

SEARCH FOR THE FLAVOR-CHANGING NEUTRAL CURRENT IN TOP  
PAIR EVENTS WITH AN ASSOCIATED PHOTON USING 13 TEV  
PROTON-PROTON COLLISION DATA COLLECTED WITH THE ATLAS  
DETECTOR

by

JASON TYLER BARKELOO

A DISSERTATION

Presented to the Department of Physics  
and the Graduate School of the University of Oregon  
in partial fulfillment of the requirements  
for the degree of  
Doctor of Philosophy

March 2020

## DISSERTATION APPROVAL PAGE

Student: Jason Tyler Barkeloo

Title: Search for the Flavor-Changing Neutral Current in Top Pair Events With an Associated Photon Using 13 TeV Proton-Proton Collision Data Collected With the ATLAS Detector

This dissertation has been accepted and approved in partial fulfillment of the requirements for the Doctor of Philosophy degree in the Department of Physics by:

David Strom

James Brau

Spencer Chang

Dev Sinha

Chair

Advisor

Core Member

Institutional Representative

and

Janet Woodruff-Borden

Vice Provost and Dean of the Graduate School

Original approval signatures are on file with the University of Oregon Graduate School.

Degree awarded March 2020

© 2020 Jason Tyler Barkeloo  
This work is licensed under a Creative Commons  
**Attribution-NonCommercial-NoDerivs (United States) License.**

## DISSERTATION ABSTRACT

Jason Tyler Barkeloo

Doctor of Philosophy

Department of Physics

March 2020

Title: Search for the Flavor-Changing Neutral Current in Top Pair Events With an Associated Photon Using 13 TeV Proton-Proton Collision Data Collected With the ATLAS Detector

Abstract for FCNC here.

This dissertation includes previously published and unpublished co-authored material.

## CURRICULUM VITAE

NAME OF AUTHOR: Jason Tyler Barkeloo

### GRADUATE AND UNDERGRADUATE SCHOOLS ATTENDED:

University of Oregon, Eugene, Oregon  
Miami University, Oxford, Ohio  
Wittenberg University, Springfield, Ohio

### DEGREES AWARDED:

Doctor of Philosophy, Physics, 2020, University of Oregon  
Master of Science, Physics, 2012, Miami University  
Bachelor of Science, Physics, 2010, Wittenberg University

### PROFESSIONAL EXPERIENCE:

**University of Oregon**, Graduate Research Assistant, ATLAS Experiment,  
June 2014-Present

**University of Oregon**, Graduate Teaching Assistant, September 2012-June  
2014

**Miami University**, Teaching and Research Assistant, August 2010 - May 2012

**Wittenberg University**, Undergraduate Researcher, August 2008 - May 2010

### GRANTS, AWARDS AND HONORS:

Weiser Senior Teaching Assistant Award – University of Oregon – 2014

American Association of Physics Teachers Outstanding Teaching Assistant –  
Miami University – 2012

### PUBLICATIONS:

J. Barkeloo, J. Brau, M. Breidenbach, R. Frey, D. Freytag, C. Gallagher, R. Herbst, M. Oriunno, B. Reese, A. Steinhebel, D. Strom. “A silicon-tungsten electromagnetic calorimeter with integrated electronics for the International Linear Collider” J. Phys.: Conf. Ser. **1162** 012016 (2019).

J. Kangara, A. Hachtel, M. C. Gillette, J. Barkeloo, E. Clements, S. Bali. “Design and construction of cost-effective fail-safe tapered amplifier systems for laser cooling and trapping experiments”, Am. J. Phys. **82**(8), 805 - 817 (2014).

A. Hachtel, J. Kleykamp, D. Kane, M. Marshall, B. Worth, J. Barkeloo, J. Kangara, J. Camenisch, M. Gillette, S. Bali. “An undergraduate lab on measurement of radiative broadening in atomic vapor”, Am. J. Phys. **81**(6), 471 (2013).

Additional ATLAS Collaboration publications can be found:

<http://inspirehep.net/search?p=exactauthor%3AJason.Barkeloo>

# TABLE OF CONTENTS

Chapter	Page
I. SIMULATION AND RECONSTRUCTION . . . . .	1
1.1. Simulation of pp Collisions . . . . .	1
1.1.1. Matrix Element Calculation and Parton Distribution Functions	2
1.1.2. Parton Shower Calculation . . . . .	4
1.1.3. Detector Simulation . . . . .	4
1.1.4. Monte Carlo Generators Used for LHC Physics . . . . .	5
1.2. Object Reconstruction . . . . .	5
1.2.1. Electrons . . . . .	7
1.2.2. Muons . . . . .	8
1.2.3. Photons . . . . .	8
1.2.4. Jets . . . . .	8
1.2.5. Missing Transverse Energy . . . . .	10
1.3. Creation of Flavor Changing Neutral Current Signal Events . . .	11
1.3.1. MadGraph5 amc@NLO . . . . .	11
II. SEARCH STRATEGY . . . . .	12
2.1. Major Backgrounds . . . . .	12
2.2. Event Reconstruction . . . . .	12
2.3. Data and Simulation Event PreSelection . . . . .	12

Chapter	Page
2.4. Control and Validation Regions . . . . .	12
2.5. Signal Region . . . . .	12
2.6. Neural Network . . . . .	12
2.6.1. Architecture and Studies . . . . .	12
III. ANALYSIS AND RESULTS . . . . .	13
3.1. Uncertainties . . . . .	13
3.2. Statistical Treatment of Results . . . . .	13
3.3. Limit on Branching Ratio $t \rightarrow q\gamma$ . . . . .	13
IV. COMPLEMENTARY SEARCHES AND OUTLOOK . . . . .	14
4.1. Comparison with Complementary Searches . . . . .	14
4.2. Future Directions . . . . .	14
4.3. Conclusion . . . . .	14
REFERENCES CITED . . . . .	15



# LIST OF FIGURES

Figure		Page
1.1.	A pictorial view of the different steps for the creation of a MC event. . .	2
1.2.	The bands are the momentum fraction, $x$ , times the unpolarized parton distribution function obtained in NNLO NNPDF3.0 global analysis at scales $\mu^2 = 10 \text{ GeV}$ and $\mu^2 = 100 \text{ GeV}^2$ . . . . .	3
1.3.	Cross section of a simulated ATLAS detector showing how various particles interact with ATLAS subsystems. . . . .	7
1.4.	C . . . . .	9
1.5.	C . . . . .	10
1.6.	C . . . . .	10

## LIST OF TABLES

Table

Page

## CHAPTER I

### SIMULATION AND RECONSTRUCTION

This chapter presents details on the simulation of various physics processes and the reconstruction of physics objects for both simulated events and data events.

#### **1.1. Simulation of pp Collisions**

To draw conclusions from ATLAS experimental data it is necessary to make accurate theoretical predictions about the processes being searched for. Having accurate background models can help identify when a data signal is behaving in a way that might suggest new physics. Due to the stochastic nature of particle physics collisions and interactions it is not practical to create exact predictions, instead the ATLAS experiment uses Monte Carlo (MC) simulations to make predictions. MC simulations are done by repeated random sampling of possible physical processes that can occur at any given time to a particle. The possibilities change based on factors such as particle energy and particle environment. A flow chart for the entire simulation chain is shown in Figure 1.1.




FIGURE 1.1. A pictorial view of the different steps for the creation of a MC event. [1]

### 1.1.1. Matrix Element Calculation and Parton Distribution Functions

Particle interactions at LHC energies do not involve the entire proton. The constituent partons that create the proton (the two up quarks, down quark, and the sea of gluons) are what interact in any given event. The gluons create many virtual quark-antiquark pairs which can interact as well. The valence quarks, the ups and down that make up the proton, are the major portion of interacting partons at low energies, mainly inelastic interactions. At LHC energies deep inelastic scattering is possible and the sea quarks play a more dominant role. Proton structure is described

by a Parton Distribution Function (PDF) which gives the probability of finding any parton with a particular momentum fraction and is shown in Figure 1.2.




FIGURE 1.2. The bands are the momentum fraction,  $x$ , times the unpolarized parton distribution function obtained in NNLO NNPDF3.0 global analysis at scales  $\mu^2 = 10 \text{ GeV}^2$  and  $\mu^2 = 100 \text{ GeV}^2$  [2]

The PDFs and hard scattering processes are included in the calculation of the Matrix Elements (ME) of any interaction. Hard scattering processes can be described by Feynman diagrams, a representation of their amplitudes. Combining the PDFs and hard scattering amplitudes gives the probability of a particular interaction occurring. Calculation of the MEs is the first stage of simulation and is done to a specified order in perturbation theory: leading order (LO), next-to-leading order (NLO), etc. Higher order calculations lead to more accurate predictions but grow in complexity exponentially making them hard to calculate both theoretically and computationally. This is what often restricts how accurate a process can be simulated.

### 1.1.2. Parton Shower Calculation

The next stage of simulating an event is the parton shower. These parton shower calculations deal with the quantum chromodynamic processes. In any interaction the particles that carry color can spontaneously emit gluons which can go on to create more gluons or quark-antiquark pairs. Depending on when this happens in the hard scattering process it is called initial state radiation (ISR) or final state radiation (FSR). The hard scattering partons as well as any additional radiated particles are used as inputs to parton shower calculations which determine how the quarks and gluons proceed through to the final state particles seen in the detector. This includes calculation of hadronization processes and further decay processes into the final state particles.

### 1.1.3. Detector Simulation

The final stage of creating a MC event is the detector simulation. The information from the event generators are processed using GEANT4 [3] and a detailed model of the ATLAS detector. GEANT4 simulates how various particles propagate through and interact with the material properties of the detector and where they leave energy which would then be measured by the ATLAS detector in an actual event. The result of this MC event construction flow is a collection of simulated data that is similar in structure to actual data collected using the ATLAS experiment. The energy deposits in both MC and real data are then reconstructed using the same software and physics objects are reconstructed. For MC this allows for comparison between the physics object reconstruction and the truth record, or the types of particles fed into the detector simulation.

#### 1.1.4. Monte Carlo Generators Used for LHC Physics

A variety of different MC generators are used in the creation of simulated events. Different generators can specialize in simulating different physics processes and handle various precision (eg., LO vs. NLO). The MC generators used in this search are summarized in this section.

MADGRAPH aMC@NLO [4]: An amplitude and event generator at LO and NLO for hard processes. Extendable to various models including effective field theory (EFT) models used in BSM searches. This generator is used to create the signal events searched for in this dissertation: discussed in Section 1.3.

POWHEG [5, 6]: **P**ositive **W**eight **H**ardest **E**mission **G**enerator is an NLO event generator that can be interfaced with other generators (i.e. PYTHIA) for showering.

PYTHIA [7]: A generator used most often for QCD final state hard processes and showering. It is commonly interfaced with other generators for showering within the ATLAS detector. SHERPA [8, 9]: A multi-parton LO generator with an emphasis on merging ME and Parton Showering.

These generators are commonly interfaced, usually with PYTHIA for showering, and this is possible due to a common file format developed at the Les Houches Accords [10]. This allows a specialty generator to be created and used to generate hard processes and then simulate the rest of the event with common showering generators that might lack the ability to simulate the process in question.

## 1.2. Object Reconstruction

After the events are simulated, or collected in case of real data, all there is is a collection of energy deposits within the detector. These energy deposits must be transformed into meaningful physics objects through reconstruction. Reconstruction

is typically done in two major parts using the specialized detectors covered in Chapter ?? . The Inner Detector and Muon System turn patterns of hits within the tracking detectors into tracks that have direction and momentum information. The calorimeter system transforms the energy deposits within the calorimeters into calibrated energy deposits with a particular position. These tracks and calorimeter deposits are used to create physics objects (electrons, muons, etc.) by using particle identification techniques to reconstruct the underlying physics event. For the analysis presented in this dissertation the final state signal particles that need to be reconstructed are one lepton (an electron or a muon), one photon, two quarks (one light flavor and one b quark), and one neutrino (missing transverse energy as it is the only particle that does not interact with the detector). Each of these particles has a particular signature in the subdetectors of the ATLAS detector, shown in Figure 1.3.






FIGURE 1.3. Cross section of a simulated ATLAS detector showing how various particles interact with ATLAS subsystems. Solid lines indicate interactions while dashed lines indicate that no interactions typically occur in that section of the detector. [11]

### 1.2.1. Electrons

Electrons interacting with the ATLAS detector leave a track in the Inner Detector as well as a cluster of energy in the electromagnetic calorimeter. The track and cluster are required to be matched together to be identified as an electron candidate[12]. As electrons move through the detector they create electromagnetic showers through bremsstrahlung which can produce electron-positron pairs. The process continues as

the particles continue to give energy to the detector. This collection of electrons, positrons, and photons creates a signature energy cluster in the calorimeter.

Electron identification algorithms are applied to the electron candidates to separate prompt and isolated electron candidates from electrons that come from backgrounds such as converted photons and misidentified jets. Non-prompt electrons are from

#### **1.2.2. Muons**

[13]

#### **1.2.3. Photons**

[14]

#### **1.2.4. Jets**

[15]




FIGURE 1.4. [15]

#### 1.2.4.1. B-Jets




FIGURE 1.5. [18]




FIGURE 1.6. [18]

### 1.2.5. Missing Transverse Energy

[19]

### **1.3. Creation of Flavor Changing Neutral Current Signal Events**

#### **1.3.1. MadGraph5 amc@NLO**

Comparison of kinematics between standard ttbar events

- param card in appendix

ATLAS Production of these events

TopQ1 Slimming/Skimmming

## CHAPTER II

### SEARCH STRATEGY

#### 2.1. Major Backgrounds

#### 2.2. Event Reconstruction

#### 2.3. Data and Simulation Event PreSelection

#### 2.4. Control and Validation Regions

#### 2.5. Signal Region

#### 2.6. Neural Network

##### 2.6.1. Architecture and Studies

## CHAPTER III

### ANALYSIS AND RESULTS

#### **3.1. Uncertainties**

#### **3.2. Statistical Treatment of Results**

#### **3.3. Limit on Branching Ratio $t \rightarrow q\gamma$**

## CHAPTER IV

### COMPLEMENTARY SEARCHES AND OUTLOOK

#### **4.1. Comparison with Complementary Searches**

#### **4.2. Future Directions**

HL-LHC and Beyond Future perspectives at Linear Colliders? - <https://www.sciencedirect.com/>

#### **4.3. Conclusion**



## REFERENCES CITED

- [1] Chaowaroj Wanotayaroj and Jim Brau. Search for a Scalar Partner of the Top Quark in the Jets+MET Final State with the ATLAS detector, Nov 2016. Presented 25 Oct 2016.
- [2] M. Tanabashi et al. Review of Particle Physics. *Phys. Rev.*, D98(3):030001, 2018.
- [3] S. Agostinelli et al. GEANT4: A Simulation toolkit. *Nucl. Instrum. Meth.*, A506:250–303, 2003.
- [4] J. Alwall, R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, O. Mattelaer, H. S. Shao, T. Stelzer, P. Torrielli, and M. Zaro. The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations. *JHEP*, 07:079, 2014.
- [5] Stefano Frixione, Paolo Nason, and Carlo Oleari. Matching NLO QCD computations with Parton Shower simulations: the POWHEG method. *JHEP*, 11:070, 2007.
- [6] Simone Alioli, Paolo Nason, Carlo Oleari, and Emanuele Re. A general framework for implementing NLO calculations in shower Monte Carlo programs: the POWHEG BOX. *JHEP*, 06:043, 2010.
- [7] Torbjorn Sjostrand, Stephen Mrenna, and Peter Z. Skands. A Brief Introduction to PYTHIA 8.1. *Comput. Phys. Commun.*, 178:852–867, 2008.
- [8] T. Gleisberg, Stefan. Hoeche, F. Krauss, M. Schonherr, S. Schumann, F. Siegert, and J. Winter. Event generation with SHERPA 1.1. *JHEP*, 02:007, 2009.
- [9] Enrico Bothmann et al. Event Generation with Sherpa 2.2. *SciPost Phys.*, 7:034, 2019.
- [10] Johan Alwall et al. A Standard format for Les Houches event files. *Comput. Phys. Commun.*, 176:300–304, 2007.
- [11] Joao Pequeno. Event Cross Section in a computer generated image of the ATLAS detector. Mar 2008.
- [12] Morad Aaboud et al. Electron reconstruction and identification in the ATLAS experiment using the 2015 and 2016 LHC proton-proton collision data at  $\sqrt{s} = 13$  TeV. *Eur. Phys. J.*, C79(8):639, 2019.
- [13] Sébastien Rettie. Muon identification and performance in the ATLAS experiment. *PoS*, DIS2018:097, 2018.

- [14] Morad Aaboud et al. Measurement of the photon identification efficiencies with the ATLAS detector using LHC Run 2 data collected in 2015 and 2016. *Eur. Phys. J.*, C79(3):205, 2019.
- [15] Matteo Cacciari, Gavin P. Salam, and Gregory Soyez. The anti- $k_t$  jet clustering algorithm. *JHEP*, 04:063, 2008.
- [16] M. Wobisch and T. Wengler. Hadronization corrections to jet cross-sections in deep inelastic scattering. In *Monte Carlo generators for HERA physics. Proceedings, Workshop, Hamburg, Germany, 1998-1999*, pages 270–279, 1998.
- [17] Selection of jets produced in 13TeV proton-proton collisions with the ATLAS detector. Technical Report ATLAS-CONF-2015-029, CERN, Geneva, Jul 2015.
- [18] Yosuke Takubo. ATLAS IBL operational experience. *PoS*, Vertex2016:004, 2017.
- [19] Morad Aaboud et al. Performance of missing transverse momentum reconstruction with the ATLAS detector using proton-proton collisions at  $\sqrt{s} = 13$  TeV. *Eur. Phys. J.*, C78(11):903, 2018.