

Detector Resolution Studies

ALCWS 2018

Jason Barkeloo

J. Brau, A. Steinhebel, E. Meyer, J. Carlson

June 1, 2018



Table of Contents

Brief Background Top FCNC

Overlap Removal Implementation

Current Status, Region Implementation

Top Quark Decays

- Standard model top branching ratio to $bW \simeq 100\%$




Figure: Leptonic final state diagram for a top decay






Figure: Top quark pair decay final states [Nature]

Top Quark Decays in the SM




- $t \rightarrow b W \approx 99.83\%$
- $t \rightarrow s W \approx 0.16\%$
- $t \rightarrow d W \approx 0.01\%$

- $t \rightarrow q_{u,c} X \approx 10^{-17} - 10^{-12}$


FCNC: What are we looking for? $t\bar{t} \rightarrow W(\rightarrow l\nu)b + q\gamma$

- ▶ Final state topology
 - ▶ One Neutrino, from W
 - ▶ One Lepton, from W
 - ▶ One B-jet, SM top
 - ▶ **One Photon, FCNC Top**
 - ▶ One Jet, FCNC Top



Background Processes

- Due to all of the processes at hadron colliders it is important to model similar event topologies well.
- Major backgrounds include $t\bar{t}$, W +Jets, Z +Jets, + processes with an associated photon



Monte Carlo Generation

- ▶ All of our MC data is put through a showering algorithm for propagation from final decay states
- ▶ Various showering algorithms are used at ATLAS - Pythia, Herwig, etc.
- ▶ All of these will add radiative photons
- ▶ These events can be contained in other samples with explicit photons originating from the hard interaction
- ▶ Need to remove these events or risk double counting events

Photon Overlap Removal

- ▶ Truth matching procedure is used to identify origin and type of truth particle corresponding to reconstructed photon
 - ▶ If reco photon is associated with a truth electron or within $R=0.2$ we can consider this $e \rightarrow \gamma$ fake
 - ▶ If origination from boson or lepton with a corresponding truth hadron: hadron fake
 - ▶ Otherwise the photon is considered coming from the hard interaction
 - ▶ The procedure rejects less than 1% of events in $t\bar{t} + \gamma$ and $V + \gamma$ samples (except $Z + \gamma$ in e+jets channel because of fake rates)

Photon Overlap Removal

- For $t\bar{t}$ and $V + jets$ samples, the prompt photon contribution is subject to large statistical uncertainty and its modelling is less trusted, it is why the $t\bar{t} + \gamma$ and $V + \gamma$ samples are used.
- For this to work phase spaces of events must be close to identical otherwise the overlap removal will take out too much




Figure: Overlapping Phase Space Regions of the photon and various objects
[Y.Li] Red: $t\bar{t} + \gamma$ Blue: $t\bar{t}$


Object Preselection

- ▶ We preselect events with objects that look like our expected topology
- ▶ Reminder that I require:
 - ▶ Exactly one lepton (e or μ) ≥ 28 GeV
 - ▶ Exactly one Good photon ≥ 25 GeV
 - ▶ Missing Transverse Energy ≥ 30 GeV
 - ▶ ≥ 2 Jets (at least one being b-tagged)
- ▶ All following plots will have signal scaled to 0.2% of nonallhadronic $\sigma_{t\bar{t}}$, MC scaled to $36.07 fb^{-1}$
- ▶ Only electron channel shown. Similar results for the muon channel are seen.


Preselection Objects

► Leading Jet p_T



No OVR





W/OVR



► Photons



► Leptons



What does this mean?

- ▶ Previous presentations have dramatically overcounted $t\bar{t}$ events
- ▶ 40 – 50% of the events in $t\bar{t}$ and $W + jets$ base samples are removed with this procedure
- ▶ The inclusion could have hidden some potential differences in ways to remove one or the other background processes




Figure: No OVR Photon ptcone20






Figure: OVR Photon ptcone20

Other Variables



► $\Delta R_{\gamma l}$

► $\gamma\theta$

No OVR



W/OVR



Signal Region



- ▶ Current Requirements:
 - ▶ Preselection Cuts
 - ▶ FCNC Top Mass: $m_{q\gamma}$ within 50GeV of m_t
 - ▶ SM Top Mass: $m_{bl\nu}$ within 50GeV of m_t
 - ▶ Z Mass: $m_{l\gamma} > 10\text{GeV}$ from m_Z
 - ▶ Photon $p_t > 50\text{GeV}$
 - ▶ $=1$ Bjet

Some SR Variables



► $\Delta R_{l\gamma}$

► $\gamma\theta$

PreSelection




SR Cuts




Some More SR Variables



► $m_{q\gamma}$



► $m_{bl\nu}$



PreSelection



SR Cuts

$t\bar{t} + \gamma$ rich region

- Preselection Cuts
- FCNC Top Mass - Orthogonal to SR
- ≥ 4 Jets




Figure: γ_e




Figure: $\Delta R_{l\gamma}$

Other Regions


- ▶ Developing more region to test the performance of MC samples
- ▶ Regions are designed to isolate various physics processes using orthogonal selections
- ▶ $t\bar{t}(+\gamma)$ is of utmost important to model extremely well, especially with the increase in cross section at 13TeV

Conclusion, Outlook

- ▶ Orthogonal validation/control regions are in development
- ▶ Data grid run complete, need to incorporate into CR/VR plots
- ▶ Next grid run will include a couple of looser regions for CR/VRs
 - ▶ 0 Photon Samples for Backgrounds with no Real Photons
 - ▶ 0 BJet Samples - possibly for WJets region
- ▶ Top Group - Pushing for MVA, want to start investigations using these techniques

Backup


Integrated Luminosity



A Couple BSM Diagrams


- R-parity-violating supersymmetric models

[arXiv:hep-ph/9705341]



- Top-color-assisted technicolor models

[arXiv:hep-ph/0303122]



Jets/AntiKT


$$d_{ij} = \min\left(\frac{1}{p_{ti}^2}, \frac{1}{p_{tj}^2}\right) \frac{\Delta_{ij}^2}{R^2}$$

$$d_{iB} = \frac{1}{p_{ti}^2}$$

$$\Delta_{ij}^2 = (\eta_i - \eta_j)^2 + (\phi_i - \phi_j)^2$$

- ▶ Find minimum of entire set of $\{d_{ij}, d_{iB}\}$
- ▶ If d_{ij} is the minimum particles i, j are combined into one particle and removed from the list of particles
- ▶ If d_{iB} is the minimum i is labelled as a final jet and removed from the list of particles
- ▶ Repeat until all particles are part of a jet with distance between jet axes Δ_{ij} is greater than R

B-tagging



$$\mathcal{L}_{tq\gamma}^{\text{eff}} = -e\bar{c}\frac{i\sigma^{\mu\nu}q_\nu}{m_t}(\lambda_{ct}^L P_L + \lambda_{ct}^R P_R)tA_\mu + H.c.$$