

# Search for Flavor Changing Neutral Currents in Top Quark Decays

B-Tagging Working Point and  $e \rightarrow \gamma$  Fakes

Jason Barkeloo

September 12, 2019



# Overview

## Brief Background

The Top Quark  
FCNC at the LHC

## B-tagging Working Point Selection



B-tagging Background  
Neural Network on B-tagging WPs

## $e \rightarrow \gamma$ Fake Rate: Initial Studies

Initial Studies  
Basic 1D Fake Rate Scale Factor  
2D Fake Rate Scale Factor

## Outlook and Conclusions

# Top Quark Decays in the SM




- ▶  $t \rightarrow bW \approx 99.83\%$
- ▶  $t \rightarrow sW \approx 0.16\%$
- ▶  $t \rightarrow dW \approx 0.01\%$

- ▶  $t \rightarrow q_{u,c}X \approx 10^{-17} - 10^{-12}$
- ▶ Limits on  $t \rightarrow \gamma q$  processes:  
[JHEP 04 (2016) 035]
  - ▶  $t \rightarrow \gamma u < 1.3 \times 10^{-4}$
  - ▶  $t \rightarrow \gamma c < 1.7 \times 10^{-3}$

## FCNC: What are we looking for? $t\bar{t} \rightarrow W(\rightarrow l\nu)b + q\gamma$


Will further investigate BJets here.

- ▶ Final state topology
  - ▶ One Neutrino, from W
  - ▶ One Lepton, from W
  - ▶ One B-jet, SM Top
  - ▶ One Photon, FCNC Top
  - ▶ One Jet, FCNC Top



## B-tagging


- ▶ B Hadrons travel a measureable distance before decay
- ▶ Tracks originate from outside of interaction point (Secondary Vertex)
- ▶ Backtracking tracks in displaced vertex gives an impact parameter
- ▶ Decay chain MVA attempts to reconstruct decay of the jet
- ▶ Outputs of these algorithms used in a BDT to determine if a Jet is from a b-quark




# Mv2c10

MV2c10 is used to tag b-jets. The c10 implies a 10% c-jet fraction in the background training sample. Can use various fixed-cut working points for b-jet identification.

Using a different working point can change which jets are identified as originating from b-quarks in the analysis.







(a)



(b)

JHEP 08 (2018) 89

# Neural Network Reminder




Branching ratio with Significance = 2:  $1.18\text{e-}5$

# Neural Network Results


Electron Channel

► 70% Working Point




BR with Sig=2:  $1.25 \times 10^{-5}$

► 77% Working Point




BR with Sig=2:  $1.23 \times 10^{-5}$

► 85% Working Point




BR with Sig=2:  $1.28 \times 10^{-5}$


Muon Channel



BR with Sig=2:  $1.31 \times 10^{-5}$




BR with Sig=2:  $1.18 \times 10^{-5}$



BR with Sig=2:  $1.19 \times 10^{-5}$



## Fake Rate Studies




Want to be able to correct the number of fake photons predicted in MC to those present in Data

## Fake Rate Object Selection

- ▶ Want to calculate fake rate in events which could enter the signal region.
- ▶ Create 2 control regions:  $Z \rightarrow ee$  and  $Z \rightarrow e\gamma$
- ▶ Require:
  - ▶ Common Object Selection (MET, Jets, Triggers, etc.)
  - ▶ Exactly 1Bjet
  - ▶  $Z \rightarrow ee$  : 2 Opposite Sign Electrons,  $86.1 \text{ GeV} < m_{e^+e^-} < 96.1 \text{ GeV}$
  - ▶  $Z \rightarrow e\gamma$  : 1 Electron,  $\geq 1$  Photon,  $86.1 \text{ GeV} < m_{e\gamma} < 96.1 \text{ GeV}$

# Truth Study / Scale Factor




Categories: Simple mis-match, mis-match to truth photon (Reco pt  $\geq 10\%$  higher than truth), non prompt photon, prompt photons


$m_{ee}, m_{e\gamma}$

## Data and MC

### ► Data



### ► Monte Carlo



# Scale Factor

$$FR^{e\text{-fake}} = \frac{N_{e,\gamma}}{N_{e,e} + N_{e,\gamma}}$$

$$SF_{FR}^{e\text{-fake}} = \frac{FR_{\text{data}}^{e\text{-fake}}}{FR_{\text{MC}}^{e\text{-fake}}}$$

Basic Scale Factor can be calculated for the entire spectrum:

$$FR_{\text{data}}^{e\text{-fake}} = 0.201$$

$$FR_{\text{MC}}^{e\text{-fake}} = 0.212$$

$$SF_{FR}^{e\text{-fake}} = 0.953$$

## Next Steps - 2D Fake Rate



# Outlook

- ▶ As always, still lots to be done
- ▶ Fake Rates  $e \rightarrow \gamma$  and  $j \rightarrow \gamma$  are being investigated,  $e \rightarrow \gamma$  shown here,  $j \rightarrow \gamma$  to be investigated soon.
- ▶ Was able to squeak an extra factor of 2 out of Neural Network since I had to redo it for working points anyway
- ▶ Questions?


# Backup



# FCNC Diagrams



# NN Input Variable Correlations



# Neural Network Model Inputs

$$\text{Separation} = \sum_i^{\text{bins}} \frac{n_{si} - n_{bi}}{n_{si} + n_{bi}}$$

## mu+jets channel

Variable	Separation
photon0iso	41.18
mqgam	28.27
photon0pt	24.07
mtSM	11.60
mlgam	7.56
deltaRjgam	5.64
deltaRbl	4.42
MWT	3.34
ST	3.30
nuchi2	3.12
jet0pt	2.81
njets	2.07
smchi2	1.89
wchi2	1.87
jet0e	1.52
deltaRlgam	1.17
leptone	0.87
deltaRjb	0.86
met	0.68
bjet0pt	0.52
leptoniso	0.27


## e+jets channel

Variable	Separation
photon0pt	23.14
mqgam	22.73
photon0iso	18.70
mtSM	11.02
mlgam	9.53
deltaRbl	5.00
deltaRjgam	4.60
ST	3.83
MWT	3.16
jet0pt	2.47
njets	1.70
nuchi2	1.59
deltaRlgam	1.40
wchi2	1.33
smchi2	1.09
deltaRjb	0.88
leptone	0.85
leptoniso	0.56
bjet0pt	0.50
met	0.47

# Input Variables

```
['photon0iso', 'photon0pt', 'mqgam', 'mlgam', 'mtSM', 'deltaRjgam', 'deltaRbl',  
'MWT', 'ST', 'njets', 'wchi2', 'jet0pt', 'deltaRlgam', 'leptone', 'met', 'bjet0pt']
```


# Integrated Luminosity



# A Couple BSM Diagrams


- R-parity-violating supersymmetric models

[arXiv:hep-ph/9705341]



- Top-color-assisted technicolor models

[arXiv:hep-ph/0303122]



# Jets/AntiKT

$$d_{ij} = \min\left(\frac{1}{p_{ti}^2}, \frac{1}{p_{tj}^2}\right) \frac{\Delta_{ij}^2}{R^2}$$

$$d_{iB} = \frac{1}{p_{ti}^2}$$

$$\Delta_{ij}^2 = (\eta_i - \eta_j)^2 + (\phi_i - \phi_j)^2$$

- ▶ Find minimum of entire set of  $\{d_{ij}, d_{iB}\}$
- ▶ If  $d_{ij}$  is the minimum particles  $i, j$  are combined into one particle and removed from the list of particles
- ▶ If  $d_{iB}$  is the minimum  $i$  is labelled as a final jet and removed from the list of particles
- ▶ Repeat until all particles are part of a jet with distance between jet axes  $\Delta_{ij}$  is greater than  $R$

$$\mathcal{L}_{tq\gamma}^{\text{eff}} = -e\bar{c}\frac{i\sigma^{\mu\nu}q_\nu}{m_t}(\lambda_{ct}^L P_L + \lambda_{ct}^R P_R)tA_\mu + H.c.$$