

SEARCH FOR THE FLAVOR-CHANGING NEUTRAL CURRENT IN TOP
PAIR EVENTS WITH AN ASSOCIATED PHOTON USING 13 TEV
PROTON-PROTON COLLISION DATA COLLECTED WITH THE ATLAS
DETECTOR

by

JASON TYLER BARKELOO

A DISSERTATION

Presented to the Department of Physics
and the Graduate School of the University of Oregon
in partial fulfillment of the requirements
for the degree of
Doctor of Philosophy

December 2019

DISSERTATION APPROVAL PAGE

Student: Jason Tyler Barkeloo

Title: Search for the Flavor-Changing Neutral Current in Top Pair Events With an Associated Photon Using 13 TeV Proton-Proton Collision Data Collected With the ATLAS Detector

This dissertation has been accepted and approved in partial fulfillment of the requirements for the Doctor of Philosophy degree in the Department of Physics by:

David Strom	Chair
James Brau	Advisor
Spencer Chang	Core Member
Dev Sinha	Institutional Representative

and

Janet Woodruff-Borden	Vice Provost and Dean of the Graduate School
-----------------------	--

Original approval signatures are on file with the University of Oregon Graduate School.

Degree awarded December 2019

© 2019 Jason Tyler Barkeloo

This work is licensed under a Creative Commons

Attribution-NonCommercial-NoDerivs (United States) License.

DISSERTATION ABSTRACT

Jason Tyler Barkeloo

Doctor of Philosophy

Department of Physics

December 2019

Title: Search for the Flavor-Changing Neutral Current in Top Pair Events With an Associated Photon Using 13 TeV Proton-Proton Collision Data Collected With the ATLAS Detector

Abstract for FCNC here.

This dissertation includes previously published and unpublished co-authored material.

CURRICULUM VITAE

NAME OF AUTHOR: Jason Tyler Barkeloo

GRADUATE AND UNDERGRADUATE SCHOOLS ATTENDED:

University of Oregon, Eugene, Oregon
Miami University, Oxford, Ohio
Wittenberg University, Springfield, Ohio

DEGREES AWARDED:

Doctor of Philosophy, Physics, 2019, University of Oregon
Master of Science, Physics, 2012, Miami University
Bachelor of Science, Physics, 2010, Wittenberg University

PROFESSIONAL EXPERIENCE:

University of Oregon, Graduate Research Assistant, ATLAS Experiment, June 2014-Present

University of Oregon, Graduate Teaching Assistant, September 2012-June 2014

Miami University, Teaching and Research Assistant, August 2010 - May 2012

Wittenberg University, Undergraduate Researcher, August 2008 - May 2010

GRANTS, AWARDS AND HONORS:

Weiser Senior Teaching Assistant Award – University of Oregon – 2014

American Association of Physics Teachers Outstanding Teaching Assistant – Miami University – 2012

PUBLICATIONS:

A. Hatchel et al., “An undergraduate lab on measurement of radiative broadening in atomic vapor”, Am. J. Phys. **81**(6), 471 (2013).

J. Kangara et al., “Design and construction of cost-effective fail-safe tapered amplifier systems for laser cooling and trapping experiments”, Am. J. Phys. **82**(8), 805 - 817 (2014).

TABLE OF CONTENTS

Chapter	Page
I. INTRODUCTION	1
1.1. The Standard Model Top Quark	1
1.2. Searching for FCNC Top Quark Decays	2
II. THEORY	4
2.1. The Standard Model	4
2.2. Particle Interactions	5
2.3. The Top Quark	5
III. THE LARGE HADRON COLLIDER AND THE ATLAS DETECTOR	6
3.1. The Large Hadron Collider	6
3.2. The ATLAS Detector	7
3.2.1. Coordinate System	8
3.2.2. Magnet Setup	9
3.2.3. Inner Detector	9
3.2.4. Middle Layers, EMCal, HCal	10
3.2.5. Muon Calorimeter	11
3.2.6. Trigger and Data Acquisition	12

Chapter	Page
IV. SIMULATION	15
4.1. Simulation of pp collisions	15
4.1.1. MC Generators Used for LHC Physics	15
4.1.2. Detector Simulation	15
4.1.3. Showering in the Detector	15
4.2. Object Reconstruction	15
4.2.1. Electrons	15
4.2.2. Photons	15
4.2.3. Jets	15
4.2.4. Muons	15
4.2.5. Creation of FCNC Signal Events	15
V. SEARCH STRATEGY	16
5.1. Major Backgrounds	16
5.2. Event Reconstruction	16
5.3. Data and Simulation Event PreSelection	16
5.4. Control and Validation Regions	16
5.5. Signal Region	16
5.6. Neural Network	16
5.6.1. Architecture and Studies	16

Chapter	Page
VI. RESULTS	17
6.1. Uncertainties	17
6.2. Statistical Treatment of Results	17
6.3. Limit on Branching Ratio $t \rightarrow q\gamma$	17
VII. COMPLEMENTARY SEARCHES AND OUTLOOK	18
7.1. Comparison with Complementary Searches	18
7.2. Future Directions	18
7.3. Conclusion	18
REFERENCES CITED	19

LIST OF FIGURES

Figure	Page
3.1. Schematic of the CERN accelerator complex.[14]	6
3.2. Schematic of the ATLAS detector.[15]	7
3.3. Cross section of a simulated ATLAS detector showing how various particles interact with ATLAS subsystems.[16]	8
3.4. Coordinate system used in the ATLAS Collaboration.[17]	9
3.5. Schematic of the ATLAS inner detector.[15]	9
3.6. Schematic of the ATLAS inner detector.[15]	10
3.7. Schematic of the structure of the ATLAS inner detector.[15]	10
3.8. Schematic of the ATLAS calorimeter system.[15]	11
3.9. Schematic of the ATLAS muon detector.[15]	12
3.10. Layout of the ATLAS trigger and data acquisition system used in Run 2.[18]	13
3.11. Total integrated luminosity as a function of time delivered by the LHC(green), recorded (yellow) and declared good for physics analysis (blue) by the ATLAS detector throughout Run 2 consisting entirely of 13 TeV pp collisions.	14
3.12. Luminosity-weighted distribution of the mean number of interactions per bunch crossing for the entirety of Run 2 shown by individual years, 2015 (yellow), 2016(orange), 2017 (purple), 2018 (green), as well as an integrated total (blue).	14

LIST OF TABLES

Table	Page
-------	------

CHAPTER I

INTRODUCTION

The Standard Model of particle physics has proven itself an exceptional and resilient theory since the combination of the electromagnetic and weak interactions in 1961[1]. Further theoretical work combined the Higgs Mechanism[2, 3] with the electroweak theory[4, 5]. The resiliency of this theoretical model has been tested to further degrees of accuracy over the decades with one of my most recent portions being the experimental confirmation of the Higgs Boson in 2012[6, 7] using the Large Hadron Collider (LHC). Further precision measurements are ongoing at various experiments at the LHC, including the ATLAS experiment.

However, the Standard Model is known to have flaws and disagreements with nature. For example, the Standard Model predicts massless neutrinos which is in conflict with experimental observation of neutrino flavor oscillation and doesn't provide an explanation for dark matter particles or their interactions with currently known particles. While these large gaps in the Standard Model are well known every precision measurement made has yet to give any significant new hints toward physics beyond the Standard Model. One new pathway to look for these hints at the LHC is through top quark decays.

1.1. The Standard Model Top Quark

The top quark was first observed at Fermilab's Tevatron in 1995[8] but the increase in energy and amount of data at the LHC has produced orders of magnitude more top quarks than previously seen opening up a pathway to precision measurements of the properties of the top quark. The top quark is the heaviest

fundamental particle with a mass of 172.51 ± 0.27 (stat) ± 0.42 (syst)[9]. This large mass also means that top quark lifetime is very short ($5 * 10^{-25}$ s) and decays before it can hadronize. This allows us to probe its' branching ratios and decay modes directly. The Standard Model predicts that the top quark decays through the charged current mode nearly 100% of the time; $t \rightarrow qW$ ($q = b, s, d$)[10]. The Standard Model also predicts a rare branching ratio of the top quark through a flavor changing neutral current (FCNC) process, to a neutral boson (photon, Z boson, Higgs Boson, or gluon) and up type quark with a branching ratio on the order of 10^{-14} .

1.2. Searching for FCNC Top Quark Decays

Precision measurements are an important litmus test for the Standard Model. Predicted branching ratios for FCNC processes in top quark decays are far beyond the experimental reach of the LHC and any observation of these decay modes would be a sure sign of new physics. Branching ratios are an important measurement due to a litany of theories for new physics beyond the Standard Model (BSM). These BSM theories predict enhancements in the top sector by many orders of magnitude such as Minimal Supersymmetric models[11], R-parity-violating Supersymmetric models[12] and two Higgs doublet models[13] can all increase this branching ratio many orders of magnitude. Even a null result to a search will set an upper limit on the branching ratio that can assist in ruling out future physical models based on their amount of large top sector enrichment.

This dissertation presents a search for top FCNCs using the entire Run 2 dataset at the LHC, containing combined 2015-2018 datasets taken by the ATLAS experiment totaling 139 fb^{-1} of integrated luminosity taken at $\sqrt{s} = 13 \text{ TeV}$. The analysis presented looks for an excess of events coming from top quark pair produced events

where one top quark decays to the most likely decay mode (a bottom quark and W boson) and the other to an up type quark (up or charm) and a photon. Chapter II a theoretical background will be presented for both the Standard Model with a closer view on the usual extensions to include the FCNC vertices. Following this Chapter III will discuss the LHC and the ATLAS experiment used in the creation of the dataset used in the analysis. In Chapter IV the special signal simulation requirements will be presented as well as the common background event simulation methodology. The search strategy including the creation of signal, control and validation regions and the training of a neural network will be examined in Chapter V. Chapter VI will discuss the results and the conclusions drawn from these results will be presented in Chapter VII.

CHAPTER II

THEORY

In this chapter a theoretical background will be presented on the Standard Model of particle physics with special attention paid to the top quark's properties and decays. This will include discussion of all of the fundamental particles and their interactions through the fundamental forces of nature: electromagnetism and the strong and weak nuclear forces.

2.1. The Standard Model

CKM Matrix[10]:

$$\begin{pmatrix} d' \\ s' \\ b' \end{pmatrix} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} = \begin{pmatrix} d \\ s \\ b \end{pmatrix}$$

$$V_{CKM} = \begin{bmatrix} |V_{ud}| & |V_{us}| & |V_{ub}| \\ |V_{cd}| & |V_{cs}| & |V_{cb}| \\ |V_{td}| & |V_{ts}| & |V_{tb}| \end{bmatrix} = \begin{bmatrix} 0.97446 \pm 0.00010 & 0.22452 \pm 0.00044 & 0.00365 \pm 0.00012 \\ 0.04214 \pm 0.00076 & & \\ 0.00896^{+0.00024}_{-0.00023} & 0.04133 \pm 0.00074 & 0.999105 \pm 0.000032 \end{bmatrix}$$

Process	SM	2HDM	QS	MSSM	RPV	XD
$t \rightarrow u\gamma$	$4 * 10^{-16}$	—	$\leq 4 * 10^{-8}$	$\leq 10^{-8}$	$\leq 10^{-9}$	—
$t \rightarrow c\gamma$	$5 * 10^{-14}$	$\leq 10^{-7}$	$\leq 4 * 10^{-8}$	$\leq 10^{-8}$	$\leq 10^{-9}$	$\leq 10^{-9}$
$t \rightarrow uZ$	$7 * 10^{-17}$	—	$\leq 6 * 10^{-4}$	$\leq 10^{-7}$	$\leq 10^{-6}$	—
$t \rightarrow cZ$	$1 * 10^{-14}$	$\leq 10^{-6}$	$\leq 6 * 10^{-4}$	$\leq 10^{-7}$	$\leq 10^{-6}$	$\leq 10^{-5}$
$t \rightarrow ug$	$4 * 10^{-14}$	—	$\leq 9 * 10^{-7}$	$\leq 10^{-7}$	$\leq 10^{-6}$	—
$t \rightarrow cg$	$5 * 10^{-12}$	$\leq 10^{-4}$	$\leq 9 * 10^{-7}$	$\leq 10^{-7}$	$\leq 10^{-6}$	$\leq 10^{-10}$
$t \rightarrow uH$	$2 * 10^{-17}$	$\leq 6 * 10^{-6}$	—	$\leq 10^{-5}$	$\leq 10^{-9}$	—
$t \rightarrow cH$	$3 * 10^{-15}$	$\leq 2 * 10^{-3}$	—	$\leq 10^{-5}$	$\leq 10^{-9}$	$\leq 10^{-4}$

2.2. Particle Interactions

CaCa

2.3. The Top Quark

DaFuck is This

CHAPTER III

THE LARGE HADRON COLLIDER AND THE ATLAS DETECTOR

This chapter details the experimental details of the collider complex at the LHC and specifically the ATLAS detector used to produce, collect and measure various particle properties.

3.1. The Large Hadron Collider

Section:LHC




FIGURE 3.1. Schematic of the CERN accelerator complex.[14]

3.2. The ATLAS Detector

Section:ATLAS




FIGURE 3.2. Schematic of the ATLAS detector.[15]




FIGURE 3.3. Cross section of a simulated ATLAS detector showing how various particles interact with ATLAS subsystems.[16]

3.2.1. Coordinate System

Coords




FIGURE 3.4. Coordinate system used in the ATLAS Collaboration.[17]

3.2.2. Magnet Setup




FIGURE 3.5. Schematic of the ATLAS inner detector.[15]

3.2.3. Inner Detector

Inner




FIGURE 3.6. Schematic of the ATLAS inner detector.[15]




FIGURE 3.7. Schematic of the structure of the ATLAS inner detector.[15]

3.2.4. Middle Layers, EMCal, HCal

Middle chunks




FIGURE 3.8. Schematic of the ATLAS calorimeter system.[15]

3.2.5. Muon Calorimeter

Muons have to get picked up I guess




FIGURE 3.9. Schematic of the ATLAS muon detector.[15]

3.2.6. Trigger and Data Acquisition

All of these subsystems lead into actual data farm




FIGURE 3.10. Layout of the ATLAS trigger and data acquisition system used in Run 2.[18]

3.2.6.1. L1Calo

3.2.6.2. HLT

3.2.6.3. Luminosity Measurements




FIGURE 3.11. Total integrated luminosity as a function of time delivered by the LHC(green), recorded (yellow) and declared good for physics analysis (blue) by the ATLAS detector throughout Run 2 consisting entirely of 13 TeV pp collisions.




FIGURE 3.12. Luminosity-weighted distribution of the mean number of interactions per bunch crossing for the entirety of Run 2 shown by individual years, 2015 (yellow), 2016 (orange), 2017 (purple), 2018 (green), as well as an integrated total (blue).

CHAPTER IV

SIMULATION

4.1. Simulation of pp collisions

4.1.1. MC Generators Used for LHC Physics

4.1.2. Detector Simulation

4.1.3. Showering in the Detector

4.2. Object Reconstruction

4.2.1. Electrons

4.2.2. Photons

4.2.3. Jets

4.2.3.1. B-Jets

4.2.4. Muons

4.2.5. Creation of FCNC Signal Events

4.2.5.1. MadGraph5 amc@NLO

Comparison of kinematics between standard ttbar events ATLAS Production of these events TopQ1 Slimming/Skimming

CHAPTER V

SEARCH STRATEGY

5.1. Major Backgrounds

5.2. Event Reconstruction

5.3. Data and Simulation Event PreSelection

5.4. Control and Validation Regions

5.5. Signal Region

5.6. Neural Network

5.6.1. Architecture and Studies

CHAPTER VI

RESULTS

6.1. Uncertainties

6.2. Statistical Treatment of Results

6.3. Limit on Branching Ratio $t \rightarrow q\gamma$

CHAPTER VII

COMPLEMENTARY SEARCHES AND OUTLOOK

7.1. Comparison with Complementary Searches

7.2. Future Directions

HL-LHC and Beyond Future prospectives at Linear Colliders?

7.3. Conclusion

REFERENCES CITED

- [1] S. L. Glashow. Partial Symmetries of Weak Interactions. *Nucl. Phys.*, 22:579–588, 1961.
- [2] Peter W. Higgs. Broken Symmetries and the Masses of Gauge Bosons. *Phys. Rev. Lett.*, 13:508–509, 1964. [,160(1964)].
- [3] F. Englert and R. Brout. Broken Symmetry and the Mass of Gauge Vector Mesons. *Phys. Rev. Lett.*, 13:321–323, 1964. [,157(1964)].
- [4] Steven Weinberg. A Model of Leptons. *Phys. Rev. Lett.*, 19:1264–1266, 1967.
- [5] Abdus Salam. Weak and Electromagnetic Interactions. *Conf. Proc.*, C680519:367–377, 1968.
- [6] Georges Aad et al. Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC. *Phys. Lett.*, B716:1–29, 2012.
- [7] Serguei Chatrchyan et al. Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC. *Phys. Lett.*, B716:30–61, 2012.
- [8] F. Abe et al. Observation of top quark production in $\bar{p}p$ collisions. *Phys. Rev. Lett.*, 74:2626–2631, 1995.
- [9] The ATLAS collaboration. Measurement of the top quark mass in the $t\bar{t} \rightarrow$ lepton+jets channel from $\sqrt{s}=8$ TeV ATLAS data. 2017.
- [10] M. Tanabashi et al. Review of Particle Physics. *Phys. Rev.*, D98(3):030001, 2018.
- [11] J. J. Cao, G. Eilam, M. Frank, K. Hikasa, G. L. Liu, I. Turan, and J. M. Yang. SUSY-induced FCNC top-quark processes at the large hadron collider. *Phys. Rev.*, D75:075021, 2007.
- [12] Junjie Cao, Zhaoxia Heng, Lei Wu, and Jin Min Yang. R-parity violating effects in top quark FCNC productions at LHC. *Phys. Rev.*, D79:054003, 2009.
- [13] G. Eilam, J. L. Hewett, and A. Soni. Rare decays of the top quark in the standard and two Higgs doublet models. *Phys. Rev.*, D44:1473–1484, 1991. [Erratum: Phys. Rev.D59,039901(1999)].
- [14] Esma Mobs. The CERN accelerator complex. Complexe des accélérateurs du CERN. Jul 2016. General Photo.

- [15] The ATLAS Experiment at the CERN Large Hadron Collider. *J. Instrum.*, 3:S08003. 437 p, 2008.
- [16] Joao Pequenao. Event Cross Section in a computer generated image of the ATLAS detector. Mar 2008.
- [17] Fabian Kuger. Signal formation processes in micromegas detectors and quality control for large size detector construction for the atlas new small wheel. 08 2017.
- [18] Moritz Backes. The ATLAS Trigger System: Ready for Run-2. *PoS*, LeptonPhoton2015:045, 2016.