

# 1 Group Actions

## 1.1 Group Actions and Permutation Representations

**Definition.** Let  $G$  be a group acting on a set  $A$

1. The *kernel* of the action is the set of elements of  $G$  that act trivially on every element of  $A$ :  $\{g \in G \mid g \cdot a = a \text{ for all } a \in A\}$ .
2. For each  $a \in A$  the *stabilizer* of  $a$  in  $G$  is the set of elements of  $G$  that fix the element  $a$ :  $\{g \in G \mid g \cdot a = a\}$  and is denoted by  $G_a$ .
3. An action is *faithful* if its kernel is the identity.

**Note.** The kernel of an action is precisely the same as the kernel of the associated permutation representation as defined in the note in section 1.7 and is rephrased below.

**Proposition 1.** For any group  $G$  and any nonempty set  $A$  there is a bijection between the actions of  $G$  on  $A$  and the homomorphisms of  $G$  into  $S_A$ .

**Definition.** If  $G$  is a group a *permutation representation* of  $G$  into the symmetric group  $S_A$  for some nonempty set  $A$ . We shall say a given action of  $G$  on  $A$  *affords* or *induces* the associated representation of  $G$ .

**Proposition 2.** Let  $G$  be a group acting on the nonempty set  $A$ . the relation on  $A$  defined by

$$a \sim b \text{ if and only if } a = g \cdot b \text{ for some } g \in G$$

is an equivalence relation. For each  $a \in A$ , the number of elements in the equivalence class containing  $a$  is  $|G : G_a|$ , the index of the stabilizer of  $a$ .

**Definition.** Let  $G$  be a group acting on the set  $A$ .

1. The equivalence class  $\{g \cdot a \mid g \in G\}$  is called the *orbit* of  $G$  containing  $a$ .
2. The action of  $G$  on  $A$  is called *transitive* if there is only one orbit, i.e., given any two elements  $a, b \in A$  there is some  $g \in G$  such that  $a = g \cdot b$ .

**Note.**

1. Every element of  $S_n$  has a unique cycle decomposition
2. Subgroups of symmetric groups are called *permutation groups*.
3. The orbits of a permutation group will refer to its orbits on  $\{1, 2, \dots, n\}$
4. The orbits of an element  $\sigma \in S_n$  will refer to the orbits of the group  $\langle \sigma \rangle$ .

## 1.2 Group Acting on Themselves by Left Multiplication - Cayley's Theorem

**Note.** In this section  $G$  is any group and we first consider  $G$  acting on itself (i.e.,  $A = G$ ) by left multiplication:

$$g \cdot a = ga \quad \text{for all } g \in G, a \in G$$

When  $G$  is a finite group of order  $n$  it is convenient to label the elements of  $G$  with the integers  $1, 2, \dots, n$  in order to describe the permutation representation afforded by this action. In this way the elements of  $G$  are listed as  $g_1, g_2, \dots, g_n$  and for each  $g \in G$  the permutation  $\sigma_g$  may be described as a permutation of the indices  $1, 2, \dots, n$  as follows:

$$\sigma_g(i) = j \quad \text{if and only if} \quad gg_i = g_j.$$

**Theorem 3.** Let  $G$  be a group, let  $H$  be a subgroup and let  $G$  act by left multiplication on the set  $A$  of left cosets of  $H$  in  $G$ . Let  $\pi_H$  be the associated permutation representation afforded by this action. Then

1.  $G$  acts transitively on  $A$
2. the stabilizer of  $G$  of the point  $1H \in A$  is the subgroup  $H$
3. the kernel of the action (i.e., the kernel of  $\pi_H$ ) is  $\cap_{x \in G} xHx^{-1}$ , and  $\ker \pi_H$  is the largest normal subgroup of  $G$  contained in  $H$ .

**Corollary 4.** (Cayley's Theorem) Every group is isomorphic to a subgroup of symmetric group. If  $G$  is a group of order  $n$ , then  $G$  is isomorphic to a subgroup of  $S_n$ .

**Corollary 5.** If  $G$  is a finite group of order  $n$  and  $p$  is the smallest prime dividing  $|G|$ , then any subgroup of index  $p$  is normal (Note that a group of order  $n$  need not have a subgroup of order  $p$ ).

## 1.3 Groups Acting on Themselves by Conjugation - The Class Equation