


COMPLEX VARIABLES

Second Edition

Stephen D. Fisher



DOVER BOOKS ON MATHEMATICS

- AN INTRODUCTION TO LINEAR ALGEBRA AND TENSORS, M. A. Akivis and V. V. Goldberg. (63545-7)
- VECTORS, TENSORS AND THE BASIC EQUATIONS OF FLUID MECHANICS, Rutherford Aris. (66110-5)
- ASYMPTOTIC EXPANSIONS OF INTEGRALS, Norman Bleistein and Richard A. Handelsman. (65082-0)
- NON-EUCLIDEAN GEOMETRY, Roberto Bonola. (60027-0)
- INTRODUCTION TO PARTIAL DIFFERENTIAL EQUATIONS, Arne Broman. (66158-X)
- AN INTRODUCTION TO ORDINARY DIFFERENTIAL EQUATIONS, Earl A. Coddington. (65942-9)
- MATRICES AND LINEAR TRANSFORMATIONS, Charles G. Cullen. (66328-0)
- INTRODUCTION TO NONLINEAR DIFFERENTIAL AND INTEGRAL EQUATIONS, H. T. Davis. (60971-5)
- SOME THEORY OF SAMPLING, W. Edwards Deming. (64684-X)
- STATISTICAL ADJUSTMENT OF DATA, W. Edwards Deming. (64685-8)
- INTRODUCTION TO LINEAR ALGEBRA AND DIFFERENTIAL EQUATIONS, John W. Dettman. (65191-6)
- LINEAR PROGRAMMING AND ECONOMIC ANALYSIS, Robert Dorfman, Paul A. Samuelson and Robert M. Solow. (65491-5)
- THEORY OF H^p SPACES, Peter L. Duren. (41184-2)
- THE THIRTEEN BOOKS OF EUCLID'S ELEMENTS, translated with an introduction and commentary by Sir Thomas L. Heath. (60088-2, 60089-0, 60090-4)
Three-volume set
- CALCULUS OF VARIATIONS WITH APPLICATIONS, George M. Ewing. (64856-7)
- DIFFERENTIAL FORMS WITH APPLICATIONS TO THE PHYSICAL SCIENCES, Harley Flanders. (66169-5)
- AN INTRODUCTION TO THE CALCULUS OF VARIATIONS, Charles Fox. (65499-0)
- FOUNDATIONS OF MODERN ANALYSIS, Avner Friedman. (64062-0)
- TECHNICAL CALCULUS WITH ANALYTIC GEOMETRY, Judith L. Gersting. (67343-X)
- INTRODUCTION TO DIFFERENCE EQUATIONS, Samuel Goldberg. (65084-7)
- PROBABILITY: AN INTRODUCTION, Samuel Goldberg. (65252-1)
- DIFFERENTIAL GEOMETRY, Heinrich W. Guggenheimer. (63433-7)
- NUMERICAL METHODS FOR SCIENTISTS AND ENGINEERS, Richard Hamming. (65241-6)
- PROBABILITY: ELEMENTS OF THE MATHEMATICAL THEORY, C. R. Heathcote. (41149-4)
- ORDINARY DIFFERENTIAL EQUATIONS, E. L. Ince. (60349-0)
- LIE ALGEBRAS, Nathan Jacobson. (63832-4)
- GREEK MATHEMATICAL THOUGHT AND THE ORIGIN OF ALGEBRA, Jacob Klein. (27289-3)
- THEORY AND APPLICATION OF INFINITE SERIES, Konrad Knopp. (66165-2)
- APPLIED ANALYSIS, Cornelius Lanczos. (65656-X)

(continued on back flap)

COMPLEX VARIABLES

Second Edition

Stephen D. Fisher
Northwestern University

 DOVER PUBLICATIONS, INC.
Mineola, New York

Copyright

Copyright © 1986, 1990 by Stephen D. Fisher
All rights reserved under Pan American and International Copyright
Conventions.

Bibliographical Note

This Dover edition, first published in 1999, is a slightly corrected,
unabridged republication of the work originally published in 1990 by
Wadsworth & Brooks, Pacific Grove, California.

Library of Congress Cataloging-in-Publication Data

Fisher, Stephen D., 1941-

Complex variables / Stephen D. Fisher. — 2nd ed.

p. cm.

"This Dover edition, first published in 1999, is a slightly corrected,
unabridged republication of the work originally published in 1990 by
Wadsworth & Brooks, Pacific Grove, California"—T.p. verso.

Includes index.

ISBN 0-486-40679-2 (pbk.)

1. Functions of complex variables. I. Title.

QA331.7.F57 1999

515'.9—dc21

99-13026

CIP

Manufactured in the United States of America
Dover Publications, Inc., 31 East 2nd Street, Mineola, N.Y. 11501

For my parents with love and appreciation

This page is intentionally left blank

Preface to the Second Edition

I was gratified by the very positive responses and reviews the first edition of this text received. Nonetheless, I wished to modify the text or exercises at a number of points; this second edition is the result. The fundamental purpose of the text has not changed—it still provides a direct route, with a minimum of prerequisites, to the most important topics in the basic theory and applications of complex variables.

The greatest changes from the first edition occur in Chapter 2, principally in the presentation of the all-important Cauchy’s Theorem. In Section 3, I give only the Green’s Theorem proof of Cauchy’s Theorem (which assumes the continuity of the derivative); the Goursat version of Cauchy’s Theorem is now in the special starred subsection 2.3.1. In the remainder of Section 3, I discuss simple connectivity (but not star-shaped domains) and derive some of the consequences of Cauchy’s Theorem. These parallel developments, one assuming continuity of the derivative and the other not, are brought together in the first theorem of Section 4, where I derive the power-series representation of an analytic function. In Sections 5 and 6 of Chapter 2, I have made other rearrangements and clarifications in the presentation of residues, Laurent series, and the use of the Residue Theorem to evaluate definite integrals.

In Chapter 4, I have made a number of alterations in the discussion of flows.

Moreover, I have modified the text and many of the exercises, throughout—adding here and deleting or rearranging there—all for added clarity. Otherwise, I have done my best to correct typographical errors that were brought to my attention by readers and teachers of the first edition. I would like to thank several people who heeded my invitation to send me suggestions for improved exposition on one topic or another; their comments and discoveries were instrumental in the development of this edition. My colleagues at Northwestern, John Franks and Sandy Zabell, were especially generous with their time in this regard. Other important comments came from Irene Erteza, Albert G. Fadell, Thomas W. Gage, Jeffrey Nunemacher, James Okon, and Richard Troxel. Additional comments from Martin Billik, R. Carmichael, F. W. Carroll, Fred Goodman, Abel Klein, Steven Krantz, Michel Lapidus, A. Matheson, Philip McCartney, Justin Peters, Walter Rudin, J. C. Taylor, John Wermer, Alvin White, Robert Whitley, and Joan Wyzkoski were also helpful. I would like to thank the manuscript reviewers: Frank Forelli,

University of Wisconsin; Tom Garrety, Williams College; B. Frank Jones, Jr., Rice University; Yue-Kuen Kwok, Hong Kong University of Science and Technology; David Minda, University of Cincinnati; Burt Rodin, University of California, San Diego; and Mitchell Tableson, Washington University.

Once again it is a pleasure to thank my editor at Wadsworth & Brooks/Cole, John Kimmel, for his sustained interest and thoroughly professional approach.

As before, I will appreciate it if readers call my attention to any typographical and mathematical errors.

Stephen D. Fisher

Preface to the First Edition

This textbook is intended for undergraduate or graduate students in science, mathematics, and engineering who are taking their first course in complex variables. Its only prerequisite is a three-semester course in calculus; no prior knowledge of Green's Theorem or line integrals is needed. A previous course in differential equations would be useful but is not necessary.

The presentation and level of rigor in this book fit this background in several different ways. First, I have not given definitions and theorems in their greatest possible generality, and I have presented only results that are of central importance to elementary complex variables. (However, many important secondary topics are to be found in the exercises that appear at the end of each section.)

Second, although virtually all of the theorems are proved in full there are a few places where I refer the reader to other sources for a fact or a complete proof (for instance, the theorem that a continuous real-valued function on a closed and bounded set attains its maximum and minimum).

Third, the presentation in Chapters 1 and 2 emphasizes areas of complex variables that have much in common with concepts that the student has studied before; for instance, limits of sequences, continuity of functions, and convergence of series. The exponential function is developed from its representation in terms of e^x , $\sin y$, and $\cos y$ so that the known continuity and differentiability of these functions can immediately be brought to bear. Furthermore, once analytic functions are defined and their basic properties are developed, I devote a section to complex power series; this includes a derivation of the fact that a convergent power series is an analytic function within its disc of convergence, whose derivative is given by the expected power series. This fact can be obtained later by a different and shorter route, but I have found it more sound pedagogically to begin in areas in which the student has had some previous experience and later move into newer areas.

Fourth, there is a plethora of solved examples (more than 220 altogether) in the text, which illustrate exactly how each concept or theorem is to be applied. In addition to the examples, there are numerous exercises at the end of each section, a total of 730 throughout the book. The easiest reinforce basic concepts, while the more challenging extend or expand upon themes in the section or relate ideas from that section to earlier sections. Sections and topics within the exercises marked with

a star can be omitted with no loss of generality; they are not needed for future developments.

There is more material in this book than can be included in a course lasting a single quarter or semester. In teaching this material at Northwestern University I have found that in a ten-week quarter I can cover all of Chapters 1, 2, and 3, and many topics from the final two chapters. The variety and extent of the applications and techniques presented in Chapters 4 and 5 allow the instructor to pick his or her favorites and permit ambitious students to read others on their own.

Complex variables is, simultaneously, a practical tool of great utility in the hands of a skilled practitioner and a mathematical structure of enormous beauty and elegance. I hope that readers of this book will retain a residue of each of these facets of the subject.

This preface would not be complete without acknowledging those people who assisted me during the writing of this book. Professor Robert B. Burckel, Kansas State University, provided me with an infinitude of improvements in the grammar and style of the first draft. Various other helpful stylistic and pedagogical suggestions were made by Professors M. D. Arthur, Michael Beals, Steven Bell, Carlos Berenstein, Bruce Berndt, James Brennan, David Colton, Michael Cullen, Abel Klein, Steven Krantz, Michael O'Flynn, James Osborn, Kent Pearce, Thomas Porschning, Glenn Schober, Daniel Shea, and Howard Swann. The computer graphics were produced by Benjamin Slivka using the Control Data CYBER 170 model 730 at Northwestern University's Vogelback Computing Center. Randall Kamien provided detailed solutions to the exercises in Chapters 1, 2 and 3; his contribution is very much appreciated. I also want to express my thanks to Professor Ezra Zeheb of the Department of Electrical Engineering, Technion, Haifa, Israel, for two very informative conversations on the Z-transform and related matters. I also had helpful talks with Professor Danny Weiss of the Department of Aeronautical Engineering, Technion, and Professor Nadav Liron of the Department of Mathematics, Technion. They all helped to increase my knowledge of the applications of complex variables.

Special thanks go to my editor at Wadsworth & Brooks/Cole, John Kimmel, who not only got me started on this book but stuck with me in what at times seemed to be an endless task. Michael Michaud of Unicorn Productions saw the book through its production; his technical expertise is in evidence throughout.

Typographical errors and mathematical slips are the bane of any author. I would appreciate any such being called to my attention.

Stephen D. Fisher

A Note to the Student

This textbook presents an introduction to the theory and applications of complex variables. The presentation has been molded by my belief that what you have already studied in calculus can be successfully applied to learning complex variables, which at its basic level is just the calculus of complex-valued functions. Where there are strong, and even obvious, analogies between the new material and calculus, I have pointed them out and arranged the presentation to emphasize these analogies. However, there are critical points at which the study of complex variables differs intrinsically from the calculus that you know, and at these points I have provided more details to explain the new material. At all times, your comprehension of the subject will be aided greatly by reading with a pencil and paper close at hand. Write things down, fill in computations that may be omitted or only partially worked out, and work through the examples by yourself. By all means, do as many exercises (assigned or not) as you can. Mathematical knowledge is not gained passively—you must be an active participant in the learning process.

This page is intentionally left blank

Contents

1	The Complex Plane	1
1.1	The Complex Numbers and the Complex Plane	1
1.1.1*	A Formal View of the Complex Numbers	10
1.2	Some Geometry	12
1.3	Subsets of the Plane	22
1.4	Functions and Limits	30
1.5	The Exponential, Logarithm, and Trigonometric Functions	43
1.6	Line Integrals and Green's Theorem	56
2	Basic Properties of Analytic Functions	77
2.1	Analytic and Harmonic Functions; The Cauchy–Riemann Equations	77
2.1.1*	Flows, Fields, and Analytic Functions	86
2.2	Power Series	93
2.3	Cauchy's Theorem and Cauchy's Formula	106
2.3.1*	The Cauchy–Goursat Theorem	119
2.4	Consequences of Cauchy's Formula	123
2.5	Isolated Singularities	135
2.6	The Residue Theorem and Its Application to the Evaluation of Definite Integrals	153
3	Analytic Functions as Mappings	171
3.1	The Zeros of an Analytic Function	171
3.1.1*	The Stability of Solutions of a System of Linear Differential Equations	183
3.2	Maximum Modulus and Mean Value	191

3.3	Linear Fractional Transformations	196
3.4	Conformal Mapping	208
3.4.1*	Conformal Mapping and Flows	219
3.5	The Riemann Mapping Theorem and Schwarz–Christoffel Transformations	224

4 Analytic and Harmonic Functions in Applications 245

4.1	Harmonic Functions	245
4.2	Harmonic Functions as Solutions to Physical Problems	254
4.3	Integral Representations of Harmonic Functions	284
4.4	Boundary-Value Problems	298
4.5	Impulse Functions and the Green's Function of a Domain	309

5 Transform Methods 318

5.1	The Fourier Transform: Basic Properties	318
5.2	Formulas Relating u and \hat{u}	335
5.3	The Laplace Transform	346
5.4	Application of the Laplace Transform to Differential Equations	356
5.5	The Z-Transform	365
5.5.1*	The Stability of a Discrete Linear System	374

Appendix 1 Locating the Zeros of a Polynomial 381

Appendix 2 A Table of Conformal Mappings 389

Appendix 3 A Table of Laplace Transforms 395

Solutions to Odd-Numbered Exercises 397

Index 425

COMPLEX VARIABLES

Second Edition

This page is intentionally left blank

1

The Complex Plane

1.1 The Complex Numbers and the Complex Plane

The theory and utility of functions of a complex variable ultimately depend in large measure on viewing the usual x - and y -coordinates in the plane as separate components of a single new variable, the complex variable z . This new variable z can then be manipulated in the same way as conventional numbers are.

The familiar numbers, such as -1 , $\frac{1}{3}$, $\sqrt{2}$, and π , which are represented by points on a line, will be referred to as **real numbers**. A **complex number** is an expression of the form

$$z = x + iy,$$

where x and y are real numbers and i satisfies the rule

$$(i)^2 = (i)(i) = -1.$$

The number x is called the **real part** of z and is written

$$x = \operatorname{Re} z.$$

The number y , despite the fact that it is also a real number, is called the **imaginary part** of z and is written

$$y = \operatorname{Im} z.$$

Thus, for instance, we have $1 = \operatorname{Re}(1 + 3i)$ and $3 = \operatorname{Im}(1 + 3i)$. The **modulus**, or **absolute value**, of z is defined by

$$|z| = \sqrt{x^2 + y^2}, \quad z = x + iy.$$

Each complex number $z = x + iy$ corresponds to the point $P(x, y)$ in the xy -plane (Fig. 1.1). The modulus of z , then, is just the distance from the point $P(x, y)$




Figure 1.1

to the origin, which in complex-number notation is 0. In this way, we see that we have three inequalities relating x , y , and $|z|$, namely

$$|x| \leq |z|, \quad |y| \leq |z|, \quad \text{and} \quad |z| \leq |x| + |y|.$$

The first two of these are obvious; the third is obtained by noting that

$$|z|^2 = x^2 + y^2 \leq x^2 + 2|x||y| + y^2 = (|x| + |y|)^2.$$

The complex conjugate of $z = x + iy$ is given by

$$\bar{z} = x - iy.$$

Occasionally in engineering books, one encounters the notation z^* for \bar{z} , as well as the use of j instead of i ; we shall not use either of these. For the specific complex number $z = 1 + 3i$,

$$\bar{z} = 1 - 3i \quad \text{and} \quad |z| = \sqrt{10}$$

(see Fig. 1.2).




Figure 1.2

Addition, subtraction, multiplication, and division of complex numbers follow the ordinary rules of arithmetic. (Keep in mind that $i^2 = -1$, and, as usual, division by zero is not allowed.) Specifically, if

$$z = x + iy \quad \text{and} \quad w = s + it,$$

then

$$z + w = (x + s) + i(y + t)$$

$$z - w = (x - s) + i(y - t)$$

$$zw = (xs - yt) + i(xt + ys)$$

$$\frac{z}{w} = \frac{\bar{w}z}{\bar{w}w} = \frac{(xs + yt) + i(ys - xt)}{s^2 + t^2}, \quad w \neq 0.$$

Here, to obtain the formula for the quotient of z and w we used the device of multiplying both numerator and denominator by $\bar{w} = s - it$.

Example 1 To illustrate these rules in a particular case, let us take

$$z = 1 + 3i \quad \text{and} \quad w = -2 + 5i.$$

Then

$$z + w = -1 + 8i$$

$$zw = [(1)(-2) - (5)(3)] + i[(1)(5) + (3)(-2)] = -17 - i$$

$$\frac{z}{w} = \frac{\bar{w}z}{\bar{w}w} = \frac{13 - 11i}{29} = \frac{13}{29} - \frac{11}{29}i.$$

Two facts are of particular importance. The first, which is used repeatedly, is that

$$z\bar{z} = (x + iy)(x - iy) = x^2 + y^2 = |z|^2.$$

The second is that z and \bar{z} have the same absolute value:

$$|z| = \sqrt{x^2 + y^2} = |\bar{z}|.$$

Another useful relation is derived with the following computation.

$$\begin{aligned} |zw|^2 &= (xs - yt)^2 + (xt + ys)^2 \\ &= x^2s^2 + y^2t^2 + x^2t^2 + y^2s^2 \\ &= (x^2 + y^2)(s^2 + t^2) \\ &= |z|^2|w|^2, \end{aligned}$$

so after taking square roots, we obtain

$$|zw| = |z||w|.$$

In a similar vein, we compute $\bar{z}\bar{w}$:

$$\begin{aligned}\bar{z}\bar{w} &= (xs - yt) - i(xt + ys) \\ &= (x - iy)(s - it) \\ &= \bar{z}\bar{w}.\end{aligned}$$
□

Polar Representation

The identification of $z = x + iy$ with the point $P(x, y)$ in the xy -plane has further interest and significance if we make use of the usual polar coordinates in the xy -plane. The polar coordinate system gives

$$x = r \cos \theta \quad \text{and} \quad y = r \sin \theta,$$

where $r = \sqrt{x^2 + y^2}$ and θ is the angle measured from the positive x -axis to the line segment from the origin to $P(x, y)$. We immediately see that $r = |z|$, so

$$z = |z|(\cos \theta + i \sin \theta).$$

This is the **polar representation** of z (see Fig. 1.3a).

Example 2 Find the polar representation $z = -1 + i$.

Solution $|z| = \sqrt{2}$ and $\theta = 3\pi/4$. Thus,

$$-1 + i = \sqrt{2} \left[\cos \frac{3\pi}{4} + i \sin \frac{3\pi}{4} \right],$$

which you can easily verify as correct, since $\cos \frac{3\pi}{4} = -\frac{1}{2}\sqrt{2}$ and $\sin \frac{3\pi}{4} = \frac{1}{2}\sqrt{2}$ (see Fig. 1.3b). □

By now you have probably noticed that θ could equally well be replaced in the formulas by $\theta + 2\pi$, by $\theta - 4\pi$, or, indeed, by $\theta + 2\pi n$, where n is any integer. This ambiguity about the appropriate angle to use in the polar representation of a complex number is not just a question of semantics. Later we shall see that this causes some fundamental problems. Putting this aside for the moment, let's proceed with other properties of the polar representation. Suppose that

$$z = |z|(\cos \theta + i \sin \theta) \quad \text{and} \quad w = |w|(\cos \psi + i \sin \psi)$$




Figure 1.3

are two complex numbers. Then

$$\begin{aligned} zw &= |z||w|\{(\cos \theta \cos \psi - \sin \theta \sin \psi) + i(\cos \theta \sin \psi + \cos \psi \sin \theta)\} \\ &= |zw|\{\cos(\theta + \psi) + i \sin(\theta + \psi)\}. \end{aligned}$$

Moreover,

$$\begin{aligned} \frac{z}{w} &= \frac{|z|(\cos \theta + i \sin \theta)}{|w|(\cos \psi + i \sin \psi)} \\ &= \left(\frac{|z|}{|w|}\right)\{\cos \theta \cos \psi + \sin \theta \sin \psi + i(\cos \psi \sin \theta - \cos \theta \sin \psi)\} \\ &= \left(\frac{|z|}{|w|}\right)\{\cos(\theta - \psi) + i \sin(\theta - \psi)\}. \end{aligned}$$

Here we have made use of the trigonometric identities for the sine and cosine of the sum and difference, respectively, of two angles. Hence, the polar representation of the product (or quotient) of two complex numbers is found by multiplying (or dividing) their respective moduli and adding (or subtracting) their respective polar angles (Fig. 1.4). In other words, multiplying w by $z = |z|(\cos \theta + i \sin \theta)$ produces a rotation of w in the counterclockwise direction of θ radians and stretches (or shrinks) $|w|$ by a factor of $|z|$.

Example 3 Find the polar representation of zw and z/w if $z = -1 + i$ and $w = \sqrt{3} + i$.

Solution From Example 2, the polar representation of z is

$$z = \sqrt{2} \left[\cos \frac{3\pi}{4} + i \sin \frac{3\pi}{4} \right].$$




Figure 1.4

The polar representation of w is

$$w = \sqrt{3} + i = 2 \left[\cos \frac{\pi}{6} + i \sin \frac{\pi}{6} \right],$$

since $|w| = 2$ and $\psi = \pi/6$. Therefore,

$$(-\sqrt{3} - 1) + i(\sqrt{3} - 1) = zw = 2\sqrt{2} \left[\cos \frac{11\pi}{12} + i \sin \frac{11\pi}{12} \right]$$

$$\frac{-\sqrt{3} + 1}{4} + \frac{i\sqrt{3} - 1}{4} = \frac{z}{w} = \frac{\sqrt{2}}{2} \left[\cos \frac{7\pi}{12} + i \sin \frac{7\pi}{12} \right]. \quad \square$$

The foregoing is now used to derive **De Moivre's Theorem***:

$$(\cos \theta + i \sin \theta)^n = \cos n\theta + i \sin n\theta,$$

for any positive integer n and any angle θ . The formula is clearly true when $n = 1$; we shall use mathematical induction to prove it true for all n . Suppose there is a positive integer m for which

$$(\cos \theta + i \sin \theta)^m = \cos m\theta + i \sin m\theta.$$

Then

$$\begin{aligned} (\cos \theta + i \sin \theta)^{m+1} &= (\cos \theta + i \sin \theta)^m (\cos \theta + i \sin \theta) \\ &= (\cos m\theta + i \sin m\theta)(\cos \theta + i \sin \theta) \\ &= \cos(m+1)\theta + i \sin(m+1)\theta, \end{aligned}$$

* Abraham De Moivre, 1667–1754.

by invoking the formula derived above for the polar representation of the product of two complex numbers. Thus, if the equality holds for m , then it holds for $m + 1$. Since we know it is true for $n = 1$, it is true for all positive integers n .

Example 4 Let $\theta = \pi/4$; then $\cos \theta = \sin \theta = \sqrt{2}/2$. Thus,

$$\left(\frac{\sqrt{2}}{2} + i\frac{\sqrt{2}}{2}\right)^4 = \left(\cos \frac{\pi}{4} + i \sin \frac{\pi}{4}\right)^4 = \cos \pi + i \sin \pi = -1. \quad \square$$

Example 5 De Moivre's Theorem can be used to derive trigonometric identities for $\cos n\theta$ and $\sin n\theta$. For instance, by cubing,

$$\begin{aligned} (\cos \theta + i \sin \theta)^3 &= \cos^3 \theta + 3i \cos^2 \theta \sin \theta - 3 \cos \theta \sin^2 \theta - i \sin^3 \theta \\ &= (\cos^3 \theta - 3 \cos \theta \sin^2 \theta) + i(3 \cos^2 \theta \sin \theta - \sin^3 \theta). \end{aligned}$$

However, De Moivre's Theorem gives $(\cos \theta + i \sin \theta)^3 = \cos 3\theta + i \sin 3\theta$. After equating the real and imaginary parts of these expressions, we find that

$$\begin{aligned} \cos 3\theta &= \cos^3 \theta - 3 \cos \theta \sin^2 \theta = 4 \cos^3 \theta - 3 \cos \theta \\ \sin 3\theta &= 3 \cos^2 \theta \sin \theta - \sin^3 \theta = 3 \sin \theta - 4 \sin^3 \theta. \end{aligned}$$

Similar formulas can be derived for $\cos 4\theta$, $\sin 4\theta$, $\cos 5\theta$, and so on. (See Exercise 19.) \square

We define an **argument** of the nonzero complex number z to be any angle θ for which

$$z = |z|(\cos \theta + i \sin \theta),$$

whether or not it lies in the range $[0, 2\pi)$; we write $\theta = \arg z$. To repeat,

$$\arg z = \theta \quad \text{is equivalent to} \quad z = |z|(\cos \theta + i \sin \theta).$$

A concrete choice of $\arg z$ is made by defining **Arg** z to be that number θ_0 in the interval $[-\pi, \pi)$ such that

$$z = |z|(\cos \theta_0 + i \sin \theta_0).$$

We may then write

$$\operatorname{Arg}(zw) = \operatorname{Arg} z + \operatorname{Arg} w \pmod{2\pi},$$

where the expression $(\pmod{2\pi})$ means that the two sides of this last formula differ by some integer multiple of 2π . For example, if $z = -1 + i$ and $w = i$, then

$zw = -1 - i$, so $\operatorname{Arg}(zw) = -3\pi/4$. However, $\operatorname{Arg} z = 3\pi/4$ and $\operatorname{Arg} w = \pi/2$, so $\operatorname{Arg} z + \operatorname{Arg} w = (5/4)\pi = -3\pi/4 + 2\pi$.

Complex Numbers as Vectors

If $z = x + iy$ and $w = s + it$ are two nonzero complex numbers, then

$$|z - w| = \sqrt{(x - s)^2 + (y - t)^2}$$

is nothing but the distance in the xy -plane from the point $P(x, y)$ to the point $Q(s, t)$. Moreover, the sum $z + w = (x + s) + i(y + t)$ and the difference $z - w = (x - s) + i(y - t)$ correspond exactly to the addition and subtraction of the vectors \mathbf{OP} and \mathbf{OQ} (Fig. 1.5).




Figure 1.5

Note also that the angle α between the vector \mathbf{OP} and the vector \mathbf{OQ} is found by using the usual dot product of two vectors:

$$\cos \alpha = (\mathbf{OP} \cdot \mathbf{OQ}) / |\mathbf{OP}| |\mathbf{OQ}|$$

$$= \frac{xs + yt}{\sqrt{x^2 + y^2} \sqrt{s^2 + t^2}}$$

$$= \frac{\operatorname{Re}(z\bar{w})}{|z| |w|}.$$

In particular, \mathbf{OP} and \mathbf{OQ} are perpendicular if and only if $\operatorname{Re}(z\bar{w}) = 0$. The relations among the lengths of the sides of the triangle formed by z , w , and $z - w$, which is just the law of cosines, is formulated here as

$$\begin{aligned} |z - w|^2 &= (x - s)^2 + (y - t)^2 \\ &= x^2 + s^2 + y^2 + t^2 - 2(xs + yt) \\ &= |z|^2 + |w|^2 - 2 \operatorname{Re}(z\bar{w}). \end{aligned}$$

In summary, the usual xy -plane has a natural interpretation as the location of the complex variable $z = x + iy$, and all the rules for the geometry of the vectors $P(x, y)$ can be recast in terms of z . Henceforth, then, we refer to the xy -plane as the **complex plane**, or simply, the plane. The x -axis will be called the **real axis**, and the y -axis will be called the **imaginary axis**.

EXERCISES FOR SECTION 1.1

- Let $z = 1 + 2i$, $w = 2 - i$, and $\zeta = 4 + 3i$. Compute (a) $z + 3w$; (b) $-2w + \bar{\zeta}$; (c) z^2 ; (d) $w^3 + w$; (e) $\operatorname{Re}(\zeta^{-1})$; (f) w/z ; (g) $\zeta^2 + 2\bar{\zeta} + 3$.
- Use the quadratic formula to solve these equations; express the answers as complex numbers. (a) $z^2 + 36 = 0$; (b) $2z^2 + 2z + 5 = 0$; (c) $5z^2 + 4z + 1 = 0$; (d) $z^2 - z = 1$; (e) $z^2 = 2z$.
- Sketch the locus of those points w with (a) $|w| = 3$; (b) $|w - 2| = 1$; (c) $|w + 2|^2 = 4$; (d) $|w + 2| = |w - 2|$; (e) $|w^2 - 2w - 1| = 0$; (f) $\operatorname{Re}[(1 - i)\bar{z}] = 0$; (g) $\operatorname{Re}[z/(1 + i)] = 0$.
- Find $\operatorname{Re}(1/z)$ and $\operatorname{Im}(1/z)$ if $z = x + iy$, $z \neq 0$. Show that $\operatorname{Re}(iz) = -\operatorname{Im} z$ and $\operatorname{Im}(iz) = \operatorname{Re} z$.
- Give the polar representation for (a) $-1 + i$; (b) $1 + i\sqrt{3}$; (c) $-i$; (d) $(2 - i)^2$; (e) $|4 + 3i|$; (f) $\sqrt{5} - i$; (g) $-2 - 2i$; (h) $\sqrt{2}/(1 + i)$; (i) $[(1 + i)/\sqrt{2}]^4$.
- Give the complex number whose polar coordinates (r, θ) are (a) $(\sqrt{3}, \pi/4)$; (b) $(1/\sqrt{2}, \pi)$; (c) $(4, -\pi/2)$; (d) $(2, -\pi/4)$; (e) $(1, 4\pi)$; (f) $(\sqrt{2}, 9\pi/4)$.
- Let a , b , and c be real numbers with $a \neq 0$ and $b^2 < 4ac$. Show that the two roots of $ax^2 + bx + c = 0$ are complex conjugates of each other.
- Suppose that A is a real number and B is a complex number. Show that

$$|z|^2 + A^2 = |z + A|^2 - 2 \operatorname{Re}(Az)$$

and

$$|z|^2 + 2 \operatorname{Re}(Bz) = |z + B|^2 - |B|^2.$$

- Show that $|z| = 1$ if and only if $1/z = \bar{z}$.
- Let z and w be complex numbers with $zw = 0$. Show that either z or w is zero.
- Show that $|z + w|^2 - |z - w|^2 = 4 \operatorname{Re}(z\bar{w})$ for any complex numbers z , w .
- Let z_1, z_2, \dots, z_n be complex numbers. Establish the following formulas by mathematical induction:
 - $|z_1 z_2 \dots z_n| = |z_1| |z_2| \dots |z_n|$
 - $\operatorname{Re}(z_1 + z_2 + \dots + z_n) = \operatorname{Re}(z_1) + \operatorname{Re}(z_2) + \dots + \operatorname{Re}(z_n)$
 - $\operatorname{Im}(z_1 + z_2 + \dots + z_n) = \operatorname{Im}(z_1) + \operatorname{Im}(z_2) + \dots + \operatorname{Im}(z_n)$
 - $\overline{z_1 z_2 \dots z_n} = \bar{z}_1 \bar{z}_2 \dots \bar{z}_n$.
- Determine which of the following sets of three points constitute the vertices of a right triangle: (a) $3 + 5i, 2 + 2i, 5 + i$; (b) $2 + i, 3 + 5i, 4 + i$; (c) $6 + 4i, 7 + 5i, 8 + 4i$.
- Show that $\cos \theta = \cos \psi$ and $\sin \theta = \sin \psi$ if and only if $\theta - \psi$ is an integer multiple of 2π .
- Show that the triangle with vertices at 0 , z , and w is equilateral if and only if $|z|^2 = |w|^2 = 2 \operatorname{Re}(z\bar{w})$.

16. Let z_0 be a nonzero complex number. Show that the locus of points tz_0 , $-\infty < t < \infty$, is the straight line through z_0 and 0.
17. Show that if $w \neq 0$, then $|z/w| = |z|/|w|$.
18. Prove the identity $1 + z + z^2 + \cdots + z^n = (1 - z^{n+1})/(1 - z)$ valid for all z , $z \neq 1$.
19. Show that $\cos n\theta$ can be expressed as a combination of powers of $\cos \theta$ with integer coefficients. (Hint: Use De Moivre's Theorem and the fact that $\sin^2 \theta = 1 - \cos^2 \theta$.)

The Schwarz* Inequality

20. Let B and C be nonnegative real numbers and A a complex number. Suppose that $0 \leq B - 2 \operatorname{Re}(\bar{\lambda}A) + |\lambda|^2 C$ for all complex numbers λ . Conclude that $|A|^2 \leq BC$. (Hint: If $C = 0$, show that $A = 0$. If $C \neq 0$, then choose $\lambda = A/C$.)
21. Let a_1, \dots, a_n and b_1, \dots, b_n be complex numbers. Establish the Schwarz inequality:

$$\left| \sum_{j=1}^n a_j \bar{b}_j \right|^2 \leq \left\{ \sum_{j=1}^n |a_j|^2 \right\} \left\{ \sum_{j=1}^n |b_j|^2 \right\}.$$

(Hint: For all complex numbers λ , we have $0 \leq \sum_{j=1}^n |a_j - \lambda b_j|^2$. Expand this and apply Exercise 20 with $A = \sum_{j=1}^n a_j \bar{b}_j$, $B = \sum_{j=1}^n |a_j|^2$, $C = \sum_{j=1}^n |b_j|^2$.

22. Verify the Schwarz inequality directly for the case $n = 2$.
23. When does equality hold in the Schwarz inequality?
24. Use the Schwarz inequality to establish that

$$\left\{ \sum_{j=1}^n |a_j + b_j|^2 \right\}^{1/2} \leq \left\{ \sum_{j=1}^n |a_j|^2 \right\}^{1/2} + \left\{ \sum_{j=1}^n |b_j|^2 \right\}^{1/2}.$$

(Hint: Expand $\sum_{j=1}^n |a_j + b_j|^2$ and apply the Schwarz inequality.)

1.1.1 A Formal View of the Complex Numbers★

The complex numbers can be developed in a formal way from the real numbers. A complex number z is defined to be an ordered pair (x, y) of real numbers; we write $z = (x, y)$. Two complex numbers $z_1 = (x_1, y_1)$ and $z_2 = (x_2, y_2)$ are *equal* when $x_1 = x_2$ and $y_1 = y_2$. The basic arithmetic operations of addition and multiplication are defined, respectively, by,

$$\text{addition: } z_1 + z_2 = (x_1 + x_2, y_1 + y_2) \tag{1}$$

$$\text{multiplication: } z_1 z_2 = (x_1 x_2 - y_1 y_2, x_1 y_2 + x_2 y_1). \tag{2}$$

* Hermann Amandus Schwarz, 1843–1921.

The **additive identity** is $\mathbf{0} = (0, 0)$, since $(0, 0) + (x, y) = (x, y) + (0, 0) = (x, y)$ for all (x, y) . The **multiplicative identity** is $\mathbf{1} = (1, 0)$, since $(1, 0)(x, y) = (x, y)(1, 0) = (x, y)$ for all (x, y) . Further, it is elementary but somewhat tedious to show that the arithmetic operations of addition and multiplication are **commutative**:

$$z_1 + z_2 = z_2 + z_1; \quad z_1 z_2 = z_2 z_1$$

and **associative**:

$$(z_1 + z_2) + z_3 = z_1 + (z_2 + z_3); \quad (z_1 z_2) z_3 = z_1 (z_2 z_3).$$

Each $z = (x, y)$ has a unique **additive inverse** $-z = (-x, -y)$, since $z + (-z) = \mathbf{0}$. A nonzero $z = (x, y)$ necessarily satisfies the condition $x^2 + y^2 > 0$, and its unique **multiplicative inverse** is

$$z^{-1} = \left(\frac{x}{x^2 + y^2}, \frac{-y}{x^2 + y^2} \right),$$

since $(z)(z^{-1}) = (1, 0) = \mathbf{1}$.

The mathematical system of the complex numbers so constructed is one example of a **field**. There are many other examples of fields besides the complex numbers; for instance, the real numbers themselves form a field, as do the rational numbers.

The complex number $(0, 1)$ has the interesting property that its square is -1 : $(0, 1)(0, 1) = (-1, 0)$. Further, $(0, 1)$ and $(0, -1)$ are the only two complex numbers with this property (see the exercises that follow). We denote $(0, 1)$ by the symbol i . Each complex number then can be written

$$\begin{aligned} z &= (x, y) = (x, 0) + (0, y) \\ &= (x, 0) + (y, 0)(0, 1) \\ &= (x, 0) + (y, 0)i. \end{aligned}$$

Complex numbers of the form $(a, 0)$ are just the real numbers with their usual rules of arithmetic:

$$(a, 0) + (b, 0) = (a + b, 0)$$

$$(a, 0)(b, 0) = (ab, 0),$$

and it is entirely natural to identify $(a, 0)$ with a . In this way we may write

$$z = (x, y) = (x, 0) + i(y, 0) = x + iy.$$

This brings us back to the point where Section 1 began.

EXERCISES FOR SECTION 1.1.1

Throughout, the usual rules of arithmetic are assumed for the real numbers; in particular, $a^2 > 0$ for any nonzero real number a .

1. Show directly from rule (2) for multiplication that $z^2 = (-z)^2$.
2. Suppose that $z = (x, y)$ and $z^2 = (-1, 0)$. Show that $z = i$ or $z = -i$.
3. Solve the equation $z^2 = (0, 1)$.
4. Suppose that z^2 is real and negative; that is, $z^2 = (a, 0)$, $a < 0$. Show that $z = (0, b)$ and find b in terms of a .
5. Show by computation that addition of complex numbers is associative: $(z_1 + z_2) + z_3 = z_1 + (z_2 + z_3)$; and commutative: $z_1 + z_2 = z_2 + z_1$.
6. Show by computation that multiplication of complex number is associative: $(z_1 z_2) z_3 = z_1 (z_2 z_3)$; and commutative: $z_1 z_2 = z_2 z_1$.
7. Define the absolute value, $|z|$, of $z = (x, y)$ by $|z| = \sqrt{x^2 + y^2}$. Show directly that $|z_1 z_2| = |z_1| |z_2|$.
8. Define the complex conjugate, \bar{z} , of $z = (x, y)$ by $\bar{z} = (x, -y)$. Show that $z\bar{z} = (|z|^2, 0)$.
9. Show that $z_1 z_2 = 0$ implies that either z_1 or z_2 is zero.
10. Let $z = (x, y)$. Show that (a) $|x| \leq |z|$; (b) $|y| \leq |z|$; (c) $|z| \leq |x| + |y|$.

1.2 Some Geometry

The Triangle Inequality

Let us begin with an important inequality that has a simple geometric interpretation. Suppose $z = x + iy$ and $w = s + it$ are two complex numbers. Then

$$\begin{aligned}|z + w|^2 &= (x + s)^2 + (y + t)^2 \\&= x^2 + s^2 + y^2 + t^2 + 2(xs + yt) \\&= |z|^2 + |w|^2 + 2 \operatorname{Re}(zw) \\&\leq |z|^2 + |w|^2 + 2|z\bar{w}| \\&= |z|^2 + |w|^2 + 2|z||w| \\&= (|z| + |w|)^2.\end{aligned}$$

Taking the square root of both sides yields the inequality

$$|z + w| \leq |z| + |w|.$$

This is the **triangle inequality**, since it simply expresses the fact that any one side of a triangle is not longer than the sum of the lengths of the other two sides (see Fig. 1.6).




Figure 1.6

If ζ and ξ are two (other) complex numbers, then by putting $z = \zeta - \xi$ and $w = \xi$ we get $|\zeta| \leq |\zeta - \xi| + |\xi|$ or

$$|\zeta| - |\xi| \leq |\zeta - \xi|.$$

Likewise,

$$|\xi| - |\zeta| \leq |\zeta - \xi|,$$

which together yield a variation of the triangle inequality,

$$||\zeta| - |\xi|| \leq |\zeta - \xi|.$$

Straight Lines

The equation of a (nonvertical) straight line, $y = mx + b$, m and b real, can be formulated as

$$0 = \operatorname{Re}((m + i)z + b).$$

More generally, if $a = A + iB$ is a nonzero complex number and b is any complex number (not just a real number), then

$$0 = \operatorname{Re}(az + b)$$

is exactly the straight line $Ax - By + \operatorname{Re}(b) = 0$; this formulation also includes the vertical lines, $x = \operatorname{Re} z = \text{constant}$. (See Fig. 1.7.)

Roots of Complex Numbers

The computation of the fractional powers of a nonzero complex number is possible with the techniques developed in Section 1 of this chapter. It was in an attempt to




Figure 1.7

find the roots of such equations as $x^2 + 1 = 0$ that the whole subject of complex numbers first arose; here, a certain “completeness” will be evidenced by the complex numbers but not by the real numbers. Suppose w is a nonzero complex number and n is a positive integer. A complex number z satisfying the equation $z^n = w$ is called an **n th root of w** . We shall determine all the distinct n th roots of w .

Let $w = |w|(\cos \psi + i \sin \psi)$ be the polar representation of w , where we specify that ψ lies in the range $[-\pi, \pi]$. Let $z = |z|(\cos \theta + i \sin \theta)$; the relation $z^n = w$ and De Moivre’s Theorem from Section 1 then yield three equations:

$$|z|^n = |w|, \quad \cos(n\theta) = \cos \psi, \quad \text{and} \quad \sin(n\theta) = \sin \psi.$$

Thus, we must have $|z| = |w|^{1/n}$; θ is not so well determined. Of course, one possibility for θ is $\theta = \psi/n$; however, there are others. We define

$$\theta_k = \frac{\psi}{n} + k \left(\frac{2\pi}{n} \right), \quad k = 0, 1, \dots, n-1.$$

Then $n\theta_k = \psi + 2\pi k$ and so $\cos n\theta_k = \cos \psi$ and $\sin n\theta_k = \sin \psi$. Complex numbers z_0, \dots, z_{n-1} are defined by the rule

$$z_k = |w|^{1/n}(\cos \theta_k + i \sin \theta_k), \quad k = 0, 1, \dots, n-1.$$

Then each of z_0, \dots, z_{n-1} is distinct (see Exercise 14, Section 1), and each satisfies

$$z_k^n = w, \quad k = 0, 1, \dots, n-1.$$

Moreover, these complex numbers z_0, \dots, z_{n-1} are the only possible roots of the equation $z^n = w$. For if

$$\cos n\theta = \cos \psi, \quad \sin n\theta = \sin \psi,$$

then (again by Exercise 14, Section 1) we have $n\theta = \psi + 2\pi j$ for some integer j . The values $j = 0, \dots, n - 1$ yield distinct numbers $\cos \theta_j + i \sin \theta_j$, whereas other values of j just give a repetition of numbers already obtained. The geometric picture of the n th roots of w is very simple: the n roots lie on the circle centered at the origin of radius $\rho = |w|^{1/n}$; the roots are equally spaced on this circle, with one of the roots having polar angle $\theta_0 = \operatorname{Arg} w/n$; for instance, see Figure 1.8.

Example 1 Find the 12th roots of 1.

Solution Since $w = 1 = \cos 0 + i \sin 0$, the modulus of all the 12th roots is 1. The roots are equally spaced on the circle of radius 1 centered at the origin. One root is $z_0 = 1$; the others have polar angles of $2\pi/12, 4\pi/12, 6\pi/12, \dots, 22\pi/12$, respectively. \square

Example 2 Find the 5th roots of $i + 1$.

Solution The polar representation of $i + 1$ is

$$1 + i = \sqrt{2} \left(\cos \frac{\pi}{4} + i \sin \frac{\pi}{4} \right),$$

so the modulus of all the 5th roots of $i + 1$ is $2^{1/10} \doteq 1.0717$, the real, positive 10th root of 2. One of the roots is located with polar angle $\pi/20$, and the others have polar angles of $\pi/20 + 2\pi/5, \pi/20 + 4\pi/5, \pi/20 + 6\pi/5$, and $\pi/20 + 8\pi/5$, respectively. (See Fig. 1.8.) \square




Figure 1.8

Example 3 Solve the equation

$$z^4 - 4z^2 + 4 - 2i = 0.$$

Solution The equation may be rearranged as

$$z^4 - 4z^2 + 4 = 2i$$

or

$$(z^2 - 2)^2 = 2i = (1 + i)^2.$$

This has solutions

$$z^2 - 2 = \begin{cases} 1 + i \\ -1 - i. \end{cases}$$

Equivalently,

$$z^2 = \begin{cases} 3 + i \\ 1 - i. \end{cases}$$

These may be solved to give the four solutions of the original equation,

$$z_1 = \sqrt[4]{10}(\cos \theta_0 + i \sin \theta_0),$$

$$z_2 = -\sqrt[4]{10}(\cos \theta_0 + i \sin \theta_0), \quad \theta_0 = \frac{1}{2} \arctan \frac{1}{3}$$

and

$$z_3 = \sqrt[4]{2}(\cos \theta_1 + i \sin \theta_1),$$

$$z_4 = -\sqrt[4]{2}(\cos \theta_1 + i \sin \theta_1), \quad \theta_1 = \frac{1}{2} \arctan(-1) = -\frac{\pi}{8}. \quad \square$$

Circles

A circle is the set of all points equidistant from a given point, the center. If z_0 is the center and r the radius, then the circle of radius r and center z_0 is described by the equation $|z - z_0| = r$. There are, however, other ways to use complex numbers to describe circles.

If p and q are distinct complex numbers, then those complex numbers z with

$$|z - p| = |z - q|$$

are equidistant from p and q . The locus of these points is precisely the straight line that is the perpendicular bisector of the line segment joining p to q . However, if ρ is a positive real number not equal to 1, those z with

$$|z - p| = \rho |z - q|$$

form a circle. To see this, suppose that $0 < \rho < 1$ (otherwise, divide both sides of the equation by ρ). Let $z = w + q$ and $c = p - q$; then the equation becomes

$$|w - c| = \rho |w|.$$

Upon squaring and transposing terms, this can be written as

$$|w|^2(1 - \rho^2) - 2 \operatorname{Re} w\bar{c} + |c|^2 = 0.$$

We complete the square of the left side and find that

$$(1 - \rho^2)|w|^2 - 2 \operatorname{Re} w\bar{c} + \frac{|c|^2}{1 - \rho^2} = \frac{|c|^2\rho^2}{1 - \rho^2}.$$

Equivalently,

$$\left| w - \frac{c}{1 - \rho^2} \right| = |c| \frac{\rho}{1 - \rho^2}.$$

Thus, w lies on the circle of radius $R = |c|\rho/(1 - \rho^2)$ centered at the point $c/(1 - \rho^2)$, and so z lies on the circle of the same radius R centered at the point

$$z_0 = \frac{p - \rho^2 q}{1 - \rho^2} = \frac{1}{1 - \rho^2} p - \frac{\rho^2}{1 - \rho^2} q.$$

Example 4 To confirm in one special case what was just done, let us look at the locus of points z with

$$|z - i| = \frac{1}{2}|z - 1|.$$

After multiplying both sides by 2 and squaring,

$$4\{|z|^2 - 2 \operatorname{Re}(z\bar{i}) + |i|^2\} = |z|^2 - 2 \operatorname{Re} z + |1|^2,$$

or after simplifying,

$$3|z|^2 - 8y + 2x = -3.$$

More algebra yields

$$3x^2 + 2x + \frac{1}{3} + 3y^2 - 8y + \frac{16}{3} = -3 + \frac{17}{3} = \frac{8}{3}.$$

Thus, the locus is

$$\left(x + \frac{1}{3}\right)^2 + \left(y - \frac{4}{3}\right)^2 = \frac{8}{9}.$$

This is a circle of radius $2(\sqrt{2}/3)$ centered at $-1/3 + 4i/3$. Now, in the notation that preceded the example, $p = i$, $q = 1$, and $\rho = \frac{1}{2}$. The radius should be

$$R = \frac{|p - q|\rho}{1 - \rho^2} = \frac{\sqrt{2}(\frac{1}{2})}{\frac{3}{4}} = \frac{2\sqrt{2}}{3},$$

and the center at

$$z_0 = \frac{p - \rho^2 q}{1 - \rho^2} = \frac{-\frac{1}{4} + i}{\frac{3}{4}} = -\frac{1}{3} + \frac{4}{3}i,$$

which, of course, is just what was previously found. Note that the center of the circle is on the line through 1 and i (Fig. 1.9). \square




Figure 1.9

We now apply the information just derived to produce a beautiful geometric pattern: two families of mutually perpendicular circles.

Let C_1 be the family of circles of the form

$$|z - p| = \rho|z - q|, \quad 0 < \rho < \infty,$$

where we include the case $\rho = 1$ (which yields a straight line) for completeness. Let L be the perpendicular bisector of the line segment from p to q . Take C_2 to be the family of circles through p and q and centered on the line L . We shall show that each circle in the family C_1 is perpendicular to each circle in the family C_2 at their two points of intersection. The computation is considerably simplified by locating the origin at the point of intersection of the line L and the line L' , which passes through p and q . L' can then be taken to be the real axis and L to be the imaginary axis; in this way, we may assume that $0 < p = -q$. A circle from the family C_1 is then centered at a point on the real axis, and because there is no loss in assuming $0 < \rho < 1$, the center of that circle is at the point $s = p(1 + \rho^2)/(1 - \rho^2)$. The center

of the circle from the family C_2 is at the point $t = i\alpha$ (α real), and this circle must pass through p and $-p$. Let $z = x + iy$ be on both circles (see Fig. 1.10).




Figure 1.10

Since z is on the circle C_1 ,

$$|z - p| = \rho|z + p|,$$

and consequently,

$$x^2(1 - \rho^2) - 2\rho x(1 + \rho^2) + p^2(1 - \rho^2) + y^2(1 - \rho^2) = 0.$$

For notational convenience, set $v = (1 + \rho^2)/(1 - \rho^2)$; the above equation then becomes

$$x^2 - 2\rho vx + p^2 + y^2 = 0.$$

On the other hand, since $z = x + iy$ is also on the circle C_2 ,

$$|z - i\alpha| = |p - i\alpha| = |-p - i\alpha| = \sqrt{p^2 + \alpha^2},$$

and so

$$x^2 + y^2 - 2\alpha y = p^2.$$

In order to prove that the segment from $i\alpha$ to z is perpendicular to the segment s to z , it must be shown that $\operatorname{Re}[(\bar{z} - i\alpha)(z - s)] = 0$; this follows from the discussion

of perpendicularity near the end of Section 1. However, using $s = p\mathbf{v}$,

$$\begin{aligned} 2 \operatorname{Re}[(z - i\alpha)(\overline{z} - s)] &= 2x(x - s) + 2y(y - \alpha) \\ &= 2(x^2 + y^2 - p\mathbf{v}x - \alpha y) \\ &= x^2 + y^2 - 2p\mathbf{v}x + x^2 + y^2 - 2\alpha y \\ &= -p^2 + p^2 = 0. \end{aligned}$$

This is the desired conclusion.

An illustration (with $p = -q = 1$) of several circles from each of these families is shown in Figure 1.11. This pattern is often called the **Circles of Appolonius**. These will come to your attention again later in the book.




Figure 1.11

EXERCISES FOR SECTION 1.2

In Exercises 1 to 10, describe the locus of points z satisfying the given equation.

- | | |
|--|--------------------------------------|
| 1. $ z + 1 = z - 1 $ | 2. $ z - 4 = 4 z $ |
| 3. $\operatorname{Re}[(4 + i)z + 6] = 0$ | 4. $\operatorname{Im}(2iz) = 7$ |
| 5. $ z + 2 + z - 2 = 5$ | 6. $ z - i = \operatorname{Re} z$ |
| 7. $\operatorname{Re}(z^2) = 4$ | 8. $ z - 1 ^2 = z + 1 ^2 + 6$ |
| 9. $ z^2 - 1 = 0$ | 10. $ z + 1 ^2 + 2 z ^2 = z - 1 ^2$ |

In Exercises 11 to 17, write the equation of the given circle or straight line in complex number notation. For example, the circle of radius 4 centered at the point $3 - 2i$ is given by the equation $|z - (3 - 2i)| = 4$.

11. The circle of radius 2 centered at $4 + i$.
12. The straight line through 1 and $-1 - i$.
13. The vertical line containing $-3 - i$.
14. The circle through $0, 2 + 2i$, and $2 - 2i$.
15. The circle through $1, i$, and 0 .
16. The perpendicular bisector of the line segment joining $-1 + 2i$ and $1 - 2i$.
17. The straight line of slope -2 through $1 - i$.
18. Show that the two lines $\operatorname{Re}(az + b) = 0$ and $\operatorname{Re}(cz + d) = 0$ are perpendicular if and only if $\operatorname{Re}(a\bar{c}) = 0$.
19. Let p be a positive real number and let Γ be the locus of points z satisfying $|z - p| = cx$, $z = x + iy$. Show that Γ is (a) an ellipse if $0 < c < 1$; (b) a parabola if $c = 1$; (c) a hyperbola if $1 < c < \infty$.
20. Let z_1 and z_2 be distinct complex numbers. Show that the locus of points $tz_1 + (1 - t)z_2$, $-\infty < t < \infty$, describes the line through z_1 and z_2 . The values $0 \leq t \leq 1$ give the line segment joining z_1 and z_2 .
21. Let α be a complex number with $0 < |\alpha| < 1$. Show that the set of all z with
 - (a) $|z - \alpha| < |1 - \bar{\alpha}z|$ is the disc $\{z: |z| < 1\}$
 - (b) $|z - \alpha| = |1 - \bar{\alpha}z|$ is the circle $\{z: |z| = 1\}$
 - (c) $|z - \alpha| > |1 - \bar{\alpha}z|$ is the set $\{z: |z| > 1\}$.

(Hint: Square both sides and simplify.)
22. Let z and w be nonzero complex numbers. Show that $|z + w| = |z| + |w|$ if and only if $z = sw$ for some positive real number s .

In Exercises 23 to 26, follow the technique outlined in the text to find all solutions of the given equation.

23. $z^5 = i$
24. $(z + 1)^4 = 1 - i$
25. $z^8 = -1$
26. $z^3 = 8$
27. Suppose that n is an odd integer and w is a negative real number. Show that one solution of the equation $z^n = w$ is a negative real number. (For instance, -2 is a root of $z^3 = -8$.)
28. Let a , b , and c be complex numbers with $a \neq 0$. Show that the solutions of $az^2 + bz + c = 0$ are $z_1, z_2 = (\frac{1}{2a})(-b \pm \sqrt{b^2 - 4ac})$, just as they are in the case when a , b , and c are real numbers.
29. Let b and c be complex numbers. Show that the roots of the quadratic equation $z^2 + bz + c = 0$ are complex conjugates of each other if and only if the quantity $b^2 - 4c$ is real and negative, b is real, and c is positive.
30. Let A be a complex number and B a real number. Show that the equation $|z^2| + \operatorname{Re}(Az) + B = 0$ has a solution if and only if $|A|^2 \geq 4B$. If this is so, show that the solution set is a circle or a single point.
31. Let C be a circle and let A and B be any two distinct points on C . Show that if P is selected on the smaller arc of C joining A to B , then the angle from the segment AP to the segment BP is independent of P . This angle is $\pi/2$ if A and B are on opposite ends of a diameter. The result remains true if “smaller” is replaced by “larger.”

32. Let z_1, \dots, z_n be complex numbers. Show by mathematical induction that $|z_1 + \dots + z_n| \leq |z_1| + \dots + |z_n|$.

Translation and Scaling*

33. Let C be a circle or a straight line. Show that the same is true of the locus of points $z + \beta$, $z \in C$, and β a fixed complex number.
34. Let C be a circle or a straight line. Show that the same is true of the locus of points αz , $z \in C$, and α a fixed nonzero complex number.

Inversion*

35. Let L be the line $y = a$, $a > 0$. Show that the locus of points $1/z$, $z \in L$, is the circle of radius $1/2a$ centered at $-i/2a$.
36. Let L be a line through the origin. Show that the locus of points $1/z$, $z \in L$, is a line through the origin. What is the relationship of the slopes of the two lines?
37. Let C be the circle $|z - c| = r$, $0 < r < c$. Show that the locus of points $1/z$, $z \in C$, is the circle centered at $c/(c^2 - r^2)$, of radius $r/(c^2 - r^2)$.
38. Let C be the circle $|z - r| = r$, $r > 0$. Show that the locus of points $1/z$, $z \in C$, is the vertical line through $1/2r$.

1.3 Subsets of the Plane

To understand the fundamentals of complex variables, it is necessary to single out several special types of subsets of the complex plane that will be used in our discussion of analytic and harmonic functions in subsequent chapters. This section gives the definitions and basic properties of these sets.

Open Sets

The set consisting of all points z satisfying $|z - z_0| < R$ is called the **open disc** of radius R centered at z_0 . A point w_0 in a set D in the complex plane is called an **interior point** of D if there is some open disc centered at w_0 that lies entirely within D (see Fig. 1.12). A set D is called **open** if all of its points are interior points.




Figure 1.12

Example 1 Each open disc $D = \{z: |z - z_0| < R\}$ is an open set. For if $r = |w_0 - z_0| < R$, choose $\varepsilon = (R - r)/3$. Then, for any z with $|z - w_0| < \varepsilon$, we have

$$|z - z_0| \leq |z - w_0| + |w_0 - z_0| < \varepsilon + r = (R - r)/3 + r < R,$$

by the triangle inequality. Thus, the open disc of radius ε centered at w_0 lies within the set D . Hence, each point of D is an interior point, so D is open. \square

Example 2 The set $R = \{z: \operatorname{Re} z > 0\}$ is an open set. To see this, let $w_0 \in R$; then $\sigma_0 = \operatorname{Re} w_0 > 0$. Let $\varepsilon = \frac{1}{2}\sigma_0$ and suppose that $|z - w_0| < \varepsilon$. Then $-\varepsilon < \operatorname{Re}(z - w_0) < \varepsilon$, so

$$\operatorname{Re} z = \operatorname{Re}(z - w_0) + \operatorname{Re} w_0 > -\varepsilon + \sigma_0 = \frac{1}{2}\sigma_0 > 0.$$

Consequently, z also lies in R . Hence, each point w_0 of R is an interior point, so R is open. \square

Example 3 The set $\{z: |\operatorname{Im} z| > 1\}$ is an open set. This can be shown in the manner of Example 2. \square

Example 4 The set of all points $z = x + iy$, with $x^2 < y$, is also an open set. Let $z_0 = x_0 + iy_0$ be in this set; then there is a positive δ with $x_0^2 + \delta^2 < y_0$. We may also assume that δ is so small that $2x_0\delta + \delta^3 < 1 + \delta$. (The latter inequality will be needed only at one rather technical point.) Suppose now that $z = x + iy$ satisfies $|z - z_0| < \delta^2$. Then $\delta^2 > |x - x_0|$, and $\delta^2 > |y - y_0|$. Therefore,

$$\begin{aligned} x^2 &< (x_0 + \delta^2)^2 = x_0^2 + 2x_0\delta^2 + \delta^4 \\ &< y_0 - \delta + 2x_0\delta^2 + \delta^4 \\ &< y - \delta^2 - \delta + 2x_0\delta^2 + \delta^4. \end{aligned}$$

However, the choice of δ gives $-\delta^2 - \delta + 2x_0\delta^2 + \delta^4 < 0$, so $x^2 < y$ whenever $z = x + iy$ satisfies $|z - z_0| < \delta^2$. \square

Example 5 The set of all z with $\operatorname{Re} z \leq 6$ is not an open set. \square

The Boundary of a Set

The set just described in Example 5 is not open because, for instance, the point $w_0 = 6$ is in it, but every open disc centered at w_0 , no matter how small the radius, must contain a point z with $\operatorname{Re} z > 6$. The point $w_0 = 6$ is called a *boundary point* because it is located on the edge of the set, at the place where the set “almost” meets its complement. The precise definition follows.

A point p is a **boundary point** of a set S if every open disc centered at p contains both points of S and points not in S . The set of all boundary points of a set S is called the **boundary** of S (Fig. 1.13).

Let us find the boundary of each of the sets in the preceding examples (see Fig. 1.13).




Figure 1.13

Example 6 The boundary of the open disc of radius R centered at z_0 is the circle of radius R centered at z_0 . \square

Example 7 The boundary of the set of those z with $\text{Re } z > 0$ is the imaginary axis. \square

Example 8 The boundary of the set of those z with $|\operatorname{Im} z| > 1$ is the set of those z with $|\operatorname{Im} z| = 1$. \square

Example 9 The boundary of the set of those $z = x + iy$ with $x^2 < y$ is the parabola $y = x^2$. \square

Example 10 The boundary of the set of those z with $\operatorname{Re} z \leq 6$ is the vertical line $\operatorname{Re} z = 6$. \square

Closed Sets

You have observed by now that the open sets just discussed contain none of their boundary. Indeed, this is an elementary theorem, which is set out below. However, what about exactly the reverse situation? A set C is called **closed** if it contains its boundary. These definitions prepare us for the following result.

THEOREM A set D is open if and only if it contains no point of its boundary. A set C is closed if and only if its complement $D = \{z: z \notin C\}$ is open.

Proof To establish the validity of this theorem, we start by supposing that D is an open set. Let p be a boundary point of D . If it happens that p is in D , then (because D is open) there is an open disc centered at p that lies within D . Thus, p is not in the boundary of D . This contradiction shows that p is not in D . Conversely, suppose D is a set that contains none of its boundary points; we must show that D is open. If $z_0 \in D$, then z_0 cannot be a boundary point of D , so there is some disc centered at z_0 that is either a subset of D or a subset of the complement of D . The latter is impossible, since z_0 itself is in D . Hence, the disc lies in D , and we have shown that each point of D is an interior point; therefore, D is open.

The second assertion of the theorem follows immediately from the fact that the boundary of a set coincides exactly with the boundary of the complement of that set; this in turn is a direct consequence of the definition of boundary point. ■

Be wary here—there are many sets that are neither open nor closed, since they contain part, but not all, of their boundary. For example, the set D of those z with $\operatorname{Re} z \leq 6$ and $\operatorname{Im} z > 2$ is neither open nor closed, since its boundary consists of those w with either $\operatorname{Re} w = 6$ and $\operatorname{Im} w \geq 2$ or $\operatorname{Im} w = 2$ and $\operatorname{Re} w \leq 6$ (Fig. 1.14).

There is also one technical point here: The complex plane itself has no boundary and so by definition is both open and closed. It happens to be the only nonempty subset of the plane that is both open and closed. (The truth of this statement depends on a fundamental property of the real numbers.) Since what follows makes no use of this fact about the plane, we will not prove it here.

Connected Sets

A **polygonal curve** is the union of a finite number of directed line segments P_1P_2 , $P_2P_3, \dots, P_{n-1}P_n$, where the terminal point of one is the initial point of the next (except for the last) (see Fig. 1.15). An open set D is **connected** if each pair p, q of




Figure 1.14




Figure 1.15

points in D may be joined by a polygonal curve lying entirely with D . That is, there are points P_2, \dots, P_{n-1} in D such that all the line segments $pP_2, P_2P_3, \dots, P_{n-1}q$ lie in D .

Example 11 An open disc is connected. □

Example 12 The set of those z with $\operatorname{Re} z > 0$ is connected. □

Example 13 The set of those $z = x + iy$ with $x^2 < y$ is connected. □

Example 14 The set of those z with $|\operatorname{Im} z| > 1$ is not connected. □

Example 15 The set of those z with $\operatorname{Re} z < 6$ is connected. □

Example 16 The set of those z with $\operatorname{Re} z \neq 0$ is not connected. □

For instance, in Example 16 the points $p = 1$ and $q = -1$ lie in the given set, but any polygonal curve joining p to q must necessarily cross the imaginary axis (where $\operatorname{Re} z = 0$), so this curve cannot lie entirely within the set $\operatorname{Re} z \neq 0$.

An open connected set is called a **domain**. Domains are the natural setting for the study of analytic and harmonic functions.

A set S is **convex** if the line segment pq joining each pair of points p, q in S also lies in S . In particular, it is immediately clear that any convex open set is connected (Fig. 1.16).




Figure 1.16

Example 17 Each open disc is convex. □

Example 18 The set of those $z = x + iy$ with $x^2 > y$ is not convex. □

Example 19 Both of the sets $\operatorname{Re} z > 0$ and $\operatorname{Re} z \leq 6$ are convex. □

An **open half-plane** is defined to be those points strictly to one side of a straight line—that is, those points z for which $\operatorname{Re}(az + b) > 0$, say. Each open half-plane is a convex set and an open set as well. A **closed half-plane** is the open half-plane plus the defining line—that is, those z with $\operatorname{Re}(az + b) \geq 0$. Each closed half-plane is convex as well as closed (Fig. 1.17).




Figure 1.17

It is instructive to see why each half-plane (open or closed) is convex. If p and q are two points, then

$$tq + (1 - t)p, \quad 0 \leq t \leq 1$$

describes the line segment from p to q ; see Exercise 20, Section 2. If p and q are in the open half-plane given by $\operatorname{Re}(az + b) > 0$, then

$$\operatorname{Re}(a(tq + (1 - t)p) + b) = t \operatorname{Re}(aq + b) + (1 - t) \operatorname{Re}(ap + b) > 0,$$

so the line segment from p to q lies in the same open half-plane. The case of a closed half-plane is almost identical.

The Point at Infinity

A helpful and frequently used convention is to add the **point at infinity** to the complex plane. This is understood in the following way. A set D contains the point at infinity in its interior if there is a large number M such that D contains all the points z with $|z| > M$. For instance, the open half-plane $\operatorname{Re} z > 0$ does not contain the point at infinity, but the open set $D = \{z: |z + 1| + |z - 1| > 1\}$ does. One “reaches” the point at infinity by letting $|z|$ increase without bound, with no restriction at all on $\arg z$. One way to visualize all this is to let $w = 1/z$ and think about $|w|$ being very small; an open set containing the point at infinity will become an open set containing $w_0 = 0$. Further, the statement “ z approaches infinity” is identical to the statement “ w converges to zero.” The point at infinity is denoted by the usual symbol for infinity: ∞ .

EXERCISES FOR SECTION 1.3

For each of the sets in Exercises 1 to 8, (a) describe the interior and the boundary, (b) state whether the set is open or closed or neither open nor closed, (c) state whether the interior of the set is connected (if it has an interior).

1. $A = \{z = x + iy: x \geq 2 \text{ and } y \leq 4\}$
2. $B = \{z: |z| < 1 \text{ or } |z - 3| \leq 1\}$
3. $C = \{z = x + iy: x^2 < y\}$
4. $D = \{z: \operatorname{Re}(z^2) = 4\}$
5. $E = \{z: z\bar{z} - 2 \geq 0\}$
6. $F = \{z: z^3 - 2z^2 + 5z - 4 = 0\}$
7. $G = \{z = x + iy: |z + 1| \geq 1 \text{ and } x < 0\}$
8. $H = \{z = x + iy: -\pi \leq y < \pi\}$
9. Let α and β be complex numbers with $\beta \neq 0$. Describe the set of points $\alpha z + \beta$ as z varies over (a) the first quadrant, $\{z = x + iy: x > 0 \text{ and } y > 0\}$; (b) the upper half-plane, $\{z = x + iy: y > 0\}$; (c) the disc $\{z: |z| < R\}$. Show that in each case the resulting set is open and connected. (**Hint:** First investigate the set αz .)
10. Describe the set of points z^2 as z varies over the second quadrant: $\{z = x + iy: x < 0 \text{ and } y > 0\}$. Show that this is an open, connected set. (**Hint:** Use the polar representation of z .)
11. A set S in the plane is **bounded** if there is a positive number M such that $|z| \leq M$ for all z in S ; otherwise, S is **unbounded**. In Exercises 1 to 8, six of the given sets are unbounded. Find them.

12. Which, if any, of the sets given in Exercises 1 to 8 contains ∞ ?
13. (a) Show that the union of two nonempty open sets is open. Do the same, replacing “open” with “closed.” Do the same replacing “union” with “intersection.”
 (b) Repeat part (a) replacing “two” with “finitely many.”
14. Let D_1 and D_2 be domains with a nonempty intersection. Show that $D_1 \cup D_2$ is a domain.
15. Let $\Omega_1 = \{z: 1 < |z| < 2 \text{ and } \operatorname{Re} z > -\frac{1}{2}\}$ and $\Omega_2 = \{z: 1 < |z| < 2 \text{ and } \operatorname{Re} z < \frac{1}{2}\}$. Show that both Ω_1 and Ω_2 are domains but $\Omega_1 \cap \Omega_2$ is not.
16. Let D be a domain and let p and q be points of D . Show that there is a polygonal curve joining p to q whose line segments are either horizontal or vertical (both types can be used). (**Hint:** Replace a “slanting” segment by (perhaps many) horizontal and vertical segments; see Fig. 1.18.)




Figure 1.18

17. Fix a nonzero complex number z_0 . Show that the set D obtained from the plane by deleting the ray $\{tz_0: 0 \leq t < \infty\}$ is a domain.
18. An open set D is **star-shaped** if there is some point p in D with the property that the line segment from p to z lies in D for each z in D . (a) Show that the disc $\{z: |z - z_0| < r\}$ is star-shaped. (b) Show that any convex set is star-shaped.
19. Determine which of the following sets are star-shaped:
 (a) $D = \{z: |z - 1| < 2 \text{ or } |z + 1| < 2\}$
 (b) $D = \{z = x + iy: x > 0 \text{ and } |z| > 1\}$
 (c) $D = \{z: |z| > 1\}$
 (d) $D = \{z = x + iy: x > 0 \text{ and } [x > y + 1 \text{ or } x > 1 - y]\}$
20. Show that each star-shaped set is connected.

Separation of a Point and Convex Set*

Let C be a closed convex set and z_0 a point *not* in C . It is a fact that there is a point p in C with $r = |z_0 - p| \leq |z_0 - q|$ for all q in C . (This last statement requires a bit of proof, but let us assume its validity.)

21. Show that the only point of C in the disc $|z_0 - z| \leq r$ is the point p .
22. Let L be the perpendicular bisector of the line segment from z_0 to p . Show that no point of C lies on L or in the half-plane, determined by L , which contains z_0 .
23. Conclude from Exercise 22 that L separates z_0 from C : z_0 and C lie in the two open half-planes determined by L , but not in the same open half-plane.
24. Show that each closed convex set is the intersection of all the closed half-planes that contain it.

Topological Properties*

25. Show that the boundary of any set D is itself a closed set.
26. Show that if $p \in D$, then p is either an interior point of D or a boundary point of D .
27. Show that a set D coincides with its boundary if and only if D is closed and D has no interior points.
28. Show that if D is a set and E is a *closed* set containing D , then E must contain the boundary of D .
29. Show that if D is a set and S is an *open* set that is a subset of D , then S must be composed entirely of interior points of D .
30. Let C be a bounded closed convex set and let D be the complement of C . Show that D is a domain.

1.4 Functions and Limits

A **function** of the complex variable z is a rule that assigns a complex number to each z within some specified set D ; D is called the **domain of definition** of the function. The collection of all possible values of the function is called the **range** of the function. Thus, a function f has as its domain of definition some subset of the complex plane and as its range some other (usually entirely different) subset of the complex plane. We frequently write $w = f(z)$ to distinguish the independent complex variable z from the dependent complex variable w .

Example 1 $f(z) = 4z^2 + 2z + 1$ has as its domain of definition the entire complex plane. Its range is also the entire complex plane, for if w is any complex number, then the equation $f(z) = w$ is nothing but the quadratic equation

$$4z^2 + 2z + 1 - w = 0.$$

This equation is solved by use of the quadratic formula. Its solutions are

$$z_j = \frac{-2 + s_j}{8}, \quad j = 1, 2,$$

where s_1, s_2 are the two square roots of $4 - 16(1 - w) = -12 + 16w$. (It is possible that $s_1 = s_2$; this occurs only when $w = \frac{3}{4}$.) \square

Example 2 $f(z) = 1/(z - 1)$ has as its domain of definition all complex numbers except $z = 1$, where it is not defined. Its range consists of all complex numbers w except $w = 0$, since $f(z) = w = 1/(z - 1)$ is solved by $z = 1 + (1/w)$; this is a complex number as long as $w \neq 0$. \square

Example 3 $g(z) = |z|^2$ has the complex plane as its domain of definition; its range consists of all nonnegative real numbers. \square

Example 4 $h(z) = i(2 - (\operatorname{Im} z)^{-1})$ is defined for all z except those on the real axis. For a pure imaginary number w , a solution of $w = h(z) = i(2 - (\operatorname{Im} z)^{-1})$ is any complex number z with $\operatorname{Im} z = (2 + iw)^{-1}$. This defines many complex numbers as long as $w \neq 2i$. Hence, the range of h is all purely imaginary numbers except $w = 2i$. \square

Example 5 Show that the range of the function $w = T(z) = (1 + z)/(1 - z)$ on the disc $|z| < 1$ is the set of those w whose real part is positive.

Solution Compute the real part of w :

$$\operatorname{Re} w = \operatorname{Re} \left(\frac{1+z}{1-z} \right) = \operatorname{Re} \frac{(1+z)(1-\bar{z})}{|1-z|^2} = \frac{1-|z|^2}{|1-z|^2}.$$

This last quantity is positive when $|z| < 1$. This shows that the range of T is a subset of those w with $\operatorname{Re} w > 0$. Now let w' be any point with $\operatorname{Re} w' > 0$; we shall show that $z' = (w' - 1)/(w' + 1)$ satisfies $|z'| < 1$. Indeed, $1 > |(w' - 1)/(w' + 1)|$ exactly when $|w' + 1|^2 > |w' - 1|^2$. We expand both $|w' + 1|^2$ and $|w' - 1|^2$ and obtain

$$|w'|^2 + 2 \operatorname{Re} w' + 1 > |w'|^2 - 2 \operatorname{Re} w' + 1.$$

This is a correct inequality because $\operatorname{Re} w' > 0$. Thus, $z' = (w' - 1)/(w' + 1)$ lies in the disc $|z| < 1$, and

$$Tz' = \frac{1+z'}{1-z'} = \frac{1 + \frac{w'-1}{w'+1}}{1 - \frac{w'-1}{w'+1}} = \frac{2w'}{2} = w'. \quad \square$$

(NOTE: $T(z) = (1 + z)/(1 - z)$ is an example of a **linear fractional transformation**, which will be studied in some detail in Section 3 of Chapter 3.)

As with functions of a real variable, the domain of definition of a function is usually easier to determine than the range. In many cases, this text will provide only a general description of certain properties of the range rather than an explicit rule

for each point in it. For instance, we may be able to state that the range is open or connected or convex.

Graphs

For real-valued functions of a real variable, like those studied extensively in calculus, the device of displaying the graph of the function is an extremely useful tool in visualizing the behavior of the function. Such a technique is not as readily available for functions of a complex variable. If the function f has only real values, its graph in 3-space (x, y, t) , $t = f(x + iy)$, can be sketched (see Fig. 1.19). However, if the function has complex values, this type of picture is not possible (at least in our world!). Moreover, almost all the functions dealt with here have complex values. One way to proceed is to graph $|f|$, but a more useful way is to use two complex planes: one for the domain variable z and a second for the range variable $w = f(z)$. For instance, we saw in Example 5 that the function $T(z) = (1 + z)/(1 - z)$ maps the disc $\{z: |z| < 1\}$ onto those w with $\operatorname{Re} w > 0$; this can be “graphed” as shown in Figure 1.20. This type of picture is very helpful in understanding the behavior of the complex-valued function of a complex variable and we will employ it frequently.




Figure 1.19




Figure 1.20

Limits

The concepts of the limit of a sequence of complex numbers, or the limit of a function of a complex variable, or even of continuity of a function of a complex variable, are almost identical to those for a real variable. Let us begin with the limit of a sequence.

Let $\{z_n\}_{n=1}^{\infty}$ be a sequence of complex numbers. We say that $\{z_n\}$ has the complex number A as a **limit**, or that $\{z_n\}$ **converges** to A , and we write

$$\lim_{n \rightarrow \infty} z_n = A \quad \text{or} \quad z_n \rightarrow A$$

if, given any positive number ε , there is an integer N such that

$$|z_n - A| < \varepsilon \quad \text{for all } n \geq N.$$

A sequence that does not converge, for any reason whatever, is called **divergent**. If $z_n = x_n + iy_n$ and $A = s + it$, then $z_n \rightarrow A$ if and only if $x_n \rightarrow s$ and $y_n \rightarrow t$. This is due to three inequalities noted in Section 1, namely

$$|x_n - s| \leq |z_n - A|, \quad |y_n - t| \leq |z_n - A|,$$

and

$$|z_n - A| \leq |x_n - s| + |y_n - t|.$$

We can equivalently state that the sequence $\{z_n\}$ of complex numbers converges to the complex number A if and only if, whenever D is any open disc centered at A , all but a finite number of the points $\{z_n\}$ lie in D .

Example 6 The sequence $z_n = 1 + (i/n)$ converges to 1. □

Example 7 The sequence $z_n = (-\frac{1}{2})^n + i(1 - (1/2n))$ converges to i . □

Example 8 The sequence $z_n = (1/n)(\cos(n\pi/4) + i \sin(n\pi/4))$ converges to zero. \square

Example 9 The sequence $z_n = n - (1/n)$ diverges. \square

Example 10 The sequence $z_n = i^n$ diverges because its terms are $i, -1, -i, 1$, in that order, repeated infinitely often. \square

One simple consequence of the definition of convergence is this:

$$\text{If } z_n \rightarrow A, \quad \text{then } |z_n| \rightarrow |A|.$$

The converse of this assertion is generally false; for example, $|(-1)^n + i/n| \rightarrow 1$, but the sequence $\{(-1)^n + i/n\}$ itself has no limit.

You will no doubt recall from calculus results about the sum, product, and quotient of convergent sequences of real numbers. Similar statements hold for sequences of complex numbers; we collect these results in the next theorem.

THEOREM 1 Let $\{z_n\}$ and $\{w_n\}$ be convergent sequences of complex numbers with limits A and B , respectively. Let λ be a complex number. The sequences $\{z_n + \lambda w_n\}$ and $\{z_n w_n\}$ then also both converge with limits $A + \lambda B$ and AB , respectively. Furthermore, if $B \neq 0$, the sequence $\{z_n/w_n\}$ converges to A/B . \square

Proof The proof of each of these results is quite direct. For example, to show that $z_n w_n \rightarrow AB$, we write $z_n w_n - AB = (z_n - A)B + (w_n - B)z_n$. Let N_1 be chosen so big that $|z_n - A| < \varepsilon'$ if $n \geq N_1$ and N_2 chosen so big that $|w_n - B| < \varepsilon'$ if $n \geq N_2$. Then $|z_n| \leq \varepsilon' + |A| < 1 + |A|$ if $\varepsilon' < 1$; and for $n \geq N = N_1 + N_2$, we obtain

$$\begin{aligned} |z_n w_n - AB| &\leq |z_n - A||B| + |w_n - B||z_n| \\ &< \varepsilon'|B| + \varepsilon'(1 + |A|) \\ &= \varepsilon'(1 + |A| + |B|) \\ &< \varepsilon, \end{aligned}$$

if ε' is chosen initially to satisfy $\varepsilon'(1 + |A| + |B|) < \varepsilon$. \blacksquare

Example 11 The sequence $z_n = 1 + i[1 - (2/n)]$ converges to $1 + i$ as $n \rightarrow \infty$. Hence, the sequence

$$z_n^2 = \frac{4}{n} - \frac{4}{n^2} + i\left(2 - \frac{4}{n}\right)$$

converges to $2i = (1 + i)^2$ as $n \rightarrow \infty$. \square

Example 12 Suppose that a_0, a_1, a_2 , and a_3 are complex numbers and $\{z_n\}$ is a sequence with $z_n \rightarrow A$. Then the sequence $\{a_0 + a_1 z_n + a_2 z_n^2 + a_3 z_n^3\}_{n=1}^\infty$ converges

to $a_0 + a_1 A + a_2 A^2 + a_3 A^3$ as $n \rightarrow \infty$, as can be seen by several applications of results in Theorem 1. \square

Suppose next that f is a function defined on a subset S of the plane. Let z_0 be a point either in S or in the boundary of S . We say that f has limit L at the point z_0 , and we write

$$\lim_{z \rightarrow z_0} f(z) = L \quad \text{or} \quad f(z) \rightarrow L \quad \text{as} \quad z \rightarrow z_0$$

if, given $\varepsilon > 0$, there is a $\delta > 0$ such that

$$|f(z) - L| < \varepsilon \quad \text{whenever} \quad z \in S \quad \text{and} \quad |z - z_0| < \delta.$$

It is worth stressing here that f has the limit L at the point z_0 exactly when the numbers $f(z)$ approach L as z approaches z_0 in any manner whatsoever from within S . This is substantially different from the case of a function of a real variable, where the real variable can approach only from the left or right. The complex variable z may approach z_0 from infinitely many directions. Once again, it will be informative to look at some examples.

Example 13 The function $f(z) = |z|^2$ has limit 4 at the point $z_0 = 2i$. \square

Example 14 The function $g(z) = 1/(1 - z)$ has limit $(1/2)(1 + i)$ at $z_0 = i$. \square

Example 15 The function $h(z) = \operatorname{Re}(z^4 + 4)$ has limit 0 at $z_0 = 1 + i$. \square

Example 16 The function $f(z) = (z^4 - 1)/(z - i)$ has limit $-4i$ at $z_0 = i$, since $z^4 - 1 = (z - i)(z + i)(z - 1)(z + 1)$, so $f(z)$ simplifies to $(z + i)(z + 1)(z - 1)$ so long as $z \neq i$. \square

Example 17 The function $f(z) = z/\bar{z}$, $z \neq 0$, has no limit at $z_0 = 0$. For if z is real, then $f(z) = 1$, while if z is purely imaginary, $z = iy$, then $f(iy) = -1$. Such a function cannot have a limit at $z_0 = 0$. \square

We say that the function f has a limit L at ∞ , and we write

$$\lim_{z \rightarrow \infty} f(z) = L$$

if, given $\varepsilon > 0$, there is a large number M such that

$$|f(z) - L| < \varepsilon \quad \text{whenever} \quad |z| \geq M.$$

Note that we only require that $|z|$ is large; there is no restriction at all on $\arg z$.

Example 18 $\lim_{z \rightarrow \infty} 1/z^m = 0$ if $m = 1, 2, \dots$. \square

Example 19 $\lim_{z \rightarrow \infty} [(z^4 + 1)/(z^4 + 5z^2 + 3)] = 1$, since the ratio can be written as

$$\frac{1 + \frac{1}{z^4}}{1 + \frac{5}{z^2} + \frac{3}{z^4}},$$

all of whose terms except the 1's go to zero as $|z| \rightarrow \infty$. \square

Example 20 $\lim_{z \rightarrow \infty} [(x + y^3)/(x^2 + y^3)]$ does not exist; for if we take $z = x$, then the expression goes to zero, but if we take $z = iy$, then the expression is identically 1. \square

The following theorem on the sum, product, and quotient of functions that have limits at z_0 is a counterpart to Theorem 1.

THEOREM 2 Suppose that f and g are functions with limits L and M , respectively, at z_0 . Let λ be a complex number. Then the functions $f + \lambda g$ and fg have the limits $L + \lambda M$ and LM , respectively, at the point z_0 ; and, if $M \neq 0$, the function f/g has the limit L/M at the point z_0 . \blacksquare

Continuity

Suppose again that f is a function defined on a subset S of the complex plane. If $z_0 \in S$, then f is **continuous** at z_0 if

$$\lim_{z \rightarrow z_0} f(z) = f(z_0).$$

That is, f is continuous at z_0 if the values of $f(z)$ get arbitrarily close to the value $f(z_0)$, so long as z is in S and z is sufficiently close to z_0 . If it happens that f is continuous at all points of S , we say f is **continuous on S** . The function f is continuous at ∞ if $f(\infty)$ is defined and $\lim_{z \rightarrow \infty} f(z) = f(\infty)$.

Let us review Examples 13 to 17 in the context of “continuity.”

Example 21 $f(z) = |z|^2$ is continuous at every point of the complex plane. \square

Example 22 $g(z) = 1/(1 - z)$ is continuous at all points of the plane except $z = 1$. \square

Example 23 $h(z) = \operatorname{Re}(z^4 + 4)$ is continuous at all points of the plane. \square

Example 24 $f(z) = (z^4 - 1)/(z - i)$ is continuous on the whole plane if we define $f(i) = -4i$. \square

Example 25 $h(z) = z/\bar{z}$ is continuous everywhere except $z = 0$. Further, there is no way to define $h(0)$ to make h continuous at $z_0 = 0$, since $h(z) = 1$ for all z that

are real and $h(z) = -1$ for z that are purely imaginary. Such a function cannot be continuous at $z_0 = 0$, the point where the real axis meets the imaginary axis. \square

Example 26 The function $g(z) = 1/(|z| + 1)$ is continuous at all points of the plane and at ∞ as well, if we set $g(\infty) = 0$.

You will no doubt recall that the sum and product of continuous functions is continuous. This follows, in fact, directly from Theorem 2. We state this formally in the next theorem.

THEOREM 3 Suppose that f and g are functions, both of which are continuous at the point z_0 . Let λ be a complex number. Then the functions $f + \lambda g$ and fg are also continuous at the point z_0 . Further, if $g(z_0) \neq 0$, the function f/g is continuous at z_0 . Finally, if h is a function continuous at each point of some disc centered at the point $w_0 = f(z_0)$, then the composition $h(f(z))$ is continuous at z_0 . ■

The proofs are left to the exercises at the end of this section. These facts allow us to conclude that each **polynomial**

$$p(z) = a_0 + a_1 z + \cdots + a_n z^n$$

is continuous on the complex plane; here a_0, \dots, a_n are complex numbers. Further, if p and q are two polynomials, their quotient $r = p/q$ is also continuous at all points at which $q(z) \neq 0$. The ratio of two polynomials is called a **rational function**.

Each complex-valued function f can be written as $f = u + iv$, where u and v are real-valued functions: $u(z) = \operatorname{Re} f(z)$, $v(z) = \operatorname{Im} f(z)$. In this way, the statements about limits, continuity, etc., of f can be recast as statements about u and v . For example, f is continuous at a point z_0 if and only if both u and v are continuous at z_0 . Some exercises on these topics are included at the end of this section.

Infinite Series

Just as the notion of the convergence of a sequence of complex numbers is almost the same as the convergence of a sequence of real numbers, so the notion of the sum of an **infinite series of complex numbers** is virtually the same as that of the sum of an infinite series of real numbers. Specifically, if the numbers z_1, z_2, \dots , are complex numbers, we define their **n th partial sum** by

$$s_n = \sum_{j=1}^n z_j = z_1 + \cdots + z_n, \quad n = 1, 2, \dots$$

We then examine the behavior of the sequence $\{s_n\}$. If the sequence $\{s_n\}$ has a limit s , then we say that the infinite series $\sum_{j=1}^{\infty} z_j$ **converges** and has sum s ; this is written

$$\sum_{j=1}^{\infty} z_j = s.$$

If for any reason the sequence $\{s_n\}$ does not have a limit, we say that the series $\sum_{j=1}^{\infty} z_j$ **diverges**. The convergence (or divergence) of the infinite series $\sum_{j=1}^{\infty} z_j$ of complex numbers can be formulated in terms of the convergence (or divergence) of two infinite series of real numbers. This follows directly by writing $z_j = x_j + iy_j$, so

$$s_n = \sum_{j=1}^n z_j = \sum_{j=1}^n x_j + i \sum_{j=1}^n y_j = \sigma_n + i\tau_n.$$

As noted earlier in this section, the sequence $\{s_n\}$ converges if and only if *both* of the sequences $\{\sigma_n\}$ and $\{\tau_n\}$ of real numbers converge, and, this being the case,

$$s = \lim s_n = \lim \sigma_n + i \lim \tau_n.$$

However, the convergence of the two sequences $\{\sigma_n\}$ and $\{\tau_n\}$ is exactly the statement that the two infinite series $\sum_{j=1}^{\infty} x_j$ and $\sum_{j=1}^{\infty} y_j$ both converge. We thus arrive at this result: Let $z_j = x_j + iy_j$; $j = 1, 2, 3, \dots$. The infinite series $\sum_{j=1}^{\infty} z_j$ converges, $z_j = x_j + iy_j$, if and only if both

$$\sum_{j=1}^{\infty} x_j \quad \text{and} \quad \sum_{j=1}^{\infty} y_j$$

converge. Furthermore, if $\sum z_j$ converges, then

$$\sum_{j=1}^{\infty} z_j = \sum_{j=1}^{\infty} x_j + i \sum_{j=1}^{\infty} y_j.$$

There is more that can be said here. Since

$$|x_j| \leq |z_j|, \quad |y_j| \leq |z_j|,$$

the convergence of the series $\sum_{j=1}^{\infty} |z_j|$ of nonnegative numbers implies, by the use of the usual comparison test for infinite series of real numbers, the convergence of both the series

$$\sum_{j=1}^{\infty} |x_j| \quad \text{and} \quad \sum_{j=1}^{\infty} |y_j|.$$

Hence, both the series

$$\sum_{j=1}^{\infty} x_j \quad \text{and} \quad \sum_{j=1}^{\infty} y_j$$

converge, so

$$\sum_{j=1}^{\infty} (x_j + iy_j)$$

also converges. Furthermore,

$$|s_n| = \left| \sum_{j=1}^n z_j \right| \leq \sum_{j=1}^n |z_j|$$

for all n . We conclude that if $\sum_{j=1}^{\infty} |z_j|$ converges, then $\sum_{j=1}^{\infty} z_j$ converges, and

$$\left| \sum_{j=1}^{\infty} z_j \right| \leq \sum_{j=1}^{\infty} |z_j|.$$

This is a useful criterion for convergence, since there are several tests available from calculus (comparison, ratio, root, integral; see the exercises) for the convergence of an infinite series of nonnegative numbers. When $\sum |z_j|$ converges, we say that the series z_j is *absolutely convergent*.

Example 27 If α is not equal to 1, the identity

$$1 + \alpha + \cdots + \alpha^n = \frac{1 - \alpha^{n+1}}{1 - \alpha}$$

is easily verified by multiplying both sides of the equation by $1 - \alpha$. If $|\alpha| < 1$, then $\alpha^{n+1} \rightarrow 0$ as $n \rightarrow \infty$, with the result that

$$1 + \alpha + \alpha^2 + \cdots = \sum_{j=0}^{\infty} \alpha^j = \frac{1}{1 - \alpha}, \quad |\alpha| < 1.$$

This is the **geometric series**, with ratio α . □

Example 28 The series

$$\sum_{j=1}^{\infty} j \left(\frac{1+2i}{3} \right)^j$$

converges, since

$$\left| j \left(\frac{1+2i}{3} \right)^j \right| = j \left(\frac{\sqrt{5}}{3} \right)^j,$$

and the series

$$\sum_{j=1}^{\infty} j \left(\frac{\sqrt{5}}{3} \right)^j$$

converges, as can be seen by use of, say, the ratio test. □

Example 29 The series

$$\sum_{k=1}^{\infty} \frac{1}{k} (i)^k$$

converges, even though the series

$$\sum_{k=1}^{\infty} \frac{1}{k}$$

diverges. Note that

$$i^k = \begin{cases} i & \text{if } k = 1, 5, 9, \dots \\ -1 & \text{if } k = 2, 6, 10, \dots \\ -i & \text{if } k = 3, 7, 11, \dots \\ 1 & \text{if } k = 4, 8, 12, \dots \end{cases}$$

Hence,

$$\sum_{k=1}^{\infty} \frac{i^k}{k} = \left(-\frac{1}{2} + \frac{1}{4} - \frac{1}{6} + \cdots \right) + i \left(1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \cdots \right),$$

and each series in the parentheses converges by the alternating-series test. \square

Example 30 The series

$$\sum_{k=1}^{\infty} \frac{1}{2^k} (1+i)^{2k}$$

diverges, since

$$\frac{1}{2^k} (1+i)^{2k} = \left\{ \frac{(1+i)^2}{2} \right\}^k = i^k,$$

so the terms of the series do not go to zero. Thus, the series must diverge (see Exercise 30). \square

Some Sums*

The identity at the beginning of Example 27 has significance beyond the geometric series. Let x be any real number not equal to 1. Then

$$1 + x + x^2 + x^3 + \cdots + x^n = \frac{1 - x^{n+1}}{1 - x}.$$

This identity can be manipulated to obtain other useful identities. For instance, differentiate both sides with respect to x , then multiply both sides by x , and add 1. The result is the identity

$$1 + x + 2x^2 + 3x^3 + 4x^4 + \cdots + nx^n = 1 - (n+1) \frac{x^{n+1}}{1-x} + x \frac{1 - x^{n+1}}{(1-x)^2}.$$

Repeating this three-step process yields another identity, this time for $1 + x + 4x^2 + 9x^3 + \dots + n^2x^n$.

EXERCISES FOR SECTION 1.4

Limits

In Exercises 1 to 8, find the limit of each sequence that converges; if the sequence diverges, explain why.

1. $z_n = \left(\frac{1+i}{\sqrt{3}}\right)^n$

2. $z_n = \left(\frac{1+i}{\sqrt{2}}\right)^n$

3. $z_n = n\left(\frac{i}{2}\right)^n$

4. $z_n = \log\left(1 + \frac{1}{n}\right)$

5. $z_n = n + \frac{i}{n}$

6. $z_n = \frac{\cos n\theta + i \sin n\theta}{n}, \quad \theta \text{ fixed}$

7. $z_n = \arg\left(1 + \frac{\alpha}{n}\right), \quad \alpha \text{ fixed}$

8. $z_n = n\left\{1 - \cos\left(\frac{\theta}{n}\right) - i \sin\left(\frac{\theta}{n}\right)\right\}, \quad \theta \text{ fixed}$

In Exercises 9 to 14, find the limit of each function at the given point, or explain why it does not exist.

9. $f(z) = |1 - z|^2$ at $z_0 = i$

10. $f(z) = \arg z$ at $z_0 = -1$

11. $f(z) = (1 - \operatorname{Im} z)^{-1}$ at $z_0 = 8$ and then at $z_0 = 8 + i$

12. $f(z) = (z - 2) \log|z - 2|$ at $z_0 = 2$

13. $f(z) = \frac{|z|^2}{z}, \quad z \neq 0, \quad \text{at } z_0 = 0$

14. $f(z) = \frac{z^3 - 8i}{z + 2i}, \quad z \neq -2i, \quad \text{at } z_0 = -2i$

Continuity

In Exercises 15 to 20, find all points of continuity of the given function.

15. $f(z) = \begin{cases} \frac{z^3 + i}{z - i}, & z \neq i \\ -3, & z = i \end{cases}$

16. $f(z) = \begin{cases} \frac{z^4 - 1}{z - i}, & z \neq i \\ 4i, & z = i \end{cases}$

17. $f(z) = (\operatorname{Im} z - \operatorname{Re} z)^{-1}$

18. $g(z) = (1 - |z|^2)^{-3}$

19. $h(z) = \begin{cases} z & \text{if } |z| \leq 1 \\ |z|^2 & \text{if } |z| > 1 \end{cases}$

20. $h(z) = \bar{z}^3$

In Exercises 21 to 24, find the limit at ∞ of the given function, or explain why it does not exist.

21. $f(z) = \frac{1}{|z| - 1}$

22. $h(z) = \frac{|z|}{z}, \quad z \neq 0$

23. $g(z) = \frac{4z^6 - 7z^3}{(z^2 - 4)^3}$

24. $h(z) = \operatorname{Arg} z, \quad z \neq 0$

25. Let f and g be continuous at z_0 . Show that $f + g$ and fg are also continuous at z_0 . If $g(z_0) \neq 0$, show that $1/g$ is continuous at z_0 .
26. Establish the following result. A function f is continuous at a point z_0 of its domain of definition S if and only if, given $\varepsilon > 0$, there is a $\delta > 0$ such that $|f(z) - f(z_0)| < \varepsilon$ for all $z \in S$ with $|z - z_0| < \delta$.
27. Let g be defined on a set containing the range of a function f . If f is continuous at z_0 and g is continuous at $f(z_0)$, then $g(f(z))$ is continuous at z_0 .
28. Suppose that $f = u + iv$ is continuous at z_0 . Show that each of the functions u , v , $u - iv$, and $(u^2 + v^2)^{1/2}$ are also continuous at z_0 . Conversely, show that if u and v are continuous at z_0 , then so is f .
29. Let u be a continuous function on a domain D . Suppose that u has this property: For each point $p \in D$, there is a disc Δ centered at p on which u is constant. Conclude that u is constant throughout D . You will have to use the fact that D is a domain.

Infinite Series

30. Suppose that $\sum_{n=1}^{\infty} a_n$ converges. Show that $\lim_{n \rightarrow \infty} a_n = 0$.

In Exercises 31 to 39, determine whether the given infinite series converges or diverges.

31. $\sum_{n=1}^{\infty} \left(\frac{1+2i}{\sqrt{6}} \right)^n$

32. $\sum_{n=1}^{\infty} n \left(\frac{1}{2i} \right)^n$

33. $\sum_{n=1}^{\infty} \left(\frac{2+i}{\sqrt{5}} \right)^n$

34. $\sum_{n=1}^{\infty} \frac{1}{2+i^n}$

35. $\sum_{n=2}^{\infty} n(n-1)\beta^{n-2}, \quad |\beta| < 1$

36. $\sum_{n=1}^{\infty} \frac{1}{n^2 + i^n}$

37. $\sum_{n=1}^{\infty} \frac{1}{n} \left(\frac{1+i}{\sqrt{2}} \right)^n$

38. $\sum_{n=0}^{\infty} (z^n - z^{n+1})$

39. $\sum_{k=1}^{\infty} \frac{k^2 + i}{(k+i)^4}$

40. Show that each of the following series converges for all z .

(a) $\sum_{n=0}^{\infty} \frac{z^n}{n!}$

(b) $\sum_{n=0}^{\infty} (-1)^n \frac{z^{2n}}{(2n)!}$

(c) $\sum_{n=1}^{\infty} \frac{z^{2n+1}}{(2n+1)!}$

41. Suppose that the series $\sum_{n=1}^{\infty} a_n$ converges. Let $|z| < 1$. Show that the series $\sum_{n=1}^{\infty} a_n z^n$ is convergent—indeed, that $\sum_{n=1}^{\infty} |a_n z^n|$ converges. (Hint: By Exercise 30, $|a_n| \leq M$ for some M and all n .)

The Root and Ratio Tests*

In Exercises 42 and 43, $\{c_n\}$ is a sequence of positive numbers.

42. (Root test) Suppose that $\lim_{n \rightarrow \infty} (c_n)^{1/n} = A$ exists. Show that the series $\sum c_n$ converges if $0 \leq A < 1$ and diverges if $A > 1$. (Hint: If $A < 1$, there is a B with $A < B < 1$. Hence, $c_n < B^n$ if $n \geq N$. (Why?) If $1 < A$, then $c_n \geq 1$ for all $n \geq N'$. (Why?))

43. (Ratio test) Suppose that

$$\lim_{n \rightarrow \infty} \frac{c_{n+1}}{c_n} = C$$

exists. Show that the series $\sum c_n$ converges if $0 \leq C < 1$ and diverges if $C > 1$. (Hint: If $C < 1$, then for D , $C < D < 1$, we have $c_{n+1} \leq Dc_n$ for all $n \geq N$. (Why?) Therefore, $c_{n+k} \leq D^k c_n$, $k = 0, 1, 2, \dots$, and $\sum c_j$ converges. If $C > 1$, then $c_{n+1} \geq c_n$ for all $n \geq N'$. (Why?) So $\{c_k\}$ does not converge to zero.)

1.5 The Exponential, Logarithm, and Trigonometric Functions

The Exponential Function

The exponential function is one of the most important functions in complex analysis. Its definition is simple:

$$e^z = e^x(\cos y + i \sin y), \quad z = x + iy.$$

The form $\exp(z)$ is used sometimes, especially if z itself has some complicated form. The definition of e^z allows us immediately to derive one of its most significant properties: For any two complex numbers z and w ,

$$e^{z+w} = e^z e^w.$$

To see this, write (as usual) $z = x + iy$ and $w = s + it$. Then, making use of two basic trigonometric identities for the sine and cosine of the sum of two numbers,

$$\begin{aligned} e^{z+w} &= e^{x+s}[\cos(y+t) + i \sin(y+t)] \\ &= e^x e^s[(\cos y \cos t - \sin y \sin t) + i(\sin y \cos t + \sin t \cos y)] \\ &= e^x(\cos y + i \sin y)e^s(\cos t + i \sin t) \\ &= e^z e^w. \end{aligned}$$

The basic definition allows several more conclusions about the exponential function. First, e^z is a continuous function of z . The functions e^x , $\cos y$, and $\sin y$ are continuous functions of x and y . Consequently, $\operatorname{Re}(e^z) = e^x \cos y$ and $\operatorname{Im}(e^z) = e^x \sin y$ are both continuous, so e^z is continuous at all points of the plane. Second,

$$|e^z| = e^{\operatorname{Re} z},$$

because

$$\begin{aligned} |e^z| &= ((e^x \cos y)^2 + (e^x \sin y)^2)^{1/2} \\ &= e^x = e^{\operatorname{Re} z}. \end{aligned}$$

In particular,

$$|e^{it}| = 1, \quad t \text{ real}.$$

Since $e^{it} = \cos t + i \sin t$, t real, we see that as t increases, e^{it} moves on the circle of radius 1 centered at the origin in a counterclockwise direction, making one complete circuit when t increases by 2π (Fig. 1.21). In particular, of course,

$$e^{2\pi im} = 1 \quad \text{for } m = 0, \pm 1, \dots$$

and

$$e^{\pi i} = -1.$$




Figure 1.21

Second, the function $f(z) = e^z$ never has the value zero, since neither e^x nor e^{iy} is ever zero. On the other hand, if w is any nonzero complex number, the equation

$$e^z = w$$

has infinitely many solutions. This can be seen, for example, by writing w in polar form:

$$w = r(\cos \psi + i \sin \psi),$$

and then setting $x = \ln r$ and $y = \psi + 2\pi m$, where m is any integer (positive or negative) and $\ln r$ is the natural logarithm of r ; \ln is log to the base e , studied in calculus. Thus,

$$\begin{aligned} e^{x+iy} &= e^{\ln r} e^{i(\psi + 2\pi m)} \\ &= r[\cos \psi + i \sin \psi] \\ &= w. \end{aligned}$$

Furthermore, every solution z of the equation $e^z = w$ has the form given above. For

if $e^z = w$, then

$$r = |w| = |e^z| = e^x,$$

so $x = \ln r$. Consequently,

$$\cos y + i \sin y = e^{iy} = \cos \psi + i \sin \psi.$$

Thus, $y = \psi + 2\pi m$ for some integer m (see Exercise 14 in Section 1).

The mapping of $f(z) = e^z$ thus carries the complex z -plane onto the complex w -plane with the origin deleted; each point w_0 has infinitely many pre-images z , each of the form $z_0 + 2\pi im$, $m = 0, \pm 1, \pm 2, \dots$, where z_0 is any solution of $e^{z_0} = w_0$. In particular, $f(z) = e^z$ carries each strip $y_0 \leq \operatorname{Im} z < y_0 + 2\pi$, $-\infty < \operatorname{Re} z < \infty$, onto the w -plane with the origin removed. The function $f(z) = e^z$ is **one-to-one** on that strip—that is, distinct points have distinct images. For $e^{z_1} = e^{z_2}$ if and only if $e^{z_1 - z_2} = 1$; this occurs exactly when $z_1 - z_2 = 2\pi ik$ for some integer k (Fig. 1.22).




Figure 1.22

The function $f(z) = e^z$ maps each horizontal line $y = c$ onto a ray from the origin to infinity—specifically, the ray $\{r \cos c + ir \sin c : 0 < r < \infty\}$, since $\exp(x + ic) = e^x \cos c + ie^x \sin c$, and e^x increases from 0 to ∞ as x increases from $-\infty$ to ∞ ; of course, e^x is always positive. Furthermore, e^z maps each vertical line $x = c$ onto the circle centered at the origin of radius e^c . This is because $\exp(c + iy) = e^c(\cos y + i \sin y)$, and $\cos y + i \sin y$ travels along the circle of radius 1 centered at the origin in a counterclockwise direction as y increases. Each point on this circle is the image of infinitely many points on the vertical line, since both $\cos y$ and $\sin y$ are periodic, with period 2π .

The Logarithm Function

The inverse of the exponential function is the logarithm function. For a nonzero complex number z , we *define* $\log z$ to be any complex number w with $e^w = z$.

The preceding discussion of the exponential function leads immediately to the relationship

$$\log z = \ln|z| + i \arg z, \quad z \neq 0.$$

However, it is obvious that this is not a single complex number, but rather a set of complex numbers, each two elements of which differ by an integer multiple of $2\pi i$. One way to be definite is to use

$$\text{Log } z = \ln|z| + i \text{Arg } z.$$

Be wary here; it may happen that $\text{Log}(z_1 z_2) \neq \text{Log } z_1 + \text{Log } z_2$ exactly because $\text{Arg}(z_1 z_2)$ need not equal $\text{Arg } z_1 + \text{Arg } z_2$. The function $\text{Log } z$ is called the **principal branch** of the logarithm of z . Other choices of $\arg z$ yield other values of $\log z$ (more on this later). To investigate the continuity of $\log z$, delete from the plane any ray beginning at the origin (for example, delete all the nonpositive real numbers) and let D be the domain remaining. Let z_0 be any point of D and choose and then fix any value for $\arg z_0$, say $\arg z_0 = \theta_0$. We then define in D a **branch** of $\log z$ by the rule $\log z = \ln|z| + i \arg z$, with the additional specification that $\log z_0 = \ln|z_0| + i\theta_0$. Then within D the values of $\arg z$ lie in a uniquely determined open interval of length 2π , which contains θ_0 ; furthermore, $\arg z$ is continuous in D with this specification. The function $\ln|z|$ is a continuous function on D as well, so the branch of $\log z$ determined by the specific choice $\arg z_0 = \theta_0$ is continuous on D . For example, if we delete the nonpositive real axis (see Fig. 1.23) and pick $z_0 = 1 + i$ and $\theta_0 = \frac{\pi}{4}$, then the branch of $\log z$ that is determined has $\arg z$ in the range $(\pi, 3\pi)$. Clearly, there are infinitely many possible branches of $\log z$ in D , each continuous.




Figure 1.23

The definition of $\log z$ allows us to complete the discussion of roots begun in Section 2. For a nonzero complex number a , we define a^z by the rule

$$a^z = e^{z \log a}.$$

This results in many values, but it does agree with the usual definition in the special case when a and z are both positive and real.

Example 1 Find the values of $(-1)^i$.

Solution Using the relation $\log(-1) = (2n + 1)\pi i$, $n = 0, \pm 1, \dots$, we obtain

$$(-1)^i = e^{i \log(-1)} = e^{-(2n+1)\pi}, \quad n = 0, \pm 1, \pm 2, \dots \quad \square$$

Example 2 Find the solutions of $z^{1+i} = 4$.

Solution Write this equation as

$$e^{(1+i)\log z} = 4,$$

so $(1+i)\log z = \ln 4 + 2\pi ni$, $n = 0, \pm 1, \dots$. Hence,

$$\begin{aligned} \log z &= (1-i)[\ln 2 + \pi ni] \\ &= (\ln 2 + \pi n) + i(\pi n - \ln 2). \end{aligned}$$

Thus,

$$\begin{aligned} z &= 2e^{\pi n}(\cos(\pi n - \ln 2) + i \sin(\pi n - \ln 2)) \\ &= 2e^{\pi n}\{(-1)^n \cos(\ln 2) + i(-1)^{n+1} \sin(\ln 2)\} \\ &= (-1)^n 2e^{\pi n}\{\cos(\ln 2) - i \sin(\ln 2)\}, \quad n = 0, \pm 1, \dots \quad \square \end{aligned}$$

Example 3 Establish the formula

$$\lim_{n \rightarrow \infty} \left(1 + \frac{z}{n}\right)^n = e^z, \quad z \text{ a complex number}$$

Solution Look at $n \operatorname{Log}(1 + z/n)$ for large n . Write

$$n \operatorname{Log}\left(1 + \frac{z}{n}\right) = n \ln \left|1 + \frac{z}{n}\right| + in \operatorname{Arg}\left(1 + \frac{z}{n}\right).$$

The real part satisfies

$$n \ln \left|1 + \frac{z}{n}\right| = \frac{1}{2}n \ln \left[1 + \frac{2x}{n} + \frac{x^2 + y^2}{n^2}\right] \rightarrow x,$$

as $n \rightarrow \infty$ by l'Hôpital's Rule, for example. Next, if $z = r(\cos \theta + i \sin \theta)$ and $\psi_n = \operatorname{Arg}(1 + z/n)$, then from Figure 1.24,




Figure 1.24

$$\tan \psi_n = \frac{\frac{r}{n} \sin \theta}{1 + \frac{r}{n} \cos \theta},$$

and so

$$n \tan \psi_n = \frac{r \sin \theta}{1 + \frac{r}{n} \cos \theta}.$$

Because $\psi_n \rightarrow 0$ as $n \rightarrow \infty$, we know that $\psi_n / (\tan \psi_n) \rightarrow 1$ (say, l'Hôpital's Rule again). Hence,

$$n \psi_n = (n \tan \psi_n) \left(\frac{\psi_n}{\tan \psi_n} \right) \rightarrow r \sin \theta = y, \quad \text{as } n \rightarrow \infty.$$

Consequently,

$$n \operatorname{Log} \left(1 + \frac{z}{n} \right) \rightarrow x + iy = z, \quad \text{as } n \rightarrow \infty,$$

so $(1 + (z/n))^n \rightarrow e^z$, as intended. □

Trigonometric Functions

The trigonometric functions of z are defined in terms of the exponential function. We begin with the cosine and the sine of z :

$$\cos z = \frac{1}{2}(e^{iz} + e^{-iz})$$

$$\sin z = \frac{1}{2i}(e^{iz} - e^{-iz}).$$

The other four trigonometric functions are now defined in terms of $\sin z$ and $\cos z$:

$$\tan z = \frac{\sin z}{\cos z}, \quad \cot z = \frac{\cos z}{\sin z}, \quad \sec z = \frac{1}{\cos z}, \quad \csc z = \frac{1}{\sin z}.$$

Of course, the definitions are only at those points at which the denominator is not zero.

If z is real, $z = x$, then $\cos z$ and $\sin z$ agree with the usual definitions of cosine and sine. Note that

$$\begin{aligned}\cos(z + 2\pi) &= \frac{1}{2}(e^{i(z+2\pi)} + e^{-i(z+2\pi)}) \\ &= \frac{1}{2}(e^{iz} + e^{-iz}) \\ &= \cos z.\end{aligned}$$

Likewise, $\sin(z + 2\pi) = \sin z$ for all z . Thus,

$$\cos(z + 2\pi k) = \cos z$$

$$\sin(z + 2\pi k) = \sin z,$$

for all z and any integer k . Furthermore, $2\pi k$ is the only number α with $\cos(z + \alpha) = \cos z$ for all z . For if

$$\cos(z + \alpha) = \cos z, \quad \text{for all } z,$$

then

$$e^{iz}e^{i\alpha} + e^{-iz}e^{-i\alpha} = e^{iz} + e^{-iz},$$

so

$$\begin{aligned}e^{iz}(e^{i\alpha} - 1) &= e^{-iz}(1 - e^{-i\alpha}) \\ &= e^{-i\alpha}e^{-iz}(e^{i\alpha} - 1).\end{aligned}$$

If $e^{i\alpha} - 1$ is not zero, it can be canceled from both sides, leaving

$$e^{iz} = e^{-i\alpha}e^{-iz}, \quad \text{for all } z,$$

but then setting $z = 0$ gives $1 = e^{-i\alpha}$, contradicting the fact that $1 \neq e^{i\alpha}$. Hence, it must be the case that $e^{i\alpha} = 1$, so $\alpha = 2\pi k$ for some integer k . Consequently, 2π is the basic period of $\cos z$ and, likewise, of $\sin z$.

In the exercises at the end of this section, you will be asked to establish the formulas

$$\cos(x + iy) = \cos x \cosh y - i \sin x \sinh y,$$

$$\sin(x + iy) = \sin x \cosh y + i \cos x \sinh y,$$

where

$$\cosh u = \frac{1}{2}(e^u + e^{-u}), \quad u \text{ real}$$

$$\sinh u = \frac{1}{2}(e^u - e^{-u}), \quad u \text{ real.}$$

Let us use the formula given above for $\sin(x + iy)$ to establish some basic properties of the function $f(z) = \sin z$.

Restrict $z = x + iy$ such that $0 \leq x < \pi/2$ and $y \geq 0$. In this restricted region, we shall show that the function $\sin z$ is one-to-one—that is, if

$$\sin(x_1 + iy_1) = \sin(x_2 + iy_2),$$

where x_1 and x_2 lie in $[0, \pi/2)$ and y_1 and y_2 are nonnegative, then $x_1 = x_2$ and $y_1 = y_2$. To see this, observe that

$$2i \sin(x + iy) = e^{ix}e^{-y} - e^{-ix}e^y.$$

Hence, if $\sin(x_1 + iy_1) = \sin(x_2 + iy_2)$, then

$$e^{ix_1}e^{-y_1} - e^{-ix_1}e^{y_1} = e^{ix_2}e^{-y_2} - e^{-ix_2}e^{y_2}.$$

Thus,

$$\begin{aligned} e^{ix_1}e^{-y_1} - e^{ix_2}e^{-y_2} &= e^{-ix_1}e^{y_1} - e^{-ix_2}e^{y_2} \\ &= e^{-ix_1}e^{-ix_2}e^{y_1}e^{y_2}[e^{ix_2}e^{-y_2} - e^{ix_1}e^{-y_1}]. \end{aligned}$$

If $e^{ix_1}e^{-y_1} - e^{ix_2}e^{-y_2} \neq 0$, we divide by it and obtain

$$1 = -e^{-ix_1}e^{-ix_2}e^{y_1}e^{y_2}.$$

The absolute value of the left side is 1, and that of the right side is $\exp(y_1 + y_2)$; this implies that $y_1 + y_2 = 0$. This in turn implies that $-1 = \exp[-ix_1 - ix_2]$, which can be true only when $x_1 + x_2 = \pi(2m + 1)$ for some integer m . However, we have restricted x_1, x_2 to lie in $[0, (\pi/2))$ and y_1, y_2 to lie in $[0, \infty)$. The only conclusion, then, is that $x_1 = x_2 = 0$ and $y_1 = y_2 = 0$. Otherwise,

$$e^{ix_1}e^{-y_1} = e^{ix_2}e^{-y_2},$$

so $y_1 = y_2$ and $x_1 - x_2$ is an integer multiple of 2π . This again implies that $x_1 = x_2$.

Let us now find the range of $f(z) = \sin z$ on the strip $0 \leq x \leq \pi/2$ and $0 \leq y < \infty$. First,

$$\operatorname{Re}(\sin z) = \sin x \cosh y \geq 0,$$

and

$$\operatorname{Im}(\sin z) = \cos x \sinh y \geq 0.$$

Hence, the values of $\sin z$ lie in the first quadrant (Fig. 1.25). Next, on the ray $0 \leq y < \infty$, $x = 0$,

$$\sin(iy) = i \sinh y = \frac{i}{2}(e^y - e^{-y}).$$




Figure 1.25

The function $\sinh(y)$ is zero when $y = 0$ and is increasing (its derivative is positive), so $\sin(iy)$ assumes all the values $i\beta$, $0 \leq \beta < \infty$, as y increases from 0 to ∞ . Next, $\sin x$ increases from 0 to 1 as x increases from 0 to $\pi/2$. Finally,

$$\sin\left(\frac{\pi}{2} + iy\right) = \cosh y,$$

and this increases from 1 to ∞ as y increases from 0 to ∞ . Thus, the boundary of the strip $0 \leq x \leq \pi/2$, $0 \leq y < \infty$ is carried by $\sin z$ onto the boundary of the first quadrant (see Fig. 1.25). We now show that the interior is mapped onto the interior; we already know it is mapped *into*. Let us write $\sin z = w = \sigma + i\tau$. The vertical segment $z = x_0 + iy$, $y \geq 0$, is mapped onto that portion of the hyperbola

$$\frac{\sigma^2}{(\sin x_0)^2} - \frac{\tau^2}{(\cos x_0)^2} = 1,$$

which lies in the first quadrant. This shows us how to solve the equation

$$\sin z = a + ib, \quad a, b \text{ positive.}$$

First choose x such that the point $a + ib$ lies on the hyperbola consisting of those $w = \sigma + it$ with

$$\frac{\sigma^2}{(\sin x)^2} - \frac{t^2}{(\cos x)^2} = 1;$$

then choose y such that $a = \sin x \cosh y$ and $b = \cos x \sinh y$.

Consequently, the function $f(z) = \sin z$ maps the semi-infinite strip $0 \leq x \leq \pi/2$ and $0 \leq y < \infty$ in a one-to-one fashion *onto* the whole first quadrant, mapping the boundary of the strip onto the boundary of the first quadrant. This type of information, and this mapping in particular, will be very important in the solution of flow and boundary-value problems. Finally, because $\sin(-\bar{z}) = -\overline{\sin z}$ (see Exercise 15), we may conclude that $f(z) = \sin z$ maps the semi-infinite strip $\{-\pi/2 < x < \pi/2, y > 0\}$ both one-to-one and onto the upper half-plane, $\{w = s + it: -\infty < s < \infty, t > 0\}$.

Inverse Trigonometric Functions

We have just seen that the function $w = \sin z$ maps the strip $\{x + iy: 0 < x < \pi/2, 0 < y < \infty\}$ onto the first quadrant $\{w = \sigma + it: \sigma, t > 0\}$, and distinct z_1 and z_2 in the strip have distinct images w_1 and w_2 in the first quadrant. Thus, given a $w = \sigma + it$ with $\sigma, t > 0$, there is one and only one z in the strip $\{x + iy: 0 < x < \pi/2, 0 < y < \infty\}$ with $w = \sin z$. Consequently, the function $\sin z$ has an inverse function, naturally called $\arcsin w$, defined at least for w in the first quadrant. Let us pursue this a little further. The formula for $\sin z$ is

$$w = \sin z = \frac{1}{2i}(e^{iz} - e^{-iz}),$$

so

$$e^{2iz} - 2iwe^{iz} - 1 = 0,$$

which is a quadratic equation in the variable e^{iz} . Solving by the quadratic formula, we obtain

$$e^{iz} = iw + \sqrt{1 - w^2},$$

so

$$z = -i \log(iw + \sqrt{1 - w^2}),$$

provided an appropriate branch of the logarithm is chosen. A careful examination of the mapping $w = \sin z$ (see the exercises) shows that $\sin z$ maps the strip

$$\left\{ x + iy : -\frac{\pi}{2} < x < \frac{\pi}{2}, -\infty < y < \infty \right\}$$

both one-to-one and onto the region D obtained from the plane by deleting the two intervals $(-\infty, -1]$ and $[1, \infty)$ (see Fig. 1.26). Thus, on D we can solve uniquely for z in terms of w , and the formula is valid. Interchanging the roles of z and w , write

$$\text{Arcsin } z = -i \text{ Log}(iz + \sqrt{1 - z^2}), \quad z \in D.$$

Similarly,

$$\text{Arccos } z = -i \text{ Log}(z + \sqrt{z^2 - 1}),$$

and

$$\text{Arctan } z = \frac{i}{2} \text{ Log}\left(\frac{1 - iz}{1 + iz}\right), \quad z \neq \pm i,$$

with appropriate interpretations of the resulting logarithms and roots.




Figure 1.26

EXERCISES FOR SECTION 1.5

Find the values(s) of the given expression in Exercises 1 to 14.

- | | | |
|-----------------------|---|--------------------------|
| 1. $e^{i\pi/4}$ | 2. $e^{5\pi i/4}$ | 3. $\log(1 + i\sqrt{3})$ |
| 4. $\log(-i)$ | 5. $(1+i)^i$ | 6. 2^{-1-i} |
| 7. $e^{-7\pi i/3}$ | 8. $\exp(\text{Log}(3 + 2i))$ | 9. $\text{Log}(4 - 4i)$ |
| 10. $\text{Log}(-1)$ | 11. $i\sqrt{3}$ | 12. $\log(\sqrt{3} - i)$ |
| 13. $\log((1 - i)^4)$ | 14. $\exp\left[\pi\left(\frac{i+1}{\sqrt{2}}\right)^4\right]$ | |

15. Establish the following relations:

- (a) $\exp(\bar{z}) = \overline{\exp(z)}$
- (b) $\sin(\bar{z}) = \overline{\sin(z)}$
- (c) $\cos(\bar{z}) = \cos(z)$

16. Establish the formulas

$$\cos(x + iy) = \cos x \cosh y - i \sin x \sinh y,$$

$$\sin(x + iy) = \sin x \cosh y + i \cos x \sinh y,$$

where

$$\cosh u = \frac{1}{2}(e^u + e^{-u}), \quad u \text{ real}$$

$$\sinh u = \frac{1}{2}(e^u - e^{-u}), \quad u \text{ real.}$$

17. Show that $\cos z = 0$ if and only if $z = \pi/2 + n\pi$, $n = 0, \pm 1, \pm 2, \dots$; show that $\sin z = 0$ if and only if $z = n\pi$, $n = 0, \pm 1, \pm 2, \dots$. That is, extending $\sin z$ and $\cos z$ from the real axis to the whole plane does not introduce any new zeros.

18. Verify that

$$\cos(z + w) = \cos z \cos w - \sin z \sin w$$

and

$$\sin(z + w) = \sin z \cos w + \cos z \sin w,$$

for all complex numbers z and w .

19. Show that both $\cos z$ and $\sin z$ are unbounded if $z = iy$ and $y \rightarrow \infty$. Also show that

$$|\cos(x + iy)| \leq e^y \quad \text{if} \quad y \geq 0, -\infty < x < \infty;$$

and

$$|\sin(x + iy)| \leq e^y \quad \text{if} \quad y \geq 0, -\infty < x < \infty.$$

20. Prove that $\cos^2 z + \sin^2 z = 1$ for all z .

21. Define $\cosh z$ and $\sinh z$ by

$$\cosh z = \frac{1}{2}(e^z + e^{-z})$$

$$\sinh z = \frac{1}{2}(e^z - e^{-z}).$$

Show that the following identities hold:

- (i) $\cosh^2(z) - \sinh^2(z) = 1$
- (ii) $\cosh z = \cos(iz)$
- (iii) $\sinh z = -i \sin(iz)$
- (iv) $|\cosh z|^2 = \sinh^2 x + \cos^2 y$
- (v) $|\sinh z|^2 = \sinh^2 x + \sin^2 y$

22. Show that $\sin(-z) = -\sin z$ and $\cos z = \cos(-z)$ for all z .

Mappings with the Exponential, Logarithm, and Trigonometric Functions

23. Show that $F(z) = e^z$ maps the strip $S = \{x + iy : -\infty < x < \infty, -\pi/2 \leq y \leq \pi/2\}$ onto the region $\Omega = \{w = s + it : s \geq 0, w \neq 0\}$ and that F is one-to-one on S (see Fig. 1.27). Furthermore, show that F maps the boundary of S onto all the boundary of Ω except $w = 0$. Explain what happens to each of the horizontal lines $\{\text{Im } z = \pi/2\}$ and $\{\text{Im } z = -\pi/2\}$.




Figure 1.27

24. Let D be the domain obtained by deleting the ray $\{x : x \leq 0\}$ from the plane, and let $G(z)$ be a branch of $\log z$ on D . Show that G maps D onto a horizontal strip of width of 2π ,

$$\{x + iy : -\infty < x < \infty, c_0 < y < c_0 + 2\pi\},$$

and that the mapping is one-to-one on D .

25. Show that $w = \sin z$ maps the strip $-\pi/2 < x < \pi/2$ both one-to-one and onto the region obtained by deleting from the plane the two rays $(-\infty, -1]$ and $[1, \infty)$ (see Fig. 1.26). (Hint: Use Exercise 22 and the fact that $\sin(\bar{z}) = \overline{\sin(z)}$.)
26. Show that the function $w = \cos z$ maps the strip $0 < x < \pi$ one-to-one and onto the region D shown in Figure 1.26. Use this to define the inverse function to $\cos z$. Derive the formula for $\arccos z$ given in the text.
27. Let $0 < \alpha < 2$. Show that an appropriate choice of $\log z$ for $f(z) = z^\alpha = \exp[\alpha \log z]$ maps the domain $\{x + iy : y > 0\}$ both one-to-one and onto the domain $\{w : 0 < \arg w < \alpha\pi\}$. Show that f also carries the boundary to the boundary (see Fig. 1.28).




Figure 1.28

28. Show that the function $w = g(z) = e^{z^2}$ maps the lines $x = y$ and $x = -y$ onto the circle $|w| = 1$. Show further that g maps each of the two pieces of the region $\{x + iy: x^2 > y^2\}$ onto the set $\{w: |w| > 1\}$ and each of the two pieces of the region $\{x + iy: x^2 < y^2\}$ onto the set $\{w: |w| < 1\}$.

Inverse Trigonometric Functions

29. Show directly that if ζ is any value of

$$-i \log(iz + \sqrt{1 - z^2}),$$

then $\sin \zeta = z$. Likewise, show that if ξ is any value of

$$\frac{i}{2} \log\left(\frac{1 - iw}{1 + iw}\right),$$

then $\tan \xi = w$.

30. Use the result in Exercise 29 and your knowledge of the branches of the logarithmic function to explain the branches of $\arcsin z$.

1.6 Line Integrals and Green's Theorem

The fundamental theorems of complex variables depend on line integrals, so this section is devoted to that topic and to formulating Green's Theorem, the basic theorem about line integrals. Several consequences of Green's Theorem are also covered.

Curves

A **curve** γ is a continuous complex-valued function $\gamma(t)$ defined for t in some interval $[a, b]$ in the real axis. The curve γ is **simple** if $\gamma(t_1) \neq \gamma(t_2)$ whenever $a \leq t_1 < t_2 \leq b$, and it is **closed** if $\gamma(a) = \gamma(b)$. The famous **Jordan* Curve Theorem** asserts that the complement of the range of a curve that is both simple and closed consists of two

* Camille Jordan, 1838–1922.

disjoint open connected sets, one bounded and the other unbounded. The bounded piece is the **inside** of the curve and the unbounded piece the **outside**. Despite the almost painful obviousness of this statement, the theorem is hard to prove.* We shall accept it as true.

Suppose γ is a curve; separate the complex number $\gamma(t)$ into its real and imaginary parts and write $\gamma(t) = x(t) + iy(t)$, $a \leq t \leq b$. The functions $x(t)$ and $y(t)$ are real-valued functions of the real variable t , so they may (or may not) be differentiable. If both $x(t)$ and $y(t)$ are differentiable at t_0 , then we say that $\gamma(t)$ is differentiable at t_0 and we set $\gamma'(t_0) = x'(t_0) + iy'(t_0)$. (This is consistent with the usual rules in calculus for differentiating a vector-valued function of a real variable.) A curve γ is **smooth** if $\gamma(t)$ has the added property that $\gamma'(t)$ not only exists but is also continuous on $[a, b]$, the derivatives at a and b being taken from the right and left, respectively. A curve is **piecewise smooth** if it is composed of a finite number of smooth curves, the end of one coinciding with the beginning of the next. That is, the curve γ is piecewise smooth if in the interval $[a, b]$ there are points t_0, t_1, \dots, t_n with $a = t_0 < t_1 < \dots < t_{n-1} < t_n = b$ such that $\gamma'(t)$ is continuous on each closed interval $[t_k, t_{k+1}]$, $k = 0, 1, \dots, n - 1$. It is not required that $\gamma'(t)$ be continuous on all of $[a, b]$.

NOTE: It is very common and convenient to refer to the range of $\gamma(t)$ as the curve γ and to $\gamma(t)$ itself as the **parametrization** of the curve. With this use of the word *curve*, a curve becomes a concrete geometric object such as a circle or a straight line segment and hence is easily visualized. The difficulty with this view is that a particular curve has many different parametrizations. The curves we will use are generally composed of straight line segments and/or arcs of circles. These have standard parametrizations, as illustrated in the examples that follow.

Example 1 Fix z_0 and z_1 in the plane and let $\gamma(t) = tz_1 + (1 - t)z_0$, $0 \leq t \leq 1$. This is a smooth simple curve whose range is the straight line segment joining z_0 to z_1 (in that order). \square

Example 2 Fix a point p in the plane and a positive number R ; the curve $\gamma(t) = p + Re^{it}$, $0 \leq t \leq 2\pi$, is a smooth simple closed curve. The range is precisely the circle of radius R centered at p ; the circle is traversed in the counterclockwise direction. \square

Example 3 The square with vertices at z_0 , iz_0 , $-z_0$, and $-iz_0$ is the range of the piecewise smooth simple closed curve $\gamma(t)$ given by the rule

$$\gamma(t) = \begin{cases} tiz_0 + (1 - t)z_0, & 0 \leq t \leq 1 \\ (t - 1)(-z_0) + (2 - t)(iz_0), & 1 \leq t \leq 2 \\ (t - 2)(-iz_0) + (3 - t)(-z_0), & 2 \leq t \leq 3 \\ (t - 3)(z_0) + (4 - t)(-iz_0), & 3 \leq t \leq 4. \end{cases}$$

* For a proof, see Newman, M. H. A. *Elements of the topology of plane sets of points*. Cambridge: Cambridge University Press, 1961.

In the process of computing certain definite integrals by means of the Residue Theorem (Section 6 of Chapter 2), we shall make use of several contours. As a way of gaining experience in the art of parametrization, we parametrize three of these contours in Examples 4 and 5.

Example 4 Parametrize each of the curves in Figure 1.29.

Solution The curve in (a) is the semicircle of radius R followed by the real segment from $-R$ to R . A parametrization is

$$\gamma: \begin{aligned} z &= Re^{i\theta}, & 0 \leq \theta \leq \pi \\ z &= x, & -R \leq x \leq R. \end{aligned}$$

The curve in (b) is a variation on the curve of (a), with the segment from $-\varepsilon$ to ε replaced by the indicated semicircle. This curve is parametrized by

$$\gamma: \begin{aligned} z &= Re^{i\theta}, & 0 \leq \theta \leq \pi \\ z &= x, & -R \leq x \leq -\varepsilon \\ z &= \varepsilon e^{i\theta}, & \pi \geq \theta \geq 0 \\ z &= x, & \varepsilon \leq x \leq R. \end{aligned}$$

Note that the third portion of the curve is parametrized by requiring that θ decrease from π to 0; this is because we travel on that portion of γ in the clockwise direction. \square




Figure 1.29

Example 5 Find a parametrization of the “keyhole” contour shown in Figure 1.30.

Solution The outer portion is on the circle $|z| = R$ and so is parametrized by $z = Re^{i\theta}$, where θ increases from δ to $2\pi - \delta$. This takes us from A to B. We go from B to C on the ray with argument $2\pi - \delta$, so that segment is parametrized by $z = te^{i(2\pi - \delta)}$, where t decreases from R to ε . The inner portion of the keyhole contour is on the circle $|z| = \varepsilon$ and so is parametrized by $z = \varepsilon e^{i\theta}$, where θ decreases from $2\pi - \delta$ to δ . This takes us from C to D. Finally, to complete the contour, we go from D to A along the ray with argument δ . This segment is parametrized by $z = te^{i\delta}$,




Figure 1.30

where t increases from ε to R . In summary, then,

$$\begin{aligned} z &= \operatorname{Re}^{i\theta}, & \delta \leq \theta \leq 2\pi - \delta \\ z &= te^{i(2\pi-\delta)}, & R \geq t \geq \varepsilon \\ \gamma: & \quad z = \varepsilon e^{i\theta}, & 2\pi - \delta \geq \theta \geq \delta \\ & \quad z = te^{i\delta}, & \varepsilon \leq t \leq R. \end{aligned}$$
□

Each curve γ is **oriented** by increasing t . The curve γ **begins** at $\gamma(a)$, is traversed as t increases from a to b , and γ **ends** at $\gamma(b)$. The **reverse orientation** is given to γ by beginning at $\gamma(b)$ and ending at $\gamma(a)$; this curve is denoted by $-\gamma$ and is given by $-\gamma(t) = \gamma(a + b - t)$, $a \leq t \leq b$. A simple closed curve γ is **positively oriented** if, for each point p on the inside of γ , the argument of $\gamma(t) - p$ increases by 2π as t increases from a to b . Equivalently, γ is positively oriented if, as you walk along γ in the direction of the orientation of γ , the inside of γ is on your left. For example, a circle is positively oriented when it is traversed counterclockwise; the same is true for a triangle or a rectangle.

Suppose $g(t) = \sigma(t) + i\tau(t)$ is a continuous complex-valued function on the interval $[a, b]$. We define the **integral of g over $[a, b]$** by

$$\int_a^b g(t) dt = \int_a^b \sigma(t) dt + i \int_a^b \tau(t) dt.$$

This definition, like the definition of the derivative of a complex-valued function of the real variable t , is consistent with the definition of the integral of a vector-valued function of the real variable t studied in calculus. Note, for instance, that

$$\operatorname{Re} \left\{ \int_a^b g(t) dt \right\} = \int_a^b \{\operatorname{Re} g(t)\} dt,$$

since both these expressions equal $\int_a^b \sigma(t) dt$.

Suppose now that γ is a smooth curve and u is a continuous function on the range of γ . We define the **line integral** of u along γ by

$$\int_{\gamma} u(z) dz = \int_a^b u(\gamma(t))\gamma'(t) dt,$$

where the right side is the integral of the complex-valued function $u(\gamma(t))\gamma'(t)$ from a to b and is computed as discussed above. For a piecewise smooth curve γ , we define the line integral of u along γ by

$$\int_{\gamma} u(z) dz = \sum_{j=0}^{n-1} \int_{t_j}^{t_{j+1}} u(\gamma(s))\gamma'(s) ds.$$

The points t_0, t_1, \dots, t_n come from the definition of “piecewise smooth”; by assumption $\gamma'(s)$ is continuous on each segment $[t_j, t_{j+1}]$, $j = 0, 1, \dots, n - 1$.

Line integrals have the familiar properties of definite integrals studied in calculus. For example,

$$\int_{\gamma} \{Au(z) + Bv(z)\} dz = A \int_{\gamma} u(z) dz + B \int_{\gamma} v(z) dz$$

if A and B are complex numbers and u, v are continuous functions on the range of γ . For the curve $-\gamma$,

$$\begin{aligned} \int_{-\gamma} u(z) dz &= \int_a^b u(\gamma(a + b - t))\{\gamma(a + b - t)\}' dt \\ &= - \int_a^b u(\gamma(a + b - t))\gamma'(a + b - t) dt; \quad \text{put } s = a + b - t \\ &= - \int_a^b u(\gamma(s))\gamma'(s) ds = - \int_{\gamma} u(z) dz. \end{aligned}$$

Suppose γ_1 and γ_2 are two curves with parameter intervals $[a_1, b_1]$ and $[a_2, b_2]$, respectively. If $\gamma_1(b_1) = \gamma_2(a_2)$, then the **sum** of γ_1 and γ_2 is the curve

$$(\gamma_1 + \gamma_2)(t) = \begin{cases} \gamma_1(t), & a_1 \leq t \leq b_1 \\ \gamma_2(t + a_2 - b_1), & b_1 \leq t \leq b_1 + b_2 - a_2. \end{cases}$$

Further, a simple computation shows that

$$\int_{\gamma_1 + \gamma_2} u(z) dz = \int_{\gamma_1} u(z) dz + \int_{\gamma_2} u(z) dz,$$

provided, of course, that u is continuous on the range of both γ_1 and γ_2 .

Let g be a complex-valued continuous function on $[a, b]$; we shall show the

inequality

$$\left| \int_a^b g(t) dt \right| \leq \int_a^b |g(t)| dt.$$

The inequality is obviously true if $\int_a^b g(t) dt = 0$, so we may assume that $\int_a^b g(t) dt \neq 0$. Let

$$\theta = \operatorname{Arg} \left(\int_a^b g(t) dt \right)$$

and define $h(t) = e^{-i\theta} g(t)$, $a \leq t \leq b$. Then

$$\begin{aligned} 0 < \left| \int_a^b g(t) dt \right| &= e^{-i\theta} \int_a^b g(t) dt \\ &= \int_a^b e^{-i\theta} g(t) dt = \int_a^b h(t) dt, \quad h(t) = e^{-i\theta} g(t). \end{aligned}$$

Hence, $\int_a^b h(t) dt$ is positive, so

$$\begin{aligned} \left| \int_a^b g(t) dt \right| &= \operatorname{Re} \int_a^b h(t) dt \\ &= \int_a^b (\operatorname{Re} h(t)) dt \\ &\leq \int_a^b |h(t)| dt = \int_a^b |g(t)| dt. \end{aligned}$$

Therefore,

$$\begin{aligned} \left| \int_{\gamma} u(z) dz \right| &= \left| \int_a^b u(\gamma(t)) \gamma'(t) dt \right| \\ &\leq \int_a^b |u(\gamma(t))| |\gamma'(t)| dt. \end{aligned} \tag{1}$$

Recall now from calculus that if $\gamma(t) = x(t) + iy(t)$, then the length of the curve that is the range of $\gamma(t)$ is given by

$$\begin{aligned} \text{length } (\gamma) &= \int_a^b \sqrt{(x'(t))^2 + (y'(t))^2} dt \\ &= \int_a^b |\gamma'(t)| dt. \end{aligned} \tag{2}$$

This makes possible the very important estimate

$$\left| \int_{\gamma} u(z) dz \right| \leq \left(\max_{z \in \gamma} |u(z)| \right) \text{length}(\gamma). \quad (3)$$

The inequality in (3) follows directly from that in (1) by replacing $|u(\gamma(t))|$ by its maximum value on the interval, $a \leq t \leq b$; this is the first factor on the right in (3). The integral that remains is that in (2), and it yields the length of the curve.

Example 6 Compute $\int_{\gamma} (z^2 - 3|z| + \operatorname{Im} z) dz$, where $\gamma(t) = 2e^{it}$, $0 \leq t \leq \pi/2$. That is, γ is the quarter-circle centered at the origin and extending from 2 to $2i$.

Solution Since $\gamma(t) = 2e^{it}$, we have $\gamma'(t) = 2ie^{it}$, so

$$\begin{aligned} \int_{\gamma} (z^2 - 3|z| + \operatorname{Im} z) dz &= \int_0^{\pi/2} (4e^{2it} - 6 + 2 \sin t) 2ie^{it} dt \\ &= \left(\frac{8}{3}e^{3it} - 12e^{it} + \frac{1}{i}e^{2it} - 2t \right) \Big|_0^{\pi/2} \\ &= \frac{28}{3} - \pi - \frac{38}{3}i. \end{aligned}$$

The second integral used the formula

$$\sin t = \frac{1}{2i}(e^{it} - e^{-it}). \quad \square$$

Example 7 Compute $\int_{\gamma} \cos z dz$, where γ is the line segment from $-(\pi/2) + i$ to $\pi + i$.

Solution The curve is given by

$$\gamma(t) = t(\pi + i) + (1 - t)\left(-\frac{\pi}{2} + i\right) = \frac{3\pi}{2}t - \frac{\pi}{2} + i, \quad 0 \leq t \leq 1.$$

Furthermore,

$$\cos(x + iy) = \cos x \cosh y - i \sin x \sinh y.$$

(See the formula in Exercise 16 of Section 5.) Hence,

$$\begin{aligned} \int_{\gamma} \cos z dz &= \int_0^1 \left\{ \cos\left(\frac{3\pi}{2}t - \frac{\pi}{2}\right) \cosh(1) - i \sin\left(\frac{3\pi}{2}t - \frac{\pi}{2}\right) \sinh(1) \right\} \frac{3\pi}{2} dt \\ &= \left\{ \cosh(1) \sin\left(\frac{3\pi}{2}t - \frac{\pi}{2}\right) + i \sinh(1) \cos\left(\frac{3\pi}{2}t - \frac{\pi}{2}\right) \right\} \Big|_0^1 \\ &= \cosh(1) - i \sinh(1). \quad \square \end{aligned}$$

Example 8 Estimate

$$\left| \int_{\gamma} \frac{1}{z^2 + 4} dz \right|,$$

where γ is the semicircle $Re^{i\theta}$, $-\pi \leq \theta \leq 0$, $R > 2$.

Solution On the semicircle,

$$\left| \frac{1}{z^2 + 4} \right| \leq \frac{1}{R^2 - 4},$$

since $|z^2 + 4| \geq |z|^2 - 4$ by the triangle inequality (see Section 2). The length of the semicircle is πR , so the absolute value of the integral cannot exceed $\pi R / (R^2 - 4)$. \square

Example 9 Estimate $|\int_{\gamma} e^{-z} dz|$, where γ is the vertical line segment from $-i + 1$ to $i + 1$.

Solution On γ , $|e^{-z}| = e^{-x} = e^{-1}$. The length of γ is 2, so the integral cannot exceed $2/e$ in absolute value. \square

Example 10 Let u be a continuous function in the disc $|z - z_0| < r$, and let γ_ε be the circle $|z - z_0| = \varepsilon$. Show that

$$\lim_{\varepsilon \rightarrow 0} \frac{1}{2\pi i} \int_{\gamma_\varepsilon} \frac{u(z)}{z - z_0} dz = u(z_0).$$

Solution The circle is the range of $\gamma_\varepsilon(t) = z_0 + \varepsilon e^{it}$, $0 \leq t \leq 2\pi$. Therefore,

$$\begin{aligned} \frac{1}{2\pi i} \int_{\gamma_\varepsilon} \frac{u(z)}{z - z_0} dz &= \frac{1}{2\pi i} \int_0^{2\pi} \frac{u(z_0 + \varepsilon e^{it})}{\varepsilon e^{it}} i \varepsilon e^{it} dt \\ &= \frac{1}{2\pi} \int_0^{2\pi} u(z_0 + \varepsilon e^{it}) dt. \end{aligned}$$

Thus,

$$\begin{aligned} \left| \frac{1}{2\pi i} \int_{\gamma_\varepsilon} \frac{u(z)}{z - z_0} dz - u(z_0) \right| &= \left| \frac{1}{2\pi} \int_0^{2\pi} \{u(z_0 + \varepsilon e^{it}) - u(z_0)\} dt \right| \\ &\leq \max_{0 \leq t \leq 2\pi} \{|u(z_0 + \varepsilon e^{it}) - u(z_0)|\}. \end{aligned}$$

This last quantity goes to zero as $\varepsilon \rightarrow 0$ exactly because u is continuous at z_0 (see Exercise 26 in Section 4). \square

We digress briefly at this point to state a differentiation formula that will be needed. Its verification is simple and is relegated to the exercises. If $z(t)$ and $w(t)$ are

two complex-valued functions on an interval $[a, b]$, both of which are differentiable, then so is their product, and

$$(zw)'(t) = z'(t)w(t) + z(t)w'(t). \quad (4)$$

Repeated applications of (4), or mathematical induction, yield

$$(z^m)'(t) = m(z(t))^{m-1}z'(t), \quad m = 1, 2, \dots. \quad (5)$$

Example 11 Let γ be a piecewise smooth closed curve. Show that

$$\int_{\gamma} z^m dz = 0, \quad m = 0, 1, 2, \dots. \quad (6)$$

Solution Let $[t_j, t_{j+1}]$ be a segment in $[a, b]$ on which $\gamma'(t)$ is continuous. From (5),

$$\frac{d}{dt} (\gamma(t))^{m+1} = (m+1)\gamma^m(t)\gamma'(t).$$

This gives

$$\int_{t_j}^{t_{j+1}} \gamma^m(t)\gamma'(t) dt = \frac{1}{m+1} \{\gamma^{m+1}(t_{j+1}) - \gamma^{m+1}(t_j)\}.$$

Hence,

$$\begin{aligned} \int_{\gamma} z^m dz &= \sum_{j=0}^{n-1} \int_{t_j}^{t_{j+1}} \gamma^m(t)\gamma'(t) dt \\ &= \sum_{j=0}^{n-1} \frac{1}{m+1} \{\gamma^{m+1}(t_{j+1}) - \gamma^{m+1}(t_j)\} \\ &= \frac{1}{m+1} \{\gamma^{m+1}(b) - \gamma^{m+1}(a)\} = 0, \end{aligned}$$

because $\gamma(b) = \gamma(a)$, since γ is a closed curve. The result in (6) will be important in Chapter 2, when we prove Cauchy's Theorem. \square

Let p and q be distinct points in the plane, and suppose that γ_1 and γ_2 are two piecewise smooth curves from p to q . Then $\gamma = \gamma_1 - \gamma_2$ is a closed piecewise smooth curve. Hence, for $m = 0, 1, 2, \dots$,

$$0 = \int_{\gamma} z^m dz = \int_{\gamma_1} z^m dz - \int_{\gamma_2} z^m dz.$$

That is, the value of the integral of z^m along a curve joining p to q does not depend

on the curve but only on p , q , and m . Indeed, a glance at the computations in Example 11 shows that

$$\begin{aligned}\int_{\gamma} z^m dz &= \int_a^b \gamma^m(t) \gamma'(t) dt = \frac{1}{m+1} \gamma^{m+1}(t) \Big|_{t=a}^{t=b} \\ &= \frac{1}{m+1} [\gamma^{m+1}(b) - \gamma^{m+1}(a)] \\ &= \frac{1}{m+1} [q^{m+1} - p^{m+1}]\end{aligned}$$

for any curve γ joining p to q .

A comment on line integrals. Recall that a curve, say an arc of a circle or a straight line segment, may have several different parametrizations. For instance, $\gamma_1(s) = s + is^2$, $0 \leq s \leq 1$, and $\gamma_2(t) = \sin t + i(1 - \cos^2 t)$, $0 \leq t \leq \pi/2$, both have the same range—namely, the graph of the parabola $y = x^2$, $0 \leq x \leq 1$. It is natural to wonder whether the value of the line integral $\int_{\gamma} f$ depends on the parametrization (that is, on the function γ whose range is the given curve). Thankfully, under very general conditions, the answer is “no.” To give a proof of this would stray too far from our present purposes, but you can find one in many calculus texts, generally under the heading of “Green’s Theorem” or “line integrals.”

Green's* Theorem

The most important result on line integrals is Green’s Theorem, which will be stated after several more definitions. Green’s Theorem is formulated for a domain Ω whose boundary Γ consists of a finite number of disjoint, piecewise smooth simple closed curves $\gamma_1, \dots, \gamma_n$ (see Fig. 1.31a). We orient the boundary Γ of Ω **positively** by requiring that Ω remain on the left as we walk along Γ . Thus, the “outer” piece of the boundary of Ω is oriented counterclockwise, and each “inner” piece of Γ (if there are any) is oriented clockwise (see Fig. 1.31b). For example, if Ω is $\{z: 1 < |z| < 2\}$, the boundary of Ω is positively oriented if the circle $\{z: |z| = 1\}$ is traversed clockwise and the circle $\{z: |z| = 2\}$ counterclockwise.

Henceforth in this section, Ω is a domain whose boundary Γ consists of a finite number of disjoint, piecewise smooth simple closed curves $\gamma_1, \dots, \gamma_n$ and Γ is positively oriented. If f is a continuous complex-valued function on Γ , we define the line integral of f over Γ by

$$\int_{\Gamma} f(z) dz = \sum_{j=1}^n \int_{\gamma_j} f(z) dz,$$

where, of course, we already know how to compute each $\int_{\gamma_j} f(z) dz$.

* George Green, 1793–1841.




Figure 1.31

Green's Theorem relates the line integral of a function f over Γ to the integral of a certain related function over Ω . In order to formulate it properly, we assume that there is some open set D that contains both Ω and Γ and, on D , f has continuous partial derivatives with respect to both x and y . That is, if $f = p + iq$, then

$$\frac{\partial f}{\partial x} = \frac{\partial p}{\partial x} + i \frac{\partial q}{\partial x}, \quad \frac{\partial f}{\partial y} = \frac{\partial p}{\partial y} + i \frac{\partial q}{\partial y},$$

where all of $\partial p / \partial x$, $\partial q / \partial x$, $\partial p / \partial y$, and $\partial q / \partial y$ are continuous on D .

With all this background, then, we can state the theorem.

THEOREM 1 Green's Theorem

$$\int_{\Gamma} f(z) dz = i \iint_{\Omega} \left\{ \frac{\partial f}{\partial x} + i \frac{\partial f}{\partial y} \right\} dx dy. \quad (7)$$

If you have studied Green's Theorem in calculus, you may not recognize this formulation, although it is one of the easiest to work with. Let us recast Green's Theorem in what may be more familiar terms. If u and v are two real-valued functions on an open set D containing Ω and Γ , and if u and v both have continuous partial derivatives with respect to x and y on D , and if on Γ we set $dx = (\operatorname{Re} \gamma'(t)) dt$ and $dy = (\operatorname{Im} \gamma'(t)) dt$, then Green's Theorem can also be stated in this way:

$$\int_{\Gamma} \{u dx + v dy\} = \iint_{\Omega} \left(\frac{\partial v}{\partial x} - \frac{\partial u}{\partial y} \right) dx dy. \quad (8)$$

The proof of equivalence of these two versions of Green's Theorem is left to the exercises. ■

The first example that follows is a verification of Green's Theorem for a triangle and some related figures. The three examples after that derive some consequences of Green's Theorem.




Figure 1.32

Example 12* Verify Green's Theorem for a triangle.

Solution Suppose first that the triangle Γ has one horizontal side (see Fig. 1.32).

Let v be a function that has continuous partial derivatives on some open set containing the triangle Γ and its inside Ω . To compute $\int_{\Gamma} v \, dy$, parametrize the edges of Γ in the following way:

$$\text{On } PQ: \quad x = t, y = b; \quad a \leq t \leq c$$

$$\text{On } QR: \quad y = t, x = c + (d - c) \frac{t - b}{e - b}; \quad b \leq t \leq e$$

$$\text{On } RP: \quad y = t, x = d + (a - d) \frac{t - e}{b - e}; \quad e \geq t \geq b.$$

(We have assumed that $b < e$; if $e < b$, just reverse the range of t and the order of the sides.) Hence,

$$dy = \begin{cases} 0 & \text{on } PQ \\ dt & \text{on } QR \\ dt & \text{on } RP. \end{cases}$$

This gives

$$\begin{aligned} \int_{\Gamma} v \, dy &= \int_{PQ} v \, dy + \int_{QR} v \, dy + \int_{RP} v \, dy \\ &= 0 + \int_b^e v(At + B, t) \, dt + \int_e^b v(Ct + D, t) \, dt, \end{aligned}$$

where for simplicity we have written

$$A = \frac{d - c}{e - b}, \quad B = c - b \frac{d - c}{e - b}, \quad C = \frac{a - d}{b - e}, \quad D = d - e \frac{a - d}{b - e}.$$

On the other hand, the area integral of $\partial v / \partial x$ over the inside Ω of the triangle is




Figure 1.33

computed in this way. Ω is described by

$$\Omega = \{(x, y) : b \leq y \leq e, Cy + D \leq x \leq Ay + B\}.$$

Consequently,

$$\begin{aligned} \iint_{\Omega} \frac{\partial v}{\partial x} dx dy &= \int_b^e \left\{ \int_{Ct+D}^{At+B} \frac{\partial v}{\partial x} dx \right\} dt \\ &= \int_b^e \{v(At + B, t) - v(Ct + D, t)\} dt \\ &= \int_{\Gamma} v dy. \end{aligned}$$

Next, any triangle Γ can be divided in two by a single horizontal line segment from one of its vertices to the opposite side so that each of the resulting triangles has one horizontal edge (Figs. 1.33a and 1.33b). Let Γ_1 and Γ_2 be the two resulting triangles, each positively oriented, and let Ω_1 and Ω_2 be their respective insides. Then

$$\int_{\Gamma_j} v dy = \iint_{\Omega_j} \frac{\partial v}{\partial x} dx dy, \quad j = 1, 2,$$

from the computations above. The integral over the segment PS appears in both $\int_{\Gamma_1} v dy$ and $\int_{\Gamma_2} v dy$, but in both cases it is zero, since PS is horizontal. Hence,

$$\begin{aligned} \int_{\Gamma} v dy &= \int_{\Gamma_1} v dy + \int_{\Gamma_2} v dy \\ &= \iint_{\Omega_1} \frac{\partial v}{\partial x} dx dy + \iint_{\Omega_2} \frac{\partial v}{\partial x} dx dy \\ &= \iint_{\Omega} \frac{\partial v}{\partial x} dx dy. \end{aligned}$$

This establishes the formula

$$\iint_{\Omega} \frac{\partial v}{\partial x} dx dy = \int_{\Gamma} v dy$$

for any triangle Γ and its inside Ω . A similar argument, using triangles with a vertical edge, establishes the formula

$$\iint_{\Omega} \frac{\partial u}{\partial y} dx dy = - \int_{\Gamma} u dx.$$

Thus, Green's Theorem is verified for a triangle. \square

Some further comments. Once Green's Theorem is proved for all triangles, it also follows for many other regions. For instance, let Ω be a bounded convex domain whose boundary consists of straight line segments (Fig. 1.34a). Select a point P in Ω and draw the line segments from P to each vertex of Ω (Fig. 1.34b). This breaks Ω up into nonoverlapping solid triangles $\Omega_1, \dots, \Omega_r$ with boundaries $\Gamma_1, \dots, \Gamma_r$, respectively. Then apply Green's Theorem to each of these triangles, obtaining

$$\iint_{\Omega_j} \left(\frac{\partial v}{\partial x} - \frac{\partial u}{\partial y} \right) dx dy = \int_{\Gamma_j} (v dy + u dx), \quad j = 1, \dots, r.$$

Each line segment joining P to a vertex of Ω is integrated over twice, once from P to the vertex and once from the vertex to P . These line integrals then cancel each other, and

$$\begin{aligned} \int_{\Gamma} (v dy + u dx) &= \sum_{j=1}^r \int_{\Gamma_j} (v dy + u dx) \\ &= \sum_{j=1}^r \iint_{\Omega_j} \left(\frac{\partial v}{\partial x} - \frac{\partial u}{\partial y} \right) dx dy \\ &= \iint_{\Omega} \left(\frac{\partial v}{\partial x} - \frac{\partial u}{\partial y} \right) dx dy. \end{aligned}$$





Figure 1.34

Thus, Green's Theorem holds for Ω . A similar sort of “triangulation” argument can be applied to show that Green's Theorem is valid for a bounded domain Ω whose boundary Γ is composed of straight line segments. (Figs. 1.35a and 1.35b show representative types.)




Figure 1.35

Example 13 Suppose that γ is a piecewise smooth positively oriented simple closed curve. The value of the integral

$$\frac{1}{2\pi i} \int_{\gamma} \frac{dz}{z - p}, \quad p \text{ not in } \gamma,$$

is a crucial factor in many of the theorems of complex variables. We shall now use Green's Theorem to show that

$$\frac{1}{2\pi i} \int_{\gamma} \frac{dz}{z - p} = \begin{cases} 0 & \text{if } p \text{ is outside } \gamma, \\ 1 & \text{if } p \text{ is inside } \gamma. \end{cases}$$

Let Ω be the domain inside γ . Suppose first that p is outside γ . Let $f(z) = (z - p)^{-1}$; then f has continuous partial derivatives everywhere except at p and, indeed, it is elementary to compute that

$$\frac{\partial f}{\partial x} = \frac{-1}{(z - p)^2}; \quad \frac{\partial f}{\partial y} = \frac{-i}{(z - p)^2}.$$

Hence, $(\partial f / \partial x) + i(\partial f / \partial y) = 0$ in Ω , so

$$\int_{\gamma} f(z) dz = i \iint_{\Omega} \left\{ \frac{\partial f}{\partial x} + i \frac{\partial f}{\partial y} \right\} dx dy = 0.$$

Suppose now that p is inside γ . Then, of course, the function $f(z) = (z - p)^{-1}$ does *not* have continuous partials at p , so Green's Theorem is not immediately applicable. However, let γ_1 be a circle centered at p of radius ϵ , where ϵ is so small that the disc of radius 2ϵ centered at p lies within Ω . Take Ω_1 to be the region inside γ but outside γ_1 and orient the boundary of Ω_1 positively, so γ is already correctly

oriented and γ_1 is oriented clockwise. Then, just as above,

$$\frac{\partial f}{\partial x} + i \frac{\partial f}{\partial y} = 0 \quad \text{on } \Omega_1,$$

so Green's Theorem implies that

$$\int_{\gamma} \frac{dz}{z-p} + \int_{\gamma_1} \frac{dz}{z-p} = 0.$$

Reversing the orientation of γ_1 so that it is now traversed counterclockwise, we conclude that

$$\int_{\gamma} \frac{dz}{z-p} = \int_{-\gamma_1} \frac{dz}{z-p}.$$

To evaluate the integral over $-\gamma_1$, set $z = p + \varepsilon e^{it}$, $0 \leq t \leq 2\pi$; then $dz = i\varepsilon e^{it} dt$, and

$$\int_{\gamma} \frac{dz}{z-p} = \int_{-\gamma_1} \frac{dz}{z-p} = \int_0^{2\pi} \frac{i\varepsilon e^{it} dt}{\varepsilon e^{it}} = 2\pi i.$$
□

Example 14 This example will derive from Green's Theorem an important formula with numerous applications. The formula is called, not surprisingly, **Green's Formula**.

Begin by defining the normal derivative of a real-valued function h on the boundary of Ω . Recall from calculus that if $\lambda = \cos \theta + i \sin \theta$ is a complex number of modulus one, then the **directional derivative** of a function h in the direction λ at a point z_0 is

$$\lim_{t \rightarrow 0} \frac{h(z_0 + t\lambda) - h(z_0)}{t}.$$

This is, in fact (as you learned in calculus), equal to the dot product of the gradient of h with the vector $(\cos \theta, \sin \theta)$ and so has the value

$$\frac{\partial h}{\partial x} \cos \theta + \frac{\partial h}{\partial y} \sin \theta.$$

Let $z_0 = z(t_0)$ be a point of Γ , and let $\gamma(t) = x(t) + iy(t)$ parameterize a portion of Γ containing z_0 . A unit vector pointing out of Ω and normal to Γ is

$$\lambda = (y'(t) - ix'(t))([x'(t)]^2 + [y'(t)]^2)^{-1/2}.$$

(We shall always assume that $[x'(t)]^2 + [y'(t)]^2$ is strictly positive.) Thus, the directional derivative of the function h in the direction normal to the boundary of

Γ , with the normal vector pointing out of Γ , is

$$\frac{\partial h}{\partial n} = \left(\frac{\partial h}{\partial x} y'(t) - \frac{\partial h}{\partial y} x'(t) \right) ([x'(t)]^2 + [y'(t)]^2)^{-1/2}.$$

Now take f and g to be two real-valued functions that have continuous partial derivatives up to order two on some open set containing both Ω and Γ . In the second formulation of Green's Theorem, formula (8), define u and v by

$$v = \frac{\partial f}{\partial x} g - \frac{\partial g}{\partial x} f \quad \text{and} \quad u = -\frac{\partial f}{\partial y} g + \frac{\partial g}{\partial y} f.$$

After a bit of rearranging, we obtain

$$\int_{\Gamma} \left[g \left(\frac{\partial f}{\partial x} dy - \frac{\partial f}{\partial y} dx \right) - f \left(\frac{\partial g}{\partial x} dy - \frac{\partial g}{\partial y} dx \right) \right] = \iint_{\Omega} [(\Delta f)g - (\Delta g)f] dx dy,$$

where

$$\Delta f = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2}$$

is the **Laplacian*** of f and (likewise) Δg is the Laplacian of g . In the line integral, multiply and divide by the expression $[(x'(t))^2 + (y'(t))^2]^{-1/2}$ and then recall from calculus that the element of **arc-length** ds along Γ is given by

$$ds = [(x'(t))^2 + (y'(t))^2]^{1/2} dt.$$

We then obtain the desired formula:

$$\int_{\Gamma} \left(g \frac{\partial f}{\partial n} - f \frac{\partial g}{\partial n} \right) ds = \iint_{\Omega} (g \Delta f - f \Delta g) dx dy. \quad \square \quad (9)$$

Example 15 In a similar fashion, if we take

$$u = f \frac{\partial f}{\partial x} \quad \text{and} \quad v = -f \frac{\partial f}{\partial y}$$

in (8), we obtain the formula

$$\int_{\Gamma} f \frac{\partial f}{\partial n} ds = \iint_{\Omega} \left\{ f \Delta f + \left(\frac{\partial f}{\partial x} \right)^2 + \left(\frac{\partial f}{\partial y} \right)^2 \right\} dx dy. \quad \square \quad (10)$$

The formulas (9) and (10) will be important in the discussion of harmonic functions. One application will suffice for now.

* Named after Pierre-Simon Laplace, 1749–1827.

DEFINITION A function $u(z) = u(x, y)$ with continuous first and second partial derivatives with respect to both x and y is **harmonic** on an open set D if

$$\Delta u = \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0 \quad \text{on } D.$$

THEOREM 2 Suppose that u is a real-valued harmonic function on an open set D and D contains a domain Ω and the boundary Γ of Ω . Assume that Γ consists of a finite number of disjoint, piecewise smooth simple closed curves. If $u = 0$ on Γ , then $u = 0$ in Ω as well.

Proof The proof comes directly from (10) with, of course, u in place of f . The integral $\int_{\Gamma} u \, \partial u / \partial n \, ds$ is zero, since $u = 0$ on Γ . On Ω , $\Delta u = 0$; hence,

$$0 = \iint_{\Omega} \left\{ \left(\frac{\partial u}{\partial x} \right)^2 + \left(\frac{\partial u}{\partial y} \right)^2 \right\} dx dy.$$

But then $\partial u / \partial x = \partial u / \partial y = 0$ on Ω , so $u(x, y)$ is constant on Ω . Since $u = 0$ on Γ , it follows that $u = 0$ on Ω . ■

EXERCISES FOR SECTION 1.6

Compute the following line integrals.

1. $\int_{\gamma} z \, dz$, where γ is the semicircle from i to $-i$, which passes through -1 .
2. $\int_{\gamma} e^z \, dz$, where γ is the line segment from 0 to z_0 .
3. $\int_{\gamma} |z|^2 \, dz$, where γ is the line segment from 2 to $3 + i$.
4. $\int_{\gamma} 1/(z + 4) \, dz$, where γ is the circle of radius 1 centered at -4 , oriented counterclockwise.
5. $\int_{\gamma} (\operatorname{Re} z) \, dz$, where γ is the line segment from 1 to i .
6. $\int_{\gamma} (z^2 + 3z + 4) \, dz$, where γ is the circle $|z| = 2$ oriented counterclockwise.
7. Let γ_1 be the semicircle from 1 to -1 through i and γ_2 the semicircle from 1 to -1 through $-i$ (Fig. 1.36).




Figure 1.36

Compute $\int_{\gamma_1} z^2 dz$ and $\int_{\gamma_2} z^2 dz$. Can you account for the fact that they are equal?

Now compute $\int_{\gamma_1} \bar{z} dz$ and $\int_{\gamma_2} \bar{z} dz$. Can you account for the fact that they are not equal?

8. Explicitly verify the conclusion of Example 11 by computing $\int_{\gamma} z dz$ and $\int_{\gamma} z^2 dz$ when γ is the square with vertices at $\pm 1 \pm i$.

9. (a) Show that $\frac{1}{2\pi} \int_0^{2\pi} e^{ik\theta} d\theta = \begin{cases} 1, & k = 0 \\ 0, & k \neq 0. \end{cases}$

- (b) Explicitly verify the conclusion of Example 13 for γ , the circle of radius 1, centered at 0. (Hint:

$$\frac{1}{z - p} = \sum_{k=0}^{\infty} \frac{p^k}{z^{k+1}} \quad \text{if } 0 \leq |p| < 1 \quad \text{and} \quad |z| = 1$$

$$\frac{1}{z - p} = - \sum_{k=0}^{\infty} \frac{z^k}{p^{k+1}} \quad \text{if } |z| = 1 \quad \text{and} \quad 1 < |p| < \infty.$$

Interchange summation and integration; then use (a).)

10. Let $f = u + iv$ be a continuous function and $\gamma(t) = x(t) + iy(t)$ be a piecewise smooth curve. Show that

$$\operatorname{Re} \left\{ \int_{\gamma} f(z) dz \right\} = \int_{\gamma} (u dx - v dy)$$

and

$$\operatorname{Im} \left\{ \int_{\gamma} f(z) dz \right\} = \int_{\gamma} (v dx + u dy).$$

Here, $dx = x'(t) dt$, $dy = y'(t) dt$.

11. Use Exercise 10 to show the equivalence of the two formulations of Green's Theorem: (7) and (8).
12. Use the result of (the extension of) Example 11 to compute the following integrals: (a) $\int_{\gamma} (z^3 - 6z^2 + 4) dz$, where γ is any curve joining $-1 + i$ to 1.
- (b) $\int_{\gamma} (z^4 + z^2) dz$, where γ is any curve joining $-i$ to $2 + i$.
13. Use Green's Theorem to derive Green's Identity:

$$\iint_{\Omega} \left\{ \frac{\partial u}{\partial x} \frac{\partial v}{\partial x} + \frac{\partial u}{\partial y} \frac{\partial v}{\partial y} \right\} dx dy = \int_{\Gamma} v \frac{\partial u}{\partial n} ds - \iint_{\Omega} v \Delta u dx dy.$$

14. Use Exercise 13 to derive Green's Formula (9):

$$\int_{\Gamma} \left(g \frac{\partial f}{\partial n} - f \frac{\partial g}{\partial n} \right) ds = \iint_{\Omega} (g \Delta f - f \Delta g) dx dy.$$

15. Let u be a continuous function on the complex plane, which is bounded:

$|u(z)| \leq C$ for all z . Let γ_R be the circle $|z| = R$. Show that

$$\lim_{R \rightarrow \infty} \int_{\gamma_R} \frac{u(z)}{(z - z_0)^2} dz = 0,$$

for each z_0 . (Hint: Use (3).)

16. Let z_0 be outside a piecewise smooth simple closed curve γ . Extend Example 13 by showing that

$$\int_{\gamma} \frac{dz}{(z - z_0)^m} = 0, \quad m = 2, 3, 4, \dots$$

Exact Differentials*

DEFINITION An expression $P(x, y) dx + Q(x, y) dy$ is an **exact differential** if there is a function $g(x, y)$ with

$$\frac{\partial g}{\partial x} = P \quad \text{and} \quad \frac{\partial g}{\partial y} = Q.$$

In Exercises 17 to 20, D is a disc and P and Q are functions on D with continuous partial derivatives with respect to x and y .

17. If $P dx + Q dy$ is an exact differential, show that

$$\int_{\gamma} \{P dx + Q dy\} = 0, \tag{11}$$

for each closed curve γ in D .

18. Use Green's Theorem to show that

$$\frac{\partial Q}{\partial x} = \frac{\partial P}{\partial y} \tag{12}$$

throughout D if and only if (11) holds.

19. Suppose that (11) holds for each closed curve γ in D . Show that $P dx + Q dy$ is an exact differential. (Hint: Let $x_0 + iy_0$ be any point in D and set

$$g(x, y) = \int_{x_0}^x P(t, y_0) dt + \int_{y_0}^y Q(x, s) ds.$$

Use (12) to show that g is the desired function.)

20. Let f be a continuous complex-valued function on D . Show that there is a complex-valued function F on D with

$$\frac{\partial F}{\partial x} = f, \quad \frac{\partial F}{\partial y} = if \tag{13}$$

if and only if

$$\int_{\gamma} f(z) dz = 0 \quad (14)$$

for each closed curve γ in D . You may need to use the results of Exercise 10.

21. Example 11 showed that $\int_{\gamma} z^m dz = 0$ for each closed curve γ , $m = 0, 1, 2, \dots$. What is the function F in Exercise 20 that corresponds to $f(z) = z^m$?

Further Reading

The paperback book by Konrad Knopp, *Elements of the Theory of Functions* (New York: Dover, 1952) is an excellent source for virtually all the material in this chapter. The sections there on linear fractional transformations can be skipped or read, depending on your taste; these functions are covered in Chapter 3 of this book.

2

Basic Properties of Analytic Functions

2.1 Analytic and Harmonic Functions; the Cauchy–Riemann Equations

Analytic functions and their close relatives, harmonic functions, are the stuff of which the subject of complex variables is built. This section introduces both of these types of functions; here and in subsequent sections, many of their significant properties are developed.

A function f defined for z in a domain D is **differentiable** at a point z_0 in D if

$$\lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0} = \lim_{h \rightarrow 0} \frac{f(z_0 + h) - f(z_0)}{h} \quad (1)$$

exists; the limit, if it exists, is denoted by $f'(z_0)$. If f is differentiable at each point of the domain D , then f is called **analytic** in D . A function analytic on the whole complex plane is called **entire**.

It is worth stressing here that the limit in (1) is required to exist *no matter how z approaches z_0* ; equivalently, no matter how h approaches 0. That is, although $|h|$ must approach zero, the argument of h can change arbitrarily. This latitude is at the root of the difference between differentiable functions of a real variable and differentiable functions of a complex variable.

Example 1 $f(z) = z^n$, $n = 1, 2, \dots$, is entire, and $f'(z) = nz^{n-1}$. For

$$\begin{aligned} (z + h)^n - z^n &= nz^{n-1}h + \frac{n(n-1)}{2}z^{n-2}h^2 + \cdots + h^n \\ &= h \left(nz^{n-1} + \frac{n(n-1)}{2}z^{n-2}h + \cdots + h^{n-1} \right), \end{aligned}$$

so

$$\frac{(z + h)^n - z^n}{h} \rightarrow nz^{n-1} \quad \text{as} \quad h \rightarrow 0. \quad \square$$

Example 2 The sum and product of functions analytic on a common domain are again analytic, and their derivatives obey the rules already familiar from calculus:

$$(f + g)' = f' + g', \quad (2)$$

and

$$(fg)' = f'g + fg'. \quad \square \quad (3)$$

Example 3 The quotient of two analytic functions is differentiable at all points z_0 at which the denominator does not vanish and

$$\left(\frac{f}{g}\right)'(z_0) = \frac{g(z_0)f'(z_0) - f(z_0)g'(z_0)}{(g(z_0))^2}, \quad g(z_0) \neq 0. \quad \square \quad (4)$$

Example 4 If f and g are differentiable and if the range of f lies within the domain of g , then $g(f(z))$ is differentiable with

$$[g(f(z))]' = g'(f(z))f'(z). \quad \square \quad (5)$$

(The proofs of formulas (2) through (5) are straightforward and are left to the exercises at the end of this section.)

From Examples 1 to 4 we can immediately give two general types of analytic functions.

Example 5 Any polynomial $p(z) = a_0 + a_1 z + \cdots + a_n z^n$ is an entire function. \square

Example 6 A rational function $r = p/q$, where p and q are polynomials, is analytic on any domain containing no zero of q . \square

Example 7 Show that the exponential function $f(z) = e^z$ is an entire function.

Solution We know that

$$e^{z+h} - e^z = e^z(e^h - 1),$$

by the properties of the exponential function derived in Section 5 of Chapter 1. Furthermore, with $h = \sigma + i\tau$,

$$\begin{aligned} e^h - 1 - h &= \{e^\sigma \cos \tau - 1 - \sigma\} + i\{e^\sigma \sin \tau - \tau\} \\ &= \{e^\sigma(\cos \tau - 1) + e^\sigma - 1 - \sigma\} + i\{e^\sigma(\sin \tau - \tau) + \tau(e^\sigma - 1)\}. \end{aligned}$$

Hence,

$$\begin{aligned} \left| \frac{e^h - 1 - h}{h} - 1 \right| &= \left| \frac{e^h - 1 - h}{h} \right| \\ &\leqslant e^\sigma \left| \frac{1 - \cos \tau}{\tau} \right| + \left| \frac{e^\sigma - 1 - \sigma}{\sigma} \right| + e^\sigma \left| \frac{\sin \tau - \tau}{\tau} \right| + |e^\sigma - 1|. \end{aligned}$$

To obtain this inequality, we used the triangle inequality, as well as the simple facts that

$$\frac{1}{|h|} \leq \frac{1}{|\tau|} \quad \text{and} \quad \frac{1}{|h|} \leq \frac{1}{|\sigma|}.$$

However, each of the four quantities within absolute value signs approaches zero as σ and τ independently approach zero (use l'Hôpital's Rule on each, if you like), so

$$\lim_{h \rightarrow 0} \frac{e^h - 1}{h} = 1.$$

This finally gives

$$\lim_{h \rightarrow 0} \frac{e^{z+h} - e^z}{h} = e^z \lim_{h \rightarrow 0} \frac{e^h - 1}{h} = e^z.$$

Consequently, e^z is differentiable at all points z , and

$$(e^z)' = e^z. \quad \square \quad (6)$$

Example 8 It follows from (5) and (6) that if $f(z)$ is analytic on a domain D , then the function

$$G(z) = \exp(f(z))$$

is also analytic on D , and

$$G'(z) = f'(z) \exp(f(z)), \quad z \in D \quad \square$$

Example 9 Let $f(z)$ be differentiable at a point z_0 in a domain D . Show that f is continuous at z_0 .

Solution For z near z_0 ,

$$|f(z) - f(z_0)| = \left| \frac{f(z) - f(z_0)}{z - z_0} \right| |z - z_0| \rightarrow |f'(z_0)| \cdot 0 = 0$$

as $z \rightarrow z_0$. Hence, $\lim_{z \rightarrow z_0} f(z) = f(z_0)$. In particular, both the real and the imaginary parts of f are continuous on any domain on which f is analytic. \square

The Cauchy*–Riemann† Equations

The essential feature of an analytic function is the fact that the limit in (1) must exist *no matter how z approaches z_0* . This leads to the pair of famous partial differential equations, which connect the real and imaginary parts of an analytic function.

* Augustine Louis Cauchy, 1789–1857.

† Bernhard Riemann, 1826–1866.

THEOREM 1 Cauchy–Riemann Equations Suppose that $f = u + iv$ is analytic on a domain D . Then throughout D ,

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \quad \text{and} \quad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}. \quad (7)$$

Proof Let us employ the definition of the derivative of f at z_0 in two different ways. First take h to be real. Then

$$\begin{aligned} f'(z_0) &= \lim_{h \rightarrow 0} \left[\frac{u(x_0 + h, y_0) - u(x_0, y_0)}{h} + i \frac{v(x_0 + h, y_0) - v(x_0, y_0)}{h} \right] \\ &= \frac{\partial u}{\partial x}(x_0, y_0) + i \frac{\partial v}{\partial x}(x_0, y_0). \end{aligned}$$

Next, take $h = ik$, where k is real. Then

$$\begin{aligned} f'(z_0) &= \lim_{k \rightarrow 0} \left[\frac{u(x_0, y_0 + k) - u(x_0, y_0)}{ik} + i \frac{v(x_0, y_0 + k) - v(x_0, y_0)}{ik} \right] \\ &= \frac{1}{i} \frac{\partial u}{\partial y}(x_0, y_0) + \frac{\partial v}{\partial y}(x_0, y_0). \end{aligned}$$

All that remains is to equate real and imaginary parts of these two expressions for $f'(z_0)$. ■

The Cauchy–Riemann equations imply that an analytic function $f = u + iv$ is actually determined by its real part u (or, equivalently, by its imaginary part v), apart from an additive constant. For if we know u , we can (presumably) find v with

$$\frac{\partial v}{\partial x} = -\frac{\partial u}{\partial y} \quad \text{and} \quad \frac{\partial v}{\partial y} = \frac{\partial u}{\partial x}.$$

For example, if $u(x, y) = x^3 - 3xy^2$, then v is to be found from

$$\frac{\partial v}{\partial x} = 6xy \quad \text{and} \quad \frac{\partial v}{\partial y} = 3x^2 - 3y^2.$$

Integrate $6xy$ with respect to x and find that $v(x, y) = 3x^2y + p(y)$, where p is a function of y alone. Then differentiate this expression for v with respect to y and compare the result with the known equality $\partial v / \partial y = 3x^2 - 3y^2$. This yields $3x^2 - 3y^2 = 3x^2 + p'(y)$. Hence, $p(y) = -y^3 + c$, where c is a constant. Consequently, $v(x, y) = 3x^2y - y^3 + c$. A word of warning is merited here: Not every function $u(x, y)$ is the real part of an analytic function. Indeed, if $f = u + iv$ is analytic on a domain D , and if we assume temporarily (and unnecessarily as it turns

out) that both u and v have continuous partial derivatives of first and second order, then

$$\begin{aligned}\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} &= \frac{\partial}{\partial x} \left(\frac{\partial u}{\partial x} \right) + \frac{\partial}{\partial y} \left(\frac{\partial u}{\partial y} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{\partial v}{\partial y} \right) + \frac{\partial}{\partial y} \left(-\frac{\partial v}{\partial x} \right) \\ &= \frac{\partial^2 v}{\partial x \partial y} - \frac{\partial^2 v}{\partial y \partial x} = 0.\end{aligned}$$

Hence, if $u = \operatorname{Re} f$, where f is analytic on D , then u must necessarily satisfy Laplace's equation:

$$\Delta u = \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0.$$

Recall from Section 6 of Chapter 1 that a continuous function u with continuous first and second partial derivatives on D is **harmonic** on D if it satisfies Laplace's equation on D . Thus, the real part u of an analytic function f is harmonic. Quite similarly, the imaginary part v of f is also harmonic. Harmonic functions arise with uncanny frequency in physical problems and provide one of the most important applications of complex variables. Many such applications are discussed in Chapter 4.

We note further that a real-valued harmonic function u determines many other harmonic functions v by the Cauchy–Riemann equations:

$$\frac{\partial v}{\partial y} = \frac{\partial u}{\partial x} \quad \text{and} \quad \frac{\partial v}{\partial x} = -\frac{\partial u}{\partial y}.$$

Such a v is called a **harmonic conjugate** of u ; any two harmonic conjugates v and v_1 of u differ by a constant, since

$$\frac{\partial}{\partial y} (v - v_1) = \frac{\partial u}{\partial x} - \frac{\partial u}{\partial x} = 0$$

and

$$\frac{\partial}{\partial x} (v - v_1) = -\frac{\partial u}{\partial y} + \frac{\partial u}{\partial y} = 0.$$

Moreover, if v is a harmonic conjugate of u in a disc Ω , then $f = u + iv$ is analytic on Ω . (The verification that f is indeed analytic depends on Theorem 3 of this section.) We will study harmonic functions further in Chapter 4.

There are other consequences of the Cauchy–Riemann equations. In each case of the following theorem, we shall draw the conclusion that the function in question is a constant.

THEOREM 2 Suppose that $f = u + iv$ is analytic on a domain D . If either u is constant on D or $u^2 + v^2$ is constant on D , then f is constant on D .

Proof Suppose that u is constant on D . Then $\partial u / \partial x = \partial u / \partial y = 0$, so by the Cauchy–Riemann equations, we also have $\partial v / \partial x = \partial v / \partial y = 0$ on D . Since $\partial v / \partial x$ is the rate of change of v on a horizontal line, it follows that v is constant on each horizontal line segment in D . Likewise, v is constant on each vertical line segment in D , since $\partial v / \partial y = 0$ in D . However, each pair of points in D can be joined by a polygonal curve consisting entirely of such segments (see Exercise 16, Section 3, of Chapter 1). Hence, v is constant in D .

Suppose that $|f| = \sqrt{u^2 + v^2}$ is equal to some constant c throughout D . If $c = 0$, then f is identically zero. Otherwise, we may divide f by c and assume that $|f| = 1$ throughout D . Hence, $\bar{f}(z) = 1/f(z)$ on D . However, $1/f(z)$ is analytic on D , being the reciprocal of a nonvanishing analytic function. Consequently, the sum $f + \bar{f}$ is analytic on D . This sum is then a real-valued analytic function; that is, $i(f + \bar{f})$ is analytic, and its real part is zero. The first part of the proof then implies that this function must be constant. Therefore, u is constant, and another application of the first part of the proof implies that f is constant. ■

One interpretation of this theorem is that an analytic function f is constant if its range lies in either a vertical line or a circle centered at the origin. We shall see later that there are, in fact, even more severe restrictions than this on the range of a nonconstant analytic function.

The Cauchy–Riemann equations, which are implied by analyticity, themselves imply analyticity with an extra hypothesis of continuity on the partial derivatives.

THEOREM 3 Suppose that $f = u + iv$ and that all of $u, v, \partial u / \partial x, \partial v / \partial x, \partial u / \partial y$, and $\partial v / \partial y$ are continuous in a disc centered at the point z_0 . If u and v satisfy the Cauchy–Riemann equations at z_0 , then f is differentiable at z_0 .

Proof The proof is based on the Taylor series expansion of u and v and the Cauchy–Riemann equations. We let $h = \delta + iv$ be a (small) complex number. The Taylor series for u near (x_0, y_0) is

$$u(x_0 + \delta, y_0 + v) = u(x_0, y_0) + \delta \left(\frac{\partial u}{\partial x} \right)(x_0, y_0) + v \left(\frac{\partial u}{\partial y} \right)(x_0, y_0) + \delta E_1 + v E_2,$$

where E_1 and E_2 depend on δ and v and approach zero as δ and v approach zero. Likewise,

$$v(x_0 + \delta, y_0 + v) = v(x_0, y_0) + \delta \left(\frac{\partial v}{\partial x} \right)(x_0, y_0) + v \left(\frac{\partial v}{\partial y} \right)(x_0, y_0) + \delta E_3 + v E_4,$$

where E_3 and E_4 depend on δ and v and approach zero as δ and v approach zero. Since the Cauchy–Riemann equations hold, replace $\partial v/\partial y(x_0, y_0)$ with $(\partial u/\partial x)(x_0, y_0)$ and $(\partial u/\partial y)(x_0, y_0)$ with $-(\partial v/\partial x)(x_0, y_0)$; then multiply the expression for v by i and add the result to that for u . This yields

$$\begin{aligned} f(z_0 + h) &= f(z_0) + \left[\left(\frac{\partial u}{\partial x} \right)(x_0, y_0) + i \left(\frac{\partial v}{\partial x} \right)(x_0, y_0) \right] h \\ &\quad + \delta(E_1 + iE_3) + v(E_2 + iE_4). \end{aligned}$$

From this and the facts that $|\delta/h| \leq 1$, $|v/h| \leq 1$, we obtain

$$\lim_{h \rightarrow 0} \frac{f(z_0 + h) - f(z_0)}{h} = \left(\frac{\partial u}{\partial x} \right)(x_0, y_0) + i \left(\frac{\partial v}{\partial x} \right)(x_0, y_0).$$

This is exactly the statement that f is differentiable at z_0 , with derivative $(\partial u/\partial x)(x_0, y_0) + i(\partial v/\partial x)(x_0, y_0)$. ■

Example 10 Let D be a domain on which there is a single-valued branch of $\log z$; for example, take D to be the complex plane with any ray from the origin to ∞ deleted. Show that $f(z) = \log z$ is differentiable on D , and find its derivative.

Solution

$$f(z) = \log z = \ln|z| + i \arg z$$

$$= \frac{1}{2} \ln(x^2 + y^2) + i \arctan \frac{y}{x},$$

where the values of $\arctan(y/x)$ are specified to lie in an interval of the form $(\theta_0, \theta_0 + 2\pi)$. Thus, the real and imaginary parts of $\log z$ may be expressed as

$$u(x, y) = \frac{1}{2} \ln(x^2 + y^2), \quad v(x, y) = \arctan \frac{y}{x}.$$

It follows that

$$\frac{\partial u}{\partial x} = \frac{x}{x^2 + y^2}, \quad \frac{\partial u}{\partial y} = \frac{y}{x^2 + y^2}, \quad \frac{\partial v}{\partial x} = \frac{-y}{x^2 + y^2}, \quad \frac{\partial v}{\partial y} = \frac{x}{x^2 + y^2}.$$

All the partial derivatives are continuous on D , and they clearly satisfy the Cauchy–Riemann equations at all points of D :

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}, \quad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}.$$

Consequently, Theorem 3 implies that $\log z$ is differentiable with derivative

$$\begin{aligned}(\log z)' &= \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = \frac{x}{x^2 + y^2} + i \frac{-y}{x^2 + y^2} \\&= \frac{\bar{z}}{|z|^2} = \frac{1}{z}.\end{aligned}$$

Thus,

$$(\log z)' = \frac{1}{z}. \quad \square \quad (9)$$

Note that $1/z$ is analytic on the whole complex plane except at $z = 0$, but the function $\log z$, whose derivative is $1/z$, is *not* analytic on the punctured plane; indeed, it is not analytic on any domain D that contains a simple closed curve that surrounds the origin (see Exercise 25).

Example 11 Let α be a real number. Show that the function

$$f(z) = z^\alpha$$

can be defined to be analytic on any domain D in the plane that omits a ray from the origin to infinity.

Solution The domain D omits a ray from 0 to infinity, so we may choose a branch of $\log z$ that is well-defined and analytic in D . For instance, if D omits the negative real axis and 0, then $\text{Log } z$ is analytic on D . With this branch of $\log z$ fixed, we define

$$z^\alpha = e^{\alpha \log z}.$$

The result of Example 8 now shows that z^α is analytic on D . \square

EXERCISES FOR SECTION 2.1

1. Establish the following differentiation formulas:

- | | |
|-------------------------------------|---------------------------------------|
| (a) $(\sin z)' = \cos z$ | (b) $(\cos z)' = -\sin z$ |
| (c) $(\sinh z)' = \cosh z$ | (d) $(\cosh z)' = \sinh z$ |
| (e) $(\tan z)' = \sec^2 z$ | (f) $(\arcsin z)' = (1 - z^2)^{-1/2}$ |
| (g) $(\arctan z)' = (1 + z^2)^{-1}$ | |

Refer to Section 5 of Chapter 1 for the definitions of these trigonometric functions.

Use the rules for differentiation, formulas (2) to (6), to find the derivative of each of the functions in Exercises 2 to 7.

- | | | | |
|--|--------------------|--------------------|-----------------------|
| 2. $z^2 + 10z$ | 3. $\exp(z^3 - z)$ | 4. $[\cos(z^2)]^3$ | 5. $(z^3 + 100)^{-4}$ |
| 6. $(\text{Log } z)^3$ on the plane minus the negative reals | | 7. $\sinh(e^z)$ | |

For each function f listed in Exercises 8 to 11, find an analytic function F with $F' = f$.

8. $f(z) = z - 2$

9. $f(z) = \frac{z^4 + 1}{z^2}$

10. $f(z) = \sin z \cos z$

11. $f(z) = \cosh(2z)$

12. Let f and g be analytic on a domain. Show that (a) $f + g$ is analytic on D and $(f + g)' = f' + g'$; (b) fg is analytic on D and $(fg)' = f'g + fg'$; (c) f/g is analytic at all points z_0 , where $g(z_0) \neq 0$ and

$$\left(\frac{f}{g}\right)' = \frac{f'g - fg'}{g^2}.$$

13. Show that if f is analytic on a domain D and g is analytic on a domain Ω containing the range of f , then $g(f(z))$ is analytic on D , and the chain rule holds: $(g(f(z)))' = g'(f(z))f'(z)$.
14. Let $P(z) = A(z - z_1)\dots(z - z_n)$, where A and z_1, \dots, z_n are complex numbers and $A \neq 0$. Show that

$$\frac{P'(z)}{P(z)} = \sum_{j=1}^n \frac{1}{z - z_j}, \quad z \neq z_1, \dots, z_n.$$

15. Let f be analytic on a domain D and suppose that $f'(z) = 0$ for all $z \in D$. Show that f is constant on D .
16. Find the derivative of the linear fractional transformation $T(z) = (az + b)/(cz + d)$, $ad \neq bc$. In what way does the condition $ad - bc \neq 0$ enter? Conclude that $T'(z)$ is never zero, $z \neq -d/c$.
17. Suppose that f is analytic on a domain D and $f'(z) = \alpha f(z)$, $z \in D$, where α is a constant. Show that $f(z) = C \exp(\alpha z)$, C a constant. (Hint: Consider $g(z) = e^{-\alpha z}f(z)$ and use Exercise 15 for g .)
18. Show that $h(z) = \bar{z}$ is not analytic on any domain. (Hint: Check the Cauchy–Riemann equations.)
19. Fill in the details to make the following argument correct: Let $w = g(z) = \text{Log } z$ and $w_0 = \text{Log } z_0$. Then

$$g'(z_0) = \lim_{z \rightarrow z_0} \frac{\text{Log } z - \text{Log } z_0}{z - z_0} = \lim_{z \rightarrow z_0} \frac{w - w_0}{z - z_0}$$

$$= \lim_{z \rightarrow z_0} \frac{w - w_0}{e^w - e^{w_0}} = \frac{1}{(e^w)'}, \quad \text{at } w = w_0$$

$$= \frac{1}{e^{w_0}} = \frac{1}{z_0}.$$

20. Let $f = u + iv$ be analytic. In each of the following, find v given u .

(a) $u = x^2 - y^2$

(b) $u = \frac{x}{x^2 + y^2}$

(c) $u = 2x^2 + 2x + 1 - 2y^2$

(d) $u = \cosh y \sin x$

(e) $u = \cosh x \cos y$

21. Let γ be a piecewise smooth simple closed curve, and suppose that F is analytic on some domain containing γ . Show that

$$\int_{\gamma} F'(z) dz = 0.$$

22. Use the conclusion of Exercise 21 and Example 13 of Section 6, Chapter 1, to prove that $f(z) = \log z$ cannot be analytic on any domain D that contains a piecewise smooth simple closed curve γ that surrounds the origin. (Hint: What is the value of $\int_{\gamma} f'(z) dz$?)
23. Show that if f is analytic on a domain D and if the range of f lies in either a straight line or a circle, then f is a constant.
24. If $z(t)$ is a differentiable function of the real variable t , $a \leq t \leq b$, and if g is an analytic function on a domain D that contains the range of $z(t)$, $a \leq t \leq b$, show that $g(z(t)) = w(t)$ is a differentiable function of t and

$$w'(t) = g'(z(t))z'(t).$$

Cauchy–Riemann Equations in Polar Coordinates*

25. Suppose $f = u + iv$ is analytic in a domain D . Show that the Cauchy–Riemann equations in polar coordinates are

$$r \frac{\partial u}{\partial r} = \frac{\partial v}{\partial \theta} \quad \text{and} \quad r \frac{\partial v}{\partial r} = -\frac{\partial u}{\partial \theta}.$$

26. Suppose that γ is a piecewise smooth simple closed curve and u is a continuous function on γ . Let D be a domain disjoint from γ , and define a function h on D by the rule

$$h(z) = \int_{\gamma} \frac{u(\zeta)}{\zeta - z} d\zeta, \quad z \in D.$$

Show that h is analytic on D .

2.1.1 Flows, Fields, and Analytic Functions*

There are two physical phenomena, each of which provides a physical setting that helps us to understand analytic functions. An analysis shows that these phenomena are really the same and that they differ only in terminology. However, the terminology is now fixed by tradition. Thus, it is worthwhile to begin by describing each of these situations separately; later we shall concentrate on just one of them.

Flows

Imagine a thin layer of incompressible liquid flowing smoothly across a domain D in the complex plane (Fig. 2.1). At each point $z \in D$, we can find the direction and speed of the liquid, which we assume does not vary with time; equivalently, we watch the path taken by a particle in the liquid and we find the tangent vector to this path. This correspondence between points of D and the value of the velocity (speed + direction) of the liquid is nothing more than a complex-valued function $f(z) = u(z) + iv(z)$, $z \in D$. Conversely, each complex-valued function f gives a flow on D by assigning to the point $z \in D$ the value $f(z)$ as the velocity of the flow at that point z . Hence, each complex-valued function on the domain D corresponds to a (time-independent) flow on D , and vice versa.





Figure 2.1

Assume now that the function f is continuous on D . Let γ be a smooth curve in D ; we examine a small segment of γ . If the segment is sufficiently small, it can be thought of as a straight line segment, and the flow is virtually constant near the segment. The flow f has a component of magnitude $f \cdot n$ perpendicular (or normal) to the segment, so the amount of the flow that crosses this small segment in a unit time is just $f \cdot n ds$, where ds is the length of the small segment. Hence, $\int_\gamma f \cdot n ds$ is the total amount of the flow that crosses γ within a unit time. Likewise, the magnitude of the component of the flow that is parallel to the small segment is $f \cdot \tau$, where τ is a unit vector tangent to (or parallel with) the small segment, so the term $f \cdot \tau ds$ represents the amount of the flow that is tangential to the small segment. Thus, $\int_\gamma f \cdot \tau ds$ is the total amount of the flow that is tangential to γ .



A flow $f = u + iv$ is termed **sourceless** in D if

$$\int_{\gamma} (f \cdot n) ds = 0, \quad ds = \text{arc length},$$

for all smooth simple closed curves γ in D , where $f \cdot n$ is the component of f normal to γ . Thus, for example, there is no *net* flow of the fluid in or out of any disc in D .

A flow $f = u + iv$ is termed **irrotational** in D if

$$\int_{\gamma} (f \cdot \tau) ds = 0, \quad ds = \text{arc length},$$

for all smooth simple closed curves γ in D , where $f \cdot \tau$ is the component of f tangential to γ . Thus, if the flow is irrotational, there is no net rotation of the fluid about any point of D .

Fields

In physics, a complex-valued function f on a domain D is thought of as representing a **field**. The term **vector-field** is also frequently used because the value $f(z)$ can be viewed as a vector whose length $|f(z)|$ gives the strength of the field at z and whose direction $\arg f(z)$ gives the orientation of the field at z . For instance, f could be the force field induced by the gravitational attraction of a collection of masses, or f could be the electrostatic field induced by a collection of charges. We assume, as with flows, that the field is in **steady state**.

Let γ be a smooth curve in D . The **work** done by the field in moving a particle along the curve γ is

$$\int_{\gamma} f \cdot \tau ds,$$

where $f \cdot \tau$ is the component of f tangent to the curve and ds is arc length along the curve. The field is termed **conservative** if the net work done when moving a particle around any closed curve is zero; that is, when

$$0 = \int_{\gamma} f \cdot \tau ds$$

for each smooth closed curve γ in D .

Again let γ be a smooth curve in D ; the **flux** of f across γ is defined to be

$$\int_{\gamma} f \cdot n ds,$$

where $f \cdot n$ is the component of f normal to γ and ds is arc length. The field f is

termed **fluxless** if the net flux of f across any smooth closed curve γ in D is zero; that is, if $\int_{\gamma} f \cdot n \, ds = 0$ for any smooth closed curve γ in D .

The reader will have noticed by now that there is a difference only in terminology between fields and flows: what is termed “irrotational” for flows is called “conservative” for fields and what is termed “sourceless” for flows is called “fluxless” for fields. For this reason, we will discuss only flows in the remainder of this section.

We now explore the relation between a sourceless/irrotational flow f and the analyticity of f . We shall assume that $f = u + iv$, where u and v have continuous partial derivatives throughout D with respect to both x and y .

Let $\gamma(t) = x(t) + iy(t)$, $a \leq t \leq b$, be a smooth curve in D ; then

$$ds = \sqrt{(x'(t))^2 + (y'(t))^2} \, dt$$

and

$$f \cdot \tau = \frac{ux' + vy'}{\sqrt{(x')^2 + (y')^2}},$$

since a unit tangent vector is $\frac{x' + iy'}{\sqrt{(x')^2 + (y')^2}}$.

Thus,

$$\int_{\gamma} f \cdot \tau \, ds = \int_{\gamma} u \, dx + v \, dy = \operatorname{Re} \left\{ \int_{\gamma} \overline{f(z)} \, dz \right\};$$

see Exercise 10, Section 6, Chapter 1. Consequently, the flow f is irrotational if and only if

$$0 = \operatorname{Re} \left\{ \int_{\gamma} \overline{f(z)} \, dz \right\}$$

for all smooth closed curves γ in D .

Just as above, we find that

$$\int_{\gamma} f \cdot n \, ds = \int_{\gamma} u \, dy - v \, dx = \operatorname{Im} \left\{ \int_{\gamma} \overline{f(z)} \, dz \right\}.$$

Hence, the flow f is sourceless exactly when

$$0 = \operatorname{Im} \left\{ \int_{\gamma} \overline{f(z)} \, dz \right\}$$

for all smooth closed curves γ in D .

Let γ be a small circle in D whose inside Ω also lies in D . Green's Theorem then gives

$$\iint_{\Omega} \left[\frac{\partial v}{\partial x} - \frac{\partial u}{\partial y} \right] dx dy = \int_{\gamma} u dx + v dy$$

and

$$\iint_{\Omega} \left[\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} \right] dx dy = \int_{\gamma} u dy - v dx.$$

Thus, for a flow f , which is either irrotational or sourceless, either the first or the second of these area integrals must be zero. By letting γ shrink down to a point, we conclude that

if a flow $f = u + iv$ is irrotational on D , then $\partial v / \partial x - \partial u / \partial y = 0$ on D ,

and

if a flow $f = u + iv$ is sourceless on D , then $\partial u / \partial x + \partial v / \partial y = 0$ on D .

Consequently, if a flow $f = u + iv$ is both sourceless and irrotational, it must satisfy

$$\frac{\partial u}{\partial x} = -\frac{\partial v}{\partial y} \quad \text{and} \quad \frac{\partial u}{\partial y} = \frac{\partial v}{\partial x}.$$

But these are precisely the Cauchy–Riemann equations for the function $\bar{f} = u - iv$, the complex conjugate of $f = u + iv$. In other words,

when a flow f is both sourceless and irrotational on D , \bar{f} must be analytic on D .

In fact, something else has actually been shown too:

A flow f on a domain D is both sourceless and irrotational on D if and only if

$$\int_{\gamma} \overline{f(z)} dz = 0$$

for all smooth closed curves γ in D .

On the other hand, the use of Green's Theorem as above shows that if \bar{f} is analytic on D , then the flow f is **locally sourceless** and **locally irrotational**. That is, each point of D is the center of some disc Ω in D , and in Ω the flow is sourceless and irrotational. Note, however, that a flow can be locally sourceless/irrotational in D but not globally sourceless/irrotational in D ; see Example 2.

Example 1 The function $f(z) = \bar{z} = x - iy$ gives a sourceless and irrotational flow on any domain. Let us find the path followed by a particle in this flow. If $x(t) + iy(t)$

is a parametrization of the path, the tangent to the path is just $dx/dt + i(dy/dt)$, so

$$\frac{dx}{dt} + i \frac{dy}{dt} = \bar{z} = x(t) - iy(t).$$

Hence $dx/dt = x$, $dy/dt = -y$, and consequently $x(t) = c_1 e^t$, $y(t) = c_2 e^{-t}$ for constants c_1 and c_2 . In particular, $xy = c_1 c_2 = \text{constant}$. The curves $xy = \text{constant}$ are thus the paths traced out by the flow. In general a streamline of the flow is the path followed by a particle in the flow. Here, then, the streamlines are the curves $xy = \text{constant}$. The direction of the flow is determined by the signs of c_1 and c_2 (Fig. 2.2). The origin is a **stagnation point** of the flow—once there, it is impossible to leave (of course, nothing actually gets there). \square




Figure 2.2

Example 2 The function $f(z) = 1/\bar{z}$ gives a *locally sourceless* and irrotational flow on any domain D that does not include the origin (where f is not defined). Note, however, that on the circle $|z| = 1$, the normal component of f is

$$f \cdot n = \cos \theta \cos \theta + \sin \theta \sin \theta = 1,$$

and so

$$\int_{|z|=1} f \cdot n \, ds = 1 \int_{|z|=1} \, ds = 2\pi.$$

Thus, the flow $f(z) = 1/\bar{z}$, although locally sourceless, is not globally sourceless (Fig. 2.3). \square

Example 3 The function $f(z) = z$ gives a flow on the plane. Since $\overline{f(z)} = \bar{z}$ is not analytic on any domain (Exercise 18, Section 1), this flow cannot be both sourceless and irrotational. Indeed, $x + iy = z = f(z) = u + iv$, so

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} = 1 \quad \text{and} \quad \frac{\partial u}{\partial y} = \frac{\partial v}{\partial x} = 0.$$




Figure 2.3

Therefore,

$$\int_{\gamma} u \, dx + v \, dy = \iint_{\Omega} \left[\frac{\partial v}{\partial x} - \frac{\partial u}{\partial y} \right] dx \, dy = 0$$

and

$$\int_{\gamma} u \, dy - v \, dx = \iint_{\Omega} \left[\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} \right] dx \, dy = 2 \text{ area}(\Omega).$$

That is, the flow is irrotational but definitely not sourceless. If you are interested, it is challenging to discover the “sources” of this particular flow. \square

The ideas of this section will prove useful later in this chapter, after some more techniques have been developed, and again in Chapters 3 and 4, in the context of conformal mapping.

EXERCISES FOR SECTION 2.1.1

In Exercises 1 to 8, decide whether or not the given function represents a locally sourceless and/or irrotational flow. For those that do, decide whether the flow is globally sourceless and/or irrotational. Sketch some of the streamlines.

1. $x^3 - 3xy^2 + i(y^3 - 3yx^2)$
2. $\frac{x}{x^2 + y^2} + i\frac{y}{x^2 + y^2}, \quad x^2 + y^2 > 0$
3. $\cos x \cosh y + i \sin x \sinh y$
4. $x^2 - y^2 + 2ixy$
5. $x^2 - y^2 - 2ixy$
6. $e^y \cos x + ie^y \sin x$
7. $e^x \cos y + ie^x \sin y$
8. $\cosh x \cos y + i \sinh x \sin y$
9. Suppose that G is analytic on a domain D and $f(z) = \overline{G'(z)}$. Show that f represents a sourceless/irrotational flow on D .

2.2 Power Series

This section is devoted to examining functions of a very special nature: power series. These functions represent a fusion of the ideas on infinite series introduced in Section 4 of Chapter 1 with the concept of an analytic function introduced and studied in Section 1 of this chapter. In Section 4 of this chapter, we shall see that there is a full and complete relationship between analytic functions and power series.

A **power series** in z is an infinite series of the special form

$$\sum_{n=0}^{\infty} a_n(z - z_0)^n, \quad (1)$$

where a_0, a_1, \dots are complex numbers, called the **coefficients** of the series; z_0 is fixed and is called the **center** of the series. One example of such a series has already appeared in Section 4 of Chapter 1, namely,

$$f(z) = \sum_{n=0}^{\infty} z^n = \frac{1}{1-z}, \quad |z| < 1.$$

Of course, any polynomial,

$$p(z) = \sum_{n=0}^{N} b_n z^n,$$

is also an example of such a series. Other examples are the power series

$$\sum_{n=0}^{\infty} \frac{z^n}{n!}, \quad \sum_{n=0}^{\infty} (-1)^n \frac{z^{2n}}{(2n)!}, \quad \sum_{n=0}^{\infty} (-1)^n \frac{z^{2n+1}}{(2n+1)!},$$

all three of which converge for all z by use of the ratio test (see Section 4 of Chapter 1).

The first major result on power series is as follows.

THEOREM 1 Suppose there is some $z_1 \neq z_0$ such that $\sum a_n(z_1 - z_0)^n$ converges. Then for each z with $|z - z_0| < |z_1 - z_0|$, the series $\sum a_n(z - z_0)^n$ is absolutely convergent.

Proof Suppose that $|z - z_0| \leq r < |z_1 - z_0|$. Since $\sum_0^{\infty} a_n(z_1 - z_0)^n$ is convergent, we know that

$$\lim_{n \rightarrow \infty} a_n(z_1 - z_0)^n = 0.$$

In particular, there is a constant M with

$$|a_n| |z_1 - z_0|^n \leq M$$

for all n . Hence,

$$|a_n|(z - z_0)^n = |a_n|(z_1 - z_0)^n \left(\frac{|z - z_0|}{|z_1 - z_0|} \right)^n \\ \leq M\rho^n,$$

where $\rho = r/|z_1 - z_0|$ is less than 1. Hence, the series $\sum a_n(z - z_0)^n$ is absolutely convergent. ■

The Radius of Convergence

For any power series $\sum_0^\infty a_n(z - z_0)^n$, there are always three mutually exclusive possibilities:

The series $\sum a_n(z - z_0)^n$ converges only for $z = z_0$. (2)

The series $\sum a_n(z - z_0)^n$ converges for all z . (3)

The series $\sum a_n(z - z_0)^n$ converges for some $z \neq z_0$, but not for all z . (4)

Let us examine the third possibility (4) more thoroughly. Suppose z' and z'' are two points with

$$\sum_{n=0}^{\infty} a_n(z' - z_0)^n \text{ convergent}, \quad \sum_{n=0}^{\infty} a_n(z'' - z_0)^n \text{ divergent}.$$

By Theorem 1,

$$|z - z_0| < |z' - z_0| \quad \text{implies that} \quad \sum_0^\infty a_n(z - z_0)^n \text{ converges},$$

and

$$|z - z_0| > |z'' - z_0| \quad \text{implies that} \quad \sum_0^\infty a_n(z - z_0)^n \text{ diverges}.$$

In particular, $|z' - z_0| < |z'' - z_0|$. That is, any point of convergence of the series lies no farther from z_0 than any point of divergence of the series. Continuing in this fashion to examine the convergence of $\sum_0^\infty a_n(z - z_0)^n$, a number R can be defined by this rule: R is the unique number such that

$$|z - z_0| < R \quad \text{implies that} \quad \sum_0^\infty a_n(z - z_0)^n \text{ converges},$$

and

(5)

$$|z - z_0| > R \quad \text{implies that} \quad \sum_0^{\infty} a_n(z - z_0)^n \text{ diverges.}$$

That is, R is the largest number with the property that whenever $|z - z_0| < R$, the series $\sum a_n(z - z_0)^n$ converges. When $|z - z_0| = R$, the series $\sum_0^{\infty} a_n(z - z_0)^n$ may converge or may diverge. The number R is the **radius of convergence** of the power series $\sum a_n(z - z_0)^n$. To make the picture complete, if the series satisfies (2), we define $R = 0$, and if (3) holds, we define $R = \infty$. To reiterate, a power series $\sum_0^{\infty} a_n(z - z_0)^n$, which converges for some $z \neq z_0$, always converges within a disc $\{z : |z - z_0| < R\}$, where R is positive or R is infinity.

The radius of convergence R of the power series $\sum_0^{\infty} a_n(z - z_0)^n$ depends in a very explicit way on the coefficients $\{a_n\}$. The general formula is a bit complicated (see Exercises 28 and 29), but two simple and useful cases are as follows.

THEOREM 2 Suppose that $\sum_0^{\infty} a_n(z - z_0)^n$ is a power series with a positive or infinite radius of convergence R .

(a) If $\lim_{n \rightarrow \infty} |a_{n+1}/a_n|$ exists, then

$$\frac{1}{R} = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right|. \quad (6)$$

(b) If $\lim_{n \rightarrow \infty} \sqrt[n]{|a_n|}$ exists, then

$$\frac{1}{R} = \lim_{n \rightarrow \infty} \sqrt[n]{|a_n|}. \quad (7)$$

Proof The proofs of (a) and (b) are nothing but the ratio and root tests, respectively. To see why (a) is true, let L be the limit of $|a_{n+1}/a_n|$. Then

$$\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}(z - z_0)^{n+1}}{a_n(z - z_0)^n} \right| = |z - z_0| \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = |z - z_0|L.$$

When $|z - z_0|L$ is less than 1, the series $\sum_0^{\infty} a_n(z - z_0)^n$ is absolutely convergent; when $|z - z_0|L$ is greater than 1, the series $\sum a_n(z - z_0)^n$ diverges; see Exercise 43, Section 4 of Chapter 1. By the definition of R , we must have $R = 1/L$ or $1/R = L$.

The proof of (b) is virtually the same, with the root test replacing the ratio test; see Exercise 42 of Section 4 of Chapter 1. ■

Example 1 The power series $\sum_0^{\infty} 5^n(z - 1)^n$ has radius of convergence $R = \frac{1}{5}$, since

$$\frac{1}{R} = \lim_{n \rightarrow \infty} \sqrt[n]{5^n} = 5.$$

Hence, the series converges within the disc $|z - 1| < \frac{1}{3}$ and diverges for all z with $|z - 1| > \frac{1}{3}$. \square

Example 2 The power series $\sum_0^{\infty} (-1)^n z^n / n!$ converges for all z , since

$$\lim_{n \rightarrow \infty} \frac{|a_{n+1}|}{|a_n|} = \lim_{n \rightarrow \infty} \frac{|(-1)^{n+1}/(n+1)!|}{|(-1)^n/n!|} = \lim_{n \rightarrow \infty} \frac{1}{n+1} = 0,$$

so $1/R = 0$, which means that $R = \infty$. \square

Example 3 Find the radius of convergence of the power series

$$\sum_{n=0}^{\infty} n^2 \frac{1}{2^{2n}} z^n.$$

Solution Use (7):

$$\begin{aligned} \frac{1}{R} &= \lim_{n \rightarrow \infty} \sqrt[n]{\frac{n^2}{2^{2n}}} = \lim_{n \rightarrow \infty} (\sqrt[n]{n})^2 (\sqrt[n]{4^{-n}}) \\ &= (1^2) \left(\frac{1}{4}\right) = \frac{1}{4}, \end{aligned}$$

because $\lim_{n \rightarrow \infty} \sqrt[n]{n} = 1$ (see Exercise 21(a) of this section). Hence, $R = 4$. \square

Example 4 Find the radius of convergence of the power series $\sum_{n=0}^{\infty} 4^n z^{3n}$.

Solution Here

$$a_k = \begin{cases} 4^{k/3} & \text{if } k = 0, 3, 6, \dots \\ 0 & \text{otherwise,} \end{cases}$$

so technically speaking, neither (6) nor (7) can be applied. However, when we write $w = z^3$, the power series becomes $\sum_0^{\infty} 4^n w^n$. This power series has radius of convergence $\frac{1}{4}$; that is, the given series

$$\begin{aligned} \text{converges if } \frac{1}{4} &> |w| = |z|^3 \\ \text{diverges if } \frac{1}{4} &< |w| = |z|^3. \end{aligned}$$

Hence, the first series $\sum_{n=0}^{\infty} 4^n z^{3n}$ has radius of convergence $R = 4^{-1/3}$. \square

Example 5 Here is another example that shows the limitations of formulas (6) and (7). Let f be the power series

$$f(z) = \sum_{n=0}^{\infty} 4^{n(-1)^n} z^n.$$

That is,

$$a_n = \begin{cases} 4^n & \text{if } n = 0, 2, 4, \dots \\ \frac{1}{4^n} & \text{if } n = 1, 3, 5, \dots \end{cases}$$

The two power series

$$f_1(z) = \sum_{n=0}^{\infty} 4^{2n} z^{2n}$$

and

$$f_2(z) = \sum_{n=0}^{\infty} \frac{1}{4^{2n+1}} z^{2n+1}$$

have radii of convergence $R_1 = \frac{1}{4}$ and $R_2 = 4$, respectively, and

$$f(z) = f_1(z) + f_2(z) \quad \text{if} \quad |z| < \frac{1}{4}.$$

Hence, the radius of convergence of f is at least $\frac{1}{4}$. But it cannot be bigger than $\frac{1}{4}$, since the series for f_1 diverges whenever $|z| > \frac{1}{4}$. Hence, $R = \frac{1}{4}$. But note that neither (6) nor (7) was available to reach this conclusion. The general formula for R , which does cover this case, depends on some special knowledge from real analysis and is presented in the exercises. \square

The Derivative of a Power Series

We can now begin the chief task of this section: to show that a function f given by a power series

$$f(z) = \sum_{n=0}^{\infty} a_n (z - z_0)^n \tag{8}$$

with a positive or infinite radius of convergence R is actually *analytic* within the disc $|z - z_0| < R$, and its derivative is given by

$$f'(z) = \sum_{n=1}^{\infty} n a_n (z - z_0)^{n-1}. \tag{9}$$

We begin by showing that the power series on the right side of (9) has radius of convergence at least R . Let $|z - z_0| = r < R$, and let s be between r and R , $r < s < R$. Then, for all integers n greater than some N ,

$$nr^{n-1} \leq s^n, \quad n \geq N, \quad (10)$$

since $\lim_{n \rightarrow \infty} n(r/s)^n = 0$; see Exercise 21(b) at the end of this section. Thus,

$$n|a_n|(z - z_0)^{n-1} \leq |a_n|s^n, \quad n \geq N,$$

and since $s < R$, the series $\sum_0^\infty |a_n|s^n$ converges. Hence, the series $\sum n a_n(z - z_0)^{n-1}$ converges whenever $|z - z_0| = r < R$.

Let g be the series on the right side of (9):

$$g(z) = \sum_{n=1}^{\infty} n a_n (z - z_0)^{n-1}.$$

It simplifies the notation to assume $z_0 = 0$; this can be done with no loss of generality. Suppose z is in the disc of radius R centered at 0; let $\delta = \frac{1}{2}(R - |z|)$, and suppose that $|h| < \delta$. Now compute

$$\begin{aligned} f(z + h) - f(z) &= \sum_1^{\infty} a_n [(z + h)^n - z^n] \\ &= a_1 h + \sum_2^{\infty} a_n [(z + h)^n - z^n]. \end{aligned}$$

This gives

$$\frac{f(z + h) - f(z)}{h} - g(z) = \sum_2^{\infty} a_n \left\{ \frac{(z + h)^n - z^n}{h} - nz^{n-1} \right\}.$$

Now the binomial theorem (Exercise 20, this section) gives

$$\frac{(z + h)^n - z^n}{h} - nz^{n-1} = \sum_{j=2}^n \binom{n}{j} z^{n-j} h^{j-1},$$

where $\binom{n}{j} = \frac{n!}{j!(n-j)!}$. Hence,

$$\left| \frac{(z + h)^n - z^n}{h} - nz^{n-1} \right| \leq \sum_{j=2}^n \binom{n}{j} |z|^{n-j} |h|^{j-1}$$

$$\leq |h| \sum_{j=2}^n \binom{n}{j} (R - 2\delta)^{n-j} \delta^{j-2}$$

$$\begin{aligned}
&= |h|\delta^{-2} \sum_{j=2}^n \binom{n}{j} (R - 2\delta)^{n-j} \delta^j \\
&< |h|\delta^{-2} \sum_{j=0}^n \binom{n}{j} (R - 2\delta)^{n-j} \delta^j \\
&= |h|\delta^{-2} [(R - 2\delta) + \delta]^n = |h|\delta^{-2}(R - \delta)^n.
\end{aligned}$$

This leads to the estimate

$$\left| \frac{f(z+h) - f(z)}{h} - g(z) \right| \leq |h|\delta^{-2} \sum_{n=2}^{\infty} |a_n|(R - \delta)^n.$$

This last quantity goes to zero as $h \rightarrow 0$ (recall that δ is fixed), so we have established that f is differentiable with derivative equal to g :

$$\left(\sum_0^{\infty} a_n z^n \right)' = \sum_1^{\infty} n a_n z^{n-1}.$$

This is what we set out to accomplish. However, even more can now be said. The series $\sum_1^{\infty} n a_n z^{n-1}$ is also a power series with radius of convergence R , so it, too, is differentiable in the disc $|z| < R$ with derivative

$$\left(\sum_1^{\infty} n a_n z^{n-1} \right)' = \sum_2^{\infty} n(n-1) a_n z^{n-2}.$$

This process can be continued indefinitely, of course, to obtain the following theorem.

THEOREM 3 If $f(z) = \sum_0^{\infty} a_n(z - z_0)^n$ has a positive or infinite radius of convergence R , then within the disc $|z - z_0| < R$, f is infinitely differentiable, each derivative being again given by a power series

$$f^{(k)}(z) = \sum_{n=k}^{\infty} n(n-1)\dots(n-k+1)a_n(z - z_0)^{n-k}, \quad k = 1, 2, \dots \quad (11)$$

In particular, by setting $z = z_0$, we have

$$\frac{f^{(n)}(z_0)}{n!} = a_n, \quad n = 0, 1, 2, \dots \quad \blacksquare \quad (12)$$

Example 6 The power series

$$\sum_{n=0}^{\infty} \frac{z^n}{n!}$$

has radius of convergence infinity (by the ratio test) and hence is an entire function. According to Theorem 3, its derivative is

$$\begin{aligned} \left(\sum_{n=0}^{\infty} \frac{z^n}{n!} \right)' &= \sum_{n=1}^{\infty} \frac{n}{n!} z^{n-1} = \sum_{n=1}^{\infty} \frac{z^{n-1}}{(n-1)!} \\ &= \sum_{j=0}^{\infty} \frac{z^j}{j!}. \end{aligned}$$

That is, the function is its own derivative. Let us denote the sum of the series by F :

$$F(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!}.$$

Then

$$\begin{aligned} (e^{-z} F(z))' &= -e^{-z} F(z) + e^{-z} F'(z) \\ &= 0, \end{aligned}$$

since $F' = F$. Thus, $F(z)e^{-z} = \lambda$ for some constant λ (by Exercise 15 of Section 1). But $1 = F(0) = \lambda$, so $F(z) = e^z$, and thus,

$$e^z = \sum_{n=0}^{\infty} \frac{z^n}{n!}. \quad \square \quad (13)$$

Example 7 Show that $\sin z$ and $\cos z$ have power series expansions valid for all z ; find the series explicitly.

Solution From Example 6, we know that

$$e^z = \sum_{n=0}^{\infty} \frac{z^n}{n!}.$$

Hence,

$$\begin{aligned} \cos z &= \frac{1}{2}(e^{iz} + e^{-iz}) = \frac{1}{2} \left(\sum_{n=0}^{\infty} \frac{(iz)^n}{n!} + \sum_{n=0}^{\infty} \frac{(-iz)^n}{n!} \right) \\ &= \sum_{n=0}^{\infty} (-1)^n \frac{z^{2n}}{(2n)!}. \end{aligned}$$

Likewise,

$$\sin z = \sum_{n=0}^{\infty} (-1)^n \frac{z^{2n+1}}{(2n+1)!}. \quad \square$$

Multiplication and Division of Powers Series*

Suppose that the two power series $f(z) = \sum_{n=0}^{\infty} a_n z^n$ and $g(z) = \sum_{n=0}^{\infty} b_n z^n$ both have radius of convergence at least R for some positive number R ; the product $f(z)g(z)$ is then defined and analytic in the disc $\{z : |z| < R\}$. Formally multiplying out the series for $f(z)$ and that for $g(z)$ and collecting equal powers of z , we obtain

$$\begin{aligned} f(z)g(z) &= (a_0 + a_1 z + a_2 z^2 + \cdots)(b_0 + b_1 z + b_2 z^2 + \cdots) \\ &= a_0 b_0 + (a_0 b_1 + a_1 b_0)z + (a_0 b_2 + a_1 b_1 + a_2 b_0)z^2 + \cdots \\ &= \sum_{n=0}^{\infty} c_n z^n, \end{aligned}$$

where

$$c_n = \sum_{k=0}^n a_k b_{n-k}.$$

These manipulations are, in fact, correct, and the power series $\sum c_n z^n$ has radius of convergence at least R . The formal statement of this result follows, and the proof is presented at the end of Section 4.

THEOREM 4 Suppose that $f(z) = \sum_{n=0}^{\infty} a_n z^n$ and $g(z) = \sum_{n=0}^{\infty} b_n z^n$ are two power series, each with radius of convergence R or more, $R > 0$. Then their product fg can also be expressed as a power series in the disc $|z| < R$:

$$(fg)(z) = \sum_{n=0}^{\infty} c_n z^n, \quad \text{where } c_n = \sum_{k=0}^n a_k b_{n-k}. \quad \blacksquare$$

Example 8 Find the power series expansion of $e^z/(1 - z)$ about $z_0 = 0$ in $|z| < 1$.

Solution We know that

$$e^z = 1 + z + \frac{z^2}{2!} + \frac{z^3}{3!} + \cdots, \quad \text{all } z$$

and that

$$\frac{1}{1 - z} = 1 + z + z^2 + z^3 + \cdots, \quad |z| < 1.$$

Hence,

$$\frac{e^z}{1 - z} = 1 + 2z + \frac{5}{2}z^2 + \frac{8}{3}z^3 + \frac{65}{24}z^4 + \cdots, \quad |z| < 1. \quad \square$$

Example 9 Find the power series for $(1 - z^2) \sin z$ about $z_0 = 0$.

Solution Again, just formally multiply out the two series

$$(1 - z^2)(\sin z) = (1 - z^2) \left(z - \frac{z^3}{6} + \frac{z^5}{120} - \dots \right)$$

$$= z - \frac{7}{6}z^3 + \frac{21}{120}z^5 - \dots \quad \square$$

Example 10 Later in this chapter, we will show that $z \csc z$ has a power series expansion near $z = 0$. Assuming this for the moment, find the first few terms of this series.

Solution Since $z \csc z = z/\sin z$, we are faced with a problem of division of series. However, every division problem can be replaced by an equivalent multiplication problem, and this is what we will do here. Write

$$z \csc z = b_0 + b_1 z + b_2 z^2 + \dots$$

Then

$$\begin{aligned} z &= (z \csc z) \sin z \\ &= (b_0 + b_1 z + b_2 z^2 + \dots) \left(z - \frac{z^3}{6} + \frac{z^5}{120} - \dots \right) \\ &= b_0 z + b_1 z^2 + \left(b_2 - \frac{b_0}{6} \right) z^3 + \left(b_3 - \frac{b_1}{6} \right) z^4 + \left(b_4 - \frac{b_2}{6} + \frac{b_0}{120} \right) z^5 + \dots \end{aligned}$$

Equate coefficients of equal powers of z and obtain a series of equations for the b_j :

$$1 = b_0, \quad 0 = b_1, \quad 0 = b_2 - \frac{b_0}{6}, \quad 0 = -\frac{b_1}{6} + b_3, \quad 0 = \frac{b_0}{120} - \frac{b_2}{6} + b_4.$$

Hence, $b_0 = 1$, $b_1 = 0$, $b_2 = \frac{1}{6}$, $b_3 = 0$, $b_4 = \frac{7}{360}$. Consequently,

$$z \csc z = 1 + \frac{z^2}{6} + \frac{7}{360}z^4 + \dots \quad \square$$

Note that this technique of dividing series works all the time, provided that it is known that the quotient does actually have a series representation. We shall look further into this issue in Section 5 of this chapter.

EXERCISES FOR SECTION 2.2

In Exercises 1 to 6, use Theorem 2 or Example 4 to find the radius of convergence of the given power series.

1. $\sum_{k=1}^{\infty} k(z-1)^k$

2. $\sum_{k=0}^{\infty} \frac{(k!)^2}{(2k)!}(z-2)^k$

3. $\sum_{j=0}^{\infty} \frac{z^{3j}}{2^j}$

4. $\sum_{k=0}^{\infty} (-1)^k z^{2k}$

5. $\sum_{n=0}^{\infty} 5^{(-1)^n} z^n$

6. $\sum_{k=1}^{\infty} \frac{(2k)(2k-2)\dots4\cdot2}{(2k-1)(2k-3)\dots3\cdot1} z^k$

In Exercises 7 to 13, find the power series about the origin for the given function.

7. e^{-z}

8. $z^2 \cos z$

9. $\frac{z^3}{1-z^3}, \quad |z| < 1$

10. $\frac{1+z}{1-z}, \quad |z| < 1$

11. $\frac{z^2}{(4-z)^2}, \quad |z| < 4$ **(Hint:** $\frac{1}{(a-z)^2} = \frac{d}{dz}[(a-z)^{-1}]$.)

12. $\sinh(z^2)$

13. $(\sin z)^2$

In Exercises 14 to 18, find a “closed form” (that is, a simple expression) for each of the given power series.

14. $\sum_{n=0}^{\infty} \frac{z^{2n}}{n!}$

15. $\sum_{n=1}^{\infty} z^{3n}$

16. $\sum_{n=1}^{\infty} n(z-1)^{n-1}$

17. $\sum_{n=0}^{\infty} (-1)^n \frac{(z-2\pi i)^n}{n!}$

18. $\sum_{n=2}^{\infty} n(n-1)z^n$ **(Hint:** Divide by z^2 .)

19. (a) Suppose that the radius of convergence of the power series $f(z) = \sum_{n=0}^{\infty} a_n(z-z_0)^n$ is R , $R > 0$. Show that the radius of convergence of

$$F(z) = \sum_{n=0}^{\infty} \frac{a_n}{n+1} (z-z_0)^{n+1}$$

is at least R .

- (b) Show that $F'(z) = f(z)$ for all z with $|z-z_0| < R$.
 (c) Use parts (a) and (b) to show that

$$\text{Log}(1-z) = -z - \frac{z^2}{2} - \frac{z^3}{3} - \frac{z^4}{4} - \cdots, \quad \text{if } |z| < 1.$$

20. Establish the **binomial formula** for any pair (a, b) of complex numbers

$$(a+b)^n = \sum_{j=0}^n \binom{n}{j} a^{n-j} b^j, \quad \binom{n}{j} = \frac{n!}{j!(n-j)!}.$$

(Hint: One technique is to prove the formula by induction on n .)

21. Show that (a) $\lim_{n \rightarrow \infty} \sqrt[n]{n} = 1$; (b) $\lim_{n \rightarrow \infty} n\rho^n = 0$ if $|\rho| < 1$;
 (c) $\lim_{n \rightarrow \infty} \sqrt[n]{M} = 1$ if M is positive.

(Hint: For (a), write

$$\sqrt[n]{n} = 1 + \varepsilon_n, \quad \varepsilon_n > 0; \quad \text{then } n = (1 + \varepsilon_n)^n > 1 + \frac{n(n-1)}{2}\varepsilon_n^2,$$

by Exercise 20.)

22. (a) If $f(z) = \sum_{n=0}^{\infty} a_n(z - z_0)^n$ has radius of convergence $R > 0$ and if $f(z) = 0$ for all z , $|z - z_0| < r \leq R$, show that $a_0 = a_1 = \dots = 0$.
 (b) If $F(z) = \sum_0^{\infty} a_n(z - z_0)^n$ and $G(z) = \sum_0^{\infty} b_n(z - z_0)^n$ are equal on some disc $|z - z_0| < r$, show that $a_n = b_n$ for all n .


23. Let a_0, a_1, a_2, \dots be complex numbers with either

$$\lim_{n \rightarrow \infty} \sqrt[n]{|a_n|} = L \quad \text{or} \quad \lim_{n \rightarrow \infty} \frac{|a_{n+1}|}{|a_n|} = L.$$

Show that the series

$$\sum_{n=0}^{\infty} \frac{a_n}{(z - z_0)^n}$$

converges absolutely for all z with $|z - z_0| > L$ and diverges for all z with $|z - z_0| < L$.



24. Let the radius of convergence of $f(z) = \sum_{n=0}^{\infty} a_n(z - z_0)^n$ be R , $0 < R \leq \infty$, and let $r < R$. Set

$$f_n(z) = \sum_{i=0}^n a_i(z - z_0)^i, \quad n = 1, 2, \dots$$

Given $\varepsilon > 0$, show that there is an N such that $|f_n(z) - f(z)| < \varepsilon$ for all z with $|z - z_0| \leq r$ and $n \geq N$.

(Hint: The series $\sum_{j=0}^{\infty} |a_j|r^j$ converges, so there is an N such that

$$\sum_N^{\infty} |a_j|r^j < \varepsilon.$$

Hence, for $n \geq N$ and $|z - z_0| \leq r$,

$$\begin{aligned} |f_n(z) - f(z)| &= \left| \sum_{n+1}^{\infty} a_j(z - z_0)^j \right| \leq \sum_{n+1}^{\infty} |a_j|r^j \\ &\leq \sum_N^{\infty} |a_j|r^j < \varepsilon. \end{aligned}$$

The Binomial Theorem for Real Exponents*

25. Suppose that α is a real number but not a nonnegative integer. Define $\binom{\alpha}{j}$ for each nonnegative integer j by

$$\binom{\alpha}{0} = 1, \quad \binom{\alpha}{1} = \alpha,$$

and in general,

$$\binom{\alpha}{j} = \frac{\alpha(\alpha - 1)\dots(\alpha - j + 1)}{j!}, \quad j = 1, 2, \dots$$

Show that $\frac{\left| \binom{\alpha}{j+1} \right|}{\left| \binom{\alpha}{j} \right|} \rightarrow 1$ as $j \rightarrow \infty$.

26. Show that the radius of convergence of the series

$$F(z) = \sum_{k=0}^{\infty} \binom{\alpha}{k} z^k$$

is 1.

27. (a) Show that the function F defined in Exercise 26 satisfies the differential equation

$$(1 + z)F'(z) = \alpha F(z).$$

(Hint: Compute F' ; then multiply it by $(1 + z)$ and add terms with equal powers of z .)

(b) Conclude from (a) that $F(z) = (1 + z)^\alpha$; that is,

$$(1 + z)^\alpha = \sum_{k=0}^{\infty} \binom{\alpha}{k} z^k, \quad |z| < 1.$$

General Formula for the Radius of Convergence of a Power Series*

Let $\{x_n\}$ be a bounded sequence of nonnegative real numbers. Define

$$\limsup_{n \rightarrow \infty} x_n = \rho$$

to be the *largest* number ρ such that *each* interval $(\rho - \varepsilon, \rho + \varepsilon)$ contains x_n for infinitely many indices n . (The fact that such a ρ exists is a fundamental property of the real numbers.)

If $\{x_n\}$ is unbounded, define $(\limsup)_{n \rightarrow \infty} x_n$ to be infinity.

28. (a) Show that there are integers $\{n_j\}_{j=1}^{\infty}$ increasing to ∞ such that

$$\lim_{j \rightarrow \infty} x_{n_j} = \rho.$$

(b) If $\{n_k\}_{k=1}^{\infty}$ is a sequence of integers increasing to ∞ and if $\lim_{k \rightarrow \infty} x_{n_k} = \sigma$, show that $\sigma \leq \rho$.

29. Let $\sum_0^{\infty} a_n(z - z_0)^n$ be a power series with radius of convergence R . Show that

$$\frac{1}{R} = \limsup_{n \rightarrow \infty} \sqrt[n]{|a_n|}.$$

Outline: Let $L = \limsup \sqrt[n]{|a_n|}$. Given $\varepsilon > 0$, there is an N such that $\sqrt[n]{|a_n|} < L + \varepsilon$ for all $n \geq N$. (Why?) Suppose that $|z - z_0| < (1 - \varepsilon)/(L + \varepsilon)$; then $|a_n| |z - z_0|^n < (1 - \varepsilon)^n$ for all $n \geq N$, so $\sum_0^{\infty} a_n(z - z_0)^n$ is absolutely convergent. Hence, $R \geq (1 - \varepsilon)/(L + \varepsilon)$ for each $\varepsilon > 0$. (Why?) Therefore, $R \geq 1/L$. Conversely, suppose that $|z - z_0| > 1/(L - \varepsilon)$ for some $\varepsilon > 0$. There are infinitely many indices n with $\sqrt[n]{|a_n|} > L - \varepsilon$. (Why?) Hence, $|a_n| |z - z_0|^n > 1$ for infinitely many indices n , so $\sum_0^{\infty} a_n(z - z_0)^n$ diverges. (Why?) Hence, $R \leq 1/(L - \varepsilon)$ for each $\varepsilon > 0$, so $R \leq 1/L$. Hence, $R = 1/L$.

2.3 Cauchy's Theorem and Cauchy's Formula

Cauchy's Theorem is the linchpin of complex variables. We shall first prove the result with the added assumption that f' is continuous. As it turns out, an analytic function automatically has a continuous derivative. However, it is somewhat technical to prove this. For that reason, we isolate most of this work as a special starred subsection 2.3.1; the final step is Theorem 1 of Section 4. Throughout this section, we shall assume that f' is continuous.

THEOREM 1 Cauchy's Theorem Suppose that f is analytic on a domain D . Let γ be a piecewise smooth simple closed curve in D whose inside Ω also lies in D . Then

$$\int_{\gamma} f(z) dz = 0.$$

Proof Recall that we will assume initially that f' is continuous. We know from Green's Theorem (Section 6 of Chapter 1) that

$$\int_{\gamma} f(z) dz = i \iint_{\Omega} \left\{ \frac{\partial f}{\partial x} + i \frac{\partial f}{\partial y} \right\} dx dy.$$

However, because $f = u + iv$ is analytic, u and v satisfy the Cauchy–Riemann equations. Hence,

$$\begin{aligned} \frac{\partial f}{\partial x} + i \frac{\partial f}{\partial y} &= \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} + i \left[\frac{\partial u}{\partial y} + i \frac{\partial v}{\partial y} \right] \\ &= \frac{\partial u}{\partial x} - \frac{\partial v}{\partial y} + i \left[\frac{\partial v}{\partial x} + \frac{\partial u}{\partial y} \right] \\ &= 0 + i(0) = 0. \end{aligned}$$
■

The incredible utility of Cauchy's Theorem is due in part to its great generality and in part to the fact that it can be conjoined with an assumption about the geometry of the underlying domain to yield additional and somewhat unexpected results. In what follows, we pursue the latter path.

DEFINITION A domain D is **simply-connected** if, whenever γ is a simple closed curve in D , the inside of γ is also a subset of D .

A less formal way of saying that D is simply-connected is to say that “ D has no holes in it.” Several examples will help make the concept more concrete.

Example 1 The disc $\{z: |z - z_0| < R\}$ is simply-connected, as is the horizontal strip $\{z: a < \operatorname{Im} z < b\}$. □

Example 2 Any convex domain Ω is simply-connected. □

Example 3 Neither the annulus $\{z: 0 < r_1 < |z - z_0| < r_2\}$ nor the punctured disc $\{z: 0 < |z - z_0| < r_2\}$ is simply-connected. This follows from the fact that for each r , $r_1 < r < r_2$, the circle $|z - z_0| = r$ lies in the annulus and the punctured disc, but the inside of this circle, which is the disc $\{z: |z - z_0| < r\}$, is not a subset of either. □

Example 4 The domain obtained by deleting the line segment $0 \leq x \leq 1$ from the open disc $\{z: |z| < 1\}$ is simply-connected, while the domain obtained by deleting the segment $0 \leq x \leq \frac{1}{2}$ is not simply-connected (see Fig. 2.4a and 2.4b). \square

Example 5 The domain obtained from the annulus $\{z: r < |z| < R\}$ by deleting those x with $r < x < R$ is simply-connected (Fig. 2.4c). \square




Figure 2.4

Our results connecting Cauchy's Theorem and simple-connectivity follow below. Recall that we assume f' to be continuous; the results of Section 2.3.1 will show this to be superfluous.

THEOREM 2 Let D be a simply-connected domain and Γ a closed curve in D that is composed of a finite number of horizontal and vertical line segments. If f is analytic in D , then

$$\int_{\Gamma} f(\zeta) d\zeta = 0.$$

This is rather intricate to prove completely, but any picture (for instance, Fig. 2.5) makes it reasonably clear. You can draw several more pictures of such




Figure 2.5

closed curves to explore their other possible configurations. The crux of the matter is that any closed loop in Γ surrounds only points of D , since D is simply-connected. Given that this is the case, repeated applications of Theorem 1 then establish Theorem 2.

We now show another connection between simple-connectivity and analyticity.

THEOREM 3 If f is analytic in a simply-connected domain D , then there is an analytic function F on D with $F' = f$ throughout D .

Proof Fix some point z_0 in D and define F at a point z of D by the rule

$$F(z) = \int_{\gamma} f(\zeta) d\zeta,$$

where γ is a polygonal curve in D joining z_0 to z , which is composed of a finite number of horizontal/vertical line segments. If γ_1 is another such curve, then $\Gamma = \gamma - \gamma_1$ is a closed curve in D made up of horizontal/vertical line segments and consequently (by Theorem 2),

$$0 = \int_{\Gamma} f(w) dw = \int_{\gamma} f(w) dw - \int_{\gamma_1} f(w) dw.$$

Thus, the value of $F(z)$ does not depend on the choice of curve γ from z_0 to z ; in mathematical language, F is *well-defined*. We now need to show that $F' = f$. Let z_1 be a point of D and let r be a number so small that the disc $\{z : |z - z_1| < r\}$ lies in D . Now let $|h| < r$, and let L be the curve joining z_1 to $z_1 + h$ which is composed of one horizontal and one vertical segment (see Fig. 2.6). Note that the total length of L is at most $2|h|$ and that

$$\int_L dz = h.$$




Figure 2.6

Let γ_1 be a polygonal curve composed of horizontal and vertical segments joining z_0 to $z_1 + h$; again refer to Figure 2.6. The curve $\Gamma = \gamma_1 - L - \gamma$ is then a closed polygonal curve consisting of horizontal and vertical segments. Hence, the line integral of f over Γ is zero. Equivalently,

$$\begin{aligned} 0 &= \int_{\gamma_1} f(\zeta) d\zeta - \int_L f(\zeta) d\zeta - \int_\gamma f(\zeta) d\zeta \\ &= F(z_1 + h) - \int_L f(\zeta) d\zeta - F(z_1). \end{aligned}$$

This gives

$$F(z_1 + h) - F(z_1) = \int_L f(\zeta) d\zeta.$$

Consequently,

$$\frac{F(z_1 + h) - F(z_1)}{h} - f(z_1) = \int_L \frac{[f(\zeta) - f(z_1)]}{h} d\zeta.$$

Given $\varepsilon > 0$, choose $\delta > 0$ so small that $|f(\zeta) - f(z_1)| < \varepsilon$ whenever $|\zeta - z_1| < \delta$. This is possible because f is continuous at z_1 . For $|h| < \delta$, inequality (3) in Section 6 of Chapter 1 and the above imply that

$$\left| \frac{F(z_1 + h) - F(z_1)}{h} - f(z_1) \right| \leq \frac{1}{|h|} \{ \max |f(w) - f(z_1)| \} 2|h| < 2\varepsilon.$$

Consequently, the derivative of F at z_1 exists and equals $f(z_1)$. ■

COROLLARY Let f be analytic on a simply-connected domain D , and let γ be a piecewise smooth closed curve in D . Then

$$\int_\gamma f(z) dz = 0.$$

Proof Let F be a function on D with $F' = f$. Then

$$\begin{aligned} \int_\gamma f(z) dz &= \int_\gamma F'(z) dz = F(\text{endpoint}) - F(\text{initial point}) \\ &= 0, \end{aligned}$$

since the curve γ is closed. ■

Cauchy's Formula

THEOREM 4 Cauchy's Formula Suppose that f is analytic on a domain D and that γ is a piecewise smooth, positively oriented simple closed curve in D whose inside Ω also lies in D . Then

$$f(z) = \frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta, \quad \text{for all } z \in \Omega.$$

Proof Since z is inside Ω , there is a positive number δ_0 so small that the disc $\{w: |w - z| < \delta_0\}$ lies within Ω also. Let δ be a positive number smaller than δ_0 , and let Ω_δ be the domain obtained by deleting the disc $\{w: |w - z| \leq \delta\}$ from Ω . The boundary of Ω_δ consists of the two curves γ , oriented positively, and the circle $\{w: |w - z| = \delta\}$, oriented negatively. We may apply Green's Theorem to the function

$$g(\zeta) = \frac{f(\zeta)}{\zeta - z}$$

on the region Ω_δ . Because g is analytic on Ω_δ , it follows (as in the proof of Theorem 1) that

$$\frac{\partial g}{\partial x} + i \frac{\partial g}{\partial y} = 0$$

throughout Ω_δ . Hence, the double integral term is zero, so

$$\begin{aligned} 0 &= \iint_{\Omega_\delta} \left[\frac{\partial g}{\partial x} + i \frac{\partial g}{\partial y} \right] dx dy = \int_{\partial\Omega_\delta} g(\zeta) d\zeta \\ &= \int_{\gamma} g(\zeta) d\zeta - \int_{|\zeta-z|=\delta} g(\zeta) d\zeta. \end{aligned}$$

Replacing g with its equivalent expression involving f and then transposing, we obtain

$$\int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta = \int_{|\zeta-z|=\delta} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

However, the right-hand side of this equation converges to $2\pi i f(z)$ as $\delta \rightarrow 0$ (see Example 10, Section 6, Chapter 1). This establishes the conclusion. ■

Cauchy's Formula has many applications, as we shall see. In the following examples, we use it to evaluate definite integrals.

Example 6 Find the value of $\int_0^{2\pi} d\theta/(2 + \sin \theta)$.

Solution The idea in this example, and others like it, is to use the substitution $z = e^{i\theta}$ effectively. We begin with three identities:

$$\cos \theta = \frac{1}{2} \left(z + \frac{1}{z} \right), \quad z = e^{i\theta}$$

$$\sin \theta = \frac{1}{2i} \left(z - \frac{1}{z} \right), \quad z = e^{i\theta}$$

$$d\theta = \frac{1}{i} \frac{dz}{z}.$$

Thus, $2 + \sin \theta = 2 + (\frac{1}{2}i)(z - 1/z) = (\frac{1}{2})(4 - iz + i/z)$, and

$$\frac{d\theta}{2 + \sin \theta} = \frac{2 dz}{iz \left(4 - iz + \frac{i}{z} \right)} = \frac{2 dz}{z^2 + 4iz - 1}.$$

Now,

$$z^2 + 4iz - 1 = [z - i(\sqrt{3} - 2)][z + i(\sqrt{3} + 2)].$$

Set $p = i(\sqrt{3} - 2)$ and $q = -i(\sqrt{3} + 2)$; p lies within the circle $|z| = 1$, while q lies outside the circle $|z| = 1$, since $|q| = \sqrt{3} + 2$. The function $(z - q)^{-1}$ is analytic in the disc $|z| < \sqrt{3} + 2$, so Cauchy's Formula (Theorem 4) gives us

$$\frac{1}{2\pi i} \int_{|z|=1} \frac{dz}{(z - q)(z - p)} = \frac{1}{p - q} = \frac{1}{2\sqrt{3}i}.$$

But the integral is just

$$\frac{1}{2\pi i} \int_{|z|=1} \frac{dz}{(z - q)(z - p)} = \frac{1}{2\pi i} \int_0^{2\pi} \frac{ie^{i\theta} d\theta}{2ie^{i\theta}(2 + \sin \theta)} = \frac{1}{4\pi i} \int_0^{2\pi} \frac{d\theta}{2 + \sin \theta}.$$

This yields

$$\int_0^{2\pi} \frac{d\theta}{2 + \sin \theta} = \frac{2\pi}{\sqrt{3}}.$$
□

Example 7 Evaluate the integral

$$\frac{1}{2\pi} \int_0^{2\pi} \frac{d\theta}{1 - 2a \cos \theta + a^2}, \quad 0 < a < 1.$$

Solution Again we use the substitution $z = e^{i\theta}$. This yields

$$1 - 2a \cos \theta + a^2 = 1 + a^2 - a\left(z + \frac{1}{z}\right),$$

so

$$\frac{d\theta}{1 - 2a \cos \theta + a^2} = \frac{dz}{iz\left(1 + a^2 - a\left(z + \frac{1}{z}\right)\right)} = \frac{dz}{i(-az^2 + (1 + a^2)z - a)}.$$

Now

$$-az^2 + (1 + a^2)z - a = -a\left(z - \frac{1}{a}\right)(z - a).$$

The point $1/a$ is outside the circle $|z| = 1$, and the point a is inside the circle $|z| = 1$. Hence,

$$\begin{aligned} \frac{1}{2\pi i} \int_{|z|=1} \frac{dz}{-az^2 + (1 + a^2)z - a} &= \frac{1}{2\pi i} \int_{|z|=1} \frac{1}{-a\left(z - \frac{1}{a}\right)} \frac{1}{z - a} dz \\ &= \frac{1}{-a\left(a - \frac{1}{a}\right)} = \frac{1}{1 - a^2}. \end{aligned}$$

But

$$\begin{aligned} \frac{1}{2\pi i} \int_{|z|=1} \frac{dz}{-az^2 + (1 + a^2)z - a} &= \frac{1}{2\pi i} \int_0^{2\pi} \frac{ie^{i\theta} d\theta}{e^{i\theta}(1 - 2a \cos \theta + a^2)} \\ &= \frac{1}{2\pi} \int_0^{2\pi} \frac{d\theta}{1 - 2a \cos \theta + a^2}. \end{aligned}$$

Thus, the integral has the value

$$\frac{1}{2\pi} \int_0^{2\pi} \frac{d\theta}{1 - 2a \cos \theta + a^2} = \frac{1}{1 - a^2}, \quad 0 < a < 1. \quad (1)$$

The function

$$P_a(\theta) = \frac{1 - a^2}{1 - 2a \cos \theta + a^2}$$

is called the **Poisson kernel** and plays an important part in the subject of harmonic functions; we will return to $P_a(\theta)$ in Chapter 4. \square

Example 8 The following is a more difficult application, this time of Cauchy's Theorem. The function $f(z) = \exp(-z^2)$ is entire. Let β be fixed, $\beta > 0$, and let R be a large positive number; let γ be the contour in Figure 2.7. Hence, $\int_{\gamma} f(z) dz = 0$, by Cauchy's Theorem. Parametrize the contour γ as indicated in Figure 2.7. Thus,

$$\begin{aligned} 0 &= \int_0^R e^{-x^2} dx + \int_0^\beta e^{-(R+it)^2} i dt + \int_R^0 e^{-(i\beta+x)^2} dx + \int_\beta^0 e^{-(it)^2} i dt \\ &= \int_0^R e^{-x^2} dx + ie^{-R^2} \int_0^\beta e^{-2iRt} e^{t^2} dt - e^{\beta^2} \int_0^R e^{-x^2} e^{-2i\beta x} dx - i \int_0^\beta e^{t^2} dt. \end{aligned}$$

As $R \rightarrow \infty$, the first integral approaches $\frac{1}{2}\sqrt{\pi}$, since

$$\int_{-\infty}^{\infty} e^{-x^2} dx = \sqrt{\pi}.$$

(This should be familiar from calculus; also see Exercise 20 in this section.)




Figure 2.7

The second integral can be estimated by

$$\left| \int_0^\beta e^{-2iRt} e^{t^2} dt \right| \leq \int_0^\beta e^{t^2} dt \leq \beta e^{\beta^2},$$

so

$$e^{-R^2} \int_0^\beta e^{-2iRt} e^{t^2} dt \rightarrow 0$$

as $R \rightarrow \infty$. (Recall that β is fixed.) The third integral converges to

$$\int_0^{\infty} e^{-x^2} [\cos(2\beta x) - i \sin(2\beta x)] dx$$

as $R \rightarrow \infty$. Setting the real and imaginary parts of the resulting expression separately equal to zero, we find that

$$e^{\beta^2} \int_0^\infty e^{-x^2} \cos(2\beta x) dx = \frac{\sqrt{\pi}}{2} \quad (2)$$

and

$$e^{\beta^2} \int_0^\infty e^{-x^2} \sin(2\beta x) dx = \int_0^\beta e^{t^2} dt. \quad \square \quad (3)$$

The next example shows that for a polynomial, Cauchy's Formula is actually an easy consequence of Taylor's Theorem and Cauchy's Theorem.

Example 9 Directly verify Cauchy's Formula for a polynomial.

Solution Let $f(\zeta) = a_0 + a_1 \zeta + \cdots + a_n \zeta^n$, $a_n \neq 0$, be a polynomial. We expand f in a Taylor series about the point z_0 :

$$f(\zeta) = b_0 + b_1(\zeta - z_0) + b_2(\zeta - z_0)^2 + \cdots + b_n(\zeta - z_0)^n,$$

where

$$b_j = \frac{f^{(j)}(z_0)}{j!}, \quad j = 0, 1, \dots, n.$$

Hence,

$$\begin{aligned} \frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{\zeta - z_0} d\zeta &= \sum_{j=0}^n b_j \frac{1}{2\pi i} \int_{\gamma} \frac{(\zeta - z_0)^j}{\zeta - z_0} d\zeta \\ &= \sum_{j=0}^n b_j \frac{1}{2\pi i} \int_{\gamma} (\zeta - z_0)^{j-1} d\zeta. \end{aligned}$$

Each of the integrals is zero except when $j = 0$ (see Exercise 16, Section 6, Chapter 1). When $j = 0$, the value of the integral is

$$\frac{1}{2\pi i} \int_{\gamma} (\zeta - z_0)^{-1} d\zeta = 1.$$

Hence,

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{\zeta - z_0} d\zeta = b_0 = f(z_0).$$

This establishes the result. \square

Example 10 Evaluate the integral

$$\int_{\gamma} \frac{z}{z+1} dz,$$

where γ is any curve in the domain $\{z : \operatorname{Im} z > 0\}$, which joins $-1 + 2i$ to $1 + 2i$.

Solution The integrand

$$f(z) = \frac{z}{z+1} = 1 - \frac{1}{z+1}$$

is the derivative of $F(z) = z - \operatorname{Log}(z+1)$. Of course, this is valid only where the function $\operatorname{Log}(z+1)$ is analytic. This is the case, however, on the complex plane with the ray $(-\infty, -1]$ deleted; this domain includes the domain $\operatorname{Im} z > 0$. Hence,

$$\begin{aligned} \int_{\gamma} \frac{z}{z+1} dz &= \int_{\gamma} f(z) dz = \int_{\gamma} F'(z) dz \\ &= F(\text{endpoint}) - F(\text{initial point}) \\ &= F(1 + 2i) - F(-1 + 2i) \\ &= [1 + 2i - \operatorname{Log}(2 + 2i)] - [-1 + 2i - \operatorname{Log}(2i)] \\ &= 2 - \left\{ \log \sqrt{8} + \frac{i\pi}{4} \right\} + \left\{ \log 2 + \frac{i\pi}{2} \right\} \\ &= 2 - \frac{1}{2} \log 2 + \frac{i\pi}{4}. \end{aligned}$$

□

EXERCISES FOR SECTION 2.3

In Exercises 1 to 4, evaluate the given integral using Cauchy's Formula or Theorem.

1. $\int_{|z|=1} \frac{z}{(z-2)^2} dz$

2. $\int_{|z|=2} \frac{e^z}{z(z-3)} dz$

3. $\int_{|z+1|=2} \frac{z^2}{4-z^2} dz$

4. $\int_{|z|=1} \frac{\sin z}{z} dz$

In Exercises 5 to 8, evaluate the definite trigonometric integral making use of the technique of Examples 6 and 7 in this section.

5. $\int_0^{2\pi} \frac{d\theta}{2 + \cos \theta}$

7. $\int_0^{2\pi} \frac{d\theta}{a + b \cos \theta}, \quad a > b > 0$

6. $\int_0^{2\pi} \frac{d\theta}{3 + \sin \theta + \cos \theta}$

8. $\int_0^\pi \frac{d\theta}{1 + \sin^2 \theta}$

In Exercises 9 to 12, evaluate the given integral using the technique of Example 10; indicate which theorem or device you used to obtain your answer.

9. $\int_{\gamma} \frac{dz}{z^2}$, where γ is any curve in $\operatorname{Re} z > 0$ joining $1 - i$ to $1 + i$.
10. $\int_{\gamma} \left(z + \frac{1}{z} \right) dz$, where γ is any curve in $\operatorname{Im} z > 0$ joining $-4 + i$ to $6 + 2i$.
11. $\int_{\gamma} e^z dz$, where γ is the semicircle from -1 to 1 passing through i .
12. $\int_{\gamma} \sin z dz$, where γ is any curve joining i to π .
13. Integrate e^{iz^2} around the contour γ shown in Figure 2.8 to obtain the **Fresnel integrals**:

$$\int_0^\infty \cos(x^2) dx = \int_0^\infty \sin(x^2) dx = \frac{\sqrt{2\pi}}{4}. \quad (4)$$

(You will also have need of the conclusion of Exercise 20.)




Figure 2.8

(Hint: On the circular arc, $z = Re^{i\theta}$, $0 \leq \theta \leq \pi/4$, you will need to know that $\sin \psi \geq (2/\pi)\psi$, $0 \leq \psi \leq \pi/2$, and hence, $e^{-R^2 \sin \psi} \leq e^{-2R^2 \psi/\pi}$. The inequality $\sin \psi \geq (2/\pi)\psi$ is proved by calculus.)

14. Specialize Theorem 4 to the case when z is the center of the circle and show that

$$f(z) = \frac{1}{2\pi} \int_0^{2\pi} f(z + re^{it}) dt. \quad (5)$$

15. (a) Use the estimate (3), Section 6, Chapter 1, and (5) above to conclude that

$$|f(z)| \leq \max_{0 \leq t \leq 2\pi} |f(z + re^{it})| \quad (6)$$

for all sufficiently small r .

- (b) Conclude from (a) that $|f|$ cannot have a strict local maximum within its domain of analyticity. That is, the graph in three-dimensional space of $(x, y, |f(x + iy)|)$ has no “peaks.”
16. Use the conclusion of Exercise 15 to establish the following result. If f is analytic and never zero on a domain D , then $|f(z)|$ has no local minima in D . That is, the graph $(x, y, |f(x + iy)|)$ has no “pits.”

17. Let $f(z) = \sum_{j=0}^{\infty} a_j(z - z_0)^j$ have a radius of convergence of $R > 0$. Show that for each $r \in (0, R)$,

$$\frac{f^{(j)}(z_0)}{j!} = a_j = \frac{1}{2\pi i} \int_{|z-z_0|=r} f(z)(z - z_0)^{-j-1} dz, \quad j = 0, 1, 2, \dots$$

(Hint: Substitute the series into the integral and interchange summation and integration.)

18. Use the result of Exercise 17 to establish these formulas:

$$(a) \frac{1}{2\pi} \int_0^{2\pi} e^{-in\theta} \sin(e^{i\theta}) d\theta = \begin{cases} 0, & n = 0, 2, 4, \dots \\ \frac{(-1)^{(n-1)/2}}{n!}, & n = 1, 3, 5, \dots \end{cases}$$

$$(b) \frac{1}{2\pi} \int_0^{2\pi} e^{-in\theta} e^{e^{i\theta}} d\theta = \frac{1}{n!}, \quad n = 0, 1, 2, \dots$$

$$(c) \frac{1}{2\pi} \int_0^{2\pi} e^{im\theta} (2e^{in\theta} - e^{i(n+1)\theta})^{-1} d\theta = \begin{cases} 0, & n < m \\ 2^{m-n-1}, & n \geq m \end{cases}$$

19. Let A be the annulus, $A = \{z : 0 \leq r < |z| < R\}$. Suppose that g is an analytic function on A (and that g' is continuous) with the property that

$$\int_{\gamma} g(z) dz = 0$$

where γ is the positively oriented circle of radius s centered at the origin, and s is some positive number in the open interval (r, R) . Show that there is an analytic function G on A with $G' = g$ throughout A . (Hint: Show that $\int_{\gamma} g(z) dz = 0$ for any piecewise smooth simple closed curve γ in A and then modify the proof of Theorem 3 to this context.)

The Integral $\int_{-\infty}^{\infty} e^{-x^2} dx$ *

20. Let $I = \int_{-\infty}^{\infty} e^{-x^2} dx$; then

$$\begin{aligned} I^2 &= \int_{-\infty}^{\infty} e^{-x^2} dx \int_{-\infty}^{\infty} e^{-y^2} dy = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-(x^2+y^2)} dx dy \\ &= \int_0^{2\pi} \int_0^{\infty} e^{-r^2} r dr d\theta = 2\pi \int_0^{\infty} r e^{-r^2} dr = \pi. \end{aligned}$$

Hence, $I = \sqrt{\pi}$.

The Poisson Kernel*

21. Verify each of the statements (a) through (e) below for $z = re^{i\theta}$ and $r < 1$.

$$(a) \operatorname{Re} \left(\frac{e^{it} + z}{e^{it} - z} \right) = \frac{1 - r^2}{1 - 2r \cos(\theta - t) + r^2}$$

- (b) $\frac{e^{it}}{e^{it} - z} + \frac{\bar{z}e^{it}}{1 - e^{it}\bar{z}} = \operatorname{Re}\left(\frac{e^{it} + z}{e^{it} - z}\right)$
- (c) $\frac{1}{2\pi} \int_0^{2\pi} f(e^{it}) \frac{e^{it}\bar{z}}{1 - \bar{z}e^{it}} dt = 0$, if $f(w)$ is analytic for $|w| < 1 + \varepsilon$
- (d) $\frac{1}{2\pi} \int_0^{2\pi} f(e^{it}) \frac{e^{it}}{e^{it} - z} dt = f(z)$
- (e) Add (c) and (d); then use (a) and (b) to conclude that

$$f(z) = \frac{1}{2\pi} \int_0^{2\pi} f(e^{it}) \frac{1 - r^2}{1 - 2r \cos(\theta - t) + r^2} dt, \quad z = re^{i\theta}.$$

2.3.1 The Cauchy–Goursat Theorem

In 1900, E. Goursat* discovered that the hypothesis “ f' is continuous” is superfluous to the conclusion of Cauchy’s Theorem. However, as might be imagined, the proof becomes more elaborate. Furthermore, the conclusion is initially established only for triangles, so that a somewhat more indirect path is needed to recover the full set of results of Section 2.3. These steps are set out here; if you do not feel the need for the extra mathematical generality gained by dropping the hypothesis of continuity on f' , you can move directly to Section 2.4.

THEOREM 1 Cauchy–Goursat Suppose f is analytic in a domain D . If Γ is a triangle in D whose inside Ω is also in D , then

$$\int_{\Gamma} f(z) dz = 0.$$

Proof Without the hypothesis of continuity on f' , we cannot apply Green’s Theorem, so we must use a more sophisticated technique. Let

$$I = \left| \int_{\Gamma} f(z) dz \right|$$

so that $I \geq 0$. We wish, of course, to show that $I = 0$. Divide the (solid) triangle $\Gamma \cup \Omega$ into four triangles by joining the midpoints of the three sides of Γ (Fig. 2.9). Orient the boundaries of all the triangles positively, call the four smaller solid triangles $\Delta_1, \dots, \Delta_4$, and let $\Gamma_1, \dots, \Gamma_4$ be the boundaries of $\Delta_1, \dots, \Delta_4$, respectively. Then

$$\int_{\Gamma} f(z) dz = \sum_{j=1}^4 \int_{\Gamma_j} f(z) dz,$$

* Edouard Goursat, 1858–1936.




Figure 2.9

so that at least one of the small triangles Δ_m satisfies

$$I = \left| \int_{\Gamma} f(z) dz \right| \leq 4 \left| \int_{\Gamma_m} f(z) dz \right| = 4I_m.$$

Rename this triangle Δ_1 . Note that the diameter of Δ_1 , the length of its longest side, is one-half the diameter of Δ . Proceed as above to divide Δ_1 into four smaller triangles by joining the midpoints of the three sides of Δ_1 . As before, we will obtain a second triangle, named Δ_2 , with boundary Γ_2 , such that

$$I_1 = \left| \int_{\Gamma_1} f(z) dz \right| \leq 4 \left| \int_{\Gamma_2} f(z) dz \right| = 4I_2$$

and $\text{diameter } (\Delta_2) = \frac{1}{2} \text{diameter } (\Delta_1) = \frac{1}{4} \text{diameter } (\Delta)$. Continuing in this manner, we obtain a sequence of triangles, $\Delta_1, \Delta_2, \Delta_3, \dots$ with respective boundaries $\Gamma_1, \Gamma_2, \Gamma_3, \dots$ such that

(i) Δ_{j+1} is a subset of Δ_j

(ii) $\text{length } (\Gamma_{j+1}) = \frac{1}{2} \text{length } (\Gamma_j)$

(iii) the diameter of $\Delta_{j+1} = \frac{1}{2} \{\text{diameter of } \Delta_j\}$

(iv) if $I_j = \left| \int_{\Gamma_j} f(z) dz \right|$, then $|I_j| \leq 4|I_{j+1}|$.

In particular, for $j = 1, 2, \dots$

(ii)' $\text{length } (\Gamma_j) = \frac{1}{2^j} \text{length } (\Gamma)$

$$(iii)' \quad \text{diameter}(\Gamma_j) = \frac{1}{2^j} \text{diameter}(\Gamma)$$

$$(iv)' \quad I \leq 4^j I_j.$$

Since the diameters of the triangles $\{\Delta_j\}$ decrease to zero and since (i) holds, there is a unique point z_0 within all the Δ_j , and z_0 lies in D . Thus, f is differentiable at z_0 so, given $\varepsilon > 0$, there is a (small) $\delta > 0$ such that

$$\left| \frac{f(z) - f(z_0)}{z - z_0} - f'(z_0) \right| < \varepsilon \quad \text{if} \quad |z - z_0| < \delta.$$

Equivalently,

$$|f(z) - [f(z_0) + f'(z_0)(z - z_0)]| < \varepsilon |z - z_0| \quad \text{if} \quad |z - z_0| < \delta. \quad (1)$$

We know that for $j \geq j_0$, the triangle Δ_j lies within the disc $\{z : |z - z_0| < \delta\}$, because $\text{diameter}(\Delta_j) \rightarrow 0$ as $j \rightarrow \infty$. Further, we know that

$$\int_{\Gamma_j} dz = \int_{\Gamma_j} z \, dz = 0, \quad (2)$$

by the version of Cauchy's Theorem proved in Section 3. The inequality given in (1), together with (2) and inequality (3) in Section 6 of Chapter 1, give

$$\begin{aligned} I_j &= \left| \int_{\Gamma_j} f(z) \, dz \right| = \left| \int_{\Gamma_j} \{f(z) - [f(z_0) + f'(z_0)(z - z_0)]\} \, dz \right| \\ &< \varepsilon \left(\max_{z \in \Gamma_j} |z - z_0| \right) (\text{length}(\Gamma_j)) \\ &\leq \varepsilon \text{diameter}(\Delta_j) \text{length}(\Gamma_j) \\ &\leq \varepsilon \frac{1}{4^j} \text{diameter}(\Delta) \text{length}(\Gamma). \end{aligned}$$

Finally, employing (iv)' above, we find that

$$0 \leq I \leq 4^j I_j \leq 4^j \left\{ \varepsilon \frac{1}{4^j} \text{diameter}(\Delta) \text{length}(\Gamma) \right\},$$

so

$$0 \leq I \leq \varepsilon \text{ diameter } (\Delta) \text{ length } (\Gamma).$$

Since ε is an arbitrary positive number, we see that the number I must be zero. ■

Extensions of Theorem 1

Although Theorem 1 appears to be very special, it can be extended to reach significantly higher levels of generality. Suppose, for instance, that γ is a simple closed curve composed of straight line segments (see Fig. 2.10a for a representative picture). Let Ω be the inside of γ , and suppose that f is analytic on a domain D that includes both γ and Ω . The region consisting of Ω and γ may be “triangulated” (see Fig. 2.10b). The line integral of f about any of the little triangles is zero, by Theorem 1. Since each edge of a triangle that is in Ω is traversed twice, once in each direction, it follows that the line integral of f over all of the edges of the triangles that lie in Ω add to zero. This leads to the conclusion that

$$\int_{\gamma} f(z) dz = 0.$$

An argument of this same nature can be used to establish the validity of the conclusion of Theorem 2 of Section 3—without, however, the hypothesis that f' is continuous. You may “triangulate” the curve in Figure 2.5 to check this for this particular example. Theorem 2 then directly implies Theorem 3 and its corollary; the proof is unchanged. We state these formally.




Figure 2.10

THEOREM 2 Let f be analytic on a simply-connected domain D .

- (a) If γ is a piecewise smooth closed curve in D , then

$$\int_{\gamma} f(z) dz = 0.$$

- (b) There is an analytic function F in D with $F' = f$ throughout D . ■

2.4 Consequences of Cauchy's Formula

Cauchy's Formula has far-reaching implications, allowing us to draw numerous nonobvious conclusions about the properties of analytic functions. We begin with one of the most important.

THEOREM 1 Suppose that f is analytic in a domain D and z_0 is a point of D . If the disc $\{z: |z - z_0| < R\}$ lies in D , then f has a power series

$$f(z) = \sum_{k=0}^{\infty} a_k (z - z_0)^k,$$

valid in this disc. Furthermore, the coefficients are given by

$$a_k = \frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{(\zeta - z_0)^{k+1}} d\zeta$$

where γ is the positively oriented circle $\{\zeta: |\zeta - z_0| = r\}$ and r is any positive number less than R .

Proof We restrict our attention to the disc $\Delta = \{z: |z - z_0| < R\}$, which by assumption lies within D . Fix any r , $0 < r < R$. We shall suppose first that f has a continuous derivative on the disc Δ . Consequently, for each z with $|z - z_0| < r$, Cauchy's Formula, Theorem 4 of Section 3 is valid:

$$f(z) = \frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta,$$

where γ is the positively oriented circle $\{\zeta: |\zeta - z_0| = r\}$. Let $s = |z - z_0|$, so that $s < r$. For this z and any ζ in γ ,

$$\frac{1}{\zeta - z} = \frac{1}{(\zeta - z_0) - (z - z_0)} = \frac{1}{\zeta - z_0} \frac{1}{1 - \left(\frac{z - z_0}{\zeta - z_0} \right)}.$$

Now

$$\left| \frac{z - z_0}{\zeta - z_0} \right| = \frac{s}{r} < 1,$$

so we can represent

$$\left(1 - \left(\frac{z - z_0}{\zeta - z_0}\right)\right)^{-1}$$

by the geometric series

$$\begin{aligned} \left(1 - \left(\frac{z - z_0}{\zeta - z_0}\right)\right)^{-1} &= 1 + \left(\frac{z - z_0}{\zeta - z_0}\right) + \left(\frac{z - z_0}{\zeta - z_0}\right)^2 + \dots \\ &= \sum_{k=0}^{\infty} \left(\frac{z - z_0}{\zeta - z_0}\right)^k. \end{aligned}$$

The series $\sum_{k=0}^{\infty} (s/r)^k$ converges, since $0 < s < r$. Consequently, given $\varepsilon > 0$, we can choose N so big that $\sum_{k=N}^{\infty} (s/r)^k < \varepsilon$. Hence, for all ζ with $|\zeta - z_0| = r$ and all $n \geq N$,

$$\left| \sum_{k=n}^{\infty} \left(\frac{z - z_0}{\zeta - z_0}\right)^k \right| \leq \sum_{k=n}^{\infty} \left(\frac{s}{r}\right)^k \leq \sum_{k=N}^{\infty} \left(\frac{s}{r}\right)^k < \varepsilon. \quad (1)$$

Now, for any n , we can write

$$\begin{aligned} \int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta &= \int_{\gamma} \frac{f(\zeta)}{\zeta - z_0} \left\{ \sum_{k=0}^{\infty} \left(\frac{z - z_0}{\zeta - z_0}\right)^k \right\} d\zeta \\ &= \sum_{k=0}^n (z - z_0)^k \left\{ \int_{\gamma} \frac{f(\zeta)}{(\zeta - z_0)^{k+1}} d\zeta \right\} + \int_{\gamma} \frac{f(\zeta)}{\zeta - z_0} \left\{ \sum_{k=n+1}^{\infty} \left(\frac{z - z_0}{\zeta - z_0}\right)^k \right\} d\zeta. \end{aligned}$$

When we employ (1) and inequality (3) from Section 6 in Chapter 1, we see that

$$\left| \int_{\gamma} \left\{ \frac{f(\zeta)}{\zeta - z_0} \right\} \left\{ \sum_{k=n+1}^{\infty} \left(\frac{z - z_0}{\zeta - z_0}\right)^k \right\} d\zeta \right| \leq \frac{M}{r} \varepsilon 2\pi r = 2\pi \varepsilon M,$$

where $M = \max\{|f(\zeta)|: |\zeta - z_0| = r\}$. From Cauchy's Formula and the previous computations, we see that for $n \geq N$

$$f(z) = \frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta = \sum_{k=0}^n (z - z_0)^k \left\{ \frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{(\zeta - z_0)^{k+1}} d\zeta \right\} + E(z; z_0),$$

where $|E(z; z_0)| < \varepsilon M$. This implies that the infinite series

$$\sum_{k=0}^{\infty} (z - z_0)^k \left\{ \frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{(\zeta - z_0)^{k+1}} d\zeta \right\}$$

converges, and its sum is $f(z)$.

To complete the proof of the theorem, we must show that its conclusion is valid when we omit the assumption that f has a continuous derivative. However, Theorem 2b of Section 2.3.1 implies that there is an analytic function F on the disc Δ with $F(z_0) = 0$ and $F' = f$ on Δ . The function F does have a continuous derivative on Δ , and consequently, by the preceding part of the proof, F has a power series valid on Δ . Theorem 2 of Section 2.2 then shows that $F' = f$ also has a power series whose radius of convergence is the same as that of F (that is, at least R). Furthermore, f' is also given by a power series and, in particular, is continuous. Hence, ipso facto, the first portion of the proof applies to f' ; we are done. ■

Theorem 1 has an immediate, and important, corollary.

If f is analytic on a domain D , then so is f' . Hence, f has derivatives of all orders, and each derivative is analytic on D . (2)

This follows because in each disc in D , f is given by a power series, and (by Theorem 3 of Section 2 of this chapter) each power series is infinitely differentiable. Note that (2) brings together the parallel threads of Section 2.3 and Section 2.3.1, in that the results of 3.1 lead to Theorem 1 of this section and hence to (2). That is to say, when we assume that f is merely once (complex) differentiable, it follows from the Cauchy–Goursat Theorem and the rest of the development in Section 2.3.1 and Theorem 1 of this section that f is infinitely (complex) differentiable. In other words, from the existence of one derivative, we are able to infer the existence of all derivatives. Contrast this with the case of real-valued functions of a real variable. For instance, the function

$$u(t) = \begin{cases} t^2, & t \geq 0 \\ -t^2, & t \leq 0 \end{cases}$$

is differentiable at all points t , $-\infty < t < \infty$, with derivative

$$u'(t) = \begin{cases} 2t & t \geq 0 \\ -2t, & t < 0, \end{cases}$$

and the derivative is even continuous. However, u' is itself not differentiable at $t = 0$.

Let us find the power series for some of the elementary analytic functions discussed in Section 1. This is accomplished by use of Theorem 1 and formula (12), Section 2 of this chapter.

Example 1 The function e^z is entire and is its own derivative. Hence, $a_n = 1/n!$ for $n = 0, 1, 2, \dots$, so

$$e^z = \sum_{k=0}^{\infty} \frac{z^k}{k!},$$

as we saw already in formula (13), Section 2. □

Example 2 The functions $\cos z$ and $\sin z$ are entire and satisfy $(\sin z)' = \cos z$ and $(\cos z)' = -\sin z$. Since $\sin(0) = 0$ and $\cos(0) = 1$, we obtain

$$\sin z = \sum_{j=0}^{\infty} (-1)^j \frac{z^{2j+1}}{(2j+1)!},$$

$$\cos z = \sum_{j=0}^{\infty} (-1)^j \frac{z^{2j}}{(2j)!}.$$
□

Example 3 The function $\text{Log}(1-z)$ is analytic in the disc $|z| < 1$ (see Example 10, Section 1 of this chapter), and its derivatives are

$$\frac{d^n}{dz^n}(\text{Log}(1-z)) = -\frac{(n-1)!}{(1-z)^n}, \quad n = 1, 2, \dots.$$

Hence, for $|z| < 1$,

$$\text{Log}(1-z) = -\sum_{n=1}^{\infty} \frac{z^n}{n}.$$
□

Example 4 The functions $\sinh(z)$ and $\cosh(z)$ are defined by

$$\sinh(z) = \frac{1}{2}(e^z - e^{-z}), \quad \cosh(z) = \frac{1}{2}(e^z + e^{-z}).$$

From the expansion of e^z above, we find that

$$\cosh(z) = \sum_{k=0}^{\infty} \frac{z^{2k}}{(2k)!}$$

and

$$\sinh(z) = \sum_{k=0}^{\infty} \frac{z^{2k+1}}{(2k+1)!}.$$

Both series converge for all z , by an application of the ratio test. □

Example 5 The function $f(z) = 1/z^3$ is analytic in the disc $D = \{z: |z-1| < 1\}$; at 1, the value of the n th derivative of f is

$$\frac{1}{2}(n+2)!(-1)^n, \quad n = 1, 2, 3, \dots;$$

thus, in D ,

$$\begin{aligned}\frac{1}{z^3} &= 1 + \sum_{n=1}^{\infty} \frac{1}{2} (n+2)! (-1)^n \frac{(z-1)^n}{n!} \\ &= 1 + \frac{1}{2} \sum_{n=1}^{\infty} (-1)^n (n+2)(n+1)(z-1)^n.\end{aligned}$$

The series is valid for $|z-1| < 1$. □

Theorem 1 has another significant implication.

Suppose that f is analytic on a domain D and, further, at some point $z_0 \in D$, $f^{(k)}(z_0) = 0$, $k = 0, 1, 2, \dots$. Then $f(z) = 0$ for all $z \in D$. (3)

The proof of (3) lies a little deeper than that of (2). To begin, let Δ be any disc in D with center at z_0 ; then by Theorem 1 and (2) of Section 2, $f(z)$ is given by a power series within Δ :

$$f(z) = \sum_{k=0}^{\infty} a_k (z - z_0)^k; \quad a_k = \frac{f^{(k)}(z_0)}{k!}, \quad k = 0, 1, \dots$$

Thus the hypothesis that $f^{(k)}(z_0) = 0$ for all k implies that $f(z) = 0$ for all $z \in \Delta$. Next, let L be a line segment in D from z_0 to another point z_1 in D . We shall show that f and all the derivatives of f vanish at all points of L . We take note of the fact that there is a positive number δ such that each disc of radius δ centered at a point of L lies within D . (This is reasonably clear from a picture, such as Fig. 2.11, but a careful proof depends on fundamental properties of real and complex numbers; we shall not prove it.) Let $\Delta_0, \Delta_1, \dots, \Delta_N$ be a “chain” of discs of radius δ centered at points $\zeta_0 = z_0, \zeta_1, \dots, \zeta_N$ in L , and with the point ζ_j in the disc Δ_{j-1} for $j = 1, 2, \dots, N$, and $z_1 \in \Delta_N$ (see Fig. 2.11).




Figure 2.11

We know already that $f = 0$ at all points of the disc Δ_0 ; hence, $f^{(k)}(\zeta_1) = 0$ for $k = 0, 1, \dots$, since $\zeta_1 \in \Delta_0$. Thus, the power-series representation for $f(z)$, which is valid in Δ_1 shows that $f = 0$ at all points of the disc Δ_1 . Consequently, $f^{(k)}(\zeta_2) = 0$ for $k = 0, 1, \dots$, since $\zeta_2 \in \Delta_1$, and we learn that $f \equiv 0$ in the disc Δ_2 . Continuing in this way, after N steps we learn that $f^{(k)}(z_1) = 0$ for $k = 0, 1, \dots$, which is what we wished to show. Hence, if $f^{(k)}(z_0) = 0$ for $k = 0, 1, 2, \dots$, then $f^{(k)}(z_1) = 0$ for $k = 0, 1, \dots$ at any point z_1 that can be reached from z_0 by a straight line segment. Since *all* points of D can be reached from z_0 by a polygonal curve consisting of a finite number of straight line segments, we see that $f(z) = 0$ for all $z \in D$.

The Order of a Zero

Suppose that f is analytic and not identically zero on a domain D , and $f(z_0) = 0$ for some $z_0 \in D$. Thus, the power series for f centered at z_0 is

$$f(z) = a_1(z - z_0) + a_2(z - z_0)^2 + \cdots.$$

We know from (3) that not all of the coefficients a_k can vanish, so there is an integer $m \geq 1$ such that

$$a_1 = \cdots = a_{m-1} = 0, \quad \text{but } a_m \neq 0.$$

Therefore,

$$f(z) = a_m(z - z_0)^m + a_{m+1}(z - z_0)^{m+1} + \cdots, \quad a_m \neq 0.$$

This means that

$$f^{(k)}(z_0) = 0 \quad \text{for } k = 0, \dots, m-1, \quad \text{but } f^{(m)}(z_0) \neq 0.$$

In this case, we say that f has a zero of order m at z_0 . Furthermore, if we set

$$g(z) = \frac{f(z)}{(z - z_0)^m},$$

then g is actually analytic in D and $g(z_0) = a_m \neq 0$. Clearly, g is analytic on $D \setminus \{z_0\}$. To see that g is analytic near z_0 , note that the series

$$a_m + a_{m+1}(z - z_0) + a_{m+2}(z - z_0)^2 + \cdots$$

converges in a disc about z_0 exactly because $(z - z_0)^m$ times this series converges to $f(z)$ in this disc. That is, if $g(z)$ is the sum of this series, then $(z - z_0)^m g(z) = f(z)$, and we are done. We state this result formally, since it will be of considerable utility for us.

If f has a zero of order m at z_0 , then $f(z) = (z - z_0)^m g(z)$, where g is analytic in D and $g(z_0) \neq 0$. (4)

Morera's* Theorem

Cauchy's Theorem, on which so much of the development of complex variables depends, has a converse. This theorem, given precisely below, states that if the integral of a continuous function f over every triangle in some domain is zero, then f must be analytic in that domain.

THEOREM 2 Morera's Theorem If f is a continuous function on a domain D and if

$$\int_{\gamma} f(z) dz = 0$$

for every triangle γ that lies, together with its interior, in D , then f is analytic on D .

Proof The proof can be adapted almost word for word from parts of Section 3. Let z_0 be a point of D and Ω be the disc $\{z : |z - z_0| < r\}$, where $r > 0$ is so small that Ω is in D . We shall show that f is analytic on the disc Ω . Define

$$F(z) = \int_{z_0}^z f(\zeta) d\zeta,$$

where the integration is along the radius joining z_0 to z . We shall show that F is analytic in Ω and that $F' = f$. It will then follow from (2) of this section that f is analytic in Ω , as well.

Let h be a small complex number; then

$$F(z + h) - F(z) = \int_z^{z+h} f(\zeta) d\zeta,$$

because the integral of f around the triangle with vertices $\{z_0, z, z + h\}$ is zero (Fig. 2.12). Hence,

$$\frac{F(z + h) - F(z)}{h} - f(z) = \int_z^{z+h} \left\{ \frac{f(\zeta) - f(z)}{h} \right\} d\zeta.$$

Suppose now that $\varepsilon > 0$ is given; choose δ so small that

$$|f(\zeta) - f(z)| < \varepsilon$$

if $|\zeta - z| < \delta$; this is possible by the continuity of f . Then for $|h| < \delta$, we have

$$\left| \int_z^{z+h} \{f(\zeta) - f(z)\} d\zeta \right| < \varepsilon|h|,$$

* Ciacinto Morera, 1856–1909.




Figure 2.12

by the estimate (3) from Section 6 of Chapter 1. Putting all this together, we find that

$$\left| \frac{F(z+h) - F(z)}{h} - f(z) \right| < \varepsilon \quad \text{if } |h| < \delta,$$

which implies that

$$\lim_{h \rightarrow 0} \frac{F(z+h) - F(z)}{h} = f(z);$$

that is, F is differentiable and $F' = f$. ■

We now give four applications of the theorems of this section.

Application 1 Liouville's Theorem

THEOREM 3 Liouville's* Theorem If F is entire and if there is a constant M such that $|F(z)| \leq M$ for all z , then F is identically constant.

Proof Set $g(z) = (F(z) - F(0))/z$; g is then an entire function. For z , $|z| = R > 0$, we may make the simple estimate

$$|g(Re^{i\theta})| \leq \frac{|F(Re^{i\theta})| + |F(0)|}{R} \leq \frac{2M}{R}.$$

Let ζ be a point of the plane, and assume R is so big that $R > |\zeta|$. By Cauchy's Formula,

$$g(\zeta) = \frac{1}{2\pi i} \int_{|z|=R} \frac{g(z)}{z - \zeta} dz,$$

so

* Joseph Liouville, 1809–1882.

$$|g(\zeta)| \leq \frac{1}{2\pi} \frac{2M}{R} \frac{2\pi R}{R - |\zeta|} \rightarrow 0 \quad \text{as } R \rightarrow \infty.$$

Consequently, $g = 0$. Thus, $F(z) = F(0)$ for all z . ■

Application 2 Analytic Logarithms

Suppose f is analytic and zero-free in a simply-connected domain D . Fix $z_0 \in D$ and define

$$h(z) = \int_{z_0}^z \frac{f'(w)}{f(w)} dw,$$

where the integral is taken over any piecewise smooth curve in D from z_0 to z . Since the integrand f'/f is analytic in D , the function h is analytic in D . As we have seen in the proof of Theorem 2, the derivative of h is the integrand; in this case, $h' = f'/f$. Thus,

$$\begin{aligned} (e^{-h(z)}f(z))' &= -h'(z)e^{-h(z)}f(z) + e^{-h(z)}f'(z) \\ &= e^{-h(z)}[-f'(z) + f'(z)] \\ &= 0. \end{aligned}$$

Consequently, $e^{-h(z)}f(z) = c$ for all $z \in D$, where c is some nonzero constant. Setting $z = z_0$ yields $c = f(z_0)$, since $h(z_0) = 0$, so

$$f(z) = f(z_0)e^{h(z)}.$$

That is to say, if f is zero-free and analytic on a simply-connected domain D , there is an analytic function g with

$$e^{g(z)} = f(z), \quad z \in D.$$

[Just set $g = h - \operatorname{Log} f(z_0)$.]

Application 3 Flows (Continued)*

Suppose that f is a sourceless and irrotational flow in a domain D . We know from the discussion in Section 2.1.1 that \bar{f} is analytic on D . Even more, we learned there that

$$\int_{\gamma} \overline{f(z)} dz = 0 \tag{5}$$

for all smooth closed curves γ in D . It is simple to see from this that (5) also holds

for all piecewise smooth closed curves γ , as well. Equation (5) actually implies that \bar{f} is the derivative of some analytic function G on D . To see this, fix a point a in D and define

$$G(z) = \int_{\Gamma} \overline{f(w)} dw,$$

where Γ is any piecewise smooth curve from a to z . $G(z)$ is independent of the choice of Γ , since if Γ' is another curve from a to z , then $\gamma = \Gamma - \Gamma'$ is a piecewise smooth closed curve so

$$0 = \int_{\gamma} \overline{f(w)} dw = \int_{\Gamma} \overline{f(w)} dw - \int_{\Gamma'} \overline{f(w)} dw.$$

Now an argument identical to the one employed in proving Theorem 2, Morera's Theorem, shows that $G' = \bar{f}$ on D . Any function G with $G' = \bar{f}$ is called a **complex potential** of the flow f . We shall see in Section 4.1 of Chapter 3 that the complex potential is intimately related to conformal mapping.

Application 4 Multiplication of Power Series

Theorem 4 of Section 2 asserts that the product of two convergent power series is again a power series, and the coefficients of the product series may be obtained from those of the original series by a rather simple rule. Theorem 1 of this section allows us to give a simple proof of that result. Let

$$f(z) = \sum_{n=0}^{\infty} a_n z^n \quad \text{and} \quad g(z) = \sum_{n=0}^{\infty} b_n z^n$$

be the two series, each with radius of convergence at least R , $R > 0$. Then the product $h(z) = f(z)g(z)$ is analytic in the disc $\{z : |z| < R\}$. Hence, by Theorem 1 of this section, h has a power series expansion in this same disc

$$h(z) = \sum_{n=0}^{\infty} c_n z^n, \quad |z| < R,$$

and the coefficients c_n are given by (12) of Section 2.2:

$$c_n = \frac{h^{(n)}(0)}{n!}.$$

However, the product rule for differentiation gives

$$h^{(n)}(0) = \sum_{k=0}^n \frac{n!}{(n-k)! k!} f^{(k)}(0) g^{(n-k)}(0).$$

Again using (12) of Section 2.2, we obtain $f^{(k)}(0) = (a_k)k!$ and $g^{(n-k)}(0) = (b_{n-k})(n-k)!$. When we substitute these into the formula above for $h^{(n)}(0)$ and cancel the factorials, we get

$$h^{(n)}(0) = n! \sum_{k=0}^n a_k b_{n-k},$$

which is just what Theorem 4 of Section 2 asserted.

EXERCISES FOR SECTION 2.4

In Exercises 1 to 8, give the order of *each* of the zeros of the given function.

1. $\frac{\sin z}{z}$

2. $(e^z - 1)^2$

3. $(z^2 + z - 2)^3$

4. $(z^2 - 4z + 4)^3$

5. $z^2(1 - \cos z)$

6. $\text{Log}(1 - z), |z| < 1$

7. $e^{2z} - 3e^z - 4$

8. $\frac{z}{z^2 + 1}$

In Exercises 9 to 16, find the power-series expansion about the given point for each of the functions; find the largest disc in which the series is valid.

9. $z(e^z - 1)$ about $z_0 = 0$

10. e^z about $z_0 = \pi i$

11. $z^3 + 6z^2 - 4z - 3$ about $z_0 = 1$

12. $\frac{z^2}{1 - z}$ about $z_0 = 0$

13. $\frac{z+2}{z+3}$ about $z_0 = -1$

14. $[\text{Log}(1 - z)]^2$ about $z_0 = 0$ (first four terms)

15. $\sin \pi z$ about $z_0 = \frac{1}{2}$

16. $\tan z$ about $z_0 = 0$ (first four terms)

17. Suppose that f is analytic on a domain D and has a zero of order m at z_0 in D . Show that (a) f' has a zero of order $m - 1$ at z_0 ; (b) f^2 has a zero of order $2m$ at z_0 .

18. Use Theorem 1 to establish the **Cauchy estimates**:

$$|f^{(n)}(z_0)| \leq \frac{n!}{r^n} \max_{|z-z_0|=r} |f(z)|, \quad n = 0, 1, 2, \dots,$$

whenever f is analytic on a domain containing the set $\{z : |z - z_0| \leq r\}$.

19. Use the Cauchy estimates (Exercise 18) for $n = 1$ to give another proof of Liouville's Theorem by showing that the derivative of a bounded entire function is identically zero.
20. Suppose that f is an entire function and $\text{Re } f(z) \leq c$ for all z . Show that f is constant. (**Hint:** Consider $\exp(f(z))$.)
21. Suppose that f is an entire function and that there are positive constants A and m with $|f(z)| \leq A|z|^m$ if $|z| \geq R_0$. Show that f is a polynomial of degree m or less. (**Hint:** Use the Cauchy estimates (Exercise 18) for $n > m$ and let $r \rightarrow \infty$.)

22. Let D be a simply-connected domain and f an analytic function on D that has no zeros in D . Let γ be a complex number, $\gamma \neq 0$. Show that there is an analytic function g on D with $f = g^\gamma$. (**Hint:** Use Application 2.)
23. Suppose that F is analytic in the region $|z| > R$, including at ∞ (that is, $G(z) = F(1/z)$ is analytic for $|z| < 1/R$). Show that F can be expressed as a “power series” in $1/z$:

$$F(z) = \sum_{k=0}^{\infty} c_k \frac{1}{z^k}, \quad |z| > R,$$

and derive a formula for c_k similar to that given in Theorem 1.

24. Use Morera’s Theorem and an interchange of the order of integration to show that each of the following functions is analytic on the indicated domain; find a power-series expansion for each function by using the known power series for the integrand and interchanging the summation and integration.

$$\begin{aligned} \text{(a)} \quad & \int_0^1 \frac{dt}{1-tz} \quad \text{on } |z| < 1 & \text{(b)} \quad & \int_0^{1/2} \operatorname{Log}(1-tz) dt \quad \text{on } |z| < 2 \\ \text{(c)} \quad & \int_{-\pi/2}^{\pi/2} \sin(z+t) dt, \quad \text{all } z. \end{aligned}$$

Differential Equations in the Complex Plane*

25. Find all solutions to the differential equation

$$f''(z) + \beta^2 f(z) = 0, \quad f \text{ is an entire function.}$$

(**Hint:** Write $f(z) = \sum_0^{\infty} a_j z^j$ and solve for the coefficients a_2, a_3, \dots in terms of a_0, a_1 , and β .)

26. Use the technique of Exercise 25 to give the solutions of these differential equations:

$$\begin{aligned} \text{(a)} \quad & f''(z) - 3f'(z) + 2f(z) = 0; \quad a_0 = 1, \quad a_1 = 2 \\ \text{(b)} \quad & f''(z) - \frac{2}{z} f'(z) + f(z) = 0; \quad z \neq 0, \quad f(0) = 1, \quad f'(0) = 0 \\ \text{(c)} \quad & f''(z) + 2f'(z) + f(z) = 0; \quad a_0 = 0, \quad a_1 = 1 \end{aligned}$$

27. If f and A are analytic in a simply-connected domain D and

$$f'(z) = A(z)f(z), \quad z \in D,$$

show that $f(z) = C \exp[\int_{z_0}^z A(w) dw]$ for a constant C , where the integral is taken over any piecewise smooth curve joining (a fixed point) z_0 to z . (**Hint:** Let

$$g(z) = \exp \left[- \int_{z_0}^z A(w) dw \right].$$

Show that $(f(z)g(z))' = 0$ in D .)

28. Use the result of Exercise 27 to give the solution to each of these differential equations:

- $f'(z) - 2zf(z) = 0$
- $f'(z) + e^z f(z) = 0$
- $f'(z) + \frac{1}{z}f(z) = 0$

Bessel Functions*

29. Let v be an integer, $v \geq 0$. Show that one solution of the differential equation

$$z^2 f''(z) + zf'(z) + (z^2 - v^2)f(z) = 0$$

is

$$J_v(z) = \sum_{n=0}^{\infty} (-1)^n \frac{z^{2n+v}}{2^{2n+v}(v+n)!n!}.$$

J_v is the Bessel function of the first kind of order v .

30. Show that the Bessel functions satisfy the recurrence relation

$$J_{v-1} - J_{v+1} = 2J'_v.$$

2.5 Isolated Singularities

An analytic function f has an **isolated singularity** at a point z_0 if f is analytic in the punctured disc $0 < |z - z_0| < r$, for some $r > 0$. The isolated singularities of an analytic function play a large role in the applications of complex variables, including analyzing flows and fields and computing definite integrals by means of the Residue Theorem. We shall analyze isolated singularities in this section; the applications will come in the next section, with the residue theorem, and in later chapters.

Three examples of isolated singularities are:

- z_0 is an isolated singularity for $f(z) = (z^2 - z_0^2)/(z - z_0)$
- z_0 is an isolated singularity for $g(z) = 7(z - z_0)^{-4}$
- z_0 is an isolated singularity for $h(z) = \exp(1/(z - z_0))$.

As these examples show, there are three possible modes of behavior for $|f|$ when $0 < |z - z_0| < r$:

$$|f(z)| \text{ remains bounded as } z \rightarrow z_0 \quad (1)$$

$$\lim_{z \rightarrow z_0} |f(z)| = \infty \quad (2)$$

$$\text{Neither (1) nor (2) holds.} \quad (3)$$

We will explore the first two of these possibilities in this section; the third we will not need. One characterization of such singularities is presented in Exercise 17.

Removable Singularities

Suppose first that (1) holds for f . Set

$$g(z) = \begin{cases} (z - z_0)^2 f(z), & 0 < |z - z_0| < r \\ 0, & z = z_0. \end{cases}$$

The function g is most certainly analytic for $0 < |z - z_0| < r$, but it is also differentiable at z_0 , since

$$\lim_{z \rightarrow z_0} \frac{g(z) - g(z_0)}{z - z_0} = \lim_{z \rightarrow z_0} (z - z_0)f(z) = 0.$$

(The last equality is because $|f|$ remains bounded as $z \rightarrow z_0$.) Since g is analytic in $|z - z_0| < r$, we know that g has a power series valid for $|z - z_0| < r$:

$$g(z) = b_0 + b_1(z - z_0) + b_2(z - z_0)^2 + \cdots$$

However, $b_0 = g(z_0) = 0$ and $b_1 = g'(z_0) = 0$, so

$$g(z) = b_2(z - z_0)^2 + b_3(z - z_0)^3 + \cdots,$$

and hence,

$$f(z) = b_2 + b_3(z - z_0) + \cdots.$$

for $0 < |z - z_0| < r$. Set $f(z_0) = b_2$. Since the power series is valid for $|z - z_0| < r$, we see that the hypothesis (1) actually implies that f is analytic in $|z - z_0| < r$. When (1) holds, we then learn that f can be extended to be analytic in the disc $|z - z_0| < r$. In this case, z_0 is called a **removable singularity** for f .

Poles

Suppose next that (2) holds; there is no harm in assuming that r is so small that $|f(z)| > 1$ for $0 < |z - z_0| < r$. Consequently, $g(z) = 1/f(z)$ is analytic on the punctured disc $\{z : 0 < |z - z_0| < r\}$ and is bounded there: $|g(z)| < 1$. But then the foregoing discussion shows that z_0 is a removable singularity for g and $g(z_0) = 0$, because (2) holds. Let $m \geq 1$ be the order of the zero of g at z_0 and write

$$g(z) = (z - z_0)^m h(z),$$

where h is analytic on the disc $\{z : |z - z_0| < r\}$ and $h(z_0) \neq 0$. Since g does not

vanish in the punctured disc $0 < |z - z_0| < r$, neither does h . Consequently, the function $H(z) = 1/h(z)$ is analytic on the disc $\{z: |z - z_0| < r\}$. This, then, leads to

$$f(z) = \frac{1}{g(z)} = \frac{1}{(z - z_0)^m} \frac{1}{h(z)} = \frac{H(z)}{(z - z_0)^m}, \quad (4)$$

where $H(z)$ is analytic on the disc $\{z: |z - z_0| < r\}$ and $H(z_0) \neq 0$. Thus, if (2) holds, f has predictable behavior near z_0 , and, in particular, $|f(z)|$ grows to infinity as $z \rightarrow z_0$, just like some power of $1/|z - z_0|$. In this case, we say that f has a **pole** at z_0 . The **order of the pole** of f is the integer m that is the order of the zero of $1/f$ at z_0 .

Essential Singularities

If (3) holds, then z_0 is termed an **essential singularity** of f . Information on the behavior of f near an essential singularity is contained in the exercises at the end of this section.

The Residue at z_0

We note here the simple but significant fact that if f is a function analytic in the punctured disc $0 < |z - z_0| < r$, then the value of the integral

$$\frac{1}{2\pi i} \int_{|\zeta - z_0|=s} f(\zeta) d\zeta \quad (5)$$

is the same for all s , $0 < s < r$. The proof is easy; if $0 < s_1 < s_2 < r$, then the region $0 < |z - z_0| < r$ is divided into four sectors by the horizontal and vertical lines through z_0 (Fig. 2.13).

As indicated in Figure 2.13, the integral over $|\zeta - z_0| = s_1$ (oriented clockwise) plus the integral over $|\zeta - z_0| = s_2$ (oriented counterclockwise) can be broken up




Figure 2.13

into the sum of four integrals (one in each sector); the line integrals over the interior line segments cancel. The integral in each sector is zero, by Cauchy's Theorem. Thus, the value of the integral in (5) does not depend on s . The value of the integral in (5) is called the **residue** of f at z_0 , and we denote it by $\text{Res}(f; z_0)$. Thus,

$$\text{Res}(f; z_0) = \frac{1}{2\pi i} \int_{|\zeta-z_0|=s} f(\zeta) d\zeta. \quad (5')$$

The residues of a function with only isolated singularities play an important role in complex variables.

The Computation of Residues

Although (5') is the definition of the residue of f at z_0 , it generally does not provide the most effective way to compute it. We set out here the most important case—when f has a pole at z_0 . The computation is based on (4):

$$f(z) = \frac{H(z)}{(z - z_0)^m}.$$

Since H is analytic on the disc $\{z: |z - z_0| < r\}$, it has a power series expansion valid there:

$$H(z) = \sum_{k=0}^{\infty} c_k (z - z_0)^k, \quad |z - z_0| < r.$$

In (4), replace H with this series, and carry out the division for $z \neq z_0$. This yields

$$f(z) = \frac{c_0}{(z - z_0)^m} + \cdots + \frac{c_{m-1}}{(z - z_0)} + c_m + c_{m+1}(z - z_0) + \cdots. \quad (6)$$

Next, replace f in (5) with this expression, and then interchange the summation and integration. Finally, recall from Exercise 16, Section 6, Chapter 1, and from Cauchy's Theorem, that

$$\frac{1}{2\pi i} \int_{|z-z_0|=s} (z - z_0)^j dz = \begin{cases} 1 & \text{if } j = -1 \\ 0 & \text{if } j \text{ is any other integer.} \end{cases}$$

This then yields the conclusion

$$\text{Res}(f; z_0) = c_{m-1} = \frac{H^{(m-1)}(z_0)}{(m-1)!}. \quad (7)$$

That is, to find the residue of a function f which has a pole at the point z_0 , write the expansion of f in positive and negative powers of $z - z_0$ in the form (6). This may always be done according to the presentation under the topic “Poles.”

The residue of f at z_0 is the coefficient of $(z - z_0)^{-1}$ in the expansion (6). (8)

In the examples that follow, we show several instances of how this is accomplished.

Example 1 Find the residue of $f(z) = (z^2 + 3z - 1)/(z + 2)$ at its pole.

Solution The pole of f is at $z_0 = -2$. Expand the numerator in powers of $z + 2$:

$$z^2 + 3z - 1 = (z + 2)^2 - (z + 2) - 3.$$

Therefore, by (8), $\text{Res}(f; -2) = -3$. □

Example 2 Find the residue of $g(z) = e^z/(z - 1)^3$ at $z_0 = 1$.

Solution Expand e^z in powers of $z - 1$:

$$e^z = ee^{z-1} = e \left(1 + (z - 1) + \frac{(z - 1)^2}{2!} + \frac{(z - 1)^3}{3!} + \dots \right).$$

Thus,

$$g(z) = \frac{e}{(z - 1)^3} + \frac{e}{(z - 1)^2} + \frac{e}{2(z - 1)} + \frac{e}{6} + \frac{e}{24}(z - 1) + \dots,$$

so

$$\text{Res}(g; 1) = \frac{e}{2}. \quad \square$$

Example 3 The function $\text{Log } z$ is analytic on the domain obtained by deleting the ray $(-\infty, 0]$ from the plane. Find the residue of $h(z) = (\text{Log } z)/(1 + z^2)^2$ at each of its poles.

Solution h has a pole of order 2 at i and at $-i$. We work first with $z_0 = i$. We may write h as

$$h(z) = \left\{ \frac{1}{(z - i)^2} \right\} \left\{ \frac{\text{Log } z}{(z + i)^2} \right\}.$$

The second factor is analytic near i . According to (7), the residue of h at i is the value of the derivative of the second factor, $(\text{Log } z)/(z + i)^2$, at $z_0 = i$. This is easily

computed to be $\pi/8 + i/4$. Therefore,

$$\text{Res}(h; i) = \frac{\pi}{8} + \frac{i}{4}.$$

Similarly,

$$\text{Res}(h; -i) = \frac{\pi}{8} + \frac{i}{4}. \quad \square$$

Example 4 Suppose that F and G are analytic functions on the disc $\{z: |z - z_0| < r_0\}$ with $G(z_0) = 0$ but $G'(z_0) \neq 0$. Show that

$$\text{Res}\left(\frac{F}{G}; z_0\right) = \frac{F(z_0)}{G'(z_0)}.$$

Solution Write $H = F/G$; now $G(z) = (z - z_0)P(z)$, where P is analytic on the disc $\{z: |z - z_0| < r_0\}$ and $P(z_0) \neq 0$. Indeed,

$$\begin{aligned} P(z_0) &= \lim_{z \rightarrow z_0} P(z) = \lim_{z \rightarrow z_0} \frac{G(z)}{z - z_0} \\ &= \lim_{z \rightarrow z_0} \frac{G(z) - G(z_0)}{(z - z_0)} \\ &= G'(z_0). \end{aligned}$$

We may write

$$H(z) = \frac{F(z)}{G(z)} = \left(\frac{1}{z - z_0}\right) \frac{F(z)}{P(z)}.$$

The function F/P is analytic on a disc centered at z_0 , and its value at z_0 is the residue of H at z_0 , by (7). Thus,

$$\text{Res}\left(\frac{F}{G}; z_0\right) = \frac{F(z_0)}{P(z_0)} = \frac{F(z_0)}{G'(z_0)}. \quad \square$$

Example 5 Find the residue of $r(z) = (z + 1)/(z^2 + 4)(z - 1)^3$ at each of its poles.

Solution r has poles of order 1 at $-2i$ and $2i$ and a pole of order 3 at 1. By Example 4, the residue of r at $-2i$ is the quotient of the value of the numerator at $-2i$ with the derivative of the denominator at $-2i$. This is easily computed to be

$$\text{Res}(r; -2i) = \frac{24 + 7i}{500}.$$

Likewise,

$$\text{Res}(r; -2i) = \frac{24 - 7i}{500}.$$

The pole at 1 is of order 3, so by (7), the residue of r at 1 is the value at 1 of one-half of the second derivative of $(z + 1)/(z^2 + 4)$. This works out to be

$$\text{Res}(r; 1) = \frac{-12}{125}. \quad \square$$

Example 6 Find the residue of $\cot \alpha z$ at $z_0 = 0$.

Solution We know that

$$\cot(\alpha z) = \frac{\cos(\alpha z)}{\sin(\alpha z)}$$

and

$$\sin(\alpha z) = \left(\frac{\sin(\alpha z)}{z} \right) z,$$

where $\lim_{z \rightarrow 0} (\sin \alpha z)/z = \alpha$. Hence, the residue of $\cot(\alpha z)$ at $z_0 = 0$ is

$$\begin{aligned} \text{Res}(\cot(\alpha z); 0) &= \text{value of } \left(\frac{\cos(\alpha z)}{\sin(\alpha z)} \right) \text{ at } z = 0 \\ &= \frac{1}{\alpha}. \end{aligned} \quad \square$$

Laurent* Series

We know that a function that is analytic in the disc $|z - z_0| < R$ may be expanded there in a power series and that, conversely, if a function has a power series valid in the disc $|z - z_0| < R$, it is analytic in this disc. But what about a function f that is analytic only in the punctured disc $0 < |z - z_0| < R$ or, even worse, analytic only in the annulus $0 \leq r < |z - z_0| < R$? We shall show here that something almost as good as a power series can be given to represent $f(z)$, $r < |z - z_0| < R$. Namely, we shall show that

$$f(z) = f_1(z) + f_2(z), \quad r < |z - z_0| < R,$$

* Pierre Alphonse Laurent, 1813–1854.

where f_1 is analytic on the disc $|z - z_0| < R$ and f_2 is analytic on the region $r < |z - z_0|$, including at ∞ . f_1 has a power series in the variable $z - z_0$ which is valid for $|z - z_0| < R$, while f_2 has a power series in the variable $(z - z_0)^{-1}$ which is valid for $r < |z - z_0|$ (see Exercise 23, Section 4). Consequently,

$$f(z) = \sum_{k=0}^{\infty} a_k(z - z_0)^k + \sum_{k=1}^{\infty} b_k(z - z_0)^{-k},$$

or

$$f(z) = \sum_{k=-\infty}^{\infty} a_k(z - z_0)^k, \quad a_{-k} = b_k, \quad k = 1, 2, \dots, \quad r < |z - z_0| < R.$$

This representation of f is its **Laurent series**. Our discussion will include formulas for the coefficients a_k . In one special case, we shall show that these coefficients are given in terms of the residues of f .

Fix a point z in the annulus (even if $r = 0$, we use this terminology), and choose r_1 and R_1 with

$$r < r_1 < |z - z_0| < R_1 < R.$$

Let Γ be the circle $|w - z_0| = R_1$ (oriented counterclockwise), and let γ be the circle $|w - z_0| = r_1$ (oriented clockwise). Now consult Figure 2.14. The closed curve beginning at P and consisting of Γ followed by the horizontal segment PQ , followed by γ , and ended by the segment QP surrounds z . We may apply Cauchy's Formula, therefore, and write

$$f(z) = \frac{1}{2\pi i} \int_{\Gamma} \frac{f(w)}{w - z} dw + \frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

(Note that the line integral over PQ is cancelled by that over QP .) For $w \in \Gamma$, we




Figure 2.14

may write

$$\begin{aligned}\frac{1}{w-z} &= \frac{1}{(w-z_0)-(z-z_0)} = \frac{1}{w-z_0} \left(\frac{1}{1 - \frac{z-z_0}{w-z_0}} \right) \\ &= \sum_{k=0}^{\infty} \frac{(z-z_0)^k}{(w-z_0)^{k+1}}.\end{aligned}$$

The series is absolutely convergent, since

$$\left| \frac{z-z_0}{w-z_0} \right| < 1.$$

In an entirely similar way, we obtain

$$\frac{1}{\zeta-z} = - \sum_{j=0}^{\infty} \frac{(\zeta-z_0)^j}{(z-z_0)^{j+1}}$$

for each $\zeta \in \gamma$; here again, the series is absolutely convergent, since

$$\left| \frac{\zeta-z_0}{z-z_0} \right| < 1.$$

In the integral representation of f given before, replace $1/(w-z)$ and $1/(\zeta-z)$ with the series obtained above. After interchanging the summation and integration, we find that

$$\begin{aligned}f(z) &= \sum_{k=0}^{\infty} (z-z_0)^k \left\{ \frac{1}{2\pi i} \int_{\Gamma} \frac{f(w)}{(w-z_0)^{k+1}} dw \right\} \\ &\quad + \sum_{j=0}^{\infty} (z-z_0)^{-j-1} \left\{ \frac{-1}{2\pi i} \int_{\gamma} f(\zeta) (\zeta-z_0)^j d\zeta \right\} \\ &= f_1(z) + f_2(z).\end{aligned}$$

The integral $\int_{|w-z_0|=s} f(w)(w-z_0)^k dw$, $k = 0, \pm 1, \pm 2, \dots$ is independent of s for $r < s < R$, since the integrand is analytic on the annulus $r < |w-z_0| < R$. Let

$$a_k = \frac{1}{2\pi i} \int_{|w-z_0|=s} \frac{f(w)}{(w-z_0)^{k+1}} dw, \quad k = 0, \pm 1, \dots \quad (9)$$

Then

$$f_1(z) = \sum_{k=0}^{\infty} a_k (z-z_0)^k, \quad f_2(z) = \sum_{k=-\infty}^{-1} a_k (z-z_0)^k$$

are analytic on the sets $\{z: |z - z_0| < R\}$ and $\{z: |z - z_0| > r\}$, respectively, and we obtain the desired formula:

$$f(z) = f_1(z) + f_2(z) = \sum_{-\infty}^{\infty} a_k(z - z_0)^k, \quad r < |z - z_0| < R. \quad (10)$$

The Laurent series can, of course, be computed using (9). But in at least one important case, it may be found by an alternate, and usually easier, technique. Specifically, if f is analytic in the punctured disc $0 < |z - z_0| < R$ and has a pole of order m at z_0 , the Laurent series for f has already been found in (6), although with different notation:

$$f(z) = \frac{c_0}{(z - z_0)^m} + \cdots + \frac{c_{m-1}}{z - z_0} + c_m + c_{m+1}(z - z_0) + \cdots,$$

$$0 < |z - z_0| < R,$$

where we have written $H(z)$ in its power series,

$$H(z) = \sum_{k=0}^{\infty} c_k(z - z_0)^k,$$

which is valid on the disc $|z - z_0| < R$ and where $c_0 \neq 0$. All that remains to do is to rewrite the coefficients in the manner $a_k = c_{k+m}$, $k = -m, -m + 1, \dots$. When a function f has a pole at z , the sum of the terms involving negative powers of $z - z_0$ is called the **principal part** of f at z_0 . This is exactly $P(1/(z - z_0))$, where P is a polynomial of degree m with constant term equal to zero.

Example 7 Find the principal part and residue of $f(z) = (z^3 + z^2)/(z - 1)^2$ at $z_0 = 1$.

Solution We expand $z^3 + z^2$ in powers of $z - 1$:

$$z^3 + z^2 = 2 + 5(z - 1) + 4(z - 1)^2 + (z - 1)^3.$$

Division then yields the Laurent series for f :

$$f(z) = \frac{2}{(z - 1)^2} + \frac{5}{z - 1} + 4 + (z - 1).$$

The principal part is $2(z - 1)^{-2} + 5(z - 1)^{-1}$, and the residue at 1 is 5. □

Example 8 Compute the Laurent series at $z_0 = 0$ for $f(z) = (\sin z)/z^3$.

Solution We use the expansion of $\sin z$ about $z_0 = 0$ and then divide by z^3 .

$$f(z) = \frac{\sin z}{z^3} = \frac{z - \frac{z^3}{3!} + \frac{z^5}{5!} - \cdots}{z^3}$$

$$= \frac{1}{z^2} - \frac{1}{3!} + \frac{z^2}{5!} - \cdots, \quad z \neq 0.$$

The principal part is $1/z^2$, and the residue at zero is 0. \square

Example 9 Find the first five terms of the Laurent series about $z_0 = 0$ for the function $g(z) = \cot z$.

Solution The function $\cot z$ is a quotient:

$$g(z) = \cot z = \frac{\cos z}{\sin z}.$$

$\sin z$ has a zero of order one at $z_0 = 0$, and $\cos 0 = 1$. Hence, g has a pole of order one at the origin. This means that the Laurent series for g has the form

$$g(z) = \frac{a_{-1}}{z} + a_0 + a_1 z + a_2 z^2 + \cdots.$$

To find the coefficients a_k , we write $(\sin z)g(z) = \cos z$, multiply out the resulting series, collect equal powers of z , and equate coefficients. Thus,

$$\begin{aligned} \{\sin z\} \{g(z)\} &= \left\{ z - \frac{z^3}{3!} + \frac{z^5}{5!} - \cdots \right\} \left\{ \frac{a_{-1}}{z} + a_0 + a_1 z + \cdots \right\} \\ &= a_{-1} + a_0 z + \left(a_1 - \frac{a_{-1}}{3!} \right) z^2 + \left(a_2 - \frac{a_0}{3!} \right) z^3 \\ &\quad + \left(a_3 - \frac{a_1}{3!} + \frac{a_{-1}}{5!} \right) z^4 + \cdots \\ &= \cos z = 1 - \frac{z^2}{2!} + \frac{z^4}{4!} - \cdots. \end{aligned}$$

After equating coefficients of equal powers of z , the resulting equations are

$$\begin{aligned} a_{-1} &= 1, \quad a_0 = 0, \quad \left(a_1 - \frac{a_{-1}}{3!} \right) = -\frac{1}{2}, \quad \left(a_2 - \frac{a_0}{3!} \right) = 0, \\ \left(a_3 - \frac{a_1}{3!} + \frac{a_{-1}}{5!} \right) &= \frac{1}{4!}, \end{aligned}$$

which may be solved successively for a_{-1} , a_0 , a_1 , a_2 , and a_3 , yielding

$$a_{-1} = 1, \quad a_0 = 0, \quad a_1 = -\frac{1}{3}, \quad a_2 = 0, \quad \text{and} \quad a_3 = -\frac{1}{45}.$$

Hence,

$$\cot z = \frac{1}{z} - \left(\frac{1}{3}\right)z - \left(\frac{1}{45}\right)z^3 + \cdots, \quad 0 < |z| < \pi. \quad \square$$

Example 10 Find the Laurent series about $z_0 = 0$ of $h(z) = z \cos(1/z)$.

Solution In this case, the origin is not a pole but rather an essential singularity. Nonetheless, the Laurent series is rather easy to find. We may, of course, use (9). However, we also know an expansion for $\cos w$ valid for any complex number w (see Example 7, Section 2). In this expansion, just replace w with $1/z$. This will give

$$\begin{aligned} h(z) &= z \cos \frac{1}{z} \\ &= z \left\{ 1 - \left(\frac{1}{2!}\right)\left(\frac{1}{z^2}\right) + \left(\frac{1}{4!}\right)\left(\frac{1}{z^4}\right) - \cdots \right\} \\ &= z - \left(\frac{1}{2!}\right)\left(\frac{1}{z}\right) + \left(\frac{1}{4!}\right)\left(\frac{1}{z^3}\right) - \cdots. \end{aligned} \quad \square$$

Example 11 Find the Laurent series on the annulus $1 < |z| < 4$ for the function $R(z) = (z + 2)/(z^2 - 5z + 4)$.

Solution The rational function R has two poles each of order one, the first at $z_1 = 1$ and the second at $z_2 = 4$. The Laurent series of R may be found by writing

$$R(z) = R_1(z) + R_2(z),$$

where R_1 is analytic for $|z| < 4$ and R_2 is analytic for $1 < |z|$. In this case, this amounts to a partial-fractions decomposition of R , a procedure familiar from calculus:

$$R(z) = \frac{z + 2}{(z - 1)(z - 4)} = \frac{A}{z + 4} + \frac{B}{z - 1}.$$

The unknowns A and B are found by clearing the denominators:

$$A(z - 1) + B(z - 4) = z + 2.$$

This yields $A = 2$, $B = -1$. We now employ the geometric series and obtain

$$\begin{aligned}\frac{z+2}{(z-1)(z-4)} &= \frac{2}{z-4} + \frac{-1}{z-1} \\ &= \left(-\frac{1}{2}\right) \sum_{k=0}^{\infty} \left(\frac{z}{4}\right)^k - \sum_{j=1}^{\infty} \left(\frac{1}{z}\right)^j \\ &= \sum_{-\infty}^{\infty} a_k z^k,\end{aligned}$$

with

$$a_k = \begin{cases} \left(-\frac{1}{2}\right) 4^{-k}, & k = 0, 1, 2, \dots \\ -1, & k = -1, -2, -3, \dots \end{cases}$$
□

We return to our discussion of the principal part of a function with poles. Suppose that f is analytic on some domain D except for a pole of order m at a point z_0 of D . Then f may be written as

$$f(z) = P\left(\frac{1}{z-z_0}\right) + g(z),$$

where P is a polynomial of degree m with zero constant term and g is analytic on D and also at z_0 . $P(1/(z-z_0))$ is the principal part of f at z_0 and is analytic on the whole plane except at z_0 ; it is also analytic at ∞ . Repeated applications of this procedure produce the following result.

If f is analytic on a domain D except for poles at the points z_1, \dots, z_n of orders m_1, \dots, m_n , respectively, then

$$f(z) = P_1\left(\frac{1}{z-z_1}\right) + \dots + P_n\left(\frac{1}{z-z_n}\right) + g(z), \quad z \in D, \quad (11)$$

where g is analytic on D and P_1, \dots, P_n are polynomials of degree m_1, \dots, m_n , respectively, each with constant term equal to zero.

The Laurent Series of a Rational Function in Powers of z and $1/z^*$

The Laurent series of a rational function is important in applications, for instance, in connection with the Z-transform discussed in Section 5 of Chapter 5. Here we discuss one particularly simple but useful case. Let R be a rational function all of whose poles in the plane have order one and which has no pole at the origin. After an initial long division, we can write

$$R(z) = S(z) + \frac{P(z)}{Q(z)},$$

where S , P , and Q are polynomials and the degree of P is strictly less than the degree of Q . Since S is already expressed as the sum of powers of z , we concentrate on expanding $f = P/Q$. Here we shall show the following.

Let f be a rational function all of whose poles z_1, \dots, z_N in the plane have order one and which has no pole at the origin and which is zero at ∞ .

Suppose that no pole of f lies in the annular region $r < |z| < R$. Then

$$f(z) = \sum_{-\infty}^{\infty} a_k z^k \quad \text{for} \quad r < |z| < R,$$

where

$$a_k = \begin{cases} \sum_{|z_j| < r} z_j^{k-1} \operatorname{Res}(f; z_j), & k \leq -1 \\ -\sum_{|z_j| > R} z_j^{-k-1} \operatorname{Res}(f; z_j), & k \geq 0. \end{cases} \quad (12)$$

$$(13)$$

The chief tool to establish (12) and (13) is (11). In that formula, we see that since f is rational, so must g be. However, g has no poles and has the value zero at ∞ ; hence, g is identically zero. Let z_1, \dots, z_m be the poles of f in the disc $\{z: |z| < r\}$, and let z_{m+1}, \dots, z_N be the remaining poles of f , which necessarily lie in the region $\{z: |z| > R\}$. From (11), we see that $f = F_1 + F_2$, where

$$F_1(z) = \sum_{j=m+1}^N P_j \left(\frac{1}{z - z_j} \right) \quad \text{and} \quad F_2(z) = \sum_{j=1}^m P_j \left(\frac{1}{z - z_j} \right).$$

Consequently, F_1 is analytic on the disc $\{z: |z| < R\}$, and F_2 is analytic on the region $\{z: |z| > r\}$, including at ∞ . Thus, $f = F_1 + F_2$ is the Laurent series of f in the annulus $\{z: r < |z| < R\}$. What remains, then, is to find the explicit formula for the coefficients a_k in the Laurent series

$$f(z) = \sum_{-\infty}^{\infty} a_k z^k.$$

This result may be obtained either by the use of the Residue Theorem or by a more elementary technique. Here we shall follow the latter path; the former is left as an exercise.

Since each of the poles of f has order one (by assumption), the polynomial P_j in (11) has degree one. Also, P_j has no constant term, so

$$P_j(z) = \frac{b_j}{z - z_j}, \quad b_j = \operatorname{Res}(f; z_j).$$

For $j = 1, \dots, m$, the point z_j satisfies $|z_j| < r$. Hence, using the geometric series for

$$\frac{1}{1 - \frac{z_j}{z}},$$

we obtain

$$\frac{1}{z - z_j} = \frac{1}{z} \cdot \frac{1}{1 - \frac{z_j}{z}} = \sum_{k=0}^{\infty} \frac{z_j^k}{z^{k+1}}.$$

In an entirely similar manner, for $j = m + 1, \dots, N$,

$$\frac{1}{z - z_j} = \frac{-1}{z_j} \cdot \frac{1}{1 - \frac{z}{z_j}} = -\sum_{k=0}^{\infty} \frac{z^k}{z_j^{k+1}}.$$

This gives (12) and (13) after summing on j .

Example 12 Find the expansion $R(z) = (z^3 - 3z^2 + 3)/(z - 1)(z - 3)$ in powers of z and $1/z$ in the regions (a) $|z| < 1$; (b) $1 < |z| < 3$; (c) $|z| > 3$.

Solution The initial long division gives $R(z) = (z + 1) + z/(z - 1)(z - 3)$. We concentrate on expanding $f(z) = z/(z - 1)(z - 3)$.

(a) In this case, $r = 0$, $R = 1$, and there are no terms to use in (12), so only (13) is used, and we obtain

$$\frac{z}{(z - 1)(z - 3)} = \sum_{k=0}^{\infty} a_k z^k, \quad a_k = -\left[-\frac{1}{2} + \frac{3}{2} \frac{1}{3^{k+1}} \right], \quad k \geq 0.$$

(b) Here, $r = 1$ and $R = 3$; thus, directly from (12) and (13),

$$\frac{z}{(z - 1)(z - 3)} = \sum_{k=-\infty}^{\infty} a_k z^k, \quad a_k = \begin{cases} -\frac{1}{2}, & k \leq -1 \\ -\frac{3}{2} \frac{1}{3^{k+1}}, & k \geq 0. \end{cases}$$

(c) Here, $r = 3$, and R does not enter the picture (or, if you prefer, is ∞), so there are no terms to use in (13). Thus,

$$\frac{z}{(z - 1)(z - 3)} = \sum_{k=-\infty}^{-1} a_k z^k, \quad a_k = -\frac{1}{2} + \frac{3}{2} 3^{k-1}, \quad k \leq -1. \quad \square$$

You may recognize that this technique for finding the Laurent series is really the partial-fractions technique, familiar from calculus.

EXERCISES FOR SECTION 2.5

In Exercises 1 to 6, locate each of the isolated singularities of the given function and tell whether it is a removable singularity, a pole, or an essential singularity (case (3)). If the singularity is removable, give the value of the function at the point; if the singularity is a pole, give the order of the pole.

1. $\frac{e^z - 1}{z}$

2. $\frac{z^2}{\sin z}$

3. $\frac{z^4 - 2z^2 + 1}{(z - 1)^2}$

4. $\pi \cot \pi z$

5. $\frac{2z + 1}{z + 2}$

6. $\frac{e^z - 1}{e^{2z} - 1}$

In Exercises 7 to 13, find the Laurent series for the given function about the indicated point. Also, give the residue of the function at the point.

7. $\frac{e^z - 1}{z^2}; \quad z_0 = 0 \quad 8. \frac{z^2}{z^2 - 1}; \quad z_0 = 1 \quad 9. \frac{\sin z}{(z - \pi)^2}; \quad z_0 = \pi$

10. $\frac{z}{(\sin z)^2}; \quad z_0 = 0$ (four terms of the Laurent series)

11. $\frac{az + b}{cz + d}; \quad z_0 = -\frac{d}{c}, \quad c \neq 0$

12. $\frac{1}{e^z - 1}; \quad z_0 = 0$ (four terms of the Laurent series)

13. $\frac{1}{1 - \cos z}; \quad z_0 = 0$ (four terms of the Laurent series)

14. If f is analytic in $|z - z_0| < R$ and has a zero of order m at z_0 , show that

$$\text{Res}\left(\frac{f'}{f}; z_0\right) = m.$$

15. If f is analytic in $0 < |z - z_0| < R$ and has a pole of order l at z_0 , show that

$$\text{Res}\left(\frac{f'}{f}; z_0\right) = -l.$$

(Hint: Write

$$f(z) = \frac{g(z)}{(z - z_0)^l},$$

where g is analytic in $|z - z_0| < R$ and $g(z_0) \neq 0$.)

16. Suppose f is analytic in $|z - z_0| < R$ and has a zero of order $m \geq 1$ at z_0 . If g is analytic in $0 < |z - z_0| < R$ and has a pole of order l , $l \leq m$, at z_0 , show that fg has a removable singularity at z_0 .
17. Let f be analytic in $0 < |z - z_0| < r$ and suppose that f has an essential singularity at z_0 . Let w be any complex number. Show that

$$g(z) = \frac{1}{f(z) - w}, \quad z \in D,$$

is *not* bounded in any punctured disc $0 < |z - z_0| < \varepsilon$. (**Hint:** if g is bounded, then show that f has a pole or a removable singularity at z_0 .) Conclude that the range of f gets arbitrarily close to all points in the complex plane. How does this differ from the situation in (1) and (2)?

18. Here is an alternate proof that

$$\frac{1}{2\pi i} \int_{|z-z_0|=s} f(z) dz = \text{Res}(f; z_0)$$

is independent of s . Assume $z_0 = 0$ with no loss of generality.

- (a) Show that

$$\frac{\partial}{\partial t} \{f(se^{it})e^{it}\} = ie^{it}f(se^{it}) + ise^{2it}f'(se^{it}).$$

- (b) Show that

$$\frac{d}{ds} \left\{ \frac{1}{2\pi} \int_0^{2\pi} sf(se^{it})e^{it} dt \right\} = \frac{1}{2\pi} \int_0^{2\pi} [f(se^{it}) + sf'(se^{it})e^{it}] e^{it} dt.$$

- (c) Conclude from (a) and (b) that

$$\frac{d}{ds} \left\{ \frac{1}{2\pi i} \int_{|z|=s} f(z) dz \right\} = \frac{1}{2\pi i} \int_0^{2\pi} \frac{\partial}{\partial t} \{e^{it}f(se^{it})\} dt.$$

Finally, calculate that the integral on the right has the value 0.

19. Suppose that the Laurent series $\sum_{-\infty}^{\infty} a_n(z - z_0)^n$ converges for $r < |z - z_0| < R$ and

$$\sum_{-\infty}^{\infty} a_n(z - z_0)^n = 0, \quad 0 < |z - z_0| < r.$$

Show that $a_n = 0$, $n = 0, \pm 1, \pm 2, \dots$ (**Hint:** Multiply the series by $(z - z_0)^{-m}$ and integrate around the circle $|z - z_0| = s$, $r < s < R$ with respect to z . The result must be zero, but it is also a_{m-1} .)

20. If f is analytic in $0 \leq r < |z - z_0| < R$, show that its Laurent series is uniquely determined.
 21. If F is analytic on $0 < |z - z_0| < R$ and if there is an analytic function G on $0 < |z - z_0| < R$ with $G' = F$, then

$$\text{Res}(F; z_0) = 0.$$

(**Hint:** Use (5).)

22. Find the Laurent series about $z_0 = 0$ for the following functions, valid in the indicated regions.
- $e^{1/z}$ in $0 < |z| < \infty$
 - $z^4 \sin\left(\frac{1}{z}\right)$ in $0 < |z| < \infty$
 - $\frac{1}{z-1} - \frac{1}{z+1}$ in $2 < |z| < \infty$
 - $\exp\left(z + \frac{1}{z}\right)$ in $0 < |z| < \infty$
 - $z \cos\left(\frac{1}{z}\right)$ in $0 < |z| < \infty$
23. Use equations (12) and (13) to find the Laurent expansions of the following rational functions in powers of z and $1/z$ in the indicated region(s).
- $f(z) = \frac{z+2}{z^2 - z - 2}$ in $1 < |z| < 2$ and then in $2 < |z| < \infty$
 - $f(z) = \frac{z^2 + 2z - 4}{(z^2 - 9)(z + 1)}$ in $|z| < 1$ and then in $1 < |z| < 3$
 - $f(z) = \frac{z+1}{(z-2)(z-3)(z-5)}$ in $2 < |z| < 3$ and then in $5 < |z| < \infty$
24. Let f be analytic in $0 < |z - z_0| < r$, and let

$$f(z) = \sum_{n=-\infty}^{\infty} a_n(z - z_0)^n$$

be its Laurent series. Show that

- z_0 is a removable singularity for f if and only if $a_n = 0$ for $n = -1, -2, \dots$
- z_0 is a pole of order $m \geq 1$ for f if and only if $a_{-m} \neq 0$ but $a_{-n} = 0$ for all $n \geq m + 1$.
- z_0 is an essential singularity for f if and only if there are infinitely many a_{-n} , $n > 0$, that are *not* zero.

Singularities at ∞ *

Suppose that f is analytic for $|z| > R$; set $g(z) = f(1/z)$ so that g is analytic for $0 < |z| < 1/R$. We classify the nature of the singularity of f at ∞ by that of g at $z = 0$; for instance, if g has a pole of order m at 0, we say f has a pole of order m at ∞ . The Laurent series for f about ∞ is just the Laurent series for $g(z) = f(1/z)$ about 0, with z replaced by $1/z$.

- Show that f has a removable singularity at ∞ if and only if $|f(z)|$ is bounded for $|z| > R_0$.
- Show that f has a pole of order m at ∞ if and only if

$$h(z) = \frac{f(z)}{z^m}$$

has a removable singularity at ∞ and $h(\infty) \neq 0$.

27. Classify the nature of the singularity at ∞ of each of the following functions. If the singularity is removable, give the value at ∞ . If the singularity is a pole, give its order; in each case find the first few terms in the Laurent series about ∞ .

$$\begin{array}{lll} \text{(a)} \quad 3z^2 + 4 - \frac{1}{z} & \text{(b)} \quad (1-z)(z-4) & \text{(c)} \quad \frac{z^2}{z-4} \\ \text{(d)} \quad (-z^2 - 2z + 3)^{-1} & \text{(e)} \quad e^{1/z} & \text{(f)} \quad e^{-z^2} \\ \text{(g)} \quad \left(\frac{1}{z} + z\right)^3 & \text{(h)} \quad \sin \frac{1}{z} & \text{(i)} \quad \sum_{n=0}^{\infty} (-1)^n \frac{z^{-2n}}{(2n)!} \end{array}$$

Bessel Functions (Continued)*

28. Let u be a complex number and let

$$G(z; u) = \exp\left[\left(\frac{u}{2}\right)\left(z - \frac{1}{z}\right)\right], \quad z \neq 0.$$

Show that $G(z; u)$ is an analytic function of z for $z \neq 0$ and has an essential singularity at $z = 0$ (unless $u = 0$).

29. Let

$$G(z; u) = \sum_{-\infty}^{\infty} J_n(u) z^n, \quad (14)$$

be the Laurent series of $G(z; u)$ about the origin, where the coefficients $\{J_n(u)\}$ are given by (9). In (9), choose $s = 1$ and conclude that

$$J_n(u) = \frac{1}{2\pi} \int_0^{2\pi} \cos(u \sin \theta - n\theta) d\theta. \quad (15)$$

(Hint: Prove (15) for u real by showing that $J_n(u)$ is real if u is real.)

30. Multiply out the series for $\exp(uz/2)$ and the series for $\exp(-u/2z)$ and then collect equal powers of z . Conclude that

$$J_n(u) = \frac{u^n}{2^n} \sum_{k=0}^{\infty} \frac{(-1)^k u^{2k}}{2^{2k} k!(n+k)!}$$

for $n \geq 0$. Compare this to the conclusion of Exercise 29, Section 4.

31. Show that $J_{-n}(u) = (-1)^n J_n(u)$. (Hint: Replace z with $-1/z$ in the Laurent series; then change the summation index n to $-n$ and compare series.)

2.6 The Residue Theorem and Its Application to the Evaluation of Definite Integrals

The Residue Theorem is of great theoretical and practical importance. It allows us to compute with ease certain definite integrals, which at first glance seem to be very

difficult and which are surely not solvable by the techniques of elementary calculus. At the same time, it provides the key in unlocking the secret to counting the zeros of an analytic function. In this section, we concentrate on its application to the evaluation of definite integrals. Later sections will contain other applications.

THEOREM 1 The Residue Theorem Suppose that f is analytic on a simply-connected domain D except for a finite number of isolated singularities at points z_1, \dots, z_N of D . Let γ be a piecewise smooth positively oriented simple closed curve in D that does not pass through any of the points z_1, \dots, z_N . Then

$$\int_{\gamma} f(z) dz = 2\pi i \sum_{z_k \text{ inside } \gamma} \operatorname{Res}(f; z_k),$$

where the sum is taken over all those singularities z_k of f that lie inside γ .

Proof Let $\gamma_1, \dots, \gamma_N$ be disjoint positively oriented circles centered at z_1, \dots, z_N , respectively, chosen to be so small that the discs they bound are disjoint. We can apply Green's Theorem to the function f on the domain Ω whose boundary is γ and $\gamma_1, \dots, \gamma_N$. Since f is analytic on Ω , the value of the line integral is zero. Therefore,

$$0 = \int_{\gamma} f(z) dz - \sum_{k=1}^N \int_{\gamma_k} f(z) dz.$$

However, $\int_{\gamma_k} f(z) dz = 2\pi i \operatorname{Res}(f; z_k)$. This finishes the proof. ■

The remainder of this section is devoted to uses of the Residue Theorem in the evaluation of definite integrals.

Integrals of Rational Functions

It is convenient to isolate here several simple facts about a polynomial

$$p(z) = a_n z^n + a_{n-1} z^{n-1} + \cdots + a_1 z + a_0, \quad a_n \neq 0.$$

If $p(\alpha) = 0$, then $p(z) = (z - \alpha)q(z)$, where q is a polynomial of degree $n - 1$. (1)

There are at most n distinct points $\alpha_1, \dots, \alpha_s$ in the plane at which $p(\alpha_j) = 0$. (2)

For $|z| = R$, R large, we have the estimate

$$\frac{1}{2} |a_n| R^n \leq |p(z)| \leq 2 |a_n| R^n. \quad (3)$$

The proofs of these three facts are all easy. For (1), expand p in powers of $(z - \alpha)$; this yields

$$p(z) = \sum_0^n c_j(z - \alpha)^j, \quad c_j = \frac{p^{(j)}(\alpha)}{j!}, \quad j = 0, \dots, n.$$

However, $c_0 = p(\alpha) = 0$, so

$$\begin{aligned} p(z) &= c_1(z - \alpha) + \cdots + c_n(z - \alpha)^n \\ &= (z - \alpha)[c_1 + \cdots + c_n(z - \alpha)^{n-1}] \\ &= (z - \alpha)q(z). \end{aligned}$$

Next, (2) follows from (1), since each time a zero is divided out, the degree of the quotient goes down by one. Finally,

$$\lim_{|z| \rightarrow \infty} \left| \frac{p(z)}{z^n} \right| = |a_n| > 0.$$

Consequently, for $|z| = R$ large enough,

$$\frac{1}{2}|a_n| \leq \left| \frac{p(z)}{z^n} \right| \leq 2|a_n|,$$

which is exactly (3).

The following proposition shows how the Residue Theorem is applied to evaluate controls of certain rational functions.

PROPOSITION Suppose P and Q are polynomials that are real-valued on the real axis and for which the degree of Q exceeds the degree of P by 2 or more. If $Q(x) \neq 0$ for all real x , then

$$\int_{-\infty}^{\infty} \frac{P(x)}{Q(x)} dx = 2\pi i \sum_U \text{Res}\left(\frac{P}{Q}; z_j\right),$$

where the sum is taken over all poles of P/Q that lie in the upper half-plane $U = \{z: \text{Im } z > 0\}$.

Proof Let γ_R be the semicircular contour in Figure 2.15. There is no loss of generality in assuming that $Q(z) \neq 0$ for $z \in \gamma_R$. On γ_R , when R is big, (3) gives




Figure 2.15

$$\left| \frac{P(z)}{Q(z)} \right| \leqslant \frac{2|a_n|R^n}{\frac{1}{2}|b_m|R^m} \leqslant 4 \left| \frac{a_n}{b_m} \right| R^{-2} = \frac{C}{R^2},$$

since $m \geq n + 2$. Hence,

$$\left| \int_{|z|=R, y>0} \frac{P(z)}{Q(z)} dz \right| \leq C \frac{2\pi R}{R^2} \rightarrow 0 \quad \text{as } R \rightarrow \infty.$$

The result now follows from the Residue Theorem. ■

Example 1 Compute

$$\int_{-\infty}^{\infty} \frac{x^2}{(1+x^2)(4+x^2)} dx.$$

Solution The polynomials are $P(z) = z^2$ and $Q(z) = (1+z^2)(4+z^2)$, respectively, and Q has zeros at $z_1 = i$ and $z_2 = 2i$ in U , the upper half-plane. Now

$$\frac{P(z)}{Q(z)} = \frac{z^2}{(z-i)(z+i)(z-2i)(z+2i)}.$$

Example 4 of Section 5 then gives

$$\operatorname{Res}\left(\frac{P}{Q}; i\right) = \frac{-1}{(2i)^3} = \frac{-1}{6i}$$

and

$$\operatorname{Res}\left(\frac{P}{Q}; 2i\right) = \frac{-4}{(-3)(4i)} = \frac{1}{3i}.$$

Hence,

$$\int_{-\infty}^{\infty} \frac{x^2}{(1+x^2)(4+x^2)} dx = 2\pi i \left[\frac{-1}{6i} + \frac{1}{3i} \right] = \frac{\pi}{3}. \quad \square$$

Example 2 Compute

$$\int_{-\infty}^{\infty} \frac{dx}{(1+x^2)^2}.$$

Solution Here $P(z) = 1$ for all z and $Q(z) = (1+z^2)^2 = (z+i)^2(z-i)^2$. Hence, Q has just one zero in the upper half-plane U , at $z_0 = i$, but that zero has order two. According to (7) of Section 5, the residue of $1/Q$ at $z_0 = i$ is the value at i of the

derivative of $g(z) = (z + i)^{-2}$. This is easily computed to be $1/4i$. Consequently,

$$\int_{-\infty}^{\infty} \frac{1}{(1+x^2)^2} dx = 2\pi i \left(\frac{1}{4i} \right) = \frac{\pi}{2}. \quad \square$$

Integrals over the Real Axis Involving Trigonometric Functions

For integrals of the form $\int_{-\infty}^{\infty} R(x) \sin x dx$ or $\int_{-\infty}^{\infty} R(x) \cos x dx$, where R is a rational function that is real-valued on the real axis, we apply the Residue Theorem to the function $f(z) = R(z)e^{iz}$ and, at the end of the computation, take the imaginary or real part of the resulting complex number.

Example 3 Compute

$$\int_{-\infty}^{\infty} \frac{\cos x}{x^2 + \alpha^2} dx, \quad \alpha > 0.$$

Solution Set

$$f(z) = \frac{e^{iz}}{z^2 + \alpha^2};$$

f has a pole at $i\alpha$ in U with residue

$$\text{Res}(f; i\alpha) = \frac{e^{-\alpha}}{2i\alpha}.$$

Let γ_R be the contour in Figure 2.15, so

$$\begin{aligned} \pi \frac{e^{-\alpha}}{\alpha} &= 2\pi i \left(\frac{e^{-\alpha}}{2i\alpha} \right) = \int_{\gamma_R} f(z) dz \\ &= \int_{-R}^R \frac{e^{ix}}{x^2 + \alpha^2} dx + i \int_0^\pi f(Re^{i\theta}) Re^{i\theta} d\theta. \end{aligned}$$

The integral over the semicircle can be estimated by

$$\begin{aligned} \left| \int_0^\pi f(Re^{i\theta}) Re^{i\theta} d\theta \right| &\leq R \int_0^\pi \frac{e^{-R \sin \theta}}{R^2 - \alpha^2} d\theta \\ &\leq \frac{\pi R}{R^2 - \alpha^2} \rightarrow 0 \quad \text{as } R \rightarrow \infty. \end{aligned}$$

Thus,

$$\frac{\pi e^{-\alpha}}{\alpha} = \int_{-\infty}^{\infty} \frac{\cos x}{x^2 + \alpha^2} dx + i \int_{-\infty}^{\infty} \frac{\sin x}{x^2 + \alpha^2} dx.$$

The integral with $\sin x$ is zero since the integrand is odd; hence,

$$\int_{-\infty}^{\infty} \frac{\cos x}{x^2 + \alpha^2} dx = \pi \frac{e^{-\alpha}}{\alpha}, \quad \alpha > 0.$$
□

Example 4 In the foregoing result, replace x with βx and set $\delta = \alpha/\beta$. This results in the formula

$$\int_{-\infty}^{\infty} \frac{\cos \beta x}{x^2 + \delta^2} dx = \left(\frac{\pi}{\delta}\right) e^{-\delta\beta}, \quad \beta, \delta > 0. \quad (4)$$

Other integral formulas may be derived formally from (4). For instance, differentiation of both sides of (4) with respect to β yields

$$\int_{-\infty}^{\infty} \frac{x \sin \beta x}{x^2 + \delta^2} dx = \pi e^{-\delta\beta}, \quad \beta, \delta > 0. \quad (5)$$

(The formalities are justifiable but will not be shown here.) A further differentiation, this time on δ , gives

$$\int_{-\infty}^{\infty} \frac{x \sin \beta x}{(x^2 + \delta^2)^2} dx = \left(\frac{\pi}{2\delta}\right) e^{-\delta\beta}.$$

Another formula comes from (5) by taking $\beta = 1$ and letting $\delta \rightarrow 0$:

$$\int_{-\infty}^{\infty} \frac{\sin x}{x} dx = \pi.$$

(Note: This last formula is only a formality at this point.)

□

In these applications of the Residue Theorem, we often need to estimate the magnitude of the line integral of e^{iz} over the semicircle $z = Re^{i\theta}$, $0 \leq \theta \leq \pi$. To do this, we note that

$$|e^{iz}| = e^{Re(iz)} = e^{-R \sin \theta}.$$

Further, elementary calculus shows that $\sin \theta \geq (2/\pi)\theta$ for $0 \leq \theta \leq \pi/2$. Hence, for $0 \leq \theta \leq \pi/2$,

$$|e^{iz}| = e^{-R \sin \theta} \leq e^{-(2R/\pi)\theta}.$$

Therefore,

$$\begin{aligned} \left| \int_{|z|=R, \operatorname{Im} z \geq 0} e^{iz} dz \right| &= \left| \int_0^\pi e^{iRe^{i\theta}} iRe^{i\theta} d\theta \right| \leq R \int_0^\pi e^{-R \sin \theta} d\theta = 2R \int_0^{\pi/2} e^{-R \sin \theta} d\theta \\ &\leq 2R \int_0^{\pi/2} e^{-(2R/\pi)\theta} d\theta = \pi(1 - e^{-R}) < \pi. \end{aligned}$$

The inequality

$$\left| \int_{\substack{|z|=R \\ \operatorname{Im} z \geq 0}} e^{iz} dz \right| < \pi$$

is often called **Jordan's Lemma**.

Example 5 Compute the value of

$$\int_{-\infty}^{\infty} \frac{x^3 \sin x}{x^4 + 16} dx.$$

Solution We set

$$f(z) = e^{iz} \frac{z^3}{z^4 + 16}$$

and integrate f over the contour in Figure 2.15. The function f has poles in U of order one at $\sqrt{2}(1+i)$ and $\sqrt{2}(-1+i)$. The residues are

$$\operatorname{Res}(f; \sqrt{2}(1+i)) = \frac{1}{4} \exp[\sqrt{2}(-1+i)]$$

and

$$\operatorname{Res}(f; \sqrt{2}(-1+i)) = \frac{1}{4} \exp[\sqrt{2}(-1+i)]$$

By Jordan's Lemma (or, better, its proof), we can make the estimate

$$\left| \int_{\substack{|z|=R \\ \operatorname{Im} z \geq 0}} f(z) dz \right| \leq \frac{R^3}{R^4 - 16} \int_0^\pi e^{-R \sin \theta} R d\theta \leq \frac{\pi R^3}{R^4 - 16} \rightarrow 0$$

as $R \rightarrow \infty$. Thus,

$$\begin{aligned} \int_{-\infty}^{\infty} \frac{x^3 \sin x}{x^4 + 16} dx &= \operatorname{Im}\{2\pi i [\operatorname{Res}(f; \sqrt{2}(1+i)) + \operatorname{Res}(f; \sqrt{2}(-1+i))]\} \\ &= \pi e^{-\sqrt{2}} \cos(\sqrt{2}). \end{aligned}$$

□

Example 6 Find the value of

$$\int_0^{\infty} \frac{\sin^2 x}{x^2} dx.$$

Solution We use the identity $2 \sin^2 x = 1 - \cos 2x = \operatorname{Re}(1 - e^{2ix})$. Set




Figure 2.16

$$f(z) = \frac{1 - e^{2iz}}{z^2}$$

and take γ to be the contour in Figure 2.16.

The integral of f over γ is zero, by Cauchy's Theorem. On the semicircle $z = Re^{it}, 0 \leq t \leq \pi$, we may estimate f by

$$|f(Re^{it})| \leq \frac{1 + e^{-2R \sin t}}{R^2} \leq \frac{2}{R^2}$$

so that the integral over the large semicircle can be bounded by

$$\left| \int_{\substack{|z|=R \\ z \in \gamma}} f(z) dz \right| \leq \frac{2}{R^2} 2\pi R \rightarrow 0 \quad \text{as } R \rightarrow \infty.$$

Furthermore, near $z = 0$, $f(z)$ has the expansion

$$\begin{aligned} f(z) &= \frac{1 - \left[1 + 2iz + \frac{(2iz)^2}{2!} + \dots \right]}{z^2} \\ &= \frac{-2i}{z} + 2 - \frac{4i}{3}z + \dots \end{aligned}$$

Hence,

$$\int_{-\pi}^0 f(\varepsilon e^{it}) i \varepsilon e^{it} dt = -2\pi + (\text{terms that go to zero as } \varepsilon \rightarrow 0).$$

Thus,

$$0 = \int_{-\varepsilon}^{\varepsilon} \frac{1 - e^{2ix}}{x^2} dx + \int_{-R}^{-\varepsilon} \frac{1 - e^{2ix}}{x^2} dx - 2\pi + E,$$

where $E \rightarrow 0$ as $R \rightarrow \infty$ and $\varepsilon \rightarrow 0$. Take the real part of the above expression:

$$2\pi = \int_{-\varepsilon}^R \frac{1 - \cos 2x}{x^2} dx + \int_{-R}^{-\varepsilon} \frac{1 - \cos 2x}{x^2} dx + \operatorname{Re}(E).$$

Now let $R \rightarrow \infty$ and $\varepsilon \rightarrow 0$ and recall that $1 - \cos 2x = 2 \sin^2 x$. The final result, then, is that

$$\frac{\pi}{2} = \int_0^\infty \left[\frac{\sin^2 x}{x^2} \right] dx. \quad \square$$

Integrals of Trigonometric Functions over $[0, 2\pi]$

Integrals of this type were discussed once already, in Section 3 of this chapter, in connection with Cauchy's Theorem. Here we can approach them more generally by allowing multiple poles or poles of order two or more.

Example 7 Compute the value of

$$\int_0^{2\pi} \frac{d\theta}{2 + \cos^2 \theta}.$$

Solution With $z = e^{i\theta}$, it follows that $d\theta = dz/iz$ and $\cos \theta = \frac{1}{2}(z + 1/z)$. This gives

$$\begin{aligned} \int_0^{2\pi} \frac{d\theta}{2 + \cos^2 \theta} &= \frac{1}{i} \int_{|z|=1} \frac{dz}{z \left(2 + \frac{1}{4} \left(z^2 + 2 + \frac{1}{z^2} \right) \right)} \\ &= 2\pi \left\{ \frac{1}{2\pi i} \int_{|z|=1} \frac{4z}{z^4 + 10z^2 + 1} dz \right\}. \end{aligned}$$

The rational function

$$R(z) = \frac{4z}{z^4 + 10z^2 + 1}$$

has two poles, each of order one, within the circle $|z| = 1$, at $z_1 = i\sqrt{5 - 2\sqrt{6}}$ and $z_2 = -i\sqrt{5 - 2\sqrt{6}}$. The residues are, respectively,

$$\begin{aligned} \operatorname{Res}(R; z_1) &= \frac{4z_1}{4z_1^3 + 20z_1} = \frac{1}{z_1^2 + 5} \\ &= \frac{1}{2\sqrt{6}} \end{aligned}$$

and

$$\begin{aligned}\text{Res}(R; z_2) &= \frac{4z_2}{4z_2^3 + 20z_2} = \frac{1}{z_2^2 + 5} \\ &= \frac{1}{2\sqrt{6}}.\end{aligned}$$

Hence,

$$\begin{aligned}\int_0^{2\pi} \frac{d\theta}{2 + \cos^2 \theta} &= 2\pi \left\{ \frac{1}{2\sqrt{6}} + \frac{1}{2\sqrt{6}} \right\} \\ &= \frac{2\pi}{\sqrt{6}}.\end{aligned}$$
□

Integrals Involving $\log x$ or Fractional Powers of x

Example 8 Find the value of

$$\int_0^\infty \frac{\log x}{(1+x^2)^2} dx.$$

Solution We take

$$f(z) = \frac{\log z}{(1+z^2)^2}$$

on the domain D obtained by deleting the negative imaginary axis; $\log z$ is determined on D by requiring the imaginary part to have values in the interval $(-\pi/2, 3\pi/2)$. We integrate f around the contour in Figure 2.16, for $R > 1$ and $0 < \varepsilon < 1$. f has a single pole at i , and it is of order two. Hence,

$$\begin{aligned}\text{Res}(f; i) &= \text{value of } \left(\frac{\log z}{(z+i)^2} \right)' \quad \text{at } z=i \\ &= \frac{(\pi+2i)}{8}.\end{aligned}$$

Thus,

$$\int_\gamma \frac{\log z}{(z^2+1)^2} dz = \frac{-\pi}{2} + \frac{i\pi^2}{4}.$$

On that part of γ that lies on the circle $|z| = R$, f can be estimated by

$$|f(z)| \leq \frac{2 \log R}{(R^2 - 1)^2}, \quad |z| = R,$$

so

$$\left| \int_{\substack{|z|=R \\ z \in \gamma}} f(z) dz \right| \leq \pi R \frac{2 \log R}{(R^2 - 1)^2} \rightarrow 0 \quad \text{as } R \rightarrow \infty.$$

Furthermore, on that part of γ that lies on the circle $|z| = \varepsilon$, we estimate f by

$$|f(z)| \leq \frac{-2 \log \varepsilon}{(1 - \varepsilon^2)^2}, \quad |z| = \varepsilon.$$

Thus,

$$\left| \int_{\substack{|z|=\varepsilon \\ z \in \gamma}} f(z) dz \right| \leq \pi \varepsilon \frac{(-2 \log \varepsilon)}{(1 - \varepsilon^2)^2} \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0.$$

On the segment of γ on the negative real axis,

$$f(z) = \frac{\log z}{(z^2 + 1)^2} = \frac{\log |x| + i\pi}{(x^2 + 1)^2}, \quad z = x, \quad -R \leq x \leq -\varepsilon.$$

All this together yields

$$\frac{-\pi}{2} + \frac{i\pi^2}{4} = \int_{-\varepsilon}^{-R} \frac{\log x}{(1 + x^2)^2} dx + \int_{-R}^{-\varepsilon} \frac{\log(-x) + i\pi}{(1 + x^2)^2} dx + E,$$

where E goes to zero as $\varepsilon \downarrow 0$ and $R \uparrow \infty$. Taking the real part of both sides, we obtain

$$\int_0^\infty \frac{\log x}{(1 + x^2)^2} dx = \frac{-\pi}{4}. \tag{6}$$

The imaginary part yields the equality

$$\int_0^\infty \frac{dx}{(1 + x^2)^2} = \frac{\pi}{4},$$

which we knew already from Example 2. □

Example 9 Find the value of $\int_0^\infty x^p / (1 + x^2) dx$, $-1 < p < 1$.

Solution Let D be the domain obtained from the complex plane by deleting the origin and the negative imaginary axis. On D we define z^p to be $\exp(p \log z)$, where $\log z$ has imaginary part in the interval $(-\pi/2, 3\pi/2)$. Then the contour in Figure 2.16 lies in D , and

$$f(z) = \frac{z^p}{1 + z^2}$$

is analytic on D except for a pole of order 1 at $z_0 = i$, with residue

$$\text{Res}(f; i) = \frac{i^p}{2i}.$$

As we have seen before, the line integral of f over the semicircle $|z| = R$ may be bounded above by

$$\frac{\pi R^{p+1}}{R^2 - 1},$$

a quantity that approaches zero as $R \rightarrow \infty$. Likewise, the line integral of f over the semicircle $|z| = \varepsilon$ may be bounded above by

$$\frac{\pi \varepsilon^{p+1}}{1 - \varepsilon^2},$$

which goes to zero as $\varepsilon \rightarrow 0$. We apply the Residue Theorem and then let $\varepsilon \rightarrow 0$ and $R \rightarrow \infty$. The conclusion is that

$$[1 + (-1)^p] \int_0^\infty \frac{x^p}{1 + x^2} dx = 2\pi i \text{Res}(f; i) = \pi i^p.$$

Simplifying and using the given formula for the p th root, we obtain

$$\int_0^\infty \frac{x^p}{1 + x^2} dx = \frac{\pi}{2} \left(\cos \frac{\pi p}{2} \right)^{-1}, \quad -1 < p < 1. \quad \square$$

REMARK It is possible to manipulate the result of Example 9 to obtain an interesting formula. Make the change of variables $x^2 = t$. Then the integral becomes

$$\frac{1}{2} \int_0^\infty \frac{t^{(p-1)/2}}{1+t} dt.$$

Set $\alpha = (p+1)/2$; with this change of parameter, we obtain the desired formula:

$$\int_0^\infty \frac{t^{\alpha-1}}{1+t} dt = \frac{\pi}{\sin(\pi\alpha)}, \quad 0 < \alpha < 1. \quad (7)$$

A number of other formulas can be obtained from this last result. In the integral in (7), replace t with t^β , $0 < \beta < \infty$. This gives

$$\int_0^\infty \frac{t^{\alpha\beta-1}}{1+t^\beta} dt = \frac{1}{\beta} \frac{\pi}{\sin(\alpha\pi)}. \quad (8)$$

For instance, with $\beta = 1/\alpha$,

$$\int_0^\infty \frac{dt}{1+t^\beta} = \frac{\pi}{\beta \sin(\pi/\beta)}, \quad 1 < \beta < \infty. \quad (9)$$

Example 10 Find the value of

$$\int_0^\infty \frac{x^{1/3}}{x^2 + 4x + 8} dx.$$

Solution Let D be the domain obtain from the complex plane by deleting the nonnegative real numbers. As in Example 9, $z^{1/3}$ is analytic on D if we choose $\log z$ to have imaginary part in the interval $(0, 2\pi)$. We set

$$f(z) = \frac{z^{1/3}}{z^2 + 4z + 8}.$$

Let γ be the “keyhole” contour pictured in Figure 2.17. In Example 5, Section 6,




Figure 2.17

Chapter 1, we gave a parametrization of γ ; this is repeated next to the figure. The poles of f are at $z_1 = -2 + 2i = 2\sqrt{2}e^{3\pi i/4}$ and $z_2 = -2 - 2i = 2\sqrt{2}e^{5\pi i/4}$. The residues of f at the poles are

$$\text{Res}(f; z_1) = \frac{\sqrt{2}e^{\pi i/4}}{4i}$$

and

$$\text{Res}(f; z_2) = \frac{\sqrt{2}e^{5\pi i/12}}{-4i}.$$

Arguments that we have employed several times now show that the line integral of f over that portion of γ on the circle $|z| = R$ goes to zero as $R \rightarrow \infty$. Likewise, the line integral of f over that portion of γ on the circle $|z| = \varepsilon$ also goes to zero. Using the parametrization of γ shown in Figure 2.17, we see that the line integral over the segment from $\varepsilon e^{i\delta}$ to $Re^{i\delta}$ is

$$\int_{\varepsilon}^R \frac{t^{1/3}e^{i\delta/3}}{t^2 e^{2i\delta} + 4te^{i\delta} + 8} e^{i\delta} dt.$$

As $\delta \rightarrow 0$, this approaches

$$\int_{\varepsilon}^R \frac{t^{1/3}}{t^2 + 4t + 8} dt.$$

Likewise, the integral over the segment from $Re^{i(2\pi-\delta)}$ to $\varepsilon e^{i(2\pi-\delta)}$ converges to

$$e^{i2\pi/3} \int_R^{\varepsilon} \frac{t^{1/3}}{t^2 + 4t + 8} dt,$$

as $\delta \rightarrow 0$. Combining these integrals with the Residue Theorem and letting $\varepsilon \rightarrow 0$ and $R \rightarrow \infty$, we obtain

$$(1 - e^{i2\pi/3}) \int_0^{\infty} \frac{t^{1/3}}{t^2 + 4t + 8} dt = \frac{\sqrt{2}\pi}{2} (e^{\pi i/4} - e^{5\pi i/12}).$$

Simplifying, we obtain

$$\int_0^{\infty} \frac{t^{1/3}}{t^2 + 4t + 8} dt = \frac{\sqrt{2}\pi}{2} \frac{\sin\left(\frac{\pi}{12}\right)}{\sin\left(\frac{\pi}{3}\right)}.$$

EXERCISES FOR SECTION 2.6

Use the method of Examples 1 and 2 to compute these integrals.

1.
$$\int_{-\infty}^{\infty} \frac{x^4}{1+x^8} dx$$

2.
$$\int_{-\infty}^{\infty} \frac{x^2}{x^4 - 4x^2 + 5} dx$$

3.
$$\int_{-\infty}^{\infty} \frac{dx}{(x^2 + a^2)(x^2 + b^2)}, \quad a, b > 0$$

Use the method of Examples 3 to 6 to compute the following integrals.

4.
$$\int_{-\infty}^{\infty} \frac{\cos \alpha x}{(x^2 + 1)(x^2 + 4)} dx$$

5.
$$\int_{-\infty}^{\infty} \frac{x \sin x}{x^4 + 1} dx$$

6.
$$\int_{-\infty}^{\infty} \frac{\sin x}{x^2 + 6x + 10} dx$$

7.
$$\int_{-\infty}^{\infty} \frac{\cos x}{(x + \alpha)^2 + \beta^2} dx$$

8.
$$\int_0^{\infty} \frac{\cos \gamma x}{(x^2 + \alpha^2)(x^2 + \beta^2)} dx$$

$$\text{Answer: } \frac{\pi}{2\alpha\beta(\alpha^2 - \beta^2)} [\alpha e^{-\gamma\beta} - \beta e^{-\gamma\alpha}]$$

Use the method of Example 7 to compute these integrals.

9.
$$\int_0^{2\pi} \frac{d\theta}{(2 - \sin \theta)^2}$$

10.
$$\int_0^{2\pi} \frac{d\theta}{(1 + \beta \cos \theta)^2}, \quad -1 < \beta < 1$$

11.
$$\int_0^{2\pi} \frac{\cos 2\theta}{1 - 2a \cos \theta + a^2} d\theta, \quad -1 < a < 1$$

12.
$$\int_0^{2\pi} \sin^{2k} \theta d\theta \quad \text{Answer: } \pi \frac{(2k)!}{(k!)^2 2^{2k-1}}$$

Use the “keyhole” contour in Figure 2.17 in the manner of Examples 9 and 10 to compute the following integrals.

13.
$$\int_0^{\infty} \frac{x^{\alpha}}{x^2 + 3x + 2} dx, \quad 0 < \alpha < 1$$

14.
$$\int_0^{\infty} \frac{\sqrt{x}}{x^2 + 2x + 5} dx \quad \text{Answer: } \frac{\pi}{2\sqrt{2}} \sqrt{\sqrt{5} - 1}$$

15.
$$\int_0^{\infty} \frac{x^{\lambda}}{x^2 + 2x \cos \omega + 1} dx, \quad -\pi < \omega < \pi, \quad 0 < \lambda < 1$$

Use the method of Example 8 in the following problems. (Note that the integrals are improper, since $\log 0$ is not defined.)

16.
$$\int_0^{\infty} \frac{\log x}{(1+x^2)(4+x^2)} dx$$

17.
$$\int_0^{\infty} \frac{\log x}{1+x^2} dx$$

18.
$$\int_0^{\infty} \frac{[\log x]^2}{1+x^2} dx \quad \text{Answer: } \frac{\pi^3}{8}$$

19.
$$\int_0^{\infty} \frac{[\log x]^4}{1+x^2} dx$$

20. (a) In the identity (7) differentiate both sides with respect to α to obtain the formula

$$\int_0^\infty \frac{t^{\alpha-1} \log t}{1+t} dt = -\pi^2 \cot(\alpha\pi) \csc(\alpha\pi), \quad 0 < \alpha < 1.$$

(b) Use the result of (a) to derive the formula

$$\int_0^\infty \frac{\log s}{1+s^\gamma} ds = -\left(\frac{\pi}{\gamma}\right)^2 \cot\left(\frac{\pi}{\gamma}\right) \csc\left(\frac{\pi}{\gamma}\right), \quad 1 < \gamma < \infty.$$

(Hint: Let $s = t^\alpha$.)

21. Show that $\int_{-\infty}^\infty \frac{\cos x}{e^x + e^{-x}} dx = \frac{\pi}{e^{\pi/2} + e^{-\pi/2}}.$

(Hint: Integrate $f(z) = e^{iz}/(e^z + e^{-z})$ over the rectangle with vertices at $\pm R$ and $\pm R + i\pi$.)

22. Use (8) or (9) to compute these integrals.

(a) $\int_0^\infty \frac{dx}{8+x^3}$

(b) $\int_0^\infty \frac{x}{x^4+16} dx$

(c) $\int_0^\infty \frac{x^\gamma}{1+x^\beta} dx, \quad 0 \leq \gamma < \beta - 1$

23. Show that (a) $\int_{-\infty}^\infty \frac{dx}{(x^2+\alpha^2)^2} = \frac{\pi}{2\alpha^3}$ (b) $\int_{-\infty}^\infty \frac{x^2}{(x^2+\alpha^2)^3} dx = \frac{\pi}{8\alpha^3}$

24. Let p be a polynomial of degree n with distinct zeros at z_1, \dots, z_n and let q be a polynomial of degree $n-2$ or less. Show that

$$\sum_{k=1}^n \operatorname{Res}\left(\frac{q}{p}; z_k\right) = 0.$$

(Hint: Integrate q/p over the circle $|z| = R$, and let $R \rightarrow \infty$.)

25. Give a proof of the Residue Theorem for functions with poles using (11) from Section 5. **(Hint:** Show that the value of the integral

$$\int_\gamma P\left(\frac{1}{z-z_j}\right) dz$$

is $2\pi i \operatorname{Res}(f; z_j)$ if z_j is inside γ and 0 otherwise.)

26. Use the Residue Theorem to calculate the following integrals.

(a) $\frac{1}{2\pi} \int_0^{2\pi} e^{e^{-i\theta}} e^{in\theta} d\theta$

(b) $\frac{1}{2\pi} \int_0^{2\pi} e^{\cos \theta} \cos n\theta d\theta \quad (\text{Hint: } g(z) = z^{n-1} \exp\left(\frac{1}{2}\left(z + \frac{1}{z}\right)\right)).$

(c) $\frac{1}{2\pi} \int_0^{2\pi} e^{\sin \theta} \sin n\theta d\theta$

The Summation of Certain Series by Means of Residues*

27. Show that the poles of $C(z) = \pi \cot \pi z$ are $z_k = k, k = 0, \pm 1, \pm 2, \dots$, that each pole has order one, and that the residue of $C(z)$ at each pole is one.

28. Show that $C(z) = \pi \cot \pi z$ is bounded on the square γ_N with vertices at $\pm(N + \frac{1}{2}) \pm i(N + \frac{1}{2})$, N a positive integer, by a bound that does *not* depend on N .
29. Let $f(z) = 1/z^2$; show that

$$\int_{\gamma_N} f(z)C(z) dz \rightarrow 0 \quad \text{as } N \rightarrow \infty.$$

Use the Residue Theorem to prove that

$$\sum_1^\infty \frac{1}{n^2} = \frac{\pi^2}{6}.$$

30. Suppose that f is analytic on the plane except for poles at ζ_1, \dots, ζ_N , none of which are integers, and suppose that $\lim_{|z| \rightarrow \infty} |zf(z)| = 0$. Show that

$$\sum_{-\infty}^{\infty} f(n) = - \sum_{j=1}^N \operatorname{Res}(f; \zeta_j); \quad C \text{ from Ex. 28} \quad (10)$$

31. Use (10) to find the sum of the following series. In every case, a is chosen such that none of the denominators vanish.

(a) $\sum_{n=1}^{\infty} \frac{1}{n^2 + a^2}$ (c) $\sum_{n=1}^{\infty} \frac{1}{(2n-1)^2}$	(b) $\sum_{-\infty}^{\infty} \frac{1}{(n+a)^2}$ (d) $\sum_{n=1}^{\infty} \frac{1}{n^4}$
---	--

Further Use of the Residue Theorem on $\pi \cot \pi z^*$

32. Let γ_N be the curve from Exercise 28. Show that

$$\frac{1}{2\pi i} \int_{\gamma_N} \frac{C(z)}{z(z-w)} dz = \frac{\pi \cot \pi w}{w} + \sum_{\substack{k=-N \\ k \neq 0}}^N \frac{1}{k(k-w)} - \frac{1}{w^2},$$

$$w \neq 0, \pm 1, \pm 2, \dots, \pm N.$$

33. Let $N \rightarrow \infty$; show that

$$C(w) = \pi \cot \pi w = \frac{1}{w} + \sum_{k=1}^{\infty} \frac{2w}{w^2 - k^2}.$$

34. Integrate $\pi \cot \pi w - (1/w)$, which is analytic at $w = 0$, from 0 to z to obtain

$$\operatorname{Log}\left(\frac{\sin \pi z}{z}\right) = \sum_{k=1}^{\infty} \operatorname{Log}\left(1 - \frac{z^2}{k^2}\right) + \log \pi,$$

z not a real number. Conclude that

$$\sin \pi z = \pi z \prod_{k=1}^{\infty} \left(1 - \frac{z^2}{k^2}\right). \quad (11)$$

Further Reading

Each of the following books gives a more sophisticated and far-reaching development of the theory of analytic functions than is presented here.

- Ahlfors, L. V. *Complex analysis*. 3rd ed. New York: McGraw-Hill, 1979.
- Burckel, R. B. *An introduction to classical complex analysis*. New York: Academic Press, 1979.
- Caratheodory, C. *Theory of functions*. Vols. 1 and 2. New York: Chelsea, 1964.
- Heins, M. *Complex function theory*. New York: Academic Press, 1968.
- Hille, E. *Analytic function theory*. Vols. 1 and 2. New York: Blaisdell, 1959.
- Knopp, K. *Theory of functions*, parts I and II; and *Problem book in the theory of functions*, parts I and II. New York: Dover, 1945; 1952.

The books by Burckel and Hille contain extensive and informative historical and biographical information on the theorems of complex analysis and their creators. These notes are well worth reading.

3

Analytic Functions as Mappings

The theme of this chapter is the “geometry” of analytic functions, in particular the nature of their range. This theme manifests itself first in several theorems in Section 1 that pinpoint the number of solutions to an equation of the form $f(z) = w_0$. The answer is formulated in several different ways, all connected closely with the behavior of the function f on a closed curve. This ability to count the number of solutions of $f(z) = w_0$ is exploited in Section 2 to prove that the range of a nonconstant analytic function is an open set. This is a profound and elegant property of analytic functions that leads in a natural way to the maximum modulus principle. Even more important, we are led directly to the study of conformal mapping, a tool of enormous beauty and power both in applications and theory.

3.1 The Zeros of an Analytic Function

Suppose that f is analytic and not identically zero in a domain D . We saw in (3) and (4) of Section 4, Chapter 2 that each zero of f has a certain **order**; that is, if $f(z_0) = 0$, there is an integer m , $m \geq 1$, such that

$$f(z) = (z - z_0)^m g(z),$$

where g is analytic in D and $g(z_0) \neq 0$. The integer m is frequently called the **multiplicity** of the zero of f at z_0 . In this section, we shall collect several results about the zeros of an analytic function. We begin with a fact closely related to (3) of Section 4, Chapter 2.

Suppose that f is analytic in a domain D . If there are distinct points z_1, z_2, \dots in D with $f(z_n) = 0$, $n = 1, 2, \dots$ and if the sequence $\{z_n\}$ converges to a point z_0 of D , then $f(z) = 0$ for all $z \in D$. (1)

To show why (1) is true, we make use of (3) in Section 4, Chapter 2; we shall show that the hypotheses in (1) imply that $f^{(k)}(z_0) = 0$ for $k = 0, 1, 2, \dots$. We note first that since $z_n \rightarrow z_0$ and since f is continuous, we must have $f(z_0) = 0$. Now $f(z)$ has a power series valid for $|z - z_0| < \delta$:

$$f(z) = a_1(z - z_0) + a_2(z - z_0)^2 + \cdots, \quad |z - z_0| < \delta.$$

Suppose we know that $a_0 = \cdots = a_{N-1} = 0$; we shall show that $a_N = 0$, too. Define a function g by

$$g(z) = \begin{cases} \frac{f(z)}{(z - z_0)^N}, & z \neq z_0 \\ a_N, & z = z_0. \end{cases}$$

By (4) of Section 4, Chapter 2, g is analytic in D . However,

$$g(z_n) = \frac{f(z_n)}{(z_0 - z_n)^N} = 0, \quad n = 1, 2, \dots \text{ and } z_n \neq z_0,$$

so $a_N = g(z_0) = \lim_{n \rightarrow \infty} g(z_n) = 0$. Thus, all the coefficients a_k of the power series for $f(z)$ about the point z_0 are zero, so $f(z) = 0$ for all z in D by (3) in Section 4, Chapter 2.

To reiterate,

if an analytic function f on a domain D vanishes on a sequence of distinct points $\{z_n\}$ of D , which converges to a point z_0 of D , then f vanishes identically in D .

Example 1 The hypothesis that z_0 lies in D is not superfluous. For instance, $f(z) = \sin(\pi/z)$ is analytic for all $z \neq 0$, and $f(1/n) = \sin(\pi n) = 0$ for $n = 1, 2, \dots$, and $z_n = 1/n \rightarrow 0$ as $n \rightarrow \infty$, yet f is not identically zero. (Obviously, f is not analytic on any disc centered at the origin.) \square

The mathematical way to phrase (1) is to say that the zeros of a nonconstant analytic function are **isolated** (from each other). This allows us to count them, as the next several theorems show.

Suppose that h is a function analytic on a domain D except for a finite number of poles. Let γ be a piecewise smooth simple closed curve in D whose inside lies in D and which passes through no zero or pole of h . Let z_1, \dots, z_N be the distinct zeros of h inside γ , and let w_1, \dots, w_M be the distinct poles of h inside γ (Fig. 3.1).




Figure 3.1

Let n_j be the order (or multiplicity) of the zero of h at z_j , $j = 1, \dots, N$. Then

$$h(z) = (z - z_j)^{n_j} G_j(z),$$

where G_j is analytic near z_j and $G_j(z_j) \neq 0$. Hence,

$$\frac{h'(z)}{h(z)} = \frac{n_j}{z - z_j} + \frac{G'_j(z)}{G_j(z)},$$

so

$$\text{Res}\left(\frac{h'}{h}; z_j\right) = n_j. \quad (2)$$

Likewise, if m_k is the order of the pole of h at w_k , $k = 1, \dots, M$, then

$$h(z) = \frac{H_k(z)}{(z - w_k)^{m_k}},$$

where H_k is analytic near w_k and $H_k(w_k) \neq 0$. Thus,

$$\frac{h'(z)}{h(z)} = \frac{-m_k}{z - w_k} + \frac{H'_k(z)}{H_k(z)}$$

for z near w_k , so

$$\text{Res}\left(\frac{h'}{h}; w_k\right) = -m_k. \quad (3)$$

We now apply the Residue Theorem (Theorem 1 from Section 6, Chapter 2) to the function (h'/h) . The result is

$$\frac{1}{2\pi i} \int_{\gamma} \frac{h'(z)}{h(z)} dz = \sum_{j=1}^N n_j - \sum_{k=1}^M m_k. \quad (4)$$

Here is the formal statement of (4).

THEOREM 1 Suppose that h is analytic in a domain D except for a finite number of poles. Let γ be a piecewise smooth positively oriented simple closed curve in D , which does not pass through any pole or zero of h and whose inside lies in D . Then,

$$\frac{1}{2\pi i} \int_{\gamma} \frac{h'(z)}{h(z)} dz = \begin{cases} \text{number of zeros} \\ \text{of } h \text{ inside } \gamma \end{cases} - \begin{cases} \text{number of poles} \\ \text{of } h \text{ inside } \gamma \end{cases}.$$

All zeros and poles are counted with multiplicities. ■

If you have been observant, you will have noticed by now that h'/h is the derivative of $\log h$, and you may well have reasoned along the following lines: $(\log h)' = h'/h$, so

$$\begin{aligned} \int_{\gamma} \frac{h'(z)}{h(z)} dz &= \int_{\gamma} (\log h(z))' dz \\ &= \log h(\gamma(b)) - \log h(\gamma(a)) \\ &= 0, \quad \text{since } \gamma(a) = \gamma(b). \end{aligned}$$

This surely runs completely contrary to the conclusion of Theorem 1. Consequently, there is a flaw in the reasoning above, and it is not overly difficult to pinpoint it—the logarithm function is not continuous on any domain that surrounds the origin. The point of Theorem 1 is, fundamentally, that the zeros and poles of h are counted by counting the number of times that the closed curve $\Gamma = h \circ \gamma$ winds around the origin. However, a lot of things are right with this fallacious computation, and all will be put right by the more careful use of the logarithm. Further, the logarithm will show the connection between the integral in (4) and the argument of h . For reasons of exposition, we will give the proof in the case when γ is a circle. More complicated curves naturally require somewhat more complicated proofs.

Let γ be the circle $\gamma(t) = z_0 + Re^{it}$, $0 \leq t \leq 2\pi$. Since h is not zero on γ , there is an annular region $R - \delta \leq |z - z_0| \leq R + \delta$ on which h is not zero. Let D be the domain $D = \{z: R - \delta < |z - z_0| < R + \delta \text{ and } 0 < \arg(z - z_0) < 2\pi\}$ (see Fig. 3.2).




Figure 3.2

The domain D is simply-connected, so there is an analytic function f on D with $f = \log h$; see Application 2, Section 4, Chapter 2. Now let ϵ be a small but positive real number. Since h'/h is continuous on the annulus $R - \delta \leq |z - z_0| \leq R + \delta$, and after parametrizing the curve γ , it follows that

$$\int_{\gamma} \frac{h'(z)}{h(z)} dz = \lim_{\epsilon \rightarrow 0} \int_{\epsilon}^{2\pi-\epsilon} \frac{h'(z_0 + Re^{it})}{h(z_0 + Re^{it})} iRe^{it} dt.$$

However, the latter integral equals

$$\begin{aligned} \int_{\varepsilon}^{2\pi-\varepsilon} f'(z_0 + Re^{it}) iRe^{it} dt &= \int_{\varepsilon}^{2\pi-\varepsilon} \frac{d}{dt} [f(z_0 + Re^{it})] dt \\ &= f(z_0 + Re^{i(2\pi-\varepsilon)}) - f(z_0 + Re^{i\varepsilon}) \\ &= \log|h(z_0 + Re^{i(2\pi-\varepsilon)})| - \log|h(z_0 + Re^{i\varepsilon})| \\ &\quad + i\{\arg h(z_0 + Re^{i(2\pi-\varepsilon)}) - \arg h(z_0 + Re^{i\varepsilon})\}. \end{aligned}$$

As $\varepsilon \rightarrow 0$, an interesting thing occurs: The real part of this expression, $\log|h(z_0 + Re^{i(2\pi-\varepsilon)})| - \log|h(z_0 + Re^{i\varepsilon})|$, goes to zero, but the imaginary part, $\{\arg h(z_0 + Re^{i(2\pi-\varepsilon)}) - \arg h(z_0 + Re^{i\varepsilon})\}$, need not. In fact, the imaginary part most clearly converges to the total change in $\arg h(z)$ as z traverses γ once. (Think of $h(z) = z^2$ on a circle centered at the origin.) This establishes, for a circle, the connection between the integral in (4) and the change in the argument of $h(z)$ as z traverses the curve γ . This relationship is called the **argument principle**. We state it formally.

THEOREM 2 The Argument Principle Suppose h is analytic on a domain D except for isolated poles. Let γ be a piecewise smooth positively oriented simple closed curve in D whose inside lies in D and which does not pass through any zeros or poles of h . Then,

$$\frac{1}{2\pi} \left\{ \begin{array}{l} \text{change in } \arg h(z) \\ \text{as } z \text{ traverses } \gamma \end{array} \right\} = \left\{ \begin{array}{l} \text{number of zeros} \\ \text{of } h \text{ inside } \gamma \end{array} \right\} - \left\{ \begin{array}{l} \text{number of poles} \\ \text{of } h \text{ inside } \gamma \end{array} \right\}. \quad \blacksquare$$

Example 2 Find the number of zeros of the function $f(z) = z^3 - 2z^2 + 4$ in the first quadrant.

Solution We examine $f(z)$ on the contour shown in Figure 3.3; R is very big. On the segment $0 \leq x \leq R$, $f(x) = x^3 - 2x^2 + 4$ is real and greater than 2. On the quarter-circle $z = Re^{it}$, $0 \leq t \leq \pi/2$,

$$f(Re^{it}) = R^3 e^{3it} \left(1 - \frac{2}{R e^{it}} + \frac{4}{R^3 e^{3it}} \right) = R^3 e^{3it} (1 + \zeta),$$




Figure 3.3

where $|\zeta| \leq 6/R < \epsilon$ for R large. Thus, $\arg f(Re^{it})$ is approximately $\arg(e^{3it}) = 3t$ for large R , so $\arg f(Re^{it})$ increases from 0 to about $3\pi/2$ as t increases from 0 to $\pi/2$. On the segment $z = iy$, $R \geq y \geq 0$,

$$f(iy) = -iy^3 + 2y^2 + 4.$$

For $R \geq y > 0$, this point lies in the fourth quadrant, since

$$\operatorname{Re}(f(iy)) = 4 + 2y^2 > 0$$

$$\operatorname{Im}(f(iy)) = -y^3 < 0.$$

Hence, as y decreases from R to 0, $f(iy)$ lies in the fourth quadrant and moves toward the point $w = 4$. Consequently, as z traverses the contour, $\arg f(z)$ increases by exactly 2π , so $f(z) = z^3 - 2z^2 + 4$ has precisely one zero in the first quadrant. \square

Example 3 Let $\lambda > 1$. Use the argument principle to show that the equation

$$z + e^{-z} = \lambda$$

has exactly one solution in the right half-plane $\{z: \operatorname{Re} z > 0\}$.

Solution We shall examine $h(z) = z + e^{-z} - \lambda$ on the curve pictured in Figure 3.4. We shall show that when R is very large, the net change in $\arg h$ over this curve is 2π . It will then follow from the argument principle that h has precisely one zero in the right half-plane. We note first that on the imaginary axis,

$$\begin{aligned} h(iy) &= iy + e^{-iy} - \lambda \\ &= (\cos y - \lambda) + i(\sin y - \sin y). \end{aligned}$$




Figure 3.4

Since $\lambda > 1$, $\operatorname{Re} h(iy)$ is always negative. Further, $\operatorname{Im} h(iy)$ decreases from a positive value at $y = R$ to a negative value at $y = -R$. That is, $h(iy)$ moves from the second to the third quadrant as y decreases from R to $-R$. On the semicircle $z = Re^{i\theta}$, $-\pi/2 \leq \theta \leq \pi/2$,

$$|e^{-Re^{i\theta}}| = e^{-R \cos \theta} \leq 1,$$

so $h(re^{i\theta}) = Re^{i\theta}\{1 + c\}$, where c is very small when R is very large. Consequently, $\arg h(Re^{i\theta})$ is approximately θ and so increases from (about) $-\pi/2$ to (about) $\pi/2$ as θ increases from $-\pi/2$ to $\pi/2$. That is, the point $h(Re^{i\theta})$ moves from the third quadrant (when $\theta = -\pi/2$ and $z = -iR$) through the fourth quadrant, then through the first quadrant, and ends in the second quadrant (when $\theta = \pi/2$ and $z = iR$). In total, then, $h(z)$ makes one full counterclockwise circuit about the origin as z traverses γ once in the positive sense, so $\arg h(z)$ increases by 2π . Hence, h has one (and only one) zero inside γ , by the argument principle. \square

Here is another theorem, closely related to the argument principle.

THEOREM 3 Rouché's* Theorem Suppose f and g are analytic on an open set containing a piecewise smooth simple closed curve γ and its inside. If

$$|f(z) + g(z)| < |f(z)| \quad \text{for all } z \in \gamma, \tag{5}$$

then f and g have an equal number of zeros inside γ , counting multiplicities.

Proof Note that the hypothesis (5) ensures that neither f nor g is zero on γ . We may assume further that all the common zeros of f and g inside γ have been canceled; this affects neither the hypotheses nor the conclusion. Let $h = g/f$; then

$$|h(z) + 1| < 1 \quad \text{for all } z \in \gamma,$$

so the range of h on γ lies in the disc of radius 1 centered at the point -1 . In particular, $\arg h(z)$ has no net change as z traverses γ (Fig. 3.5).

By Theorem 2, the number of zeros of h inside γ equals the number of poles of h inside γ . But the number of zeros of h is just the number of zeros of g , and the number of poles of h just the number of zeros of f . \blacksquare

REMARK Suppose that f and g satisfy the hypotheses of Theorem 3 except that

$$|f(z) - g(z)| < |f(z)|, \quad z \in \gamma. \tag{6}$$

It is then still true that f and g have equally many zeros inside γ . The reasoning is elementary: According to Theorem 3, the hypothesis (6) implies that f and $-g$ have equally many zeros inside γ . However, a point is a zero of $-g$ exactly when it is a zero of g . We are done.

* Eugene Rouché, 1832–1910.




Figure 3.5

Rouché's Theorem has a number of useful applications, as the following examples show.

Example 4 Show that all the zeros of

$$p(z) = 3z^3 - 2z^2 + 2iz - 8$$

lie in the annulus $1 < |z| < 2$.

Solution On the circle $|z| = 1$,

$$|p(z) + 8| \leq 3 + 2 + 2 = 7 < 8,$$

so $p(z)$ and $f(z) \equiv 8$ have the same number of zeros within $|z| = 1$, by Rouché's Theorem; thus, $p(z)$ does not vanish in the disc $|z| \leq 1$. Furthermore, on $|z| = 2$,

$$|p(z) - 3z^3| \leq 2(4) + 2(2) + 8 = 20 < 24 = |3z^3|,$$

so $p(z)$ and $f(z) = 3z^3$ have an equal number of zeros within the circle $|z| = 2$; that is, $p(z)$ has three of its zeros within $|z| = 2$. Consequently, all the zeros of p lie in the annulus $1 < |z| < 2$. \square

Example 5 Find the number of roots of the equation

$$\frac{z^2 - 4}{z^2 + 4} + \frac{2z^2 - 1}{z^2 + 6} = 0 \quad (7)$$

that lie within the unit circle $|z| = 1$.

Solution When we clear the fractions, we see that we must determine the number of roots within the circle $|z| = 1$ of the equation

$$(z^2 - 4)(z^2 + 6) = (z^2 + 4)(1 - 2z^2).$$

Let $p(z) = (z^2 - 4)(z^2 + 6) = z^4 + 2z^2 - 24$, and let $q(z) = (z^2 + 4)(1 - 2z^2) = -2z^4 - 7z^2 + 4$. Then, on $|z| = 1$,

$$|p(z)| = |24 - 2z^2 - z^4| \geq 24 - 2 - 1 = 21$$

and

$$|q(z)| = |2z^4 + 7z^2 - 4| \leq 2 + 7 + 4 = 13.$$

Hence, $|p(z)| > |q(z)| = |\{p(z) - q(z)\} - \{p(z)\}|$, which implies that p and $p - q$ have an equal number of zeros within the circle $|z| = 1$. But clearly p has no zeros within $|z| < 1$, so (7) has no roots within $|z| < 1$. \square

Rouché's Theorem also provides a direct and simple proof of the following result.

THEOREM 4 The Fundamental Theorem of Algebra A polynomial of degree n has exactly n zeros, counting multiplicities.

Proof We do not affect the solutions of $p(z) = 0$ by dividing $p(z)$ by the coefficients of z^n ; thus, we may suppose that

$$p(z) = z^n + a_{n-1}z^{n-1} + \cdots + a_1z + a_0.$$

For large values of $|z| = R$,

$$\begin{aligned} \left| \frac{p(z) - z^n}{z^n} \right| &= |a_{n-1}z^{-1} + \cdots + a_0z^{-n}| \\ &\leq \frac{|a_0| + \cdots + |a_{n-1}|}{R} \leq \frac{1}{2}. \end{aligned}$$

Hence, $|p(z) - z^n| \leq \frac{1}{2}|z|^n < |z^n|$ for $|z| = R$, so by Rouché's Theorem (Theorem 3), we learn that $p(z)$ and z^n have an equal number of zeros within the circle $|z| = R$. But z^n has exactly n zeros within $|z| = R$, so the same is true for $p(z)$. \blacksquare

EXERCISES FOR SECTION 3.1

Use the technique of Example 2 to determine the number of zeros of f in the first quadrant.

- | | |
|---|--|
| 1. $f(z) = z^2 - z + 1$
3. $f(z) = z^3 - 3z + 6$
5. $f(z) = z^9 + 5z^2 + 3$ | 2. $f(z) = z^4 - 3z^2 + 3$
4. $f(z) = z^2 + iz + 2 + i$
6. $f(z) = z^7 + 6z^3 + 7$ |
|---|--|

In Exercises 7 and 8, use the technique of Example 3 to determine the number of zeros of the given function in the upper half-plane.

7. $z^4 + 3iz^2 + z - 2 + i$
8. $2z^4 - 2iz^3 + z^2 + 2iz - 1$
9. Show that the equation $z^4 - 5z^2 + 3 = e^{-z}$ has no solutions on the imaginary axis and precisely two solutions in the half-plane $\operatorname{Re} z > 0$.
10. Show that there is no entire function F with $F(x) = 1 - \exp[2\pi i/x]$ for $1 \leq x \leq 2$.
11. Show that there is no entire function F that satisfies the equation $F^2(z) = \sin z$ for all z in the complex plane. Show that the same conclusion holds if “all z in the complex plane” is replaced by “ z in some open set,” or even by “for some sequence of z that have a limit point in the plane.”

In Exercises 12 to 15, use the technique of Example 4 to determine how many zeros of the given function lie in the given annulus.

12. $z^3 - 3z + 1$ in $1 < |z| < 2$
13. $z^4 - 2z - 2$ in $\frac{1}{2} < |z| < \frac{3}{2}$
14. $ze^z - \frac{1}{4}$ in $0 < |z| < 2$
15. $4z^3 - 12z^2 + 2z + 10$ in $\frac{1}{2} < |z - 1| < 2$
16. Let f and g be analytic on a domain containing a simple closed curve γ and its inside. Show that if $|f(z)| > |g(z)|$ for all $z \in \gamma$, then the two equations $f(z) = g(z)$ and $f(z) = 0$ have an equal number of solutions inside γ .
17. Use the result of Exercise 16 to determine how many solutions each of the following equations has in the given region.
 - (a) $z^3 - 10 = z^2 + z + 6$ in $|z| < 1$
 - (b) $z^3 = z^2 + z + 6$ in $|z| < 1$
 - (c) $z + a = e^z$ for $a > 1$, in $\operatorname{Re} z < 0$. (Hint: let γ be the semicircle of radius R from iR to $-iR$ through $-R$ plus the segment from $-iR$ to iR on the imaginary axis.)
18. Extend formula (4) to prove the following. Let g be analytic on a domain containing γ and its inside. Then

$$\frac{1}{2\pi i} \int_{\gamma} \frac{h'(z)}{h(z)} g(z) dz = \sum_{i=1}^N g(z_i) - \sum_{j=1}^M g(w_j),$$

where z_1, \dots, z_N are the zeros of h and w_1, \dots, w_M are the poles of h inside γ , each listed according to its multiplicity.

19. Suppose that f is analytic and one-to-one on a domain D . In this exercise, we shall show that the inverse function of f is analytic on the domain $\Omega = f(D)$. Let γ be a piecewise smooth simple closed curve in D whose inside lies in D , and let w be a point of Ω that lies inside the curve $\Gamma = f \circ \gamma$. In Exercise 18, take $g(z) = z$ and $h(z) = f(z) - w$. Conclude that f^{-1} , the inverse of f , is given by

$$f^{-1}(w) = \frac{1}{2\pi i} \int_{\gamma} \frac{zf'(z)}{f(z) - w} dz.$$

The integral expression on the right is an analytic function of w ; see exercise 26 in Section 1 of Chapter 2.

20. Suppose that f is analytic on a domain containing $\{z: |z| \leq 1\}$ and that $|f(e^{i\theta})| < 1$, $0 \leq \theta \leq 2\pi$. Show that f has exactly one **fixed point** in the disc $|z| < 1$; that is, the equation $f(z) = z$ has precisely one solution in the disc $|z| < 1$.
21. Let p and q be polynomials of degree n . If $p(z) = q(z)$ at $n + 1$ distinct points of the plane, then $p(z) = q(z)$ for all z .
22. Use Liouville's Theorem to prove the Fundamental Theorem of Algebra. (**Hint:** If p is a polynomial with no zero (and degree p is 1 or more), then $1/p$ is an entire function. Use (3) of Section 6, Chapter 2, to show that $1/p$ is bounded.)
23. Let P be a polynomial with $|P(e^{it})| = 1$ for all t , $0 \leq t \leq 2\pi$. Show that $P(z) = \lambda z^N$, $|\lambda| = 1$. (**Hint:** Let N be the degree of P and set

$$Q(z) = z^N P\left(\frac{1}{\bar{z}}\right).$$

Show that Q is a polynomial and that $P(e^{it})Q(e^{it}) = e^{iNt}$, $0 \leq t \leq 2\pi$. Conclude that Q is constant.)

24. Let p be a polynomial of degree n all of whose zeros are real. Let $x_1 < x_2 < \dots < x_r$ be the distinct zeros of p with respective multiplicities m_1, m_2, \dots, m_r .
- Show that p' is not zero in $(-\infty, x_1)$ or in (x_r, ∞) .
 - Show that p' has precisely one zero in (x_j, x_{j+1}) , $j = 1, \dots, r - 1$.

Rational Functions on the Extended Plane*

Suppose that f is analytic on a plane except for a finite number of poles; suppose, in addition, that f either is analytic at ∞ or has a pole at ∞ . Complete the following exercises to show that f is, in fact, a rational function.

Let m_0 be the order of the pole of f at ∞ with $m_0 = 0$ if f is analytic at ∞ . Let w_1, \dots, w_r be the distinct poles of f in the plane with respective orders m_1, \dots, m_r . Set

$$g(z) = (z - w_1)^{m_1} \cdots (z - w_r)^{m_r} f(z).$$

25. Show that g is an entire function.

26. Show that

$$|g(z)| \leq C|z|^m, \quad |z| > R.$$

where C is a constant and $m = m_0 + m_1 + \cdots + m_r$.

27. Conclude that g is a polynomial of degree m or less, and thus f is a rational function. (**Hint:** Use Exercise 21, Section 4, Chapter 2.)

Mapping Properties of Rational Functions*

Let P and Q be polynomials of respective degrees N and M and assume that P and Q have no common zero. Let $R = P/Q$ and define $d = \text{degree of } R$ to be the maximum of N and M .

28. Suppose that $M \geq N$. (a) Show that R has a removable singularity at ∞ . (b) If $M > N$, then R has a zero of order $M - N$ at ∞ . (c) If $M = N$, then $R(\infty) \neq 0$.

29. Suppose that $M < N$. Show that R has a pole at ∞ of order $N - M$.
30. Counting multiplicities and counting the zeros or poles at ∞ , if any, show that R has d zeros and d poles.
31. Let α be a complex number. Show that $R - \alpha$ is a rational function with the same degree as R .
32. Show that the equation $R(z) = \alpha$ has exactly d solutions, counting multiplicities, for each complex number α . Conclude that R maps the complex plane plus ∞ onto itself and each point is “covered” exactly d times.

Legendre* Polynomials*

We define the n th Legendre polynomial P_n by

$$P_n(z) = \frac{1}{2^n n!} \frac{d^n}{dz^n} (z^2 - 1)^n. \quad (8)$$

33. Let γ be a smooth simple closed curve around z . Show that

$$P_n(z) = \frac{1}{2\pi i} \frac{1}{2^n} \int_{\gamma} \frac{(w^2 - 1)^n}{(w - z)^{n+1}} dw. \quad (9)$$

34. In Exercise 33, take γ to be the circle of radius $\sqrt{|z^2 - 1|}$ centered at z ; show that

$$P_n(z) = \frac{1}{2\pi} \int_0^{2\pi} (z + \sqrt{z^2 - 1} \cos \theta)^n d\theta. \quad (10)$$

35. Prove the generating formula:

$$\frac{1}{\sqrt{1 - 2zt + t^2}} = \sum_{n=0}^{\infty} P_n(z)t^n.$$

(Hint: In the series on the right, substitute for P_n the formula (10). You will also have need for Exercise 7, Section 3, Chapter 2.)

36. Let $F(z, t) = (1 - 2zt + t^2)^{-1/2}$.

(a) Show that F satisfies the partial differential equation

$$(1 - z^2) \frac{\partial^2 F}{\partial z^2} - 2z \frac{\partial F}{\partial z} + \frac{\partial^2 (tF)}{\partial t^2} = 0.$$

(b) Use (a) and the conclusion of Exercise 35 to show that P_n satisfies Legendre's differential equation:

$$(1 - z^2)f''(z) - 2zf'(z) + n(n + 1)f(z) = 0.$$

37. Let $F(z, t)$ be the function in Exercise 36. (a) Show that F also satisfies the partial

* Adrien Marie Legendre, 1752–1833.

differential equation

$$(1 - 2zt + t^2) \frac{\partial F}{\partial t} - (z - t)F = 0.$$

(b) Use (a) and Exercise 35 to show that the Legendre polynomials satisfy

$$(n + 1)P_{n+1}(z) - (2n + 1)zP_n(z) + nP_{n-1}(z) = 0,$$

and, using (8),

$$(2n + 1)P_n(z) = P'_{n+1}(z) - P'_{n-1}(z).$$

These are called **recursion relations**.

3.1.1 The Stability of Solutions of a System of Linear Differential Equations*

The operation of many physical processes is described by a system of ordinary differential equations, together with a set of initial conditions. It is a matter of great relevance whether small perturbations in the initial conditions will, over the long run, produce substantial or even unbounded changes in the corresponding solution. As an elementary example, imagine placing a ball on the curve $y = x^2$. If placed at the origin $(x_0, y_0) = (0, 0)$, the ball remains stationary; if placed at some point (x_0, y_0) , $x_0 \neq 0$, the ball rolls back and forth about $(0, 0)$ indefinitely (or, if there is friction, tends to $(0, 0)$ as time goes to infinity). On the other hand, if the ball is placed on the curve $y = -x^2$, it moves farther and farther from $(0, 0)$ as times goes to infinity, *unless* it was placed (with a steady hand!) exactly at $(0, 0)$ to begin with. Obviously, the first situation is what we can call “stable,” while the second is “unstable.” In this section, we shall show that if the zeros of a certain polynomial associated with the system of differential equations all have negative real parts, then the solutions of the differential equations are stable, in the sense that small perturbations in the initial conditions produce only small perturbations in the solution. A general knowledge of the solution of a linear first-order system of differential equations would be helpful but is not strictly necessary in reading this section.

The system we shall investigate is

$$\dot{\mathbf{x}} = A\mathbf{x} + \mathbf{b}, \quad (1)$$

where A is an $n \times n$ matrix with constant entries, \mathbf{b} is an $n \times 1$ column vector with constant entries, and

$$\mathbf{x} = \mathbf{x}(t) = \begin{pmatrix} x_1(t) \\ \vdots \\ x_n(t) \end{pmatrix}.$$

The solutions of this system involve the zeros of the polynomial p given by

$$p(z) = \det(zI - A), \quad (2)$$

where I is the $n \times n$ identity matrix and “det” stands for determinant. The polynomial p has degree n and so has n zeros in the plane, counting multiplicities. Let $\lambda_1, \dots, \lambda_r$ be the *distinct* zeros and let m_1, \dots, m_r be their respective multiplicities so that

$$p(z) = \prod_1^r (z - \lambda_j)^{m_j}.$$

The study of linear first-order systems of which (1) is representative tells us that each solution of the homogeneous system $Ax = \dot{x}$ has the form

$$x_l(t) = \sum_{j=1}^r p_{jl}(t) e^{\lambda_j t}, \quad l = 1, \dots, n, \quad (3)$$

where each p_{jl} is a polynomial of degree at most $m_j - 1$. Furthermore, given a set of initial conditions $x(0) = x_0$, there is a unique solution $x(t)$ of (1) with $x(0) = x_0$.

We shall now examine the difference between the solution $x_1(t)$ corresponding to initial conditions $x_1(0) = x_1$ and the solution $x_0(t)$ and watch how this difference behaves as $t \rightarrow \infty$. Note that the linearity of the system implies that

$$A(x_0 - x_1) = (\dot{x}_0 - b) - (\dot{x}_1 - b) = \dot{x}_0 - \dot{x}_1$$

and

$$x_0(0) - x_1(0) = x_0 - x_1.$$

Hence, if we write $w = x_0 - x_1$, then w solves the homogeneous equation

$$Aw = \dot{w} \quad \text{and} \quad w(0) = w_0 = x_0 - x_1.$$

We shall now show that if all the zeros of the polynomial p in (2) lie in the left half-plane, then any solution $w(t)$ of the homogeneous linear system $Aw = \dot{w}$ tends to zero as $t \rightarrow \infty$. Indeed, the proof is elementary. Let $w_l(t)$ be the l th component of the solution. From (3) we have

$$|w_l(t)| \leq \sum_{k=1}^r |p_{kl}(t)| e^{t \operatorname{Re} \lambda_k}.$$

Now $p_{kl}(t)$ has degree at most $m_k - 1$, so we know that

$$|p_{kl}(t)| \leq A_{kl} t^{m_k - 1}, \quad t \geq 1,$$

where A_{kl} is a positive constant. Furthermore, our assumption about the zeros of p

implies that there is a positive number σ with

$$\operatorname{Re} \lambda_j \leq -\sigma, \quad j = 1, \dots, r,$$

because $\operatorname{Re} \lambda_j < 0$ for $j = 1, \dots, r$. Assuming $t \geq 1$, as we can do, we find that

$$|w_i(t)| \leq C_i t^{n-1} e^{-\sigma t}, \quad t \geq 1,$$

where C_i is some positive constant. However, it is elementary that

$$\lim_{t \rightarrow \infty} (t^{n-1} e^{-\sigma t}) = 0,$$

so $w_i(t) \rightarrow 0$ as $t \rightarrow \infty$, as we wished to show.

Since no eigenvalue of A is zero, we know that A is invertible, so there is a (constant) vector v_0 with $Av_0 = -b$. Set $x_0(t) = v_0$ for all t . Then $x_0(t)$ solves (1) and so *any* other solution of (1) tends to v_0 as $t \rightarrow \infty$. The precise nature of how $x(t)$ converges to v_0 depends, of course, on the properties of the matrix A and is an interesting topic in its own right.

It is interesting to note that if the polynomial p given in (2) has all its zeros in the half-plane $\operatorname{Re} z < 0$, then *any* set of initial conditions produces a solution that, over the long run, converges to the constant $x(t) = v_0$. That is, if the operation of a mechanical system is determined by (1) and if the associated polynomial in (2) has all its roots in the half-plane $\operatorname{Re} z < 0$, then no matter how the system starts, it will (eventually) stabilize at the same “steady state.”

Stable Polynomials

A polynomial p whose zeros all lie in the half-plane $\operatorname{Re} z < 0$ is called **stable**. The choice of terminology and the importance of such polynomials have been made clear above. Here we shall look for conditions that ensure that p is stable. We give a full solution if the degree of p is two or three. We also give the result (but not the justification) for general polynomials. These results go under the general heading of the **Routh–Hurwitz criteria**.

PROPOSITION 1 Suppose that A and B are real numbers. The real parts of both the roots of

$$z^2 + Az + B = 0$$

are negative if and only if both A and B are positive.

Proof The roots of the equation are

$$z_1 = \frac{1}{2}(-A + \sqrt{A^2 - 4B})$$

and

$$z_2 = \frac{1}{2}(-A - \sqrt{A^2 - 4B}).$$

If $\operatorname{Re} z_1$ and $\operatorname{Re} z_2$ are negative, then

$$-A = \operatorname{Re}(z_1 + z_2) < 0.$$

Furthermore, if $A^2 - 4B \leq 0$, then $4B \geq A^2 > 0$, and therefore $B > 0$. Otherwise, $A^2 - 4B > 0$, and

$$0 > 2 \operatorname{Re} z_1 = -A + \sqrt{A^2 - 4B},$$

so $A > \sqrt{A^2 - 4B}$, which implies that $B > 0$.

Conversely, if both A and B are positive, then virtually the same arguments show that $\operatorname{Re} z_1$ and $\operatorname{Re} z_2$ are both negative. ■

Example 1 The polynomial $p(z) = z^2 + z + 6$ is stable, but the polynomial

$$q(z) = z^2 + 2z - \varepsilon$$

is not stable, no matter how small the positive number ε is made. (One of its zeros lies at $z = -1 + \sqrt{1 + \varepsilon}$, which is positive.) □

The situation for cubic polynomials is not quite as simple as it is for quadratic polynomials, but it is not terribly difficult either.

PROPOSITION 2 Let A , B , and C be real numbers. The real part of each of the roots of the cubic polynomial

$$z^3 + Az^2 + Bz + C = 0$$

is negative if and only if A , B , and C are all positive and $AB > C$.

Proof Let $p(z) = z^3 + Az^2 + Bz + C = (z - z_1)(z - z_2)(z - z_3)$. Then

$$A = -z_1 - z_2 - z_3$$

$$B = z_1 z_2 + z_1 z_3 + z_2 z_3$$

$$C = -z_1 z_2 z_3.$$

We first examine the case when the root z_1 is real and the other two roots z_2 and z_3 are complex conjugates of each other. We write

$$z_1 = a, \quad z_2 = b + ic, \quad z_3 = b - ic,$$

where a , b , and c are all real. Thus, we find that

$$A = -a - 2b, \quad B = 2ab + b^2 + c^2, \quad C = -a(b^2 + c^2).$$

Suppose first that a and b are negative; it clearly follows from the formulas above that A , B , and C are all positive. Furthermore,

$$C - AB = 2b(a^2 + 2ab + b^2 + c^2) < 0,$$

so $AB > C$. Conversely, suppose that all of A , B , C , and $AB - C$ are positive; we must show that a and b are negative. Since

$$0 < C = -a(b^2 + c^2),$$

clearly $a < 0$. Furthermore, from above

$$0 < AB - C = -2b(a^2 + 2ab + b^2 + c^2),$$

so b must also be negative.

The other case to be examined is when all the roots z_1 , z_2 , z_3 are real; for simplicity we denote them by a , b , and c , respectively. Clearly,

$$A = -a - b - c$$

$$B = ab + bc + ac$$

$$C = -abc,$$

so if a , b , and c are all negative, then, A , B , and C are all positive. Furthermore,

$$AB - C = -b(a^2 + c^2) - c(a^2 + b^2) - a(b^2 + c^2) - 2abc > 0$$

if a , b , and c are all negative.

Conversely, suppose all four numbers A , B , C , and $AB - C$ are positive. First of all, since $0 < C = -abc$, either a , b , and c are all negative or one of these numbers is negative and the other two are positive. Hence, in order to reach a contradiction, let us assume that $a < 0$ and that b and c are positive. But then

$$-a = A + b + c,$$

so

$$\begin{aligned} 0 < B &= a(b + c) + bc \\ &= -(A + b + c)(b + c) + bc \\ &= -A(b + c) - b^2 - c^2 - bc < 0, \end{aligned}$$

a contradiction. This completes the proof. ■

Example 2 The polynomial

$$p(z) = 6 + 4z + 2z^2 + z^3$$

is stable, since $A = 2$, $B = 4$, and $C = 6$ are all positive and $AB = 8 > 6 = C$. \square

Example 3 The polynomial

$$p(z) = 12 + 5z + 2z^2 + z^3$$

is not stable, since $C = 12 > 10 = (5)(2) = AB$. \square

The general case is set forth in the following theorem.

THEOREM 1 Routh*-Hurwitz† Condition Suppose that a_1, a_2, \dots, a_n are real numbers; set $a_j = 0$ if $j > n$. All the roots of the polynomial $p(z) = a_n + a_{n-1}z + a_{n-2}z^2 + \dots + a_1z^{n-1} + z^n$ have negative real part if and only if for each $k = 1, \dots, n$, the determinant of the $k \times k$ matrix

$$M_k = \begin{bmatrix} a_1 & a_3 & a_5 & \dots & a_{2k-1} \\ 1 & a_2 & a_4 & \dots & a_{2k-2} \\ 0 & a_1 & a_3 & \dots & a_{2k-3} \\ 0 & 1 & a_2 & & \\ \vdots & \vdots & \vdots & & \\ 0 & 0 & 0 & \dots & a_k \end{bmatrix}$$

is positive. \blacksquare

Example 4 Let the polynomial in question be

$$p(z) = z^4 + 4z^3 + z^2 + 2z + \frac{1}{4}.$$

According to Theorem 1, p is stable if and only if the determinant of each of these four matrices is positive:

$$M_1 = [4], \quad M_2 = \begin{bmatrix} 4 & 2 \\ 1 & 1 \end{bmatrix}, \quad M_3 = \begin{bmatrix} 4 & 2 & 0 \\ 1 & 1 & \frac{1}{4} \\ 0 & 4 & 2 \end{bmatrix},$$

and

* Edward John Routh, 1831–1907.

† Adolf Hurwitz, 1859–1919.

$$M_4 = \begin{bmatrix} 4 & 2 & 0 & 0 \\ 1 & 1 & \frac{1}{4} & 0 \\ 0 & 4 & 2 & 0 \\ 0 & 1 & 1 & \frac{1}{4} \end{bmatrix}.$$

A computation shows that the determinants of M_1 and M_2 are positive, but those of M_3 and M_4 are zero. Hence, p is not stable. \square

EXERCISES FOR SECTION 3.1.1

1. Use the Routh–Hurwitz criteria to test the following polynomials for stability.

(a) $3 - 4z + 5z^2$ (c) $-\frac{1}{4} + 2z - \frac{1}{4}z^2$ (e) $1 + z + z^3$ (g) $3 + 2z + 2z^2 + 5z^3$ (i) $z^4 + z^3 + 2z^2 + z + \frac{1}{2}$ (k) $z^5 + z^4 + 5z^3 + 7z^2 + 4z + 8$	(b) $2 + z + \frac{1}{2}z^2$ (d) $1 + z - 3z^2 + z^3$ (f) $3 + 4z + z^2 + z^3$ (h) $1 + 4z^2 + 4z^4$ (j) $z^4 + 2z^3 + 3z^2 + 3z + 4$ (l) $z^4 + 4z^3 + 9z^2 + 8z + 5$
--	---
2. Let γ_R be the curve consisting of the semicircle Re^{it} , $-\pi/2 \leq t \leq \pi/2$, followed by the segment iy , $R \geq y \geq -R$. Show that a polynomial p , which does not vanish on the imaginary axis, is stable if and only if

$$\int_{\gamma_R} \frac{p'(z)}{p(z)} dz = 0,$$

for all large R .

3. Determine whether or not the systems determined by the following matrices are stable.

$$(a) A = \begin{bmatrix} -1 & 2 \\ 1 & -3 \end{bmatrix}$$

$$(b) A = \begin{bmatrix} 0 & -3 \\ 1 & -2 \end{bmatrix}$$

$$(c) A = \begin{bmatrix} 1 & 2 \\ 2 & -6 \end{bmatrix}$$

$$(d) A = \begin{bmatrix} 1 & w \\ -w & 1 \end{bmatrix}, \quad w \text{ complex}$$

$$(e) A = \begin{bmatrix} 1 & -1 & 1 \\ 1 & -2 & 1 \\ -2 & -4 & 0 \end{bmatrix}$$

$$(f) A = \begin{bmatrix} 1 & 3 & 1 \\ 1 & 2 & 3 \\ 2 & 0 & -1 \end{bmatrix}$$

4. Suppose the polynomial p given in (2) has all its zeros in the closed half-plane $Re w \leq 0$, and any zeros that lie on the imaginary axis are of order one. Show that any solution of the system (1) remains bounded as $t \rightarrow \infty$.

Work out in detail the solution of the system with

$$A = \begin{bmatrix} 0 & \omega \\ -\omega & 0 \end{bmatrix}, \quad \omega > 0,$$

to verify what happens in one special case.

5. Show by means of an example that if p given in (2) has a zero in the half-plane

$\operatorname{Re} z > 0$, then the homogeneous system $Ax = \dot{x}$ has unbounded solutions, and hence small perturbations in the initial conditions can result in unboundedly large variations in the resulting solution.

6. Examine the behavior as $t \rightarrow \infty$ of the solutions of the two equations

$$y'' = -y; \quad y(0) = x_0, y'(0) = 0$$

and

$$y'' = y; \quad y(0) = x_0, y'(0) = 0.$$

The first corresponds to the system

$$\begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} x = \dot{x},$$

and the second to the system

$$\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} x = \dot{x}.$$

The matrix $\begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$ has eigenvalues of $\pm i$, while the matrix $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ has eigenvalues of ± 1 .

7. In the theory of **feedback-control systems**, the question of the stability of the system takes the form of determining whether a function of the form $1 + (1/c)F$ has any zeros in the right half-plane; here F is a rational function and c is a nonzero constant. One answer to this question is provided by the Routh–Hurwitz condition. Write $F = P/Q$, where P and Q are polynomials with no common zero. Then $1 + (1/c)F$ has no zeros in the right half-plane if and only if the polynomial $cQ + P$ is stable. Another answer is provided by the **Nyquist stability criterion**, which is a variation on the argument principle. Let γ_R be the curve shown in Figure 3.6a, and let Γ_R be its image under F , shown in Figure 3.6b.




Figure 3.6

(Note that γ_R is oriented negatively; this is done because it is traditional to evaluate $F(iy)$ as y increases.) Show that if Γ_R winds around $-c$ a net positive number of times in the clockwise direction, then $1 + (1/c)F$ has a zero in the right half-plane (and so the system is unstable). In Figure 3.6b, those areas marked by slashes are values of c for which $1 + (1/c)F$ has a zero. More on the derivation of the stability criterion for feedback-control systems can be found in the book by A. Kyrala cited in the references at the end of Chapter 4.

3.2 Maximum Modulus and Mean Value

Suppose that f is an analytic function on a domain D . We saw in the beginning of Chapter 2 that if the range of f lies in a circle or on a straight line, then f is necessarily constant. We shall see now that a great deal more is true; either f is constant on D or the range of f is an open set.

Suppose that f is not identically constant, and let $w_0 = f(z_0)$ be an arbitrary point in the range of f . The function $f(z) - w_0$ is not identically zero, since f is not identically constant, and so $f(z) - w_0$ has a zero of order $m \geq 1$ at z_0 . Choose $r > 0$ so small that $f(z) - w_0$ has no zero in the region $0 < |z - z_0| \leq r$; this is possible because the zeros of a nonconstant analytic function are isolated (see Section 1). Let δ be the minimum value of $|f(z) - w_0|$ for all z with $|z - z_0| = r$, and let w be any point with $|w - w_0| < \delta$. Then on the circle $|z - z_0| = r$,

$$|(f(z) - w) - (f(z) - w_0)| = |w - w_0| < \delta \leq |f(z) - w_0|. \quad (1)$$

Therefore, Rouché's Theorem (Theorem 3 of Section 1) implies that the two functions $f - w$ and $f - w_0$ have an equal number of zeros within the circle $|z - z_0| = r$. However, $f - w_0$ has exactly m zeros, and hence so does $f - w$. This shows that each point w_0 in the range of f lies at the center of a small disc, which is also within the range of f . Thus, we have established this theorem. ■

THEOREM 1 Suppose that f is a nonconstant analytic function on a domain D . Then the range of $f(z)$, as z varies over D , is an open set. ■

We can draw an immediate conclusion from the work that preceded the statement of Theorem 1.

Suppose that f is a nonconstant analytic function on a domain D and that $f - f(z_0)$ has a zero of order m at z_0 . Then f is m -to-1 near z_0 ; in particular, if $f'(z_0) = 0$, then f is not one-to-one in any disc containing z_0 . (2)

Another very important consequence of Theorem 1 follows.

COROLLARY 1 The Maximum-Modulus Principle If f is a nonconstant analytic function on a domain D , then $|f|$ can have no local maximum on D .

Proof If $|f(z_0)| \geq |f(z)|$ for all z with $|z - z_0| < r$, then $f(z_0)$ lies on the boundary of the open set $W = \{f(z) : |z - z_0| < r\}$. This is in contradiction to the fact that W is an open set containing $f(z_0)$. ■

The maximum-modulus principle expressed in Corollary 1 implies, for example, that if we observe a nonconstant, sourceless, irrotational flow on a domain D , then at no point of D is the speed of the flow largest, or even locally largest. No matter where we look, the speed is always strictly greater at some nearby point.

The maximum-modulus principle (Corollary 1) has ramifications for the real and imaginary parts of an analytic function. Suppose that f is a nonconstant analytic function on a domain D ; let $u = \operatorname{Re} f$ and $g = e^f$. Then both g and $1/g$ are analytic on D , so neither of the functions

$$|g| = e^u \quad \text{and} \quad \left| \frac{1}{g} \right| = e^{-u}$$

has a local maximum in D ; consequently, u has no local maxima and no local minima in D . This is stated formally in this way:

If f is analytic and nonconstant on a domain D , then $\operatorname{Re} f$ has no local maxima and no local minima on D . (3)

Let us continue this line of thought a bit further. Let D be a bounded domain, and let B be the boundary of D . The set $D \cup B$ is both closed and bounded. There is a result from real variables that asserts that a continuous real-valued function on a closed and bounded set actually *attains* its maximum value.* Let us apply this to each of the three functions $|f|$, $\operatorname{Re} f$, and $-\operatorname{Re} f$, where f itself is analytic on D and continuous on $D \cup B$. We know that each of these three functions is continuous on $D \cup B$ and so must attain its maximum somewhere on $D \cup B$. If f is nonconstant, then the maximum can not be attained on D itself. Hence, the maximum must be attained on B , the boundary of D . Quite obviously, the same conclusion holds if, in fact, f is constant on D . Thus, in all cases, $|f|$, $\operatorname{Re} f$, and $-\operatorname{Re} f$ attain their maximum values on B , the boundary of D .

If f is analytic on a bounded domain D and continuous on $D \cup B$, where B is the boundary of D , then each of $|f|$, $\operatorname{Re} f$, and $-\operatorname{Re} f$ attains its maximum value on B . (4)

In particular, one consequence of (4) is that if $\operatorname{Re} f$ is zero everywhere on B , then $\operatorname{Re} f$ is zero throughout D , so f itself is constant on all of D . It should be stressed that D 's being bounded is not a superfluous hypothesis here. For instance, $f(z) = iz$ is analytic and nonconstant on the whole plane, $\operatorname{Re} f = 0$ on the real axis, and yet f is not constant on the half-plane $\{\operatorname{Im} z > 0\}$, whose boundary is the real axis.

One of the best-known applications of the maximum-modulus principle is the following result, known for historical reasons as Schwarz's Lemma.

* See, for instance, W. Rudin. *Principles of mathematical analysis*. 3rd ed. New York: McGraw-Hill, 1976.

THEOREM 2 Schwarz's Lemma Suppose that f is analytic in the disc $|z| < 1$, that $f(0) = 0$, and that $|f(z)| \leq 1$, for all z in the disc. Then

$$|f(z)| \leq |z|, \quad |z| < 1.$$

Equality can hold for some $z \neq 0$ only if $f(z) = \lambda z$, where λ is a constant of absolute value 1.

Proof Since $f(0) = 0$, we know that $g(z) = f(z)/z$ is also analytic on $|z| < 1$. For $|z| = r$,

$$|g(z)| = \frac{|f(z)|}{r} \leq \frac{1}{r}.$$

By the maximum-modulus principle, the inequality $|g(z)| \leq 1/r$ is true for $|z| < r$ as well. Since r can be made as near 1 as we like, we must have $|g(z)| \leq 1$ if $|z| < 1$; thus, $|f(z)| \leq |z|$ for $|z| < 1$. Furthermore, if $|f(z_0)| = |z_0|$ for some $z_0 \neq 0$, then $1 = |g(z_0)|$; consequently, $|g(z)|$ has an interior maximum. This implies that $g(z)$ is a constant λ , $|\lambda| = 1$, and gives the conclusion that $f(z) = \lambda z$. ■

You may contrast the result in Theorem 2 to the case of real-valued functions of a real variable. For instance, the function

$$u(x) = \frac{2x}{x^2 + 1}, \quad -1 \leq x \leq 1,$$

satisfies $-1 \leq u(x) \leq 1$, $u'(x)$ is continuous, $u(0) = 0$, yet $|u(x)| > |x|$ if $1 > |x| > 0$.

Mean Value

Theorem 4 of Section 3, Chapter 2 (Cauchy's Formula) gives us

$$f(z_0) = \frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{\zeta - z_0} d\zeta,$$

when γ is a circle and z_0 is inside γ . If we take z_0 to be the center of the circle, then $\zeta = z_0 + re^{it}$, $0 \leq t \leq 2\pi$, $d\zeta = ire^{it} dt$, and we obtain

$$f(z_0) = \frac{1}{2\pi} \int_0^{2\pi} f(z_0 + re^{it}) dt. \tag{5}$$

The result expressed in (5) is the **Mean-Value Theorem** for analytic functions. This result can also be obtained by replacing $f(z_0 + re^{it})$ with the power-series expansion for f about the point z_0 and integrating term by term.

In formula (5), take the real part of both sides and write $u = \operatorname{Re} f$. We find that

$$u(z_0) = \frac{1}{2\pi} \int_0^{2\pi} u(z_0 + re^{it}) dt. \quad (6)$$

As we will see in Chapter 4, those functions u , which are the real parts of some analytic function f , are of great importance in applications. Formula (6), which shows us how to “recover” $u(z_0)$ from the values of u on the circle $|z - z_0| = r$, is but a special case of a more general formula that allows us to recover $u(\zeta)$ for all ζ inside $|z - z_0| = r$ from the values of u on the circle $|z - z_0| = r$.

EXERCISES FOR SECTION 3.2

- Let $f(z) = z^2/(z + 2)$; find the maximum value of $|f(z)|$ as z varies over the disc $|z| \leq 1$.
- Let $f(z) = ze^z$; find the maximum value of $|f(z)|$ as z varies over the region $D = \{x + iy: x^2 + y^2 \leq 4, x \geq 0, y \geq 0\}$.
- Find the maximum value of $|g(z)|$,

$$g(z) = \frac{z}{4z^2 - 1},$$

as z varies over the region $\{z: |z| \geq 1\}$.

- (a) If $|z| = r < 1$, show that

$$|e^z - 1| \geq r - \frac{r^2}{2-r}.$$

(Hint:

$$e^z - 1 = z + \frac{z^2}{2} + \frac{z^3}{3!} + \dots,$$

so

$$|e^z - 1| = |z| \left| 1 + \frac{z}{2} + \frac{z^2}{3!} + \dots \right| \geq r \left\{ 1 - \sum_{k=1}^{\infty} \frac{r^k}{2^k} \right\}.$$

- (b) Use (a) to show that the equation $z(e^z - 1) = w$ has exactly two solutions z with $|z| < \frac{1}{2}$ whenever $|w| < \frac{1}{6}$. (Hint: Look at the work around (1), preceding the statement of Theorem 1.)
- The function

$$S(z) = \exp \left[\frac{1+z}{1-z} \right]$$

is analytic on the disc $\{z: |z| < 1\}$ and $|S(z)| = 1$ if $|z| = 1$, $z \neq 1$. However, $S(0) = e > 1$. Why does this not contradict the maximum-modulus principle?

6. Let D be a bounded domain with boundary B . Suppose that f and g are both analytic on D and continuous on $D \cup B$, and suppose further that $\operatorname{Re} f(z) = \operatorname{Re} g(z)$ for all $z \in B$. Show that $f = g + i\alpha$ in D , where α is a real constant.
7. Let F be analytic and nonconstant on the disc $|z - z_0| < R$, and suppose that $\operatorname{Re}(F(z_0)) = 0$. Show that on every circle $|z - z_0| = r$, $0 < r < R$, $\operatorname{Re} F$ assumes both positive and negative values.
8. Let f be analytic on a bounded domain D and continuous on $D \cup B$, where B is the boundary of D . Show that if f is never zero on D , then the minimum of $|f|$ is assumed on B . You will need to use the fact that $|f|$ does, indeed, assume a minimum somewhere on $D \cup B$.
9. Suppose that f is analytic on a domain D , which contains a simple closed curve γ and the inside of γ . If $|f|$ is constant on γ , then either f is constant or f has a zero inside γ .
10. Let f be a nonconstant entire function and U be an open set in the plane. Show that there is a z_0 such that $f(z_0) \in U$.
11. Suppose that p is a polynomial of degree n and that $|p(z)| \leq M$ if $|z| = 1$. Show that $|p(z)| \leq M|z|^n$ if $|z| \geq 1$. (Hint: Apply the maximum-modulus principle to $f(z) = p(z)/z^n$ on the domain $|z| > 1$, including the point “ ∞ .”)

In Exercises 12 to 14, f is analytic on the disc $|z| < 1$ and satisfies $|f(z)| \leq M$ if $|z| < 1$.

12. Suppose that $f(\alpha) = 0$ for some α , $|\alpha| < 1$. Show that

$$|f(z)| \leq M \left| \frac{z - \alpha}{1 - \bar{\alpha}z} \right|, \quad |z| < 1.$$

13. Show that

$$M \left| \frac{f(z) - f(z_0)}{M^2 - \overline{f(z_0)f(z)}} \right| \leq \left| \frac{z - z_0}{1 - \bar{z}_0 z} \right| \quad \text{for all } z, z_0 \text{ in } \{w: |w| < 1\}.$$

14. Suppose that $f^{(k)}(0) = 0$ for $k = 0, \dots, N$. Show that $|f(z)| \leq M|z|^{N+1}$ for all z , $|z| < 1$.
15. Let g be analytic and nonconstant on a simply-connected domain D ; for instance, D could be a disc. Show that the set $\{z: \operatorname{Re} g(z) = c\}$ contains no simple closed curve. (Hint: Use Exercise 9 with $f = \exp(g)$.)
16. Use (6) to show that the real part of a nonconstant analytic function has no strict local maxima or local minima.
17. Let f and g be analytic on a domain that contains a simple closed curve γ and the inside of γ . Assume further that f is never zero inside γ . Show that if $|f(z)| \geq |g(z)|$ for all z in γ , then $|f(z)| \geq |g(z)|$ for all z inside γ as well. Give an example to show that the hypothesis “ $f \neq 0$ inside γ ” is essential for the validity of the conclusion.

18. Suppose that f is analytic in $\{z: 0 < |z - z_0| < R\}$ and that f has a pole of order m , $m \geq 1$, at z_0 . Show that there is a large number M with this property: For each w_0 with $|w_0| > M$, the equation $f(z) = w_0$ has exactly m solutions z , z near z_0 .

Positive Rational Functions and Stable Polynomials*

If p is a polynomial, define $\tilde{p}(z) = \overline{p(-\bar{z})}$.

19. Show that $\tilde{\tilde{p}} = p$. For which polynomials Q is $\tilde{Q} = Q$?
20. Suppose that p is stable (see Section 3.1.1). Show that p and \tilde{p} have no common zero.
21. Suppose that p is stable. Show that the rational function $r = \tilde{p}/p$ maps the right half-plane, $\operatorname{Re} z > 0$, into the disc $\{w: |w| < 1\}$. Conversely, show that if $r = \tilde{p}/p$ maps $\operatorname{Re} z > 0$ into $|w| < 1$ and if p and \tilde{p} have no common zero, then p is stable.
22. A rational function R is positive if $\operatorname{Re}\{R(z)\} > 0$ whenever $\operatorname{Re} z > 0$. Show that
 - (a) R is positive if and only if $1/R$ is positive.
 - (b) If R is positive, then R has no zero or pole in the half-plane $\operatorname{Re} z > 0$, and any zero or pole on the imaginary axis has order one. (Hint: By (a), it suffices to consider just zeros.)
23. Suppose that p is stable. Show that $R = (p - \tilde{p})/(p + \tilde{p})$ is positive. Conversely, show that if p and \tilde{p} have no common zero and if $R = (p - \tilde{p})/(p + \tilde{p})$ is positive, then p is stable.

3.3 Linear Fractional Transformations

A linear fractional transformation T is a rational function of the special form

$$T(z) = \frac{az + b}{cz + d},$$

where a , b , c , and d are complex numbers and $ad - bc \neq 0$. The restriction $ad - bc \neq 0$ is essential, for otherwise

$$T'(z) = \frac{ad - bc}{(cz + d)^2} = 0 \quad \text{for all } z,$$

so T is identically constant.

A linear fractional transformation is a one-to-one function. For suppose that

$$\frac{az_1 + b}{cz_1 + d} = T(z_1) = T(z_2) = \frac{az_2 + b}{cz_2 + d}.$$

Then

$$(az_1 + b)(cz_2 + d) = (az_2 + b)(cz_1 + d),$$

and, after some cancellation,

$$bcz_2 + adz_1 = adz_2 + bcz_1,$$

or

$$(ad - bc)z_1 = (ad - bc)z_2.$$

Consequently, $z_1 = z_2$. Hence, the function T maps distinct points onto distinct images. Note also that T has a pole of order one at $-d/c$ and $\lim_{|z| \rightarrow \infty} T(z) = a/c$.

Since T is one-to-one, there is a function T^{-1} that is the inverse of T in the sense of the composition of functions:

$$T^{-1}(T(z)) = z$$

for all z . T^{-1} is also a linear fractional composition, as this computation shows.

$$T(z) = w = \frac{az + b}{cz + d}$$

implies that $czw - az = b - dw$, and thus,

$$T^{-1}(w) = z = \frac{-dw + b}{cw - a}.$$

Hence, a linear fractional transformation is a one-to-one mapping of the complex plane plus the point at ∞ onto itself. Conversely, a one-to-one (analytic) mapping of the complex plane plus “ ∞ ” onto itself is a linear fractional transformation. (See Exercise 3, Section 4.)

Example 1 Show that a linear fractional transformation T of the form

$$T(z) = \lambda \frac{a - z}{1 - \bar{a}z}, \quad |\lambda| = 1, |a| < 1 \tag{1}$$

maps the disc $\Delta = \{z: |z| < 1\}$ onto itself.

Solution We begin by noting that

$$|T(z)|^2 = \frac{|a - z|^2}{|1 - \bar{a}z|^2} = \frac{|a|^2 - 2 \operatorname{Re}(\bar{a}z) + |z|^2}{1 - 2 \operatorname{Re}(\bar{a}z) + |a|^2|z|^2},$$

so $1 > |T(z)|^2$ exactly when

$$1 + |a|^2|z|^2 > |a|^2 + |z|^2.$$

This last inequality is equivalent to

$$(1 - |a|^2)(1 - |z|^2) > 0,$$

which is true exactly when $z \in \Delta$. Hence, T does map Δ into itself. The inverse function of T , in the sense of composition of functions, is

$$T^{-1}(w) = \bar{\lambda} \frac{a\bar{\lambda} - w}{1 - \bar{a}\bar{\lambda}w}.$$

(See above.) This transformation has the same general form as T , with a replaced by $a\bar{\lambda}$ (a point which also lies in Δ) and $\bar{\lambda}$ in place of λ . Hence, T^{-1} also maps Δ into itself. To see that T maps Δ onto itself, let w_0 be any point of Δ ; then $z_0 = T^{-1}(w_0)$ lies in Δ by the comments above and so $w_0 = T(T^{-1}(w_0)) = T(z_0)$, showing that the range of T is all of Δ . \square

Fixed Points and Triples to Triples

A linear fractional transformation that is not identically equal to z has at most two distinct **fixed points**; that is, points z for which

$$T(z) = z.$$

This is evident from the fact that z is a solution of the equation $T(z) = z$ exactly when z is a root of the quadratic equation

$$0 = cz^2 + (d - a)z - b,$$

and, of course, a quadratic equation has at most two distinct roots. It is a consequence of this that if T and S are two linear fractional transformations that are equal at three distinct points, say $T(z_j) = S(z_j)$ for $j = 1, 2, 3$, then $T(z) = S(z)$ for all z . This is because the linear fractional transformation $S^{-1}(T(z))$ has three distinct fixed points:

$$S^{-1}(T(z_j)) = z_j, \quad j = 1, 2, 3.$$

Hence, $S^{-1}(T(z)) = z$ for all z , and thus, $T(z) = S(z)$ for all z .

On the other hand, if three distinct complex numbers z_1, z_2 , and z_3 are given and if any other three distinct complex numbers w_1, w_2 , and w_3 are chosen, there is a necessarily unique linear fractional transformation L with $L(z_j) = w_j, j = 1, 2, 3$. We derive its form in this way. Set

$$T(z) = \left(\frac{z - z_1}{z - z_3} \right) \left(\frac{z_2 - z_3}{z_2 - z_1} \right).$$

Then $T(z_1) = 0$, $T(z_2) = 1$, and $T(z_3) = \infty$. Let

$$S(w) = \left(\frac{w - w_1}{w - w_3} \right) \left(\frac{w_2 - w_3}{w_2 - w_1} \right),$$

so that $S(w_1) = 0$, $S(w_2) = 1$, and $S(w_3) = \infty$. L is given by

$$L(z) = S^{-1}(T(z)).$$

We can carry out a similar sort of computation if one of z_1 , z_2 , and z_3 or w_1 , w_2 , and w_3 is ∞ ; this is left for the exercises at the end of this section.

Example 2 Find the linear fractional transformation that sends $0, 1, 2$ to $-1, 0, 4$, respectively.

Solution We use the formulas to obtain

$$T(z) = \left(\frac{z - 0}{z - 2} \right) \left(\frac{1 - 2}{1 - 0} \right) = \frac{z}{2 - z}$$

and

$$S(w) = \left(\frac{w - (-1)}{w - 4} \right) \left(\frac{0 - 4}{0 - (-1)} \right) = -4 \frac{w + 1}{w - 4}.$$

Hence,

$$S^{-1}(z) = 4 \frac{z - 1}{z + 4},$$

and so

$$L(z) = S^{-1}(T(z)) = \frac{8z - 8}{-3z + 8}. \quad \square$$

Lines and Circles

It is an important property of a linear fractional transformation that it maps each circle onto another circle or onto a straight line and each straight line onto another straight line or a circle. We show this below.

First, if $T(z) = az + b$, $a \neq 0$, then the assertion that T maps circles and straight lines to the same type of figure is quite clear. The circle

$$C: \{z: |z - z_0| = r\}$$

is transformed to the circle

$$C': \{w: |w - (az_0 + b)| = |a|r\},$$

and the straight line

$$L: \{z: \operatorname{Re}(Az + B) = 0\}$$

is transformed to the straight line

$$L': \{w: \operatorname{Re}[(A/a)w + B - b(A/a)] = 0\}.$$

Thus, we henceforth assume that

$$T(z) = \frac{az + b}{cz + d}, \quad c \neq 0.$$

Now

$$T(z) = \frac{az + b}{cz + d} = \frac{1}{c} \left\{ \frac{bc - ad}{cz + d} + a \right\}, \quad c \neq 0,$$

and we see from this that T is actually the composition of several simpler linear fractional transformations. Precisely, set

$$U(z) = cz + d, \quad V(w) = \frac{1}{w}, \quad \text{and} \quad W(\zeta) = \frac{1}{c}[(bc - ad)\zeta + a].$$

Then

$$T(z) = W(V(U(z))).$$

From the foregoing discussion, we know that both U and W send circles to circles and lines to lines. Consequently, the desired conclusion will be reached if we can show that V maps circles and lines to circles and lines. The equation

$$\alpha(x^2 + y^2) + \beta x + \gamma y = \delta,$$

where α, β, γ , and δ are real and not all of α, β, γ are zero, represents either a circle (iff $\alpha \neq 0$ and $\beta^2 + \gamma^2 + 4\alpha\delta > 0$) or a straight line (iff $\alpha = 0$). Now

$$\frac{1}{z} = \frac{x}{x^2 + y^2} + i \frac{-y}{x^2 + y^2} = u + iv,$$

so replacing z by $1/z$ yields the equation

$$\delta(u^2 + v^2) - \beta u + \gamma v = \alpha.$$

This is again the equation (in u, v coordinates) of either a line or a circle. You will have no difficulty verifying that a circle or line through the origin is transformed by inversion to a line, and that a circle or a line *not* through the origin is transformed




Figure 3.7

by inversion to a circle (which goes through the origin when and only when a line was inverted) (Fig. 3.7).

There is an interesting connection between certain linear fractional transformations and “rigid motions” of the extended plane with respect to the chordal distance; see Exercises 24 to 27.

The Circulation Produced by a Linear Fractional Transformation*

Let

$$T(z) = \frac{z+2}{2z+1};$$

this linear fractional transformation fixes the two points 1 and -1 . We shall show that if C is a circle centered on the imaginary axis that passes through both 1 and -1 , then T maps C onto itself. (Of course, 1 and -1 are the only fixed points of T .) Let $i\alpha$ be the center of C , α real. Then the radius of C is $r = \sqrt{\alpha^2 + 1}$, so a point z lies on C precisely when $|z - i\alpha|^2 = 1 + \alpha^2$. Hence, the equation of C is

$$x^2 + (y - \alpha)^2 = 1 + \alpha^2$$

or

$$x^2 + y^2 - 2\alpha y = 1.$$

Now

$$\begin{aligned} |T(z) - i\alpha|^2 &= \left| \frac{z+2}{2z+1} - i\alpha \right|^2 \\ &= \frac{(x+2+2\alpha y)^2 + (y-\alpha-2\alpha x)^2}{(2x+1)^2 + (2y)^2}, \end{aligned}$$

and a modest amount of computation, using the fact that $x^2 + y^2 - 2xy = 1$, shows that

$$|T(z) - i\alpha|^2 = 1 + \alpha^2,$$

which, of course, is exactly the statement that $T(z)$ lies in the circle C if z is in C . Set

$$T_1(z) = T(z) = \frac{z+2}{2z+1},$$

and then

$$T_2(z) = T_1(T_1(z)) = \frac{5z+4}{4z+5},$$

and, in general,

$$T_{n+1}(z) = T_n(T_1(z)), \quad n = 1, 2, 3, \dots$$

For instance,

$$T_3(z) = \frac{13z+14}{14z+13}, \quad T_4(z) = \frac{41z+40}{40z+41}.$$

In general, we see that $T_n(z) \rightarrow 1$ as $n \rightarrow \infty$ so long as $z \neq -1$. Thus, T sets up a circulation in the plane with a source at -1 and a sink at $+1$: each point z in the plane is on precisely one circle C through z , 1 , and -1 (the real axis counts as a circle for this purpose); T carries C onto itself and pushes z toward 1 along the circle. Thus, the points in the plane move toward 1 and away from -1 under the action of T . We note that $T'(1) = -1/3$ and $T'(-1) = -3$. Thus, the sink is located at that fixed point at which the modulus of the derivative is less than 1 (Fig. 3.8).

You should convince yourself that the critical issue in the foregoing discussion is that the linear fractional transformation fixes the two points 1 and -1 , and not its special form. A similar phenomenon occurs for a linear fractional transformation that fixes the two distinct points p and q (see Exercise 9 at the end of this section). You should also look back at the Circles of Appolonius (Fig. 1.11), with the foregoing in mind.

Example 3 This example continues the preceding discussion. We shall investigate what happens when the two fixed points move toward each other. Set

$$T_\varepsilon(z) = \frac{z+\varepsilon^2}{z+1}, \quad 0 < \varepsilon < 1.$$

T_ε fixes ε and $-\varepsilon$, so (exactly as above) T_ε sets up a circulation using the family of circles centered on the imaginary axis and passing through ε and $-\varepsilon$ (see Fig. 3.9a). As $\varepsilon \rightarrow 0$, $T_\varepsilon(z) \rightarrow T_0(z) = z/(z+1)$ and we see that this T_0 sets up a circulation for





Figure 3.8


Circulation of $T_\epsilon(z) = \frac{z + \epsilon^2}{z + 1}$
with fixed points $\pm \epsilon$



(a)

A dipole located at the origin;
the circulation is given by

$$T(z) = \frac{z}{z + 1}$$



(b)

Figure 3.9

the family of circles centered on the imaginary axis and passing through the origin. This circulation is termed a **dipole**. Note that $T_0(z) = z/(z + 1)$ has only one fixed point and that it is a double root of $T_0(z) = z$ at $z = 0$. A similar dipole will appear as the circulation of any linear fractional transformation with a single (distinct) fixed point p , when p is a double root of $T(z) = z$ (Fig. 3.9b).

EXERCISES FOR SECTION 3.3

1. Let z_1, z_2, z_3 and w_1, w_2, w_3 be two triples of distinct complex numbers. Set

$$\alpha = \frac{z_2 - z_3}{z_2 - z_1} \quad \text{and} \quad \beta = \frac{w_2 - w_3}{w_2 - w_1}.$$

Show that the linear fractional transformation that maps z_j to w_j for $j = 1, 2, 3$ is given explicitly by

$$L(z) = \frac{z(\alpha w_3 - \beta w_1) + (\beta w_1 z_3 - \alpha z_1 w_3)}{z(\alpha - \beta) + (\beta z_3 - \alpha z_1)}.$$

2. Verify that the linear fractional transformation

$$S(z) = \frac{z - z_1}{z_2 - z_1}$$

maps z_1 to 0, z_2 to 1, and ∞ to ∞ .

3. Verify that the linear fractional transformation

$$T(z) = \frac{z_2 - z_1}{z - z_1}$$

maps z_1 to ∞ , z_2 to 1, and ∞ to 0.

4. In each case, find the linear fractional transformation that maps the first triple (z_1, z_2, z_3) onto the second triple (w_1, w_2, w_3) .
- $(1, i, -1)$ onto $(-1, i, 1)$
 - $(1, 4, \infty)$ onto $(0, 1 - i, 1 + i)$
 - $(1, 0, i)$ onto $(1, 0, 1 + i)$
 - $(\infty, -1, i)$ onto $(1, 0, 1 - i)$
 - $(i, 2, -i)$ onto $(-i, 3, i)$
5. Find a linear fractional transformation that carries
- the circle $|z| = 1$ onto the line $\operatorname{Re}((1 + i)w) = 0$
 - the circle $|z| = 1$ onto the circle $|w - 1| = 1$
 - the real axis onto the line $\operatorname{Re} w = \frac{1}{2}$
 - the circle $|z - z_0| = r$ onto the circle $|w| = 1$
 - the line $\operatorname{Re} z = \frac{1}{2}$ onto the circle $|w - 4i| = 4$
6. Show that a linear fractional transformation T that maps the circle $|z| = 1$ onto itself has the form

$$T(z) = \lambda \frac{z - \gamma}{1 - \bar{\gamma}z}, \quad |\lambda| = 1, \quad |\gamma| \neq 1, \quad \text{or} \quad Tz = \frac{\lambda}{z}, \quad |\lambda| = 1.$$

7. In each of the following, find a linear fractional transformation T with the given property.

- (a) T fixes the points 0 and 1 and $T(i) = \infty$.
- (b) T maps the real axis onto itself and the imaginary axis onto the circle $|w - \frac{1}{2}| = \frac{1}{2}$.
- (c) T maps the real axis onto itself and the imaginary axis onto the circle $|w - \frac{5}{4}| = \frac{3}{4}$.
- (d) T maps the real axis onto itself and the line $y = x$ onto the circle

$$|w + i| = \sqrt{2}.$$

8. (a) In Exercise 7(c), determine what happens to the first quadrant under the action of T .
 (b) In Exercise 7(d), determine what happens to the sector $\{z: 0 \leq y \leq x\}$ under the action of T .
9. Discuss the circulation set up by each of the following linear fractional transformations; find the fixed points (if any) and which way the circulation rotates.
 - (a) $T(z) = z/(z + 2)$
 - (b) $T(z) = (z + 4)/(z + 1)$
 - (c) $T(z) = 3z - 2$
 - (d) $T(z) = -\frac{1}{2}z$
 - (e) $T(z) = z + 1$ (no fixed points)
10. (a) If T is a linear fractional transformation that maps the circle C_1 onto the circle C_2 , shows that T carries the inside of C_1 onto either the inside of C_2 or onto the outside of C_2 . (**Hint:** Suppose a point z_0 inside C_1 goes to a point w_0 inside C_2 . Let z_1 be any other point inside C_1 and L the line segment joining z_0 and z_1 . T carries L to an arc of a circle (or a straight line segment) joining w_0 and $w_1 = T(z_1)$ and this arc (or segment) does not meet C_2 . [Why?])
 (b) T maps the point $z_0 = -d/c$ to ∞ ; tell how this decides whether T maps the inside of C_1 onto the inside of C_2 or onto the outside of C_2 .

Fixed Points

11. Show that a linear fractional transformation that fixes the two points 1 and -1 has the form

$$T(z) = \frac{z + \xi}{\xi z + 1}$$

- if $\xi = T(0) \neq \infty$. (If $Tz = 1/z$, then $T(1) = 1$, $T(-1) = -1$.)
12. Suppose that T fixes the two points p and q , $p \neq q$. Let S be any linear fractional transformation with $S(1) = p$, $S(-1) = q$, and let $U = S^{-1} \circ T \circ S$. Show that U fixes 1 and -1 .
 13. Suppose that U is a linear fractional transformation that fixes 1 and -1 . Show that $U'(1) = \{U'(-1)\}^{-1}$.
 14. Let T be a linear fractional transformation fixing p and q , $p \neq q$. Use Exercise 13 to show that $T'(p) = \{T'(q)\}^{-1}$.
 15. Show that the linear fractional transformation

$$T(z) = \frac{1-z}{1+z}$$

satisfies $T(T(z)) = z$ for all z . Find the general form of those linear fractional transformations S with $S(S(z)) = z$. (Equivalently, $S(z) = S^{-1}(z)$.)

Cross-Ratio

The **cross-ratio** of four distinct complex numbers z_0, z_1, z_2 , and z_3 is defined to be the complex number

$$(z_0, z_1, z_2, z_3) = \frac{z_0 - z_1}{z_0 - z_2} \frac{z_3 - z_2}{z_3 - z_1}.$$

16. Show that if T is a linear fractional transformation with $Tz_j = w_j$ for $j = 1, 2, 3$, then

$$(z, z_1, z_2, z_3) = (T(z), w_1, w_2, w_3) \quad \text{for all } z.$$

17. Let z_1, z_2, z_3 lie in a circle or on a straight line C . Show the point z_0 is also on C if and only if the cross-ratio (z_0, z_1, z_2, z_3) is real. (**Hint:** By Exercise 16, it is enough to do this for $C =$ real axis and $z_1 = 1, z_2 = 0, z_3 = -1$. The remainder is a computation: The cross-ratio of $(z, 1, 0, -1)$ is real if and only if $\zeta = (z-1)/z$ is real. Show that ζ is real if and only if z is real.)
18. Use Exercises 16 and 17 to give another demonstration of the fact that a linear fractional transformation T carries circles and lines to circles and lines.

Reflection*

Let L be a straight line and z be a point in the complex plane. The **reflection** of z over L is the unique point z^* located on the line L' through z and perpendicular to L , lying on the other side of L from z , and at a distance from L equal to that of z from L (Fig. 3.10a). Let C be the circle $|\zeta - \zeta_0| = r$ and z a point of the complex plane, $z \neq \zeta_0$. The **reflection** of z over C is the unique point z^* located on the ray from ζ_0 to ∞ , which passes through z , lying on the other side of C from z and satisfying $|z - \zeta_0||z^* - \zeta_0| = r^2$.




Figure 3.10

19. (a) If z is a point in L or C , show that $z^* = z$.
 (b) If L is the real axis, show that $z^* = \bar{z}$, the complex conjugate of z .
 (c) Show that $(z^*)^* = z$ for any L , C , and z .
20. If C is the circle $|\zeta - \zeta_0| = r$, show that

$$z^* = \frac{r^2}{\bar{z} - \bar{\zeta}_0} + \zeta_0.$$

21. Let z_1, z_2 , and z_3 be points on the circle C of Exercise 20; fill in the reasons for each of the following equalities (all but one use the invariance of the cross-ratio under linear fractional transformations).

$$\begin{aligned} (z^*, z_1, z_2, z_3) &= \left(\frac{r^2}{\bar{z} - \bar{\zeta}_0} + \zeta_0, z_1, z_2, z_3 \right) \\ &= \left(\frac{r^2}{\bar{z} - \bar{\zeta}_0}, z_1 - \zeta_0, z_2 - \zeta_0, z_3 - \zeta_0 \right) \\ &= \left(\bar{z} - \bar{\zeta}_0, \frac{r^2}{z_1 - \zeta_0}, \frac{r^2}{z_2 - \zeta_0}, \frac{r^2}{z_3 - \zeta_0} \right) \\ &= (\overline{z - \zeta_0}, \overline{z_1 - \zeta_0}, \overline{z_2 - \zeta_0}, \overline{z_3 - \zeta_0}) \\ &= (\overline{z}, \overline{z_1}, \overline{z_2}, \overline{z_3}). \end{aligned}$$

22. (a) Conclude from Exercise 21 that z^* is the reflection of z over the circle C if and only if

$$(z, z_1, z_2, z_3) = (\overline{z^*}, \overline{z_1}, \overline{z_2}, \overline{z_3}) \quad (2)$$

for any triple z_1, z_2, z_3 of distinct points in C .

- (b) Let L be a straight line through z_1 and z_2 ; choose z_3 to be ∞ and modify Exercises 20 and 21 to conclude that (2) also characterizes z^* for reflection over L .
23. Show that the net result of two successive reflections (over lines/circles) is a linear fractional transformation.

Chordal Distance and Linear Fractional Transformations

The **chordal distance** between two points z and w is

$$\chi(z, w) = \frac{|z - w|}{(1 + |z|^2)^{1/2}(1 + |w|^2)^{1/2}}$$

if neither of the points is “ ∞ ”; the chordal distance from z to ∞ is

$$\chi(z, \infty) = (1 + |z|^2)^{-1/2}.$$

(The chordal distance is another way of measuring how far it is from z to w ; for z and w close together, it is almost a multiple of the usual distance $|z - w|$. Hence, any sequence that converges with respect to the usual distance also converges with respect to the chordal distance. However, no points are further apart than 1 in chordal distance. The name *chordal* comes from the fact that z and w can be viewed as points on a sphere in three-dimensional space of radius $\frac{1}{2}$ centered at $(0, 0, \frac{1}{2})$; then $\chi(z, w)$ is the length of the chord joining z to w .)

The next several problems delineate an interesting connection between linear fractional transformations and the chordal distance.

24. Let a and b be complex numbers with $|a|^2 + |b|^2 = 1$, and set

$$Tz = \frac{az + b}{-\bar{b}z + \bar{a}}. \quad (3)$$

- Show that T preserves chordal distance; that is, $\chi(Tz, Tw) = \chi(z, w)$, for all z, w .
25. If α is a given point in the extended plane, show that there is a linear fractional transformation of the form (3) with $T\alpha = 0$.
26. Suppose that f and g are two functions mapping the extended plane into itself, both of which preserve chordal distance. Show that their composition $f \circ g$ also preserves chordal distance.
27. Let f be a function that is analytic on the extended plane except possibly for isolated poles, and suppose that f preserves chordal distance: $\chi(f(z), f(w)) = \chi(z, w)$. Show that f is a linear fractional transformation of the form (3). (Hint: Compose f with a T of the form (3) chosen such that $T(f(0)) = 0$. Then determine the form of $T \circ f$.)

3.4 Conformal Mapping

Conformal maps are functions that preserve angles between curves in a sense that we will make precise below. They are indispensable tools in studying flows and fields and in solving boundary-value problems, as we shall see in Section 3.4.1 and Chapter 4. In this section and the next, we shall study some of the basic facts about conformal mapping.

Suppose that γ is the range of a smooth curve $z(t)$, $a \leq t \leq b$, and γ passes through the point $z_0 = z(t_0)$, $a < t_0 < b$. The curve γ has a tangent vector $z'(t_0)$ at z_0 , and we suppose that $z'(t_0)$ is not zero. We now investigate what happens to the curve γ when we apply an analytic function f to it. The curve γ is transformed into a new curve Γ in the w -plane; Γ is given by

$$\Gamma: w(t) = f(z(t)), \quad a \leq t \leq b.$$

Furthermore, by the familiar chain rule, we find that the tangent vector to Γ at $w_0 = f(z_0)$ is

$$w'(t_0) = f'(z_0)z'(t_0).$$




Figure 3.11

See Figure 3.11. In particular, we obtain the two relations

$$|w'(t_0)| = |f'(z_0)| |z'(t_0)| \quad (1)$$

and

$$\arg w'(t_0) = \arg(f'(z_0)) + \arg(z'(t_0)). \quad (2)$$

We make the natural assumption that $f'(z_0) \neq 0$ and then note that the transformation of γ into Γ by f has these two characteristics: The tangent vector is scaled in length by a factor $|f'(z_0)|$, and the tangent vector is rotated through an angle $\psi_0 = \arg(f'(z_0))$.

Suppose now that γ_1 and γ_2 are two smooth curves that intersect at the point $z_0 = z_1(t_0) = z_2(s_0)$. We define the **angle between γ_1 and γ_2 at z_0** to be the angle θ measured counterclockwise from the tangent vector $z'_1(t_0)$ to the tangent vector $z'_2(s_0)$, if neither of these tangent vectors is zero. In this case,

$$\theta = \arg(z'_2(s_0)) - \arg(z'_1(t_0)),$$

for appropriate determinations of the argument of the tangent vectors $z'_2(s_0)$ and $z'_1(t_0)$. See Figure 3.12. (This definition of the angle between two curves breaks down if one of the vectors $z'_1(t_0)$ or $z'_2(s_0)$ is zero; this will not be a matter of great concern to us.)

Suppose now that ϕ is a function, perhaps not analytic, defined in the disc $|z - z_0| < r$ and satisfying $\phi(z) \neq \phi(z_0)$ if $0 < |z - z_0| < r$. We say that ϕ is **conformal** at z_0 if, whenever two curves γ_1 and γ_2 meet at z_0 , the angle from Γ_1 to Γ_2 (their images under ϕ) is equal to the angle from γ_1 to γ_2 . That is, both the magnitude and the sense of the angle is preserved by ϕ . We shall now show that if f is analytic in the disc $|z - z_0| < r$ and if $f'(z_0) \neq 0$, then f is conformal at z_0 .

If γ_1 and γ_2 intersect at z_0 , then Γ_1 and Γ_2 intersect at $w_0 = f(z_0)$, and equation (2) shows that the angle from Γ_1 to Γ_2 given by




Figure 3.12

$$\arg(w_2'(s_0)) - \arg(w_1'(t_0)),$$

is equal to

$$\arg(z_2'(s_0)) - \arg(z_1'(t_0)).$$

But the latter is, of course, just the angle from γ_1 to γ_2 . Thus, we have established the following result.

If f is analytic in the disc $|z - z_0| < r$ and if $f'(z_0) \neq 0$, then f is conformal at z_0 . (3)

The conclusion expressed in (3) is false whenever $f'(z_0) = 0$, but all is not lost. If $f - f(z_0)$ has a zero of order $m \geq 2$ at z_0 , it can be shown that f increases angles between intersecting curves by a factor of m ; the details are presented in the exercises at the end of this section.

In particular, since we have shown in (2) of Section 2 that the derivative of a one-to-one analytic function is never zero, we obtain this result.

If f is both analytic and one-to-one on a domain D , then f is conformal at all points of D . (4)

Many times the word *conformal* is used in complex variables to mean a one-to-one analytic function, although technically speaking it should be used to refer to an analytic function that is just locally one-to-one.

Example 1 The function $f(z) = e^z$ has a nonzero derivative at all points of the plane, so it is a conformal mapping at all points. Note that e^z is one-to-one on any strip of the form $\{x + iy: a \leq y \leq b\}$, $0 < b - a < 2\pi$, but is not one-to-one if $b - a \geq 2\pi$. \square

Example 2 The function $h(z) = \sin z$ is conformal at all points except $z = \pi/2 + n\pi$, $n = 0, \pm 1, \dots$, which are the zeros of $h'(z) = \cos z$. \square

Example 3 The function $g(z) = 1/z$ is conformal at all points z except $z = 0$, where it is not defined. In fact, a nonconstant linear fractional transformation $T(z) = (az + b)/(cz + d)$, $ad \neq bc$, is conformal at all points z except $z = -d/c$, where it is not defined. This follows immediately from (4), since we showed in Section 3 that T is one-to-one. \square

Example 4 Show that each one-to-one analytic function f that maps the disc $\Delta = \{z : |z| < 1\}$ onto itself is a linear fractional transformation of the form

$$f(z) = \lambda \frac{a - z}{1 - \bar{a}z}, \quad |\lambda| = 1, \quad |a| < 1.$$

Solution Example 1 of Section 3 shows that any function of the form of f is a one-to-one mapping of Δ onto itself. Now let f be any one-to-one analytic function mapping Δ onto Δ . Let $f(0) = a$ and set

$$g(z) = \frac{a - f(z)}{1 - \bar{a}f(z)} = \phi(f(z)),$$

where $\phi(z) = (a - z)/(1 - \bar{a}z)$. Then g is a one-to-one analytic function mapping Δ onto Δ , and $g(0) = \phi(f(0)) = \phi(a) = 0$. Therefore, $|g(z)| \leq |z|$, by Schwarz's Lemma (Theorem 2 of Section 2). The inverse of g , g^{-1} , is also a one-to-one analytic function of Δ onto Δ , which sends 0 to 0; so, again by Schwarz's Lemma, we find that $|g^{-1}(z)| \leq |z|$. Equivalently, $|\zeta| \leq |g(\zeta)|$ for all $\zeta \in \Delta$. Hence, $|z| = |g(z)|$ for all $z \in \Delta$, and this implies, again by Schwarz's Lemma, that $g(z) = \lambda z$, where λ is a constant of absolute value 1. Therefore,

$$\lambda z = \frac{a - f(z)}{1 - \bar{a}f(z)}, \quad |\lambda| = 1,$$

and hence

$$f(z) = \lambda \frac{\beta - z}{1 - \bar{\beta}z}, \quad \beta = \bar{\lambda}a \in \Delta. \quad \square$$

Example 5 Show that the function $\varphi(z) = \frac{1}{2}(z + z^{-1})$ is a one-to-one conformal mapping of the region $1 < |z| < \infty$ onto the exterior of the closed interval $[-1, 1]$.

Solution φ is one-to-one, since the equality $\varphi(z_1) = \varphi(z_2)$ implies that

$$z_1 + \frac{1}{z_1} = z_2 + \frac{1}{z_2}$$

or

$$z_1 - z_2 = \frac{1}{z_2} - \frac{1}{z_1} = \frac{z_1 - z_2}{z_1 z_2}.$$

If $z_1 \neq z_2$, then the term $z_1 - z_2$ may be canceled from both sides of this last equation, giving

$$\frac{1}{z_1 z_2} = 1 \quad \text{or} \quad z_1 z_2 = 1.$$

This is clearly impossible, since both z_1 and z_2 have modulus greater than 1.

φ is conformal, because $\varphi'(z) = \frac{1}{2}(1 - z^{-2})$, which is never zero, since $|z| > 1$. To determine the range of φ , let w be any complex number that is not in the interval $[-1, 1]$. The equation $\varphi(z) = w$ is equivalent to the quadratic equation

$$z^2 - 2zw + 1 = 0.$$

Every quadratic has two roots (which may coincide). Note that if z_1 is one root, then $z_2 = 1/z_1$ is the other. Hence, one of the roots will satisfy $|z| > 1$ unless both lie on the circle $|z| = 1$. If that were the case, then a root would be $z_1 = e^{i\theta}$ for some real θ , and the above quadratic becomes

$$e^{2i\theta} - 2we^{i\theta} + 1 = 0.$$

After dividing by $e^{i\theta}$, this yields

$$0 = e^{i\theta} - 2w + e^{-i\theta} = 2(\cos \theta - w).$$

Since w is not in the interval $[-1, 1]$, this is obviously not possible. Hence, one (and only one) of the roots of $z^2 - 2zw + 1 = 0$ lies in the region $|z| > 1$; this z is the solution we have been seeking. \square

Level Curves

Let p be a real-valued function on a domain D . The set of z that are solutions of the equation $p(z) = c$, where c is a constant, is called a **level set** or **level curve** of the function p . Level curves are important in describing flows (and thus fields), because the path followed by a particle in the flow is just a level curve of a certain analytic function, as we shall see in Section 3.4.1. We shall now show that for an analytic function $f = u + iv$, the level curves of u are “almost always” perpendicular to the level curves of v .

Suppose that $f = u + iv$ and $f'(z_0) \neq 0$. We know that f is one-to-one in some small disc $D = \{z : |z - z_0| < r\}$; let $g = \sigma + i\tau$ be the inverse function to f on $\Omega = f(D)$, so $g(f(z)) = z$ if $|z - z_0| < r$. We know from Exercise 19, Section 1, that g is analytic. Let γ_1 consist of those points z with $|z - z_0| < r$ and $u(z) = u(z_0)$. Then

$$\begin{aligned}\gamma_1 &= \{z: u(z) = u(z_0)\} = \{z: \operatorname{Re} f(z) = u(z_0)\} \\ &= \{g(w): \operatorname{Re} w = u(z_0), w \in \Omega\},\end{aligned}$$

so γ_1 is precisely the range of the function $g(w)$ on the set $\{w \in \Omega: \operatorname{Re} w = u(z_0)\}$ and hence is a smooth arc. Likewise, the set γ_2 , consisting of those points z with $|z - z_0| < r$ and $v(z) = v(z_0)$, can be described by

$$\begin{aligned}\gamma_2 &= \{z: v(z) = v(z_0)\} = \{z: \operatorname{Im} f(z) = v(z_0)\} \\ &= \{g(w): \operatorname{Im} w = v(z_0), w \in \Omega\}.\end{aligned}$$

Hence, γ_2 is precisely the range of $g(w)$ on the set $\{w \in \Omega: \operatorname{Im} w = v(z_0)\}$, and it, too, is a smooth arc. We wish to show that γ_1 and γ_2 meet at a right angle at z_0 . But this is now immediate, since the function g is conformal and the lines $\operatorname{Re} w = u(z_0)$ and $\operatorname{Im} w = v(z_0)$ meet at a right angle at the point $u(z_0) + iv(z_0) = f(z_0)$ (Fig. 3.13). Exercise 15 gives another proof of this fact.






Figure 3.13 The level curves γ_1 , where $u(x, y)$ is constant, and γ_2 , where $v(x, y)$ is constant, meet at a right angle.

In this manner, we see that each analytic function f whose derivative does not vanish gives an orthogonal “coordinate” system by means of the curves $u(z) = \text{constant}$, $v(z) = \text{constant}$. The simple case $f(z) = z$ gives the usual coordinate system. A few others are given in the following examples. It is worth mentioning again that not only do we obtain lovely pictures, but the curves $\operatorname{Im} f = \text{constant}$ represent the paths followed by a particle in the flow given by \bar{f}' , so these curves are of great importance in applications. More on this is in Section 3.4.1 of this chapter and in Chapter 4.

Example 6 $f(z) = \operatorname{Log} z$ on the plane minus the ray $(-\infty, 0]$. Here the level curves are $\log |z| = \text{constant}$ and $\operatorname{Arg} z = \text{constant}$, which are, respectively, circles centered at the origin and rays emanating from the origin (Fig. 3.14). \square

Example 7 $f(z) = z^2 = x^2 - y^2 + i2xy$; the level curves sketched in Figure 3.15 are two families of mutually perpendicular hyperbolas. \square

Figure 3.14 The curves $\text{Arg } z = c_2$, $\text{Log } |z| = c_1$.Figure 3.15 The curves $\text{Im}(z^2) = c_1$, $\text{Re}(z^2) = c_2$.

Example 8 $f(z) = \log((z - 1)/(z + 1))$; here the level curves of the real part of f are exactly the circles $|z - 1| = \rho|z + 1|$ discussed in Section 2, Chapter 1, while the level curves of the imaginary part of f (see Figure 3.16) are the circles $\text{Arg}((z - 1)/(z + 1)) = \text{constant}$. \square

Another way to find the level curves is as follows. The function $w = (z - 1)/(z + 1)$ maps circles through 1 and -1 into straight lines through the origin and maps the circles $|(z - 1)/(z + 1)| = c$ to the circles $|w| = c$. The function $\zeta = \log w$ maps the rays $\arg z = \text{constant}$ onto the straight lines $\text{Im } \zeta = \text{constant}$ and maps the circles $|w| = c$ onto the circles $|\zeta| = \log c$. Hence, $\zeta = \log[(z - 1)/(z + 1)] = f(z)$ has as its level curves $\text{Im } f(z) = \text{constant}$ exactly the circles through 1 and -1 .




Figure 3.16 The circles $\operatorname{Arg}\left(\frac{z-1}{z+1}\right) = k_1$ and $\left|\frac{z-1}{z+1}\right| = k_2$.

Example 9 $f(z) = \log(z + \sqrt{z^2 - R^2})$, on the plane minus the segment $[-R, R]$. The function $g(z) = z + \sqrt{z^2 - R^2} = w$ has as its inverse function

$$z = \frac{1}{2} \left(w + \frac{R^2}{w} \right) = G(w).$$

A simple computation shows that $G(w)$ carries rays $\operatorname{Arg} w = c$ onto the hyperbolas

$$\left(\frac{x}{\cos c} \right)^2 - \left(\frac{y}{\sin c} \right)^2 = R^2, \quad z = x + iy$$

and the circles $|w| = s$, $s > R$, onto the ellipses

$$\left(\frac{4x^2}{s + \frac{R^2}{s}} \right)^2 + \left(\frac{4y^2}{s - \frac{R^2}{s}} \right)^2 = 1, \quad z = x + iy.$$

Hence, $g(z)$ carries these hyperbolas and ellipses onto the rays $\operatorname{Arg} w = c$ and the circles $|w| = s$, respectively. Thus, $f(z) = \log g(z)$ has these ellipses and hyperbolas as its level curves; see Figure 3.17 for the case $R = 1$, $c' = (2/\pi)c$. \square

Example 10 Fix a , $|a| < 1$, and consider

$$f(z) = \log \left(\frac{z-a}{1-\bar{a}z} \right), \quad |z| < 1.$$




Figure 3.17 The curves $\left(\frac{x}{\cos c}\right)^2 - \left(\frac{y}{\sin c}\right)^2 = 1$, $\frac{4x^2}{\left(s + \frac{1}{s}\right)^2} + \frac{4y^2}{\left(s - \frac{1}{s}\right)^2}$

The function $g(z) = (z - a)/(1 - \bar{a}z)$ has constant modulus on the family of circles determined by

$$|z - a| = \rho \left| z - \frac{1}{\bar{a}} \right|.$$

The value $\rho = |a|$ gives the circle $|z| = 1$; the argument of $g(z)$ is constant on the family of circles passing through a and $1/\bar{a}$. Thus, the level curves of $f(z)$ are exactly these two families of mutually perpendicular circles, one family lying entirely within the disc $|z| < 1$ (Fig. 3.18).

Example 11 $f(z) = \sqrt{z}$ on the plane minus the ray $(-\infty, 0]$. Here

$$u(re^{i\theta}) = \operatorname{Re} f(re^{i\theta}) = \sqrt{r} \cos\left(\frac{\theta}{2}\right)$$

$$v(re^{i\theta}) = \operatorname{Im} f(re^{i\theta}) = \sqrt{r} \sin\left(\frac{\theta}{2}\right),$$

for $0 < r < \infty$ and $-\pi < \theta < \pi$. Using the half-angle formula from trigonometry, we find that the level curves of $u(re^{i\theta})$ and $v(re^{i\theta})$ are given by the polar equations

$$r(1 + \cos \theta) = \text{constant}$$

$$r(1 - \cos \theta) = \text{constant},$$

respectively. A few of these curves are sketched in Figure 3.19. \square




Figure 3.18 The curves $|z - a| = \rho |1 - \bar{a}z|$, $\text{Arg}((z - a)/(1 - \bar{a}z)) = c$, $-\pi < c < \pi$.




Figure 3.19 The curves $r(1 + \cos \theta) = c_1$ and $r(1 - \cos \theta) = c_2$.

You might review the preceding six examples to note the following two phenomena. (1) The level curves $\operatorname{Re} f(z) = c_1$ never touch the boundary of D , whereas (2) the level curves $\operatorname{Im} f(z) = c_2$ always meet the boundary of D .

EXERCISES FOR SECTION 3.4

1. Show that $g(z) = z^2$ is one-to-one on the half-plane $\operatorname{Re} z > 0$ but not on any larger open set.
2. Find all one-to-one analytic functions that map the upper half-plane U onto itself. (**Hint:** $\phi(z) = i(1+z)/(1-z)$ maps the unit disc onto U and ϕ is one-to-one.)
3. (a) Show that each one-to-one entire function has the form $Az + B$, $A \neq 0$. (**Hint:** Check the behavior at “ ∞ .”)
 (b) Extend (a) to show that a one-to-one rational function on the plane plus “ ∞ ” must be a linear fractional transformation.
4. Let f be a one-to-one analytic function on a domain D and suppose that there is an analytic function h on D with $h^2 = f$. Show that h is also one-to-one.
5. Suppose that w is not in the interval $[-R, R]$. Show that the equation $z + R^2/z = 2w$ has one solution z with $|z| < R$ and one solution z with $|z| > R$.
6. Use the basic results on conformality to show that each of the circles $|z - a| = \rho|z - b|$, $0 < \rho < \infty$, is perpendicular to each of the circles $\arg((z - a)/(z - b)) = c$, $0 \leq c < 2\pi$. (**Hint:** Consider $f(z) = \log((z - a)/(z - b))$.)
7. Sketch some of the level curves of the real and imaginary parts of the following functions. (a) $f(z) = \log((z - 2)/(1 - 2z))$. (b) $g(z) = \sqrt{z - 1}$ on the plane with the interval $(-\infty, 1]$ deleted. (**Hint:** Set $z = 1 + w^2$ with $w = s + it$. Then solve for z when $s = \text{constant}$ or $t = \text{constant}$.) (c) $h(z) = (1 + z)/(1 - z)$. (d) $f(z) = z^2 - 2z$.
8. Let f and g be conformal analytic functions with the range of f a subset of the domain of g . Show that the composition $g(f(z))$ is also conformal.
9. Let p be a polynomial of degree 1 or more with r distinct zeros. Let $\gamma_\varepsilon = \{z : |p(z)| = \varepsilon\}$; show that γ_ε consists of r disjoint closed curves if ε is small and one closed curve if ε is very large. These curves are called **lemniscates**. For the specific polynomial $p(z) = z^2 - 1$, show that the lemniscate is a pair of disjoint simple closed curves if $0 < \varepsilon < 1$, a single simple closed curve if $\varepsilon > 1$, and a “figure eight” if $\varepsilon = 1$.
10. Let f be a one-to-one analytic mapping of the unit disc $|z| < 1$ onto itself with two fixed points in $|z| < 1$. Show that $f(z) = z$.
11. Let f be analytic on a domain D , let z_0 be a point of D , and set $w_0 = f(z_0)$. Suppose that $f(z) - w_0$ has a zero of order m , $m \geq 2$, at z_0 . Follow the outlined steps to show that f “multiplies angles by m ” at z_0 .
 - (a) The angle from the ray $\gamma_1 = \{z : \arg(z - z_0) = \psi\}$ to the ray $\gamma_2 = \{z : \arg(z - z_0) = \psi + \theta\}$ is θ .
 - (b) In a disc $|z - z_0| < r$, we have $f(z) - w_0 = (z - z_0)^m g(z)$, where $g(z_0) \neq 0$.
 - (c) If $z_1 \in \gamma_1$ and $z_2 \in \gamma_2$ then $\arg(f(z_1) - w_0) = m\psi + \arg g(z_1)$, and $\arg(f(z_2) - w_0) = m(\psi + \theta) + \arg g(z_2)$.

- (d) Let $\Gamma_1 = f(\gamma_1)$ and $\Gamma_2 = f(\gamma_2)$. The angle from Γ_1 to Γ_2 is

$$\lim_{\substack{z_1 \rightarrow z_0 \\ z_2 \rightarrow z_0}} [\arg(f(z_1) - w_0) - \arg(f(z_2) - w_0)] = m\theta.$$

12. Suppose that f is analytic on a convex domain D and that $\operatorname{Re}(f'(z)) > 0$ for all $z \in D$. Show that f is one-to-one on D . (**Hint:** $f(z_2) - f(z_1) = \int_\gamma f'(w) dw$, where γ is the line segment joining z_1 to z_2 .)
13. Use Exercise 12 to show that the function $g(z) = z + e^z$ is one-to-one on the strip $\{x + iy: -\infty < x < \infty, 0 < y < \pi\}$. (**Hint:** Show that the conclusion of Exercise 12 remains valid if $\operatorname{Im}(f')$ is positive on D .)
14. Let F be a one-to-one analytic function from a domain D_1 onto a domain D_2 . Suppose that $\{z_k\}$ is a sequence of points in D_1 with $z_k \rightarrow p$, where p is in the boundary of D_1 . Show that if $\{f(z_k)\}$ converges to a point q , then q must necessarily lie in the boundary of D_2 . (**Hint:** If $q \in D_2$, then $q = f(z_0)$ for some $z_0 \in D_1$. Show that f carries a small disc centered at z_0 into a small disc centered at q and so derive a contradiction.)
15. Let $f = u + iv$ be analytic, and suppose that $f'(z_0) \neq 0$. Show that the level curves $u = u(z_0)$ and $v = v(z_0)$ are orthogonal at z_0 by showing that the dot product of the gradient of u at z_0 with that of v at z_0 is zero. You will need to use the Cauchy–Riemann equations.

3.4.1 Conformal Mapping and Flows*

One of the most significant applications of conformal mapping is to flows. This section is devoted to the first steps in this application by making the connection between level curves, conformal mapping, and flows. This will not only provide us with an important application of complex variables but will also make it apparent why we want to search for specific conformal mappings. Thus it leads into the material of the next section.

We already noted in Section 2.1.1 and Section 2.4, Application 3, Chapter 2 that if f is a sourceless and irrotational flow on some domain D , there must be an analytic function G defined on D , called the **complex potential** of the flow, with

$$G'(z) = \overline{f(z)}, \quad z \in D.$$

We shall now show that the level curves $\operatorname{Im} G = \text{constant}$ represent the paths followed by particles in the flow; that is, these level curves are the **streamlines** for the flow.

Let z_0 be a point of D at which $f(z_0) \neq 0$. Then $G'(z_0) = \overline{f(z_0)} \neq 0$, so $G(z)$ is one-to-one in some small disc $\{z: |z - z_0| < \delta\}$, centered at z_0 . This implies that there is an analytic function $H(w)$ defined at least for $w \in \Omega = \{G(z): |z - z_0| < \delta\}$ with $H(G(z)) = z$ if $|z - z_0| < \delta$. The level curve

$$\Gamma_0 = \{z: \operatorname{Im} G(z) = \operatorname{Im} G(z_0), |z - z_0| < \delta\}$$

can also be described, therefore, as

$$\begin{aligned}\Gamma_0 &= \{H(w): \operatorname{Im} w = \operatorname{Im} G(z_0), w \in \Omega\} \\ &= \{H(\tau + ic_0): \tau + ic_0 \in \Omega, c_0 = \operatorname{Im} G(z_0)\}.\end{aligned}$$

The tangent vector to the curve Γ_0 is just the derivative of $H(\tau + ic_0)$ with respect to τ and this in turn is equal to the derivative of H with respect to w , since H is analytic. By the chain rule, then

$$1 = H'(G(z))G'(z) = H'(w)\overline{f'(z)}, \quad w = G(z).$$

Consequently,

$$\frac{H'(w)}{|H'(w)|} = \frac{|f(z)|}{\overline{f(z)}} = \frac{f(z)}{|f(z)|}.$$

Thus, the unit tangent vector to Γ_0 is parallel to $f(z)$ at each point z of Γ_0 . This implies that the level curve Γ_0 is the path followed by a particle in a flow (see also Exercise 11 at the end of this section).

Streamlining

Flow problems are typically of two types. The first, illustrated in Figures 3.20a and 3.20b, is when an impermeable object is placed in a flow of uniform velocity across the plane. The resulting flow *around* the object is to be found, assuming a (virtually) uniform velocity far from the object.

The second type of flow, illustrated in Figures 3.21a and 3.21b is, for instance, the flow through a channel or around a corner, or generally in a simply-connected region whose boundary consists of straight lines except for a few irregularities or corners. The boundary is impermeable, and the flow is assumed to be (virtually) uniform “at ∞ ”—that is, far from the irregular part of the boundary. Both types of




Figure 3.20




Figure 3.21

flow problem can be handled by consideration of appropriate conformal mappings, as we shall now show.

Our objective is to find the path described by a particle as it moves through the domain D when the motion is governed by a sourceless, irrotational flow f that remains in D ; that is, the particle does not cross the boundary of D . We let G be an analytic function on D with $G'(z) = \bar{f(z)}$, $z \in D$; we then know from the foregoing discussion that the path followed by a particle in the flow is just one of the level curves, $\operatorname{Im} G = \text{constant}$. The particles never cross the boundary of D , so the boundary of D is one of the streamlines of the flow; that is, $\operatorname{Im} G$ is constant on the boundary D . Consequently, to find the path taken by a particle in the flow, we need to find an analytic function G on D whose imaginary part is constant on the boundary of D and which has the correct behavior at “ ∞ .” This is where conformal mapping enters the picture.

Suppose that G is a one-to-one analytic mapping of the simply-connected domain D onto the upper half-plane $U = \{x + iy: y > 0\}$ and G maps the boundary of D onto the real axis. Then $\operatorname{Im} G(z)$ is zero if z is in the boundary of D . If G^{-1} has the correct behavior at ∞ , then G is the complex potential that we want. That is, the level curves $\operatorname{Im} G = \text{constant}$ are the paths followed by the particles in the flow, and the function $f(z) = \bar{G'(z)}$ is the velocity of the flow. We write $w = G(z)$ and $H(w) = z$; that is, $H(w)$ is the inverse function to $G(z)$; $H(w)$ maps the upper half-plane U onto D . The paths followed by particles in the flow are given by

$$\Gamma_c(t) = \{H(t + ic): -\infty < t < \infty\}, \quad 0 < c < \infty.$$

For example, the domain D pictured in Figure 3.21b is mapped onto U by the function $w = G(z)$, where

$$z = (w^2 - 1)^{1/2},$$

as we shall see in Example 10 of Section 5. Hence, the curves Γ_c , given by

$$\Gamma_c(t) = [(t + ic)^2 - 1]^{1/2}, \quad -\infty < t < \infty, \quad c > 0$$

are exactly the paths followed by the particles in this flow. For c very large, these are virtually the same as the horizontal lines $\operatorname{Im} z = c$. For $c = 0$, the curve is just

the boundary of the domain D . (The square root must be correctly interpreted, however.)

The search for the function G is called “streamlining” the domain D , and the function $V(x, y) = \operatorname{Im} G(x + iy)$ is called the **stream function** of D .

Example 1 The function $G(z) = z^2$ maps the first quadrant conformally onto the upper half-plane U . Its inverse is the function $H(w) = \sqrt{w}$. The stream function is $\operatorname{Im}(z^2) = 2xy$. \square

Example 2 The function $G(z) = \sin z$ maps the strip $\{x + iy : |x| < \pi/2, 0 < y < \infty\}$ conformally onto U . Its inverse is $H(w) = \arcsin w$ (see Section 5, Chapter 1). The stream function is $\cos x \cosh y = \operatorname{Im}(\sin z)$. \square

Example 3 Fix a point a , $0 \leq a < 1$. The function

$$G(z) = i \frac{(1-a)(1+z)}{(1-z)(1+a)}$$

maps the disc $\{|z| < 1\}$ conformally onto U , sending a to i . Its inverse is

$$H(w) = \frac{w(1+a) - i(1-a)}{w(1+a) + i(1-a)}.$$

The stream function is

$$\frac{(1-a)}{(1+a)} \frac{(1-|z|^2)}{|1-z|^2}.$$
 \square

EXERCISES FOR SECTION 3.4.1

1. Show that the function $G(z) = \frac{1}{2}(z^2 + 1/z^2)$ maps the region D , which is in the first quadrant and also exterior to the circle $|z| = 1$ conformally onto the upper half-plane (Fig. 3.22). Use the function $H = G^{-1}$ to streamline D .




Figure 3.22

2. Show that the function $H(z) = z^\alpha$, $0 < \alpha < 2$, maps the upper half-plane U onto the plane minus a “wedge” of angle $\pi(2 - \alpha)$ (Fig. 3.23). Use H to streamline this region.




Figure 3.23

3. Show that the function $K(z) = z/(1 - z)^2$ is a conformal mapping of the disc $\{z: |z| < 1\}$ onto the domain D obtained by deleting from the plane the interval $(-\infty, -\frac{1}{4}]$ (Fig. 3.24). Show that $z = (\zeta - i)/(\zeta + i)$ is a conformal mapping of U onto the disc $\{z: |z| < 1\}$. Use these two functions to streamline D .




Figure 3.24

4. Use the function H from Example 3 to streamline the disc $\{z: |z| < 1\}$.
 5. Show that the function $G(z) = -(1 - z^2)^2$ maps the region D shown in Figure 3.25 conformally onto U . Use this to streamline D .




Figure 3.25

6. Show that the function $\text{Log}(z - a)$ represents the complex potential of a flow in the plane with a single source at the point a . What are the streamlines for this flow?
7. Show that the function $\text{Log}((z - 1)/(z + 1))$ represents the complex potential of a flow in the plane with a source at -1 and a sink (= negative source) at $+1$. Find the streamlines for this flow.
8. Show that the function

$$\text{Log}\left(\frac{z - a}{1 - \bar{a}z}\right), \quad |a| < 1,$$

represents a complex potential of a flow *within* the disc $|z| < 1$, which has a sink at $z = a$ (see Example 10, Section 3.4).

9. Sketch some of the streamlines for the flow

$$f(z) = \frac{(\bar{p} - \bar{q})}{(\bar{z} - \bar{p})(\bar{z} - \bar{q})}.$$

(**Hint:** An indefinite integral of \bar{f} is $G(z) = \text{Log}((z - p)/(z - q))$.)

10. The function $G(z) = e^z$ is a conformal mapping of the strip $S = \{x + iy: 0 < y < \pi, -\infty < x < \infty\}$ onto U with $\text{Im } G = 0$ on the edges of S . However, the curves $\text{Im } G = \text{constant}$ are *not* the horizontal lines $y = c$, which are the lines of a uniform flow in S . Explain this apparent discrepancy.
11. Let f be a sourceless, irrotational flow in a domain D , and let G be an analytic function on D with $G' = \bar{f}$. Let z_0 be a point of D at which $f(z_0) \neq 0$, and let δ be a small positive number for which $G'(z) \neq 0$ if $|z - z_0| < \delta$. Let $\Omega = \{G(z): |z - z_0| < \delta\}$ and let $H(w)$ be the analytic function on Ω with $H(G(z)) = z, |z - z_0| < \delta$.

Let $u(t), a < t < b$, be a continuously differentiable function with values in the line $L: \{\tau + ic_0: \tau + ic_0 \in \Omega, c_0 = \text{Im } G(z_0)\}$, which satisfies the differential equation

$$u'(t) = |H'(u(t))|^{-2}, \quad a < t < b,$$

and set $\gamma(t) = H(u(t)), a < t < b$. (That such a function exists is a result from the theory of ordinary differential equations.) Show that the range of the curve $\gamma(t)$ is the level curve

$$\Gamma_0 = \{z: \text{Im } G(z) = c_0\}$$

and that $\gamma'(t) = f(\gamma(t))$ for $a < t < b$. (**Hint:** Look back to the discussion at the beginning of this section.)

3.5 The Riemann Mapping Theorem and Schwarz–Christoffel Transformations

The theorem of greatest theoretical importance in the subject of conformal mapping is the famous theorem of Bernhard Riemann.

THEOREM 1 Riemann Mapping Theorem Suppose D is a simply-connected domain with at least two points in its boundary; let p be a point of D . Then there is a one-to-one analytic function ϕ that maps D onto the open unit disc $\Delta = \{w: |w| < 1\}$ and $\phi(p) = 0$. Furthermore, ϕ is uniquely determined by the requirement that $\phi'(p)$ be positive. ■

It is a consequence of Theorem 1 that *any* two simply-connected domains (each with two or more boundary points) can be linked by a one-to-one analytic function. For if D_1 and D_2 are two such domains and if ϕ_2 maps D_2 onto Δ with $\phi_2(p_2) = 0$, then the analytic function $\psi = \phi_2^{-1} \circ \phi_1$ maps D_1 onto D_2 , $\psi(p_1) = p_2$, and ψ is one-to-one. In the language of complex variables, D_1 and D_2 are **conformally equivalent**; the mapping is uniquely determined by the requirements that $\psi(p_1) = p_2$ and $\psi'(p_1)$ be positive. See Figure 3.26.




Figure 3.26 Two simply-connected domains that are necessarily conformally equivalent.

The proof of Theorem 1 requires concepts not covered in this book; in any event, the usual proof provides no clue to how you actually write down the conformal mapping; it is an “existence” proof only (see the notes at the end of the chapter). Theorem 1 is of great utility nonetheless, since it provides us with the knowledge that a conformal map actually exists and hence might be found explicitly (there is, obviously, little point in searching for an object that does not exist). We begin our search for useful conformal mappings by reviewing some of the conformal maps that we have already run across. Then we show that a more-or-less explicit rule can be given for constructing conformal maps of the upper half-plane $U = \{w: \text{Im } w > 0\}$ onto a **polygon**. The latter construction produces the **Schwarz–Christoffel** transformations, which are useful tools in analyzing flows and other physical phenomena, as we saw in Section 3.4.1.

Examples of Conformal Mappings

Example 1 $f(z) = i(1 + z)/(1 - z)$ maps the disc $\Delta = \{z: |z| < 1\}$ onto the half-plane $U = \{w: \text{Im } w > 0\}$ (Fig. 3.27). □

Example 2 $g(z) = e^z$ maps the strip $\{-\pi \leq y < \pi\}$ onto the punctured plane, those w with $w \neq 0$ (Fig. 3.28). (See Section 5, Chapter 1.) □




Figure 3.27




Figure 3.28




Figure 3.29

Example 3 $h(z) = \text{Log } z$ maps the region Ω obtained by deleting the ray $(-\infty, 0]$ from the plane onto the strip $\{w: |\text{Im } w| < \pi\}$ (Fig. 3.29). (See Section 5, Chapter 1.) \square

Example 4 $F(z) = \sin z$ maps the strip $\{z = x + iy: 0 < x < \pi/2 \text{ and } y > 0\}$ onto the first quadrant, $\{w: \text{Re } w > 0 \text{ and } \text{Im } w > 0\}$ (Fig. 3.30). (See Section 5, Chapter 1.) \square




Figure 3.30




Figure 3.31

Example 5 $f(z) = z^p$, $0 < p < 2$, maps the upper half-plane U onto the region described by $\{w = re^{i\psi} : 0 < \psi < \pi p; 0 < r < \infty\}$ (Fig. 3.31). \square

Schwarz*-Christoffel[†] Transformations

A Schwarz–Christoffel transformation is an analytic conformal mapping of the upper half-plane onto a polygon. The key to understanding it is the examination of the behavior at the point x_0 of the function f given by

$$f(z) = A(z - x_0)^\beta + B,$$

where x_0 and β are real numbers, $0 < \beta < 2$, and A and B are complex numbers. The root is determined by choosing $\arg(z - x_0)$ to lie in the interval $(-\pi/2, 3\pi/2)$; that is, we delete from the plane the vertical ray from x_0 down.

To begin, suppose $z = x$ is real and $x > x_0$. Then $\arg f'(x) = (\beta - 1)(0) + \arg A$, so the curve parametrized by f has a tangent vector of constant slope, $\arg A$; that is, it is a straight line segment. On the other hand, if $x < x_0$, then $\arg f'(x) =$

* Hermann Amandus Schwarz, 1843–1921.

† Elwin Bruno Christoffel, 1829–1900.




Figure 3.32

$\pi(\beta - 1) + \arg A$, so $f(x)$ lies on a straight line making an angle $\pi(\beta - 1) + \arg A$ with the positive real axis (Fig. 3.32).

Let us set $\alpha = \beta - 1$, so that $-1 < \alpha < 1$; we see that the mapping f carries the real axis into a polygonal curve of only two pieces, and the pieces meet at B with angle of $\pi\alpha = \pi(\beta - 1)$. Let us continue this idea, concentrating on f' instead of f , since it is the value of $\arg(f')$ that determines the slope of the various pieces of the image curve. Let $x_1 < \dots < x_N$ be points on the real axis, and let $\alpha_1, \dots, \alpha_N$ be real numbers, all in the interval $(-1, 1)$. We shall examine the behavior of the function f , whose derivative is given by

$$f'(z) = A(z - x_1)^{\alpha_1} \dots (z - x_N)^{\alpha_N}.$$

Once again, the roots are determined by the requirement that $\arg(z - x_j)$ lie in the range $(-\pi/2, 3\pi/2)$. If x is real and near x_j , but is slightly more than x_j , then

$$\arg f'(x) = \arg A + \pi\alpha_{j+1} + \dots + \pi\alpha_N.$$

However, if x is again real and near x_j , but is slightly less than x_j , then

$$\arg f'(x) = \arg A + \pi\alpha_j + \pi\alpha_{j+1} + \dots + \pi\alpha_N.$$

If we add to this the facts that $\arg f'(x) = \arg A$ whenever $x > x_N$ and that $\arg f'(x) = \arg A + \pi\alpha_1 + \dots + \pi\alpha_N$ whenever $x < x_1$, we see that f maps the real axis onto a polygonal curve with $N + 1$ pieces (one or two of which may be infinite in length). See Figure 3.33.

Now we shall apply this knowledge to the mapping of the upper half-plane onto a given polygon. Let us suppose the polygon has $N + 1$ sides with vertices at points w_0, \dots, w_N , arranged in the usual counterclockwise order around the polygon.

Let $\theta_0, \theta_1, \dots, \theta_N$ be the exterior angles at w_0, \dots, w_N , respectively (Fig. 3.34). The angles $\theta_0, \dots, \theta_N$ lie in the range $(-\pi, \pi)$, and since the polygonal curve $\overrightarrow{w_0 w_1}, \overrightarrow{w_1 w_2}, \dots, \overrightarrow{w_N w_0}$ is a simple closed positively oriented curve, we see that




Figure 3.33 An illustration of the case where $N = 4$.




Figure 3.34 A polygon with 6 sides ($N = 5$).

$$\theta_0 + \theta_1 + \cdots + \theta_N = 2\pi.$$

Set $\alpha_j = -\theta_j/\pi$ so that

$$-1 < \alpha_j < 1 \quad \text{and} \quad \sum_{j=0}^N \alpha_j = -2.$$

These preliminary comments then prepare the way for the next theorem.

THEOREM 2 Schwarz-Christoffel Let P be a polygon in the plane with vertices w_0, \dots, w_N and corresponding exterior angles $\theta_0, \dots, \theta_N$; set $\alpha_j = -\theta_j/\pi$. Then there are real numbers x_1, \dots, x_N with $x_1 < \cdots < x_N$ and a constant A such that the function $f(z)$ whose derivative is

$$f'(z) = A(z - x_1)^{\alpha_1} \cdots (z - x_N)^{\alpha_N} \quad (1)$$

gives a one-to-one analytic mapping of the upper half-plane $\operatorname{Im} z > 0$ onto the polygon P ; f maps z_j to w_j for $j = 1, \dots, N$ and $f(\infty) = \lim_{x \rightarrow \pm\infty} f(x) = w_0$. ■

A full proof of Theorem 2 rests on Theorem 1 and on the reflection principle, which will be covered in Section 3 of Chapter 4. Here we shall content ourselves with several comments and a number of examples. First, we have arranged the mapping f so that $f(\infty) = w_0$, but this is not necessary; there will be times that we will want to exploit some obvious symmetries in the polygon and not require that $f(\infty) = w_0$. In general, we can select any three of the vertices w_j and any three points x_j on the real line or at ∞ and require that $f(x_j) = w_j$ for these three values of j . For example, if we demand that $f(-1) = w_2$, $f(0) = w_5$, and $f(2) = w_6$, then we have $x_2 = -1$, $x_5 = 0$, and $x_6 = 2$, so that $x_1, x_3, x_4, x_7, \dots$, necessarily satisfy

$$x_1 < -1 < x_3 < x_4 < 0 < 2 < x_7 < \dots.$$

Second, Theorem 2 is stated for bounded polygons, but it holds as well for unbounded polygons—which will be the most useful cases. This can be seen most easily by looking again at the behavior of the function whose derivative is given in (1). Another technique is to obtain the unbounded polygon as a limit of bounded ones. This is sometimes useful to determine the angles $\alpha_1, \dots, \alpha_N$ and will be illustrated in several of the examples.

Example 6 Find the Schwarz–Christoffel transformation of the upper half-plane U onto the equilateral triangle shown in Figure 3.35.




Figure 3.35

Solution The exterior angles of an equilateral triangle are all equal to $\frac{2}{3}\pi$, so that

$$\alpha_j = -\frac{\theta_j}{\pi} = -\frac{2}{3}, \quad j = 0, 1, 2.$$

We select $x_1 = -1$ and $x_2 = 1$. Then,

$$\begin{aligned} f'(z) &= A(z + 1)^{-2/3}(z - 1)^{-2/3} \\ &= A(z^2 - 1)^{-2/3}. \end{aligned}$$

Hence,

$$f(z) = A \int_1^z \frac{dw}{(w^2 - 1)^{2/3}} + B,$$

where we have selected 1 as the initial point for the integration. (Another choice would produce another B .) To find A and B , we note that

$$a = f(1) = B$$

and

$$i\sqrt{3}a = A \int_1^\infty \frac{dt}{(t^2 - 1)^{2/3}} + B.$$

If we denote by β the value of the integral

$$\beta = \int_1^\infty \frac{dt}{(t^2 - 1)^{2/3}},$$

then we find that

$$A = \frac{a(i\sqrt{3} - 1)}{\beta}$$

and

$$B = a.$$

□

Example 7 Find the Schwarz–Christoffel transformation of the upper half-plane U onto the region shown in Figure 3.36.




Figure 3.36

Solution The exterior angle at w_0 is θ where

$$\theta = \arctan\left(\frac{\tau_0}{\sigma_0}\right) \in (-\pi, 0),$$

and the exterior angle at w_1 is just $-\theta$. If we select $x_0 = -1$ and $x_1 = 1$, then the derivative of the mapping function is

$$f'(z) = A(z+1)^\alpha(z-1)^{-\alpha}, \quad \alpha = -\frac{\theta}{\pi} \in (0, 1).$$

The function $f(z)$ cannot usually be given explicitly. However, in the special case when $\sigma_0 = 0$, we have $\theta = -\pi/2$, so $\alpha = \frac{1}{2}$ and

$$f'(z) = A \left(\frac{z+1}{z-1} \right)^{1/2}.$$

Thus,

$$f(z) = A \{(z^2 - 1)^{1/2} + \text{Log}(z + (z^2 - 1)^{1/2})\} + B.$$

The constants A and B are found from

$$i\tau_0 = f(-1) = A\{i\pi\} + B$$

$$0 = f(1) = A\{0\} + B.$$

Hence, $B = 0$ and $A = \tau_0/\pi$. The mapping is thus

$$f(z) = \frac{\tau_0}{\pi} \{(z^2 - 1)^{1/2} + \text{Log}(z + (z^2 - 1)^{1/2})\}.$$

Some of the streamlines for the flow through the region (when $\sigma_0 = 0$ and $\tau_0 = 1$) are shown in Figure 3.37 below. \square




Figure 3.37 The flow over a vertical step. The streamlines are

$$\Gamma_c(t) = \frac{1}{\pi} \{[(t+ic)^2 - 1]^{1/2} + \text{Log}(t+ic + [(t+ic)^2 - 1]^{1/2})\}, \quad -\infty < t < \infty; c > 0.$$




Figure 3.38

Example 8 Find the Schwarz-Christoffel transformation that maps the upper half-plane onto the region in Figure 3.38.

Solution The exterior angles at w_0 , w_1 , and w_2 are, respectively,

$$\theta_0 = \theta_2 = -\arctan\left(\frac{a}{b}\right) \in \left[-\frac{\pi}{2}, 0\right)$$

and

$$\theta_1 = -2\theta_0 - \pi \in (-\pi, 0).$$

Hence, $\alpha_0 = \alpha_2 = -\theta_0/\pi \in (0, \frac{1}{2}]$ and $\alpha_1 = (2\theta_0/\pi) + 1 \in (0, 1)$. Set $\alpha_0 = \alpha_2 = \alpha$, so that $\alpha_1 = -2\alpha + 1$. With the choices $x_0 = -1$, $x_1 = 0$, and $x_2 = 1$, the derivative of the mapping function is

$$\begin{aligned} f'(z) &= A(z+1)^\alpha z^{-2\alpha+1}(z-1)^\alpha \\ &= A(z^2-1)^\alpha z^{-2\alpha+1}. \end{aligned}$$

Once again, we cannot find $f(z)$ explicitly, in general. The special case when $b = 0$ gives $\alpha = \frac{1}{2}$ and

$$f'(z) = A(z^2-1)^{1/2},$$

so

$$f(z) = A\{z\sqrt{1-z^2} + \arcsin z\} + B.$$

A and *B* are found by the requirements that

$$-ia = f(-1) = A \left\{ -\frac{\pi}{2} \right\} + B$$

$$0 = f(0) = B.$$

Thus, the mapping is

$$f(z) = i \frac{2a}{\pi} \{z\sqrt{1-z^2} + \arcsin z\}. \quad \square$$

Example 9 Find the Schwarz–Christoffel mapping of the upper half-plane *U* onto the region in Figure 3.39.

Solution We find this mapping as the limit of mappings onto the region shown in Figure 3.40. We select $x_0 = -1$, $x_1 = 0$, and $x_2 = 1$ and think of b as being very large. Thus, the exterior angles at w_0 and w_2 are almost $-\pi/2$, and the exterior angle at w_1 is almost π . Hence, the derivative of the mapping function onto the desired region will be




Figure 3.39




Figure 3.40

$$\begin{aligned}f'(z) &= A(z+1)^{1/2} z^{-1}(z-1)^{1/2} \\&= A\left(\frac{z^2-1}{z^2}\right)^{1/2}.\end{aligned}$$

Thus,

$$f(z) = A \left\{ \sqrt{z^2 - 1} + \arcsin\left(\frac{1}{z}\right) \right\} + B.$$

We find the constants A and B from the conditions

$$-a = f(-1) = A \left\{ -\frac{\pi}{2} \right\} + B$$

$$a = f(1) = A \left\{ \frac{\pi}{2} \right\} + B.$$

Consequently,

$$f(z) = \frac{2a}{\pi} \left\{ \sqrt{z^2 - 1} + \arcsin\left(\frac{1}{z}\right) \right\}.$$

Figure 3.41 shows some of the streamlines and some of the equipotentials for the




Figure 3.41 The flow over a deep hole in a streambed. The streamlines are

$$\Gamma_c(t) = \frac{2a}{\pi} \left[\sqrt{(t+ic)^2 - 1} + \arcsin\left(\frac{1}{t+ic}\right) \right], \quad -\infty < t < \infty, c > 0.$$

flow through this region. This models the flow over a deep hole in a streambed. \square

Example 10 Find the mapping of the upper half-plane U onto the region R obtained by deleting from U the segment $\{it: 0 \leq t \leq a\}$ (Fig. 3.42).

Solution We obtain the given region as a limit of the regions shown in Figure 3.43 as $\varepsilon \rightarrow 0$. The exterior angle at w_0 is almost $\pi/2$, as is the exterior angle at w_2 ; the exterior angle at w_1 is almost $-\pi$. Hence, as $\varepsilon \rightarrow 0$, we find that the derivative of the desired mapping is

$$\begin{aligned} f'(z) &= A(z+1)^{-1/2} z(z-1)^{-1/2} \\ &= A \frac{z}{(z^2 - 1)^{1/2}}, \end{aligned}$$

where, as before, we have selected $x_0 = -1$, $x_1 = 0$, and $x_2 = 1$. Hence,

$$f(z) = A(z^2 - 1)^{1/2} + B.$$

A and B are found by the conditions

$$0 = f(-1) = B, \quad ia = f(0) = Ai.$$

Consequently, the mapping is

$$f(z) = a(z^2 - 1)^{1/2}.$$




Figure 3.42




Figure 3.43




Figure 3.44

Some of the streamlines for the flow through this region are shown in Figure 3.44. The streamlines are given by $z(t) = \sqrt{(t + ic)^2 - 1}$, $-\infty < t < \infty$. This models the flow over an obstacle in a streambed. \square

Example 11 Show that the function $\phi(z) = z + e^z$ maps the strip $\{-\pi < y < \pi\}$ one-to-one and onto the entire complex plane minus the two rays

$$R_1 = \{\sigma + i\pi : \sigma \leq -1\}$$

$$R_2 = \{\sigma - i\pi : \sigma \leq -1\}.$$

Show further that ϕ maps the two lines $y = \pi$ and $y = -\pi$ onto the two rays R_1 and R_2 , respectively, covering each point twice (except the points $-1 \pm i\pi$, which are covered once) (Fig. 3.45).

Solution Since $\phi(\bar{z}) = \overline{\phi(z)}$, we will consider only the region where $0 \leq y < \pi$. The function $g(z) = e^z$ maps the strip $\{x + iy : 0 < y < \pi\}$ onto the upper half-plane U , so we shall now show by use of the Schwarz–Christoffel transformations that the function $f(z) = z + \operatorname{Log} z$ maps U onto the region Ω obtained by removing from U




Figure 3.45




Figure 3.46




Figure 3.47

the ray R_1 (Fig. 3.46). The composition $\phi(z) = f(g(z)) = z + e^z$ will then map the strip $\{x + iy : 0 < y < \pi\}$ onto Ω , as desired.

Consider the region shown in Figure 3.47. As $R \rightarrow \infty$, the region shown approaches the region Ω . We select $x_0 = -1$ and $x_1 = 0$ and note that the exterior angle at w_0 is almost $-\pi$ and the exterior angle at w_1 is almost π , if R is very big. Hence, as $R \rightarrow \infty$, we find that the derivative of the mapping function is

$$\begin{aligned}f'(z) &= A(z + 1)^1 z^{-1} \\&= A\left(1 + \frac{1}{z}\right).\end{aligned}$$

Thus, $f(z) = A(z + \text{Log } z) + B$. To determine A and B , we note that

$$\pi = \text{Im } f(x) = \text{Im}(Ax + \log|x| + i\pi + B)$$

for any $x \leq -1$. Thus,

$$0 = x \text{Im } A + \text{Im } B, \quad x \leq -1,$$

which implies that A and B are real. Furthermore,

$$-1 + i\pi = f(-1) = A(-1 + i\pi) + B,$$

which then gives $A = 1$ and $B = 0$. \square




Figure 3.48




Figure 3.49

Example 12 Find the Schwarz–Christoffel transformation of the upper half-plane U onto the region pictured in Figure 3.48.

Solution Once again, we find the desired mapping as a limit of mappings onto regions that “grow” to the desired region. We start by considering the region pictured in Figure 3.49. The exterior angle at w_0 is $\pi/2$; the exterior angles at w_1 , w_2 , and w_3 are nearly π , $-\pi/2$, and π , respectively. We let w_0 correspond to -1 , w_1 to 0 , w_2 to some point a^2 , $a > 0$, and w_3 to ∞ . The derivative of the desired mapping is then

$$f'(z) = A(z+1)^{-1/2}z^{-1}(z-a^2)^{1/2},$$

where A is a constant. Our first task is to find a^2 in terms of σ_0 and τ_0 , the coordinates of the corner at w_0 . Consider Figure 3.50.

Let z lie on the semicircle $\{\varepsilon e^{i\theta} : 0 \leq \theta \leq \pi\}$, ε small; as z traverses this semicircle clockwise, the imaginary part of $f(z)$ goes from 0 to τ_0 . Hence,

$$\tau_0 = \operatorname{Im} \int f'(z) dz = \operatorname{Im} \left\{ \int_{\pi}^0 A \left(\frac{\varepsilon e^{i\theta} - a^2}{\varepsilon e^{i\theta} + 1} \right)^{1/2} id\theta \right\}.$$

Letting $\varepsilon \rightarrow 0$, we find that




Figure 3.50

$$\tau_0 = \operatorname{Im} \{Ai(ia)(-\pi)\} = \pi a \operatorname{Im} A.$$

Likewise, if z traverses the semicircle $\{\operatorname{Re}^{it}: 0 \leq t \leq \pi\}$ counterclockwise, the real part of $f(\operatorname{Re}^{it})$ decreases from σ_0 to 0. Hence,

$$-\sigma_0 = \operatorname{Re} \int f'(z) dz = \operatorname{Re} \left\{ \int_0^\pi A \left(\frac{\operatorname{Re}^{it} - a^2}{\operatorname{Re}^{it} + 1} \right)^{1/2} idt \right\}.$$

Letting $R \rightarrow \infty$, we find that

$$-\sigma_0 = \operatorname{Re} \{iA\pi\} = -\pi \operatorname{Im} A.$$

Combining these two equations yields

$$a = \frac{\tau_0}{\sigma_0}, \quad \operatorname{Im} A = \frac{\sigma_0}{\pi}, \quad A = \frac{i\sigma_0}{\pi},$$

and so

$$f'(z) = i \frac{\sigma_0}{\pi} \frac{1}{z} \left(\frac{z - a^2}{z + 1} \right)^{1/2}, \quad a = \frac{\tau_0}{\sigma_0}.$$

Thus, the derivative of the Schwarz–Christoffel transformation is completely known, and all that remains is to integrate.

To find f , we may either consult integral tables or proceed in this way. Let

$$w = \left(\frac{z+1}{z-a^2} \right)^{1/2}$$

and define g by the relation $g(w) = f(z)$. The chain rule then gives

$$\frac{dg}{dw} = \frac{df}{dz} \frac{dz}{dw} = A \frac{1}{z} \left(\frac{z - a^2}{z + 1} \right)^{1/2} \frac{dz}{dw}, \quad A = \frac{i\sigma_0}{\pi}.$$

However,

$$z = \frac{1 + a^2 w^2}{w^2 - 1},$$

so

$$\begin{aligned} \frac{dg}{dw} &= A \frac{w^2 - 1}{1 + a^2 w^2} \frac{1}{w} \frac{2w(1 + a^2)(-1)}{(w^2 - 1)^2} \\ &= 2A(1 + a^2) \frac{1}{(1 - w^2)(1 + a^2 w^2)}. \end{aligned}$$

This last expression yields immediately to the partial-fraction technique of elementary calculus:

$$\frac{2(1 + a^2)}{(1 - w^2)(1 + a^2 w^2)} = \frac{1}{1 - w} + \frac{1}{1 + w} + \frac{a^2}{1 - iaw} + \frac{a^2}{1 + iaw}.$$

Integration then gives

$$f(z) = g(w) = A \left[\operatorname{Log} \left(\frac{1 + w}{1 - w} \right) - ia \operatorname{Log} \left(\frac{1 + iaw}{1 - iaw} \right) \right] + B,$$

where

$$w = \left(\frac{z + 1}{z - a^2} \right)^{1/2}, \quad a = \frac{\tau_0}{\sigma_0}, \quad A = \frac{i\sigma_0}{\pi}.$$

B is zero, since $g(0) = f(1) = 0$. □

EXERCISES FOR SECTION 3.5

In Exercises 1 to 6, use conformal maps or combinations of conformal maps that you already know such as linear fractional transformations, powers, roots, $\sin z$, $\log z$, etc., to find a one-to-one analytic function mapping the given region D onto the upper half-plane U . In each case, tell as much as possible about the image of the boundary of D under the mapping.

1. $D = \{z = x + iy: x \text{ and } y \text{ are positive}\}$
2. $D = \{z: |\operatorname{Arg} z| < \alpha\}, \alpha \leq \pi$
3. $D = \{z = x + iy: |y - 1| < 2\}$
4. $D = \{z: |z - z_0| < r_0\}$
5. $D = \{z: |z| > R \text{ and } \operatorname{Im} z > 0\}$
6. $D = \{z: z \notin (-\infty, -1] \cup [1, \infty)\}$
7. Find a conformal map of the infinite strip $0 < y < \pi$ onto the semi-infinite strip $\{\sigma + it: 0 < \sigma < 1, t > 0\}$.
8. Find a conformal map of the region between the two circles $|z| = 1$ and $|z - \frac{1}{2}| = \frac{1}{2}$ onto the disc $|w| < 1$. (Hint: First apply $z \rightarrow 1/(z - 1)$.)
9. Find a conformal map of the first quadrant onto the strip $|\operatorname{Im} w| < 1$.

In Exercises 10 to 14, find a Schwarz–Christoffel transformation of the upper half-plane onto the given domain D .

10. $D = \{z : 0 < \operatorname{Arg} z < 4\pi/3\}$

11. $D = \{z = x + iy : -\pi/2 < x < \pi/2 \text{ and } y > 0\}$

12. D is pictured in Figure 3.51a. (**Hint:** Map onto the region in Figure 3.51b and then let $b \rightarrow 0$.)




Figure 3.51

13. D is pictured in Figure 3.52.




Figure 3.52

14. D is pictured in Figure 3.53. (**Hint:** Obtain this region as the limit of the regions in Figure 3.54.) Evaluate the constants as in Example 12.




Figure 3.53




Figure 3.54

15. D is pictured in Figure 3.55. (**Hint:** Obtain this region as the limit of the regions shown in Figure 3.56.) Let -1 correspond to ia , 0 to $-R$; then let $R \rightarrow \infty$ to find $f'(z)$. f may be obtained as in Example 12.




Figure 3.55




Figure 3.56

16. Let C_1 and C_2 be two circles, with C_2 inside C_1 . Show that there is a linear fractional transformation ϕ such that $\phi(C_1)$ and $\phi(C_2)$ are concentric circles. (**Hint:** An initial translation, dilation, and rotation reduce the result to the case when C_1 is the circle $|z| = 1$ and C_2 is centered on the positive real axis. Show that there is an $\alpha \in (0, 1)$ such that

$$\psi(z) = \frac{z - \alpha}{1 - \alpha z}$$

carries C_2 to a circle centered at the origin. The key is that ψ will carry a diameter of C_2 to a diameter of $\psi(C_2)$.

17. Suppose that f is a one-to-one analytic function mapping the disc $|z| < 1$ onto a *bounded* domain D . Show that the **area** of D is given by

$$A(D) = \iint_{|z|<1} |f'(z)|^2 dx dy, \quad z = x + iy.$$

You will need to know the “change-of-variables” formula for double integrals.

18. Using the formula in Exercise 17, show that the area of D is given by

$$A(D) = \pi \sum_{n=1}^{\infty} n |a_n|^2,$$

where $\sum_{n=0}^{\infty} a_n z^n$ is the power series for f in $|z| < 1$.

19. Let D be a simply-connected domain and ϕ the Riemann mapping of D onto the disc $\{w: |w| < 1\}$. Let $\{z_k\}$ be a sequence of points in D with $z_k \rightarrow p$, p in the boundary of D . Show that the sequence $\{|\phi(z_k)|\}$ converges to 1. (See Exercise 14 in Section 3.4.)

Further Reading

A lovely treatment of the geometry of complex numbers, including an extensive coverage of linear fractional transformations, can be found in Schwerdtfeger, H. *Geometry of complex numbers*. New York: Dover, 1979. Linear fractional transformations are frequently called Möbius transformations, after August Ferdinand Möbius, 1790–1868.

More on conformal mapping, especially of multiply connected domains, and a development of the theory of analytic functions on such domains, is in Nehari, Z. *Conformal mapping*. New York: Dover, 1975. Another possible source is Bieberbach, L. *Conformal mapping*. New York: Chelsea, 1953. The references listed at the end of Chapter 2 also all have more to say on the topics of this chapter and are good places to seek further information. The books by Ahlfors, Burckel, Hille, and Nehari all contain a proof of the Riemann Mapping Theorem. That in Burckel is the somewhat “constructive” proof of the theorem developed by P. Koebe. To understand it, or any of the proofs of this theorem, you must go into certain aspects of function theory not touched on here.

Further information on the stability of solutions to a linear system of ordinary differential equations can be found in Sanchez, D. *Ordinary differential equations*. San Francisco: W.H. Freeman, 1968, or in Lefschetz, S. *Differential equations: geometric theory*. New York: Dover, 1977.

4

Analytic and Harmonic Functions in Applications

4.1 Harmonic Functions

A continuous complex-valued function u with continuous first and second partial derivatives on an open set D in the plane is **harmonic** on D if it satisfies **Laplace's equation**:

$$\Delta u = \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0 \quad \text{on } D.$$

Harmonic functions are critical components in the solution of numerous physical problems, such as the flow of water through an underground aquifer, steady-state temperature distribution, electrostatic field intensity, elasticity, and diffusion. Many of these physical problems are discussed in Section 2 of this chapter, and techniques for their solution are derived in Sections 3 and 4. In this section, we review the intimate connection between harmonic and analytic functions and use this connection to establish a number of fundamental properties of harmonic functions.

Suppose first that f is an analytic function on a domain D ; we write $f = u + iv$. Recall that u and v satisfy the Cauchy–Riemann equations:

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}, \quad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}.$$

Both u and v are harmonic, since

$$\begin{aligned}\Delta u &= \frac{\partial}{\partial x} \left(\frac{\partial u}{\partial x} \right) + \frac{\partial}{\partial y} \left(\frac{\partial u}{\partial y} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{\partial v}{\partial y} \right) + \frac{\partial}{\partial y} \left(-\frac{\partial v}{\partial x} \right) \\ &= \frac{\partial^2 v}{\partial x \partial y} - \frac{\partial^2 v}{\partial y \partial x} = 0,\end{aligned}$$

and likewise, $\Delta v = 0$. Thus, the real and imaginary parts of an analytic function are harmonic. Moreover, the converse is true, at least locally. If $u(z)$ is a real-valued harmonic function on a disc, $\{z : |z - z_0| < r\}$, then define $f(z)$ by

$$f(x + iy) = u(x, y) - i \int_{x_0}^x \frac{\partial u}{\partial y}(t, y) dt + i \int_{y_0}^y \frac{\partial u}{\partial x}(x_0, s) ds. \quad (1)$$

From the Fundamental Theorem of Calculus,

$$\frac{\partial f}{\partial x} = \frac{\partial u}{\partial x} - i \frac{\partial u}{\partial y}.$$

Furthermore,

$$\begin{aligned} \frac{\partial f}{\partial y} &= \frac{\partial u}{\partial y} - i \int_{x_0}^x \frac{\partial^2 u}{\partial y^2}(t, y) dt + i \frac{\partial u}{\partial x}(x_0, y) \\ &= \frac{\partial u}{\partial y} + i \int_{x_0}^x \frac{\partial^2 u}{\partial x^2}(t, y) dt + i \frac{\partial u}{\partial x}(x_0, y) \\ &= \frac{\partial u}{\partial y} + i \left\{ \frac{\partial u}{\partial x}(x, y) - \frac{\partial u}{\partial x}(x_0, y) \right\} + i \frac{\partial u}{\partial x}(x_0, y) \\ &= \frac{\partial u}{\partial y} + i \frac{\partial u}{\partial x} = i \frac{\partial f}{\partial x}. \end{aligned}$$

Hence, if we write $f = u + iv$ (that is, $v(x, y)$ is the two terms in (1) involving the integrals), then

$$\frac{\partial v}{\partial x} = -\frac{\partial u}{\partial y} \quad \text{and} \quad \frac{\partial v}{\partial y} = \frac{\partial u}{\partial x}.$$

These are the Cauchy–Riemann equations, so f is analytic on this disc and $u = \operatorname{Re} f$ there. We summarize this information here.

A real-valued function u on a domain D is harmonic on D if and only if on every disc in D , u is the real part of some analytic function. (2)

Note that a harmonic function is locally the real part of some analytic function, but it need not be the real part of an analytic function on all of D . An example of such a function is

$$u(x, y) = \frac{1}{2} \log(x^2 + y^2).$$

u is harmonic on the annulus, $0 \leq r < x^2 + y^2 < R$, but, because locally $u(x, y)$ is

the real part of $\log z$, there is no analytic function on all of the annulus whose real part equals u .

The function v that is the imaginary part of f (that is, $u + iv$ is analytic) is determined completely by u except for adding a purely real constant. For if $u + iv_1$ is also analytic, then $(u + iv) - (u + iv_1)$ is again analytic. But this difference is just $i(v - v_1)$, so $i(v - v_1)$ must be constant, since an analytic function with purely imaginary values is constant. Such a function v is called a **harmonic conjugate** of u .

Example 1 Find a harmonic conjugate of (a) $2xy$; (b) $e^x \cos y$; (c) $(e^x + e^{-x}) \cos y$.

Solution One way to proceed is to use the rule for v given in (1), but it is just as easy to use the Cauchy–Riemann equations:

$$\frac{\partial v}{\partial x} = -\frac{\partial u}{\partial y} \quad \frac{\partial v}{\partial y} = \frac{\partial u}{\partial x}.$$

For (a):

$$\frac{\partial v}{\partial x} = -\frac{\partial u}{\partial y} = -2x;$$

thus, $v(x, y) = -x^2 + p(y)$. However,

$$p'(y) = \frac{\partial v}{\partial y} = \frac{\partial u}{\partial x} = 2y,$$

so $p'(y) = 2y$. Hence, $v(x, y) = -x^2 + y^2 + c$, where c is a constant.

For (b):

$$\frac{\partial v}{\partial x} = -\frac{\partial u}{\partial y} = e^x \sin y;$$

thus, $v(x, y) = e^x \sin y + p(y)$. Since

$$e^x \cos y + p'(y) = \frac{\partial v}{\partial y} = \frac{\partial u}{\partial x} = e^x \cos y,$$

we find that $p'(y) = 0$. Hence, $v(x, y) = e^x \sin y + c$.

For (c):

$$\frac{\partial v}{\partial x} = -\frac{\partial u}{\partial y} = (e^x + e^{-x}) \sin y$$

and

$$\frac{\partial v}{\partial y} = \frac{\partial u}{\partial x} = (e^x - e^{-x}) \cos y.$$

Thus, $v(x, y) = (e^x - e^{-x}) \sin y + c$. □

Maximum Modulus

With (2) in mind, we can look back at (4) of Section 2, Chapter 3 and conclude that

If u is a real-valued nonconstant harmonic function on a domain D , then u has no local maximum and no local minimum in D . (3)

This is the **maximum principle** for harmonic functions. In fact, a variation of (3) holds for any harmonic function u whether real- or complex-valued.

If $u(x, y)$ is harmonic and nonconstant on a domain D , then $|u(x, y)|$ has no local maximum in D . (4)

The proof of (4) is left as an exercise.

A theorem from real variables asserts that a real-valued continuous function on a closed and bounded set attains both its maximum and minimum. This, and (3), give the next result.

Suppose that u is a real-valued harmonic function on a bounded domain D and that u is also continuous on the union of D and the boundary, B , of D . Then u attains both its maximum value and its minimum value over $D \cup B$ on B . In particular, if $u \equiv 0$ on B , then $u \equiv 0$ on D , as well. (5)

Mean Value

Suppose that u is a complex-valued harmonic function on a domain D . Then we can write $u = u_1 + iu_2$, where both u_1 and u_2 are real-valued and harmonic* on D . Hence, in a disc $\{z : |z - z_0| < R\}$, which lies in D , we know that u_1 and u_2 are each the real part of some analytic functions, say f_1 and f_2 . Thus, from (5) of Section 2, Chapter 3, we find that for any r , $0 \leq r < R$,

$$\begin{aligned} u(z_0) &= u_1(z_0) + iu_2(z_0) = \operatorname{Re} f_1(z_0) + i \operatorname{Re} f_2(z_0) \\ &= \operatorname{Re} \left\{ \frac{1}{2\pi} \int_0^{2\pi} f_1(z_0 + re^{it}) dt \right\} + i \operatorname{Re} \left\{ \frac{1}{2\pi} \int_0^{2\pi} f_2(z_0 + re^{it}) dt \right\} \\ &= \frac{1}{2\pi} \int_0^{2\pi} \{\operatorname{Re} f_1(z_0 + re^{it}) + i \operatorname{Re} f_2(z_0 + re^{it})\} dt \\ &= \frac{1}{2\pi} \int_0^{2\pi} u(z_0 + re^{it}) dt. \end{aligned}$$

This is the **mean-value property** of harmonic functions:

$$u(z_0) = \frac{1}{2\pi} \int_0^{2\pi} u(z_0 + re^{it}) dt. \quad (6)$$

* The fact that the real part of a harmonic function is again harmonic, but the real part of an analytic function is not analytic, points to a significant property of the Laplacian operator Δ .

In the exercises, you are asked to show how (6) can be used to derive (4). Property (6) actually characterizes harmonic functions in the sense that any continuous function that satisfies (6) must be harmonic; we shall not prove this.

Composition

Suppose that u is a harmonic function for z in some domain D . If $\phi(\zeta) = z$ is an analytic function of ζ , which maps a domain Ω into D , then

$$w(\zeta) = u(\phi(\zeta))$$

is a harmonic function of ζ , as ζ varies over Ω (Fig. 4.1). One way to show this is the following. Let ζ_0 be a point of Ω and $z_0 = \phi(\zeta_0)$. Let W be a small disc centered at z_0 in D , and let V be a small disc centered at ζ_0 with the property that $\phi(\zeta) \in W$ for all $\zeta \in V$; this is possible because ϕ is continuous at ζ_0 . We assume that u is real-valued; thus, there is an analytic function f in the disc W with $u = \operatorname{Re} f$; the function $g(\zeta) = f(\phi(\zeta))$ is then analytic in the disc V , and $\operatorname{Re} g = \operatorname{Re}(f \circ \phi) = u \circ \phi = w$, so $w(\zeta)$ is harmonic on V , being the real part of the analytic function $g(\zeta)$.




Figure 4.1

The Graph of a Harmonic Function and Strain Energy

Suppose that D is a bounded domain and $u(x, y)$ is a real-valued continuously differentiable function on D whose values on B , the boundary of D , are those of some fixed real-valued function f : $u(x, y) = f(x, y)$ if $(x, y) \in B$. The **energy integral** of u is

$$E(u) = \iint_D \left\{ \left(\frac{\partial u}{\partial x} \right)^2 + \left(\frac{\partial u}{\partial y} \right)^2 \right\} dx dy.$$

This integral has the physical interpretation as the total strain energy of a thin membrane stretched in the shape S of the surface in three-dimensional space, which is the graph of the function u :

$$S = \{(x, y, u(x, y)): x + iy \in D\}.$$

The derivation of this formula goes as follows. The strain energy in a small piece of surface is proportional to the change in the surface area from the undeformed to

the deformed state. That is,

$$dE = \text{new surface area} - \text{old surface area}$$

$$\begin{aligned} &= \sqrt{1 + \left(\frac{\partial u}{\partial x}\right)^2 + \left(\frac{\partial u}{\partial y}\right)^2} dA - dA \\ &\approx \left[1 + \frac{1}{2} \left\{ \left(\frac{\partial u}{\partial x}\right)^2 + \left(\frac{\partial u}{\partial y}\right)^2 \right\} \right] dA - dA \\ &= \left(\frac{1}{2} \right) \left\{ \left(\frac{\partial u}{\partial x}\right)^2 + \left(\frac{\partial u}{\partial y}\right)^2 \right\} dA. \end{aligned}$$

When we “sum” all the strain energy, we obtain the energy integral given above, up to a constant that multiplies the integral. (In passing from the second line to the




Figure 4.2 The graph of $x^3 - 3xy^2$.

third, we made use of the simple approximation $\sqrt{1+s} \approx 1 + \frac{1}{2}s$. The accuracy of this approximation increases as s decreases, so the formula is “better” if the deformations are small.)

We now pose the problem of minimizing $E(u)$ over all functions $u(x, y)$ with $u = f$ on B ; that is, we wish to find that surface with a prescribed edge whose strain energy is minimal. Suppose that u_0 produces this minimum:

$$E(u_0) \leq E(u), \quad \text{for all } u \text{ with } u = f \text{ on } B.$$

Let v be any continuously differentiable function on D that is zero on B . Then $u_0 + \epsilon v$ equals f on B for any number ϵ , and because u_0 has the minimum energy integral, $E(u_0) \leq E(u_0 + \epsilon v)$. Hence,

$$E(u_0) \leq E(u_0 + \epsilon v)$$

$$= E(u_0) + 2\epsilon \iint_D \left\{ \frac{\partial u_0}{\partial x} \frac{\partial v}{\partial x} + \frac{\partial u_0}{\partial y} \frac{\partial v}{\partial y} \right\} dx dy + \epsilon^2 E(v).$$

Cancel the $E(u_0)$ term from both sides, divide by ϵ , and then let $\epsilon \rightarrow 0$. If ϵ is always positive, we conclude that

$$\iint_D \left\{ \frac{\partial u_0}{\partial x} \frac{\partial v}{\partial x} + \frac{\partial u_0}{\partial y} \frac{\partial v}{\partial y} \right\} dx dy \geq 0,$$

and just the reverse inequality if ϵ is always negative. Thus, we must have

$$\iint_D \left\{ \frac{\partial u_0}{\partial x} \frac{\partial v}{\partial x} + \frac{\partial u_0}{\partial y} \frac{\partial v}{\partial y} \right\} dx dy = 0.$$

We now suppose that D , B , v , and u_0 satisfy the hypotheses of Green's Theorem, and we apply Green's Identity, from Exercise 13, Section 6, Chapter 1. Thus,

$$\begin{aligned} 0 &= \iint_D \left\{ \frac{\partial u_0}{\partial x} \frac{\partial v}{\partial x} + \frac{\partial u_0}{\partial y} \frac{\partial v}{\partial y} \right\} dx dy \\ &= \int_B v \frac{\partial u_0}{\partial n} ds - \iint_D (v)(\Delta u_0) dx dy. \end{aligned}$$

However, $v(x, y) = 0$ if $x + iy$ is in B , so we finally have

$$0 = \iint_D (v)(\Delta u_0) dx dy$$

for any (smooth) function $v(x, y)$ that is zero on B . This, of course, implies that $\Delta u_0 = 0$ on D (see Exercise 17). That is, the energy integral is minimized exactly by the function u_0 , which is the unique harmonic function on D with values f on B .

Figure 4.2 shows the surface $S = \{(x, y, x^3 - 3xy^2), -1 \leq x, y \leq 1\}$. This

then depicts a thin membrane having minimum strain energy stretched across the frame whose equation is the edge of this surface.

EXERCISES FOR SECTION 4.1

1. Decide whether each of the following functions u is harmonic; if so, find a harmonic conjugate.
 - (a) $u(x, y) = x^4 - 6x^2y^2 + y^4$;
 - (b) $u(x, y) = [\cos(2xy)] \exp(x^2 - y^2)$;
 - (c) $u(x, y) = \frac{x^2}{x^2 + y^2 + 1}$;
 - (d) $u(x, y) = \sin(x^2 - y^2) \cosh(2xy)$;
 - (e) $u(x, y) = \arctan x$.
2. Let f be analytic and never zero in a domain D ; show that $\ln |f|$ is harmonic in D .
3. Use Exercise 2 to show that $\log r$ is harmonic if $0 < r < R$.
4. Use the chain rule to derive this expression for the Laplace operator in polar coordinates:

$$\Delta u = \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = \frac{1}{r} \frac{\partial u}{\partial r} + \frac{\partial^2 u}{\partial r^2} + \frac{1}{r^2} \frac{\partial^2 u}{\partial \theta^2}.$$

5. Use the result in Exercise 4 to answer these two questions.
 - (a) When is a function of θ alone harmonic?
 - (b) When is a function of r alone harmonic?
6. Suppose that u is harmonic on a domain D and $u(x, y) = 0$ for all points $x + iy$ in some open set V in D . Show that $u = 0$ on all of D .
7. Let $\Phi = u + iv$ be a nonconstant analytic function on the domain D with range in the domain Ω . Suppose that f is a smooth function on Ω and g is defined on D by $g(z) = f(\Phi(z))$. Show by direct (and tedious) use of the chain rule that

$$\frac{\partial^2 g}{\partial x^2} + \frac{\partial^2 g}{\partial y^2} = |\Phi'(x + iy)|^2 \left\{ \frac{\partial^2 f}{\partial u^2} + \frac{\partial^2 f}{\partial v^2} \right\}.$$

Use this to conclude that if f is harmonic in Ω , then g is harmonic in D .

8. Use formula (6) to establish (4). (Hint: If $|u(z_0)|$ is greater than $|u(z)|$ for $|z_0 - z| < \delta$, then

$$|u(z_0)| = \left| \frac{1}{2\pi} \int_0^{2\pi} u(z_0 + re^{it}) dt \right|, \quad 0 < r < \delta$$

$$\leq \frac{1}{2\pi} \int_0^{2\pi} |u(z_0 + re^{it})| dt \leq \frac{1}{2\pi} \int_0^{2\pi} |u(z_0)| dt$$

$$= |u(z_0)|.$$

What is wrong with this?)

9. Show that the function

$$\frac{1 - r^2}{1 - 2r \cos \theta + r^2}$$

is harmonic for $0 \leq r < 1, 0 \leq \theta < 2\pi$.

10. Show that the second-degree polynomial in x and y

$$p(x, y) = ax^2 + bxy + cy^2 + dx + ey + f$$

(a, b, \dots, f are constants) is harmonic if and only if $a + c = 0$.

11. Show that the cubic polynomial in x and y

$$q(x, y) = Ax^3 + Bx^2y + Cxy^2 + Dy^3 + Ex^2 + Fxy + Gy^2 + Hx + Iy + J$$

is harmonic if and only if

$$3A + C = 0,$$

$$3D + B = 0,$$

$$E + G = 0.$$

12. Suppose that u is a real-valued harmonic function on a domain D and that u^2 is also harmonic on D . Conclude that u is constant. Show that the same conclusion holds if u^n is harmonic for some integer $n, n \geq 2$. Note how this differs from the situation for analytic functions.
13. Let $u(x, y)$ be harmonic and bounded for all $x + iy$ with $y > 0$. Suppose further that $u(x, y)$ is continuous for all $x + iy$ with $y \geq 0$ and that $u(x, 0) = 0$ for all x . Show that $u(x, y) = 0$ for all $x + iy$. By means of an example, show that the hypothesis that $u(x, y)$ is bounded cannot be omitted.
14. Find the harmonic conjugate v on the disc $\{z: |z| < 1\}$ of

$$u(z) = \operatorname{Arg} \left(\frac{1+z}{1-z} \right).$$

Conclude that u is bounded but v is unbounded.

15. Let $u(x, y)$ be harmonic and real-valued in the disc $|z - x_0| < \delta$, x_0 real, and suppose that $u(x, 0) = 0$, $x_0 - \delta < x < x_0 + \delta$. Show that $u(x, y)$ satisfies the relation $u(x, y) = -u(x, -y)$, $z = x + iy$ and $|z - x_0| < \delta$. (Hint: Let $f(z)$ be the analytic function on the disc $|z - x_0| < \delta$ with $f(x_0) = 0$ and $u = \operatorname{Re} f$. Show that $f(z)$ satisfies the relation $\overline{f(\bar{z})} = -f(z)$.)
16. Let u be a real-valued harmonic function that is bounded above on the whole plane; that is, $u(z) \leq M$ for all z . Show that u is constant.

Vanishing Integrals

17. Suppose that H is a continuous bounded real-valued function on a domain D such that $\iint H v \, dx \, dy = 0$ for every smooth real-valued function v on D , which is zero on B , the boundary of D . Conclude that $H = 0$ on D . (Hint: Suppose that

$H(z_0) > 0$ at some point $z_0 \in D$. Then there are (small) positive numbers ε and δ such that $H(z) \geq \delta$ if $|z - z_0| \leq \varepsilon$. (Why?) Define

$$v(z) = \begin{cases} (\varepsilon^2 - |z - z_0|^2)^2, & \text{if } |z - z_0| \leq \varepsilon \\ 0, & \text{if } |z - z_0| > \varepsilon, z \in D. \end{cases}$$

Show that for this choice of v , we have $\iint vH \, dx \, dy > 0$.)

4.2 Harmonic Functions as Solutions to Physical Problems

It is a matter of considerable significance that the theory of functions of a complex variable can be applied to a great number of “real-world” problems. That is, the solution to the physical problem can be formulated as a function that, by means of the physical laws operating within the given context, can be shown to be harmonic or analytic. We have already seen that this is the case in the investigation of an irrotational, sourceless flow of an ideal fluid, where we learned that the complex conjugate of the flow must be an analytic function; indeed, it must be the derivative of an analytic function. Once the solution of the given problem is known to be analytic or harmonic, then the arsenal of results we have collected to this point (or that we will develop later) can be applied and the solution can be analyzed or given explicitly.

This section contains an analysis of three types of physical problems: steady-state heat flow; electrostatic fields; and the flow of an ideal fluid, including Bernoulli’s Law and a derivation of the lift generated by a flow. There is also a short discussion of diffusion and elasticity.

Steady-State Temperature Distribution

We imagine a thin plate across which there is a flow of heat; we assume that the temperature at a point $z = x + iy$ in the plate depends on the location but not on time—that is, the temperature is in steady state. Let $T(x, y)$ denote the temperature at a point $z = x + iy$ in the plate; we shall show that T is a harmonic function of x and y .

We envision a (very) small rectangle in the plate with sides of length h and k and lower left corner at $z = x + iy$ (Fig. 4.3). In a unit time period, there is a flux or flow of heat in the amount of $-ck(\partial T / \partial x)(x, y)$ across the left-hand




Figure 4.3

vertical side of the rectangle. Here c is a constant that incorporates the **thermal conductivity** of the material. Likewise, the flux across the right-hand vertical side is $-ck(\partial T/\partial x)(x + h, y)$. The difference

$$-ck \left[\frac{\partial T}{\partial x}(x, y) - \frac{\partial T}{\partial x}(x + h, y) \right]$$

represents the net flux of heat out of the rectangle in the x -direction. Similarly,

$$-ch \left[\frac{\partial T}{\partial y}(x, y) - \frac{\partial T}{\partial y}(x, y + k) \right]$$

represents the net flux of heat out of the rectangle in the y -direction. Of course, both these expressions are only approximately correct, but the accuracy of the approximation increases as h and k approach zero. Since there is no net flux of heat out of the rectangle (because there are no heat sources and no heat sinks within the region), we find that

$$0 = ckh \left\{ \frac{\frac{\partial T}{\partial x}(x + h, y) - \frac{\partial T}{\partial x}(x, y)}{h} + \frac{\frac{\partial T}{\partial y}(x, y + k) - \frac{\partial T}{\partial y}(x, y)}{k} \right\},$$

and after dividing by ckh , and letting h and k both approach zero, we obtain

$$0 = \frac{\partial^2 T}{\partial x^2} + \frac{\partial^2 T}{\partial y^2},$$

which tells us that $T(x, y)$ is a harmonic function of x and y .

Electrostatics

Most electrostatic phenomena involve three independent space variables, but certain important situations are basically just two-dimensional. For instance, a very long charged wire, a group of parallel charged wires, and a very long charged cylinder each produces an electric field in the surrounding space. This electric field, it turns out, is the same in each plane that is perpendicular to the wires or cylinder (we shall derive this fact). By considering such a cross-sectional plane, we can use complex variables to understand the electric field in this plane and hence in all space.

The basic tool is **Coulomb's Law**, which asserts that two charged particles of charges q and Q , respectively, exert a force on one another whose magnitude is

$$\frac{CqQ}{r^2},$$

where r is the distance between the particles; the force is directed along the line

through the two points and is attractive if q and Q have opposite signs and repulsive if q and Q have the same sign. C is a constant depending on the choice of units; we shall take $C = 1$.

Imagine now a very long, straight wire carrying a uniform charge of λ coulombs per unit length. We may suppose that the wire is perpendicular to the usual xy -plane (the complex plane) and cuts through the plane at the point z_1 . Let r be the distance from z to z_1 : $r = |z - z_1|$, and let h be the coordinate along the wire (Fig. 4.4).




Figure 4.4

The electrostatic force at z , due to the wire, has two components—one is in the xy -plane and is directed toward the wire and the other is perpendicular to this plane. The first we shall temporarily term the planar force and the second the vertical force. We envision the wire as being made up of many small pieces of equal length dh , each therefore carrying a total charge of λdh . The magnitude of the planar and vertical forces at z due to this short segment are, respectively,

$$dm_p = q \cos \theta \frac{\lambda dh}{s^2}$$

and

$$dm_v = q \sin \theta \frac{\lambda dh}{s^2},$$

by Coulomb's Law, where q is the charge at z and s is the distance from z to the small segment on the wire. Simple trigonometry produces the relations

$$h = r \tan \theta$$

$$s = r \sec \theta.$$

Hence, $dh = r \sec^2 \theta d\theta$, so

$$dm_p = q \cos \theta \frac{\lambda r \sec^2 \theta d\theta}{r^2 \sec^2 \theta} = q\lambda \cos \theta \frac{d\theta}{r}$$

and

$$dm_v = q\lambda \sin \theta \frac{d\theta}{r}.$$

If the wire is very long, then θ varies from $-\pi/2$ to $\pi/2$, so the total magnitude of the planar force at the point z is

$$m_p = \int_{-\pi/2}^{\pi/2} dm_p = q\lambda \frac{1}{r} \int_{-\pi/2}^{\pi/2} \cos \theta d\theta = \frac{2q\lambda}{r}.$$

The vertical force is

$$m_v = \int_{-\pi/2}^{\pi/2} dm_v = \frac{q\lambda}{r} \int_{-\pi/2}^{\pi/2} \sin \theta d\theta = 0.$$

That is, the charged wire produces an electric field that is directed perpendicular to the axis of the wire and whose strength is directly proportional to the product of the charge at z and the charge per unit length on the wire and inversely proportional to the distance between the point and the wire. Note how this differs from the situation when just point charges are involved—there, the force depends (inversely) on the square of the distance between the charges.

The electric force thus has magnitude

$$\frac{2\lambda q}{|z - z_1|}$$

and is directed along the line joining z to z_1 . The force is repulsive if q and λ have the same sign and attractive if q and λ have opposite signs. Hence, the force is

$$F(z) = 2\lambda q \frac{z_1 - z}{|z_1 - z|^2} = \frac{2\lambda q}{\bar{z}_1 - \bar{z}}.$$

The real-valued function

$$\phi(x, y) = -q\lambda \log[(x - x_1)^2 + (y - y_1)^2]$$

has the property that its gradient is F ; that is,

$$\frac{\partial \phi}{\partial x} + i \frac{\partial \phi}{\partial y} = F.$$

Furthermore, a simple computation shows that $\phi(x, y)$ is a harmonic function of x and y , for

$$\frac{\partial^2 \phi}{\partial x^2} = q\lambda \frac{(x - x_1)^2 - (y - y_1)^2}{[(x - x_1)^2 + (y - y_1)^2]^2}$$

and

$$\frac{\partial^2 \phi}{\partial y^2} = q\lambda \frac{(y - y_1)^2 - (x - x_1)^2}{[(x - x_1)^2 + (y - y_1)^2]^2}.$$

Adding, we find that

$$\frac{\partial^2 \phi}{\partial x^2} + \frac{\partial^2 \phi}{\partial y^2} = 0.$$

The harmonic function ϕ , which carries all the information about the electric field, is called the **electrostatic potential** of the field F .

After understanding the situation of one wire, we can easily pass on to other, more complicated, configurations. First, suppose there are many long wires, all perpendicular to the xy -plane. Let z_1, \dots, z_n be the points at which the wires meet the xy -plane, and let $\lambda_1, \dots, \lambda_n$ be their charge densities, which we assume to be uniform all along the length of each. By superposition of forces, we find the force at z to be

$$F(z) = 2q \sum_{j=1}^n \lambda_j \frac{z_j - z}{|z_j - z|^2} = 2q \sum_{j=1}^n \frac{\lambda_j}{\bar{z}_j - \bar{z}}.$$

Note that

$$\bar{F}(z) = 2q \sum_{j=1}^n \frac{\lambda_j}{z_j - z}$$

is a rational function of z of degree n and has therefore exactly n zeros. Since one zero is at infinity, $\bar{F}(z)$ has at most $n - 1$ zeros in the plane. Each of these zeros is an **equilibrium point** of the field, since there the force vanishes. In the case that $\lambda_1, \dots, \lambda_n$ are all positive integers, we also have

$$\bar{F}(z) = -\frac{2qP'(z)}{P(z)},$$

where $P(z)$ is the polynomial

$$P(z) = (z - z_1)^{\lambda_1} \cdots (z - z_n)^{\lambda_n}.$$

Thus, here we can apply the information studied in Section 1 of Chapter 3 about the location of the zeros of an analytic function. Also, there is considerable information on the zeros of a polynomial; see Appendix 1.

The potential function for this field is


$$\phi(x, y) = -q \sum_{j=1}^n \lambda_j \log[(x - x_j)^2 + (y - y_j)^2];$$

$\phi(x, y)$ is the sum of harmonic functions and so is harmonic. Furthermore, as before, the gradient of ϕ is F .


Next, imagine a very long charged cylinder with its axis parallel to the vertical axis (see Fig. 4.5). We assume the charge is uniform along any line in the cylinder parallel to the axis of the cylinder but not necessarily uniformly distributed in the cross-sectional cuts. Each cross section is identical and is nothing but a curve γ ; γ is not necessarily a closed curve. Suppose that $z(t)$ is a parametrization of γ , $a \leq t \leq b$. Let $u(z)$ be the strength of the charge at the point z on γ ; that is, the line through z and parallel to the axis of the cylinder carries a uniform charge of $u(z)$ coulombs per unit length. To find the electric field produced by the charged curve γ (or rather by the cylinder), we imagine γ broken up into many small segments (see Fig. 4.6). Let ds be the arc length along γ :

$$\begin{aligned} ds &= \sqrt{x'(t)^2 + y'(t)^2} dt \\ &= |z'(t)| dt. \end{aligned}$$

The total charge due to the segment of length ds is (about) $d\lambda = u(z) ds$, so the field it produces is



A section of a “closed” cylinder



An “open” cylinder

Figure 4.5




Figure 4.6

$$\begin{aligned} dF &= \frac{2q}{\bar{z}(t) - \bar{z}} d\lambda \\ &= \frac{2q}{\bar{z}(t) - \bar{z}} u(z(t)) |z'(t)| dt. \end{aligned}$$

Thus, the electric field is just

$$F(z) = -2q \int_a^b \frac{u(z(t))}{\bar{z} - \bar{z}(t)} |z'(t)| dt.$$

The electrostatic potential of this field is

$$\phi(x, y) = -2q \int_a^b \log\{|z - z(t)|\} u(z(t)) |z'(t)| dt.$$

Equivalently,

$$\phi(z) = -2q \int_{\gamma} \{\log |z - w|\} u(w) d|w|,$$

where $d|w|$ is short for $|z'(t)| dt$; $u(w)d|w|$ is the **charge density** along the curve γ . If we write $d\rho(w)$ for the charge density, then (except for the factor of $2q$).

$$\phi(z) = \int_{\gamma} \log \frac{1}{|z - w|} d\rho(w).$$

This is an example of a **potential function**; the study of such functions and the potentials they produce is called **potential theory**.

Flow of an Ideal Fluid

The analysis of the flow of an ideal fluid provides a setting in which we can apply many of the theorems and formulas derived earlier, especially results from conformal mapping.

We begin by envisioning an incompressible, frictionless fluid flowing through a region in space. If all cross sections perpendicular to some axis are the same, the flow can be described by just two space variables, which, as usual, we take to be the regular xy -coordinates. Situations such as this arise, for example, in the flow of a fluid through a wide channel with identical cross sections (Fig. 4.7a) or the flow of a fluid past a long object, each cross section of which is the same (Fig. 4.7b). The latter model is applicable, for instance, in analyzing the flow past the wing of an aircraft in order to determine the lift (though air is compressible, the model is valid at relatively low speeds).




Figure 4.7

In Sections 1.1 and 4 of Chapter 2 and 4.1 of Chapter 3, we have already discussed some aspects of the flow of an ideal fluid. To summarize, we saw that if the velocity of the flow was given by $f = u + iv$, the twin assumptions that the flow is irrotational and sourceless–sinkless imply that $\bar{f} = u - iv$ is analytic on the domain and, even more, \bar{f} is the derivative of another analytic function G on the domain. We write $G = \phi + i\psi$; ϕ is called the **potential function** of the flow, and ψ is the **stream function** of the flow. The terminology results from the fact that the path followed by a particle within the flow is just $\psi(x, y) = \text{constant}$. Furthermore, ϕ is the potential of the flow in just the same sense that this word was used earlier when describing the electrostatic potential: the gradient of ϕ is just $(u, -v)$. Hence, the level lines $\phi(x, y) = \text{constant}$ are the curves where the flow has the same potential. Furthermore, since $G' = u - iv$, we see that whenever $f \neq 0$, it follows that the level curves of ϕ and ψ are perpendicular. Finally, both ϕ and ψ are harmonic functions, so we can apply to them any results that we know for harmonic functions in general.

We begin our discussion of flows by examining flows in several regions, making use of conformal mapping as needed. We then elaborate on sources and sinks for flows and illustrate how the ideas are applied. Next, we derive Bernoulli's

Law, which relates the pressure and the speed of a flow. Finally, we examine the circulation of a flow and show how a circulation around an object can produce a force on it.

The simplest flow is the uniform flow $f(z) = A$, A real and not equal to zero, in the upper half-plane $U = \{z: \operatorname{Im} z > 0\}$. This flow is sourceless and irrotational in U ; a complex potential is $G(z) = Az$. The streamlines are the curves $\operatorname{Im} G = c$. In this case, these are the horizontal lines $Ay = c$ or $y = c/A$. If $A > 0$, the flow is from left to right; the reverse holds if $A < 0$. The function $f(z) = A$ also gives a flow in any horizontal strip $S = \{x + iy: a < y < b\}$. The flux of this flow in S is $A(b - a)$; this is the total quantity crossing any arc connecting the line $y = a$ to the line $y = b$ in unit time. We show here how this simple flow can be combined with conformal mapping to find flows in other regions that are more complicated in shape.




Figure 4.8

Example 1 Find the flow out of the end of a very long channel.


Solution We take the channel to have sides $\{z = x + iy: y = \pm\pi\}$ and $-\infty < x < -1\}$. We know from Example 11, Section 5, Chapter 3 that the function $\phi(w) = w + e^w$ is a conformal mapping of the strip $\{w = \sigma + it: -\pi < t < \pi, -\infty < \sigma < \infty\}$ onto the complex plane with the two rays $\{x \pm i\pi: -\infty < x < -1\}$ deleted. (See Fig. 4.8.) The uniform flow $(Q/2\pi)w$ of strength Q in the strip $\{\pi > |\tau|\}$ is carried by $\phi^{-1}(z) = w$ to the desired flow out of the channel. Thus, the curves $\operatorname{Im} w = \text{constant}$, which are the streamlines in the strip, are carried to the streamlines of the flow out of the channel; the latter, then, are

$$\Gamma_c = \{\sigma + ic + e^\sigma e^{ic}: -\infty < \sigma < \infty\}, \quad c = \text{constant}, \quad -\pi < c < \pi. \quad \square$$

Some of these are sketched in Figure 4.9.

Example 2 Find the flow through a horizontal aperture of width L .

Solution The aperture may be taken to be the segment $(-\sigma, \sigma)$, $2\sigma = L$, on the real axis, and the domain D to be the complex plane with the two rays $(-\infty, -\sigma]$



$$g(z) = z + e^z$$

Figure 4.9 The flow out the end of a long channel. The streamlines are

$$\Gamma_c(x) = x + ic + e^x, \quad -\infty < x < \infty; c \in (-\pi, \pi).$$




Figure 4.10

and $[\sigma, \infty)$ deleted (see Fig. 4.10a). The function $z = \Phi(\zeta) = \sigma \sin \zeta$ maps the vertical strip $-\pi/2 < \operatorname{Re} \zeta < \pi/2$ onto D . The uniform flow $g(\zeta) = iQ\zeta/\pi$ in the strip has total flux Q across any arc joining the right-hand side to the left-hand side of the strip. Hence,

$$f(z) = g(\Phi^{-1}(z)) = \frac{iQ}{\pi} \arcsin \frac{z}{\sigma}$$

gives the desired flow in the domain D . The streamlines of g are the vertical lines $\operatorname{Re} \zeta = c$, $-\pi/2 < c < \pi/2$. Hence, their images under Φ are the streamlines of f . These are the hyperbolas

$$\frac{x^2}{(\sin c)^2} - \frac{y^2}{(\cos c)^2} = \sigma^2;$$

see Section 5, Chapter 1. Some of these curves are shown in Figure 4.11. \square




Figure 4.11 The flow from the lower half-plane to the upper half-plane through a horizontal aperture.

Point Sources and Sinks for Flows

Suppose that z_0 is a point in a domain D . A flow f in $D \setminus \{z_0\}$ has a source of strength Q at z_0 if

$$\operatorname{Im} \left\{ \int_{\gamma} \overline{f(z)} dz \right\} = Q,$$

for every piecewise smooth, positively oriented simple closed curve γ in D that contains z_0 in its interior. If $Q < 0$, it is conventional to refer to z_0 as a sink of strength $-Q$.

The function $f(z) = q/\bar{z}$, $q > 0$, produces a flow with a source of strength $2\pi q$ at the origin, since

$$\begin{aligned} \operatorname{Im} \left\{ \int_{\gamma} \overline{f(z)} dz \right\} &= \operatorname{Im} \left\{ \int_{\gamma} \frac{q}{z} dz \right\} \\ &= \operatorname{Im} \{2\pi i q\} \\ &= 2\pi q, \end{aligned}$$

for any piecewise smooth positively oriented simple closed curve γ that contains the origin in its interior; see Example 13, Section 6, Chapter 1. Similarly, the function $q/(\bar{z} - \bar{z}_0)$ produces a flow with a source of strength $2\pi q$ at the point z_0 .

A flow in the upper half-plane U has a source of strength Q at a point x_0 in the real axis if

$$\operatorname{Im} \left\{ \int_{\Gamma} \overline{f(z)} dz \right\} = Q,$$

for every simple piecewise smooth arc Γ in the closed upper half-plane $\{x + iy : y \geq 0\}$ that connects a point on the real axis to the right of x_0 to a point on the real axis to the left of x_0 (see Fig. 4.12a). As above, if $Q < 0$, then x_0 is called a sink of strength $-Q$.




Figure 4.12

Example 3 Show that $f(z) = q/\bar{z}$, $q > 0$, gives a flow in the upper half-plane U with a source at the origin of strength πq .

Solution A complex potential for f is $G(z) = q \operatorname{Log} z$. The imaginary part of G is $q \operatorname{Arg} z$; the curves $q \operatorname{Arg} z = c$ are the rays emanating from the origin, making an angle c/q with the positive real axis. These all lie in U provided $0 < c/q < \pi$. The cases $c = 0$ and $c = \pi$ give the positive and negative half-lines, respectively. Thus, f is indeed a (sourceless, irrotational) flow in U .

To see that f has a source at $z_0 = 0$, let Γ be any arc as in Figure 4.12a. Let γ be the piecewise smooth simple closed curve shown in Figure 4.12b, where ε is so small that Γ does not meet the circle $|z| = \varepsilon$. Since \bar{f} is analytic on and inside γ , the integral of \bar{f} over γ is zero, by Cauchy's Theorem. It follows that

$$\begin{aligned} 0 &= \operatorname{Im} \left\{ \int_{\gamma} \overline{f(z)} dz \right\} \\ &= \operatorname{Im} \left\{ \int_{\Gamma} \overline{f(z)} dz + q \int_S \frac{dz}{z} + q \int_L \frac{dz}{z} \right\}, \end{aligned}$$

where S is the semicircle $\{ee^{i\theta} : \pi \geq \theta \geq 0\}$ and L stands for the two line segments in the real axis that form a part of γ . On these line segments, $z = x$, so the integrals are real-valued. Consequently, their imaginary parts are zero, and we find that

$$\begin{aligned} \operatorname{Im} \left\{ \int_{\Gamma} \overline{f(z)} dz \right\} &= -\operatorname{Im} \left\{ q \int_{\pi}^0 \left(\frac{ie^{i\theta}}{\varepsilon e^{i\theta}} \right) d\theta \right\} \\ &= q\pi. \end{aligned}$$
□

Example 4 It is evident from the results of Example 3 that

$$f(z) = \frac{q_1}{z - x_1} + \frac{q_2}{z - x_2}$$

gives a flow in the upper half-plane with sources at x_1 and x_2 of strengths q_1 and q_2 , respectively. (If either or both of q_1, q_2 are negative, then of course the corresponding point x_1, x_2 is a sink.) This idea extends to any number of sources on the real axis:

$$f(z) = \sum_{k=1}^N \frac{q_k}{z - x_k}$$

gives a flow in the upper half-plane with sources at x_1, \dots, x_N of respective strengths q_1, \dots, q_N . □

Example 5 A source of strength Q is located on one of the long sides of a semi-infinite strip. Find the resulting flow.

Solution We take the strip to be the domain $D = \{x + iy : -\sigma < x < \sigma, y > 0\}$ and locate the source at the point $\sigma + i\tau$, $\tau \geq 0$ (Fig. 4.13a). The function $\psi(z) = \sin(\pi z/2\sigma)$ maps the strip D onto the upper half-plane, $\operatorname{Im} w > 0$, with $\psi(\sigma) = 1$ and $\psi(-\sigma) = -1$ (see Section 5, Chapter 1). The point $\sigma + i\tau$ is carried by ψ to the point

$$A = \frac{1}{2} \left(\exp \left\{ \frac{\pi\tau}{2\sigma} \right\} + \exp \left\{ -\frac{\pi\tau}{2\sigma} \right\} \right) = \cosh \left(\frac{\pi\tau}{2\sigma} \right).$$




Figure 4.13

The function

$$g(w) = \left(\frac{Q}{\pi}\right) \operatorname{Log}(w - A)$$

is the complex potential of a flow in the upper half-plane U with a source at A of strength Q ; (see Example 3). Hence,

$$G(z) = \left(\frac{Q}{\pi}\right) \operatorname{Log}\left(\sin\left(\frac{\pi z}{2\sigma}\right) - A\right)$$

is the complex potential for the desired flow in the semi-infinite strip. The velocity of this flow is

$$\overline{G'(z)} = \frac{Q}{2\sigma} \frac{\cos\left(\frac{\pi\bar{z}}{2\sigma}\right)}{\sin\left(\frac{\pi\bar{z}}{2\sigma}\right) - A}.$$

Write $z = x + iy$, with $-\pi/2 < x < \pi/2$ and $y > 0$; this leads to

$$\overline{G'\left(\frac{2\sigma z}{\pi}\right)} = \left(\frac{Qi}{2\sigma}\right) \frac{e^{ix} + e^{-ix}e^{-2y}}{e^{ix} - e^{-ix}e^{-2y} - 2iAe^{-y}}.$$

This quantity approaches

$$v_\infty = \frac{iQ}{2\sigma}$$

as y approaches ∞ . Hence, the velocity of this flow is virtually uniform far from the source. Furthermore, since the width of the strip is 2σ , the total flux out of the “end” of the strip (that is, at ∞) is exactly Q , the amount entering at $\sigma + i\tau$. Thus, the amount flowing into the strip per unit time equals the amount flowing out, as it must. Some streamlines of this flow are shown in Figure 4.14. \square

Example 6 Find the flow in a long narrow channel if there is a source of strength Q at a point on one side.

Solution This problem is actually a special case of Example 5, despite the different geometric configuration. We take the channel to be the strip $\{x + iy: -\sigma < x < \sigma, -\infty < y < \infty\}$ and locate the source at the point $x_0 = \sigma, y_0 = 0$ (see Fig. 4.15). The flow has obvious up-down symmetry about the x -axis and hence may be considered as a flow in the semi-infinite strip $\{x + iy: -\sigma < x < \sigma, 0 < y < \infty\}$ resulting from a source at $x_0 = \sigma, y_0 = 0$ of strength $Q/2$. However, this is a special case of Example 5. The complex potential of this flow is




Figure 4.14 The flow in a semi-infinite channel resulting from a source on one of the long sides.




Figure 4.15 The flow in a long channel resulting from a source on one side.

$$G(z) = \left(\frac{Q}{2\pi}\right) \text{Log} \left(\sin \left(\frac{\pi z}{2\sigma} \right) - 1 \right).$$

(The constant A from Example 5 equals 1, since $\tau = 0$.) Our problem is therefore solved. The stagnation points of this flow occur where its velocity is zero. These are found from

$$0 = G'(z) = \left(\frac{Q}{4\sigma}\right) \frac{\cos\left(\frac{\pi z}{2\sigma}\right)}{\sin\left(\frac{\pi z}{2\sigma}\right) - 1}.$$

Such points occur, then, only when $\cos(\pi z/2\sigma) = 0$ and $\sin(\pi z/2\sigma) \neq 1$. Thus,

$$\frac{\pi z}{2\sigma} = \left(-\frac{\pi}{2}\right) + (\text{any multiple of } 2\pi).$$

The only such point in the channel is $\pi z/2\sigma = -\pi/2$ or $z = -\sigma$. Several of the streamlines of this flow are sketched in Figure 4.15. \square

Digression At this point, it is worth spending a minute to elaborate on the use of the words *source* and *sink* in connection with the circulation produced by a linear fractional transformation (see Section 3.3). Specifically, the linear fractional transformation

$$T(z) = \frac{z+2}{1+2z}$$

sets up a circulation from -1 to 1 with a point z moving progressively closer to 1 along the circle centered on the imaginary axis and passing through -1 , 1 , and z as T is iterated. That is, the streamlines are the circles centered on the imaginary axis and passing through both -1 and 1 . Recall now from Section 2 of Chapter 1 that the family of curves $|z-1| = \rho|z+1|$, $0 < \rho < \infty$, are all circles centered on the real axis. In the notation of that section, this is the family C_1 . The orthogonal family C_2 is the collection of circles we are interested in. However, C_1 is just the family of circles given by

$$\operatorname{Re} \left\{ \operatorname{Log} \left(\frac{z+1}{z-1} \right) \right\} = \log \left| \frac{z+1}{z-1} \right| = -\log \rho = \text{constant}.$$

Consequently, by Example 7 of Section 3.4, the orthogonal family C_2 is just the family of curves described by

$$c = \operatorname{Im} \left\{ \operatorname{Log} \left(\frac{z+1}{z-1} \right) \right\} = \operatorname{Arg} \left(\frac{z+1}{z-1} \right).$$

The complex potential for the flow is, therefore,

$$G(z) = \operatorname{Log} \left(\frac{z+1}{z-1} \right) = \operatorname{Log}(z+1) - \operatorname{Log}(z-1).$$

This flow has a source at -1 and a sink at $+1$, just as we already knew.




Figure 4.16

Bernoulli's* Law

Suppose that $f = u + iv$ represents the velocity of a locally sourceless and locally irrotational flow in a domain D ; we shall use the knowledge that $\bar{f} = u - iv$ is analytic on D to derive a fundamental relation between the pressure and the speed of the flow. We begin by imagining a small rectangle in D , as in Figure 4.16. Let $p(x, y)$ be the pressure at the point (x, y) . The net force acting on this rectangle in the x -direction is (almost)

$$\{p(x, y) - p(x + h, y)\}k.$$

However, by Newton's Law, force is mass times acceleration. The mass of the little rectangle is its (constant) density ρ , multiplied by its area hk . Furthermore, the acceleration in the x -direction is the time derivative of u , since u is the x -component of the velocity. This yields

$$\begin{aligned} (\text{mass})(\text{acceleration}) &= \rho hk \frac{du}{dt} \\ &= \rho hk \left\{ \frac{\partial u}{\partial x} \frac{dx}{dt} + \frac{\partial u}{\partial y} \frac{dy}{dt} \right\}. \end{aligned}$$

However, $dx/dt = u$ and $dy/dt = v$, which then gives

$$\{p(x, y) - p(x + h, y)\}k = \rho hk \left\{ \frac{\partial u}{\partial x} u + \frac{\partial u}{\partial y} v \right\}.$$

Divide both sides by hk , then let h and k tend to 0. We obtain

$$-\frac{\partial p}{\partial x} = \rho \left\{ \frac{\partial u}{\partial x} u + \frac{\partial u}{\partial y} v \right\}.$$

In an entirely similar manner, working with the y -component of the forces acting on the rectangle, we obtain

$$-\frac{\partial p}{\partial y} = \rho \left\{ \frac{\partial v}{\partial x} u + \frac{\partial v}{\partial y} v \right\}.$$

* Daniel Bernoulli, 1700–1782.

However, $\partial u / \partial x = -\partial v / \partial y$ and $\partial u / \partial y = \partial v / \partial x$, by the Cauchy–Riemann equations, since $u - iv$ is analytic. This gives us two equations:

$$\rho \left(\frac{\partial u}{\partial x} u + \frac{\partial v}{\partial x} v \right) + \frac{\partial p}{\partial x} = 0$$

and

$$\rho \left(\frac{\partial u}{\partial y} u + \frac{\partial v}{\partial y} v \right) + \frac{\partial p}{\partial y} = 0.$$

Integration of the first with respect to x and the second with respect to y yields

$$\frac{1}{2} \rho \{u^2 + v^2\} + p = A(y)$$

and

$$\frac{1}{2} \rho \{u^2 + v^2\} + p = B(x),$$

where $A(y)$ and $B(x)$ are functions of y alone and of x alone, respectively. Since the left-hand sides are identical, it is clear that $A(y)$ must also equal $B(x)$, so both are (the same) constant. Thus, the equations are

$$\frac{1}{2} (u^2 + v^2) + \frac{1}{\rho} p = \text{constant}.$$

This says that for an ideal fluid, half the square of the speed plus the pressure divided by the (constant) density adds up to a constant; this is **Bernoulli's Law**.

Circulation and Lift

We now investigate the flow around an object—for instance, the situation pictured in Figure 4.17. Let Γ be the boundary of the object, which we take to be a simple




Figure 4.17 The flow around a solid object with boundary Γ .

closed piecewise smooth curve. We assume that the flow f is both locally irrotational and locally sourceless in the domain exterior to Γ ; that is, we assume that $\bar{f} = u - iv$ is analytic in the domain exterior to Γ . Let γ be any smooth simple closed positively oriented curve that contains Γ in its inside. The value of the integral

$$c = \int_{\gamma} f \cdot \tau \, ds$$

is called the **circulation** of the flow around Γ ; here $f \cdot \tau$ is the component of f tangent to the curve γ , and ds is arc length. As we showed in Section 1.1, Chapter 2,

$$c = \operatorname{Re} \left\{ \int_{\gamma} \bar{f}(z) \, dz \right\},$$

so it follows from the fact that \bar{f} is analytic that the integral itself is not dependent on the choice of curve γ . Hence, the value of the circulation depends on the flow f only. Furthermore, because the flow is around Γ (that is, Γ is a streamline), we know that the component of f that is normal to Γ must be zero. (For if $f \cdot n$ is, say, positive at a point p of Γ , then there is a net flow over Γ in a small segment of Γ near p , contradicting the fact that Γ is a streamline of the flow.) Hence,

$$0 = \int_{\Gamma} f \cdot n \, ds = \operatorname{Im} \left\{ \int_{\Gamma} \bar{f}(z) \, dz \right\}.$$

Once again, the analyticity of \bar{f} implies that

$$0 = \operatorname{Im} \left\{ \int_{\gamma} \bar{f}(z) \, dz \right\}$$

for any piecewise smooth closed curve γ surrounding Γ .

We now calculate the force exerted on Γ by the flow f . For this purpose, we make the reasonable assumption that the flow is “uniform at ∞ ”; that is, the limit

$$\lim_{|z| \rightarrow \infty} \bar{f}(z) = a$$

exists. Since \bar{f} is analytic in the region exterior to Γ and bounded at ∞ , ∞ is a removable singularity for \bar{f} . Hence, for $|z|$ large, \bar{f} has the expansion

$$\bar{f}(z) = a + \frac{c}{2\pi i} \frac{1}{z} + \sum_{k=2}^{\infty} \frac{b_k}{z^k}, \quad (1)$$

where c is the circulation and b_2, b_3, \dots are constants.

To begin the analysis of the force exerted on Γ by the flow, we let $p(x, y)$ be the pressure at the point (x, y) in Γ (Fig. 4.18). By a simple resolution of forces argument, we find that the horizontal component of the force on a little segment in Γ is $-p \, dy$, so the total *horizontal* component of the force is




Figure 4.18

$$H = - \int_{\Gamma} p \, dy.$$

Likewise, the total *vertical* component of the force on Γ is

$$V = \int_{\Gamma} p \, dx.$$

From Bernoulli's Law, derived earlier,

$$p = -\frac{1}{2}\rho(u^2 + v^2) + \text{constant},$$

where ρ is the (constant) density. Hence,

$$\begin{aligned} V + iH &= \int_{\Gamma} p(dx - i \, dy) \\ &= -\frac{1}{2}\rho \int_{\Gamma} (u^2 + v^2)(dx - i \, dy), \end{aligned}$$

since the integral of the constant term is zero. A bit of manipulation produces the relation

$$(u^2 + v^2)(dx - i \, dy) - (u - iv)^2(dx + i \, dy) = -2i(u - iv)(-v \, dx + u \, dy).$$

However, $-v \, dx + u \, dy$ is exactly the component of the flow f that is normal to Γ . Since Γ is a streamline of the flow, this quantity is zero. This results in

$$\begin{aligned} V + iH &= -\frac{1}{2}\rho \int_{\Gamma} (u - iv)^2(dx + i \, dy) \\ &= -\frac{1}{2}\rho \int_{\Gamma} [\overline{f(z)}]^2 \, dz. \end{aligned}$$

The function \bar{f}^2 is analytic, so the integral above is identical with the integral of \bar{f}^2 over the circle $|z| = R$, where R is so large that (1) is valid. Therefore, after squaring the expression for \bar{f} in (1) and replacing \bar{f}^2 with the resulting series, we obtain

$$\begin{aligned} V + iH &= -\frac{1}{2}\rho \int_{|z|=R} [\bar{f}(z)]^2 dz \\ &= -\frac{1}{2}\rho \int_{|z|=R} \left(a^2 + \frac{ca}{\pi i} \frac{1}{z} + \sum_{k=2}^{\infty} \frac{d_k}{z^k} \right) dz \\ &= -\rho c a, \end{aligned}$$

where $a = \lim_{z \rightarrow \infty} \bar{f}(z)$ is the complex conjugate of the velocity of the flow at ∞ .

This result, called the **Kutta–Joukowski Theorem**, is what we have been aiming for. It says that the flow f produces a force on Γ that is directed perpendicular to a , the velocity at ∞ , and of magnitude $\rho|c||a|$. In particular, if a is real, the force is vertical and thus is a **lifting force**.

Example 7 We begin our analysis of the lift generated by a flow with the flow around a circle. We make use of the observation in Section 4.1, Chapter 3 that it is enough to find an appropriate function that has the circle as one of the level curves of its imaginary part and the correct behavior at ∞ . Specifically, let the circle be centered at the origin and of radius R and let

$$G_1(z) = s \left(\lambda z + \frac{\bar{\lambda} R^2}{z} \right), \quad s > 0, \quad |\lambda| = 1$$

and

$$G_2(z) = -\frac{ic}{2\pi} \log z, \quad c \text{ real.}$$

G_1 is just a scaling of the mapping discussed in Example 5, Section 4, Chapter 3. Here s , λ , and c are all constants. On the circle $|z| = R$,

$$\operatorname{Im} G_1(Re^{it}) = 0$$

and

$$\operatorname{Im} G_2(Re^{it}) = -\frac{c}{2\pi} \log R.$$

Consequently, the circle $|z| = R$ is a streamline for $G_1 + G_2$. Although $G_2(z)$ is not well-defined (because $\log z$ has multiple values), its derivative is. Set

$$\begin{aligned}\overline{f(z)} &= G'_1(z) + G'_2(z) \\ &= s \left(\lambda - \frac{\bar{\lambda}R^2}{z^2} \right) - \frac{ic}{2\pi} \frac{1}{z}.\end{aligned}$$

Then \bar{f} is analytic on the plane minus the origin and f is also analytic at ∞ , where

$$\lim_{|z| \rightarrow \infty} \overline{f(z)} = \lambda s.$$

Furthermore, the circulation of the flow f is

$$\operatorname{Re} \left\{ \int_{|z|=r} \overline{f(z)} dz \right\} = c, \quad r \geq R.$$

Hence, \bar{f} is exactly the flow with circulation c and velocity λs at ∞ ; see Exercise 3.

The points z at which $f(z) = 0$ are called **stagnation points** of the flow. For this flow, there are stagnation points when

$$s \left(\lambda - \frac{\bar{\lambda}R^2}{z^2} \right) = \frac{ic}{2\pi z}.$$

Multiplying through by z^2 , we obtain a quadratic equation in z , which has roots

$$z_1, z_2 = \frac{ic \pm \sqrt{16\pi^2 s^2 R^2 - c^2}}{4\pi s \lambda}.$$

If $0 \leq c \leq 4\pi s R$, then both roots lie on the circle $|z| = R$; they coincide if $c = 4\pi s R$. Several streamlines of this flow are sketched in Figure 4.19. \square

Example 8 The Joukowski Airfoil Let C be a circle with center z_0 which passes through the point $-R$ and contains the point R in its inside, $R > 0$. The function


$$\phi(z) = z + \frac{R^2}{z}$$

maps C onto a simple closed curve Γ (Fig. 4.20). The image curve Γ , which resembles the cross section of a wing, is called a **Joukowski*** airfoil.


Our goal is to find the flow f with Γ as a streamline and with uniform velocity λs at ∞ , $s > 0$ and $|\lambda| = 1$; for this we make use of conformal mapping and the results of Example 7.

Our first task is to prove that the mapping ϕ given above is a one-to-one mapping of the exterior of the circle C onto the exterior of the curve Γ . To this end, let Γ_0 be the (unique) circle through both the points R and $-R$, which is tangent to


* N. J. Joukowski, 1842–1921.



$$\sin \Theta (r - 1/r) - a \ln(R) = c$$




Two stagnation
points on the circle



One stagnation
point on the circle

Figure 4.19



$$\phi(z) = z + \frac{R^2}{z}$$

Figure 4.20




Figure 4.21

C at $-R$ (Fig. 4.21). The center of Γ_0 lies at the point $i\alpha_0$, α_0 real. It is easy to prove (see Exercise 1 at the end of this section) that a point $z = x + iy$ lies inside, on, or outside Γ_0 if and only if the quantity

$$|z|^2 - 2\alpha_0 y - R^2$$

is negative, zero, or positive, respectively. Thus, z is outside Γ_0 if and only if $w = R^2/z$ is inside Γ_0 (also, see Exercise 2). From this it follows easily that $\phi(z) = z + R^2/z$ is one-to-one on the outside of Γ_0 . For if $\phi(z_1) = \phi(z_2)$, then

$$z_1 - z_2 = R^2 \left(\frac{z_1 - z_2}{z_1 z_2} \right).$$

If $z_1 \neq z_2$, then $z_2 = R^2/z_1$, so z_1 and z_2 lie on opposite sides of Γ_0 , a contradiction. Since the outside of C is within the outside of Γ_0 , ϕ is one-to-one outside C .

Now we are ready to analyze the flow around the Joukowski airfoil. Let $G(\zeta)$ be the complex potential of this flow, and let H be the complex potential of the flow around the circle C ; we know H from Example 7. After taking into account the change of origin and scale, we find that

$$H(z) = s \left[\lambda(z - z_0) + \bar{\lambda} \frac{|z_0 + R|^2}{z - z_0} \right] - \frac{ic}{2\pi} \log(z - z_0).$$

However, the function $\zeta = \phi(z)$ carries the exterior of C onto the exterior of Γ , as we have just shown, so $H(z)$ and $G(\zeta)$ must be related by $H(z) = G(\phi(z))$. Consequently,

$$H'(z) = G'(\phi(z))\phi'(z) = G'(\phi(z)) \left(1 - \frac{R^2}{z^2} \right).$$

Thus, the flow $H'(z)$ has a stagnation point at $z = -R$. Some simple arithmetic then yields

$$c = -4\pi s \operatorname{Im}(\lambda(R + z_0)),$$

so the circulation is determined by three factors: (1) the point R ; (2) the center, z_0 , of the circle C ; and (3) λs , the velocity of the flow at ∞ . Thus, in turn, the circulation of the flow determines the lift in accordance with the Kutta–Joukowski formula derived earlier. If $\lambda = 1$ (that is, the flow is horizontal at ∞), then the lift is

$$\text{Lift} = 4\pi s^2 \operatorname{Im} z_0, \quad s = \text{speed of flow at } \infty.$$

□

Other Physical Phenomena in Which Harmonic Functions Arise

Diffusion

Imagine a level channel connecting a very large body of fresh water to a very large body of salt water, say the ocean (Fig. 4.22). The water in the channel will have various degrees of salinity, depending on location; that is, the salt from the ocean will diffuse through the channel to the fresh water, which being large, will absorb it. If we let $c(x, y)$ be the concentration of salt at the point $x + iy$ in the channel, then an analysis exactly like that for temperature distribution shows that $c(x, y)$ is a harmonic function of x and y . More generally, if a substance diffuses from a region of high concentration to one of low or zero concentration, then the concentration function $c(x, y)$ is a harmonic function in any domain in which there are no sources or sinks.




Figure 4.22 The diffusion of salt through a channel joining the ocean with a lake.

Planar Stress

In an isotropic, homogeneous body, there are nine stress components: the normal stress component and two shear stress components in each of the x , y , and z directions. There are compatibility relations among these stress components, which, in the case of plane strain with no body forces acting, reduce to

$$\frac{\partial X_x}{\partial x} + \frac{\partial X_y}{\partial y} = 0 \quad (2)$$

$$\frac{\partial X_y}{\partial x} + \frac{\partial Y_y}{\partial y} = 0 \quad (3)$$

$$\left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) (X_x + Y_y) = 0. \quad (4)$$

Here X_x , X_y , and Y_y are the normal and shear stress components in the x -direction and the normal stress component in the y -direction, respectively. We now assume that the cross-section of the body is a simple closed curve together with the domain D inside it. From equation (2), we deduce that there is a function F with

$$\frac{\partial F}{\partial x} = -X_y \quad \text{and} \quad \frac{\partial F}{\partial y} = X_x.$$

From equation (3), it follows that there is a function G with

$$\frac{\partial G}{\partial x} = Y_y \quad \text{and} \quad \frac{\partial G}{\partial y} = -X_y.$$

From the relations among the partial derivatives of F and G , we further conclude that there is a function U with

$$\frac{\partial U}{\partial x} = G \quad \text{and} \quad \frac{\partial U}{\partial y} = F.$$

Thus,

$$\frac{\partial^2 U}{\partial x^2} = Y_y, \quad \frac{\partial^2 U}{\partial y^2} = X_x, \quad \text{and} \quad \frac{\partial^2 U}{\partial x \partial y} = -X_y.$$

U is called the **stress function** or the **Airy function**, after G. B. Airy (1801–1892), who worked with it (circa 1860). Thus, from equation (4),

$$0 = \Delta(X_x + Y_y) = \Delta(\Delta U) = \Delta^2 U.$$

The stress function U is therefore **biharmonic**—that is, it satisfies the equation

$$\frac{\partial^4 U}{\partial x^4} + 2 \frac{\partial^4 U}{\partial x^2 \partial y^2} + \frac{\partial^4 U}{\partial y^4} = 0.$$

The famous nineteenth-century physicist J. C. Maxwell (1831–1879) was the first to note that U must be biharmonic.

The Representation of Biharmonic Functions

If U is biharmonic, then $P = \Delta U$ is harmonic. Let ϕ be an analytic function on D with

$$4\phi' = P + iQ,$$

where Q is a harmonic conjugate of P on D ; such a ϕ exists because D is simply connected. We write $\phi = p + iq$. A simple computation shows that

$$U - xp - yq$$

is harmonic on D . Hence, this is the real part of some analytic function f on D , again because D is simply-connected. Consequently,

$$U(z) = \operatorname{Re}\{\bar{z}\phi(z) + f(z)\}, \quad (5)$$

where ϕ and f are analytic and

$$4 \operatorname{Re} \phi' = \Delta U.$$

(5) is the most general representation of a biharmonic function U .

EXERCISES FOR SECTION 4.2

- Let Γ_0 be the circle of radius $\sqrt{R^2 + \alpha_0^2}$ centered at the point $i\alpha_0$, $\alpha_0 > 0$. Show that a point z is outside, on, or inside Γ_0 exactly when the quantity $|z|^2 - 2\alpha_0 y - R^2$ is positive, zero, or negative.
- Let Γ_0 be the circle from Exercise 1. Show that z is inside Γ_0 if and only if R^2/z is outside Γ_0 .
- Let Γ be a simple closed curve. Show that a flow outside Γ (including at ∞) with Γ as a streamline is completely determined by specifying the velocity at ∞ and the circulation. That is, show that there is at most *one* analytic function g outside Γ with
 - $g(z) = a + \frac{c}{2\pi i} \frac{1}{z} + \dots$; a, c fixed, $|z|$ large,
 - $\operatorname{Im} G = 0$ on Γ , where $G' = g$.
- Find the stagnation points of the flow in Example 5.

5. Discuss the flow within a disc caused by placing a source and a sink of equal strengths at distinct points on the edge of the disc. Find the streamlines for the flow and sketch a few.
6. Extend Example 4 by finding the potential of the flow in the upper half-plane U resulting from a source uniformly distributed on the interval $[-1, 1]$.
7. Find the flow in the semi-infinite strip $\{x + iy: y > 0 \text{ and } -\pi < x < \pi\}$ established by placing two sources across from each other (on the long sides).
8. Redo Exercise 7 with one of the sources changed to a sink.
9. Find the flow in the strip of Exercise 7 if there is a source on the short edge (where $y = 0$).
10. Let D be a region bounded by a smooth simple closed curve Γ and let f be a sourceless flow in D ; f is not assumed to be irrotational. The kinetic energy of the flow is

$$K = \frac{1}{2} \rho \iint_D |f(z)|^2 dx dy,$$

where ρ is the density (which we assume to be constant). Show that among all sourceless flows in D with given normal velocity at the boundary of D (that is, the normal derivative $\partial f / \partial n$ is specified at Γ), the flow of *smallest* kinetic energy is *irrotational*. This result is known as **Kelvin's* Minimum-Energy Theorem**; you will need to use Green's Identity (Exercise 13, Section 6, Chapter 1).

11. (a) Show that the function $G(z) = \operatorname{Log}(z^2 + 1)$ is a conformal mapping of the first quadrant onto the horizontal strip $S = \{s + it: -\infty < s < \infty, 0 < t < \pi\}$.
 (b) Show that the flow f in the first quadrant whose potential function is $G(z) = \operatorname{Log}(z^2 + 1)$ is the sourceless, irrotational flow resulting from a source $z_0 = i$ on the boundary of the first quadrant. What is the strength of the source?
 (c) Extend the result in (b) to the case when the source is located at some other point in the boundary of the first quadrant. See Figure 4.23.
12. Find the sourceless, irrotational flow in the upper half-plane that results from a source at the point i . (**Hint:** The flow will be symmetric about the imaginary axis; you may now use the result of Exercise 11b.)
13. Find the electrostatic potential due to a uniformly distributed charge on a straight wire. (**Hint:** Take the wire to be the interval $[-1, 1]$ and the charge density to be $d\rho = c_0 dx$.)
14. Show that the electrostatic potential inside a long hollow uniformly charged tube is constant and that outside the tube the potential is zero.
15. Find the flow past a long cylinder with an elliptical cross section (Fig. 4.24).

* Sir William Thompson, Baron Kelvin of Largs, 1824–1907.




Figure 4.23 The flow in the first quadrant from a source on the imaginary axis.




Figure 4.24 The flow around the cylinder with elliptical cross section $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$.

16. Show that the function

$$f(z) = \frac{K}{2\pi i} \frac{1}{\bar{z} - \bar{z}_0}, \quad K > 0,$$

is the velocity of a flow that rotates clockwise about the point z_0 . This flow has a vortex at z_0 of strength K . Show that the streamlines are circles centered at z_0 .

17. Show that the function

$$g(z) = \frac{K_1}{2\pi i} \frac{1}{z - z_1} - \frac{K_2}{2\pi i} \frac{1}{z - \bar{z}_2},$$

K_1 and K_2 both positive, is the velocity of a flow with two vortices, one at z_1 of strength K_1 with clockwise rotation, the other at z_2 of strength K_2 with counterclockwise rotation. What is the circulation of this flow on a curve γ , which is very far from both z_1 and z_2 ? (Some streamlines of this flow are shown in Fig. 4.25.)




Figure 4.25

18. The flow

$$h(z) = \frac{Q + iK}{2\pi} \frac{1}{z - z_0}$$

produces a **spiral vortex** at z_0 (Fig. 4.26). Sketch some of the streamlines and decide how the signs of Q and K affect the streamlines; that is, when will the streamlines spiral into z_0 clockwise and when will they spiral in counterclockwise? See Figure 4.26.





Figure 4.26 A spiral vortex produced by the flow $\frac{Q+iK}{2\pi i} \frac{1}{z}$, $QK \neq 0$.

19. Use Example 2 to find the electrostatic potential produced by two parallel, horizontal charged plates whose edges are separated by a distance 2σ (see the diagram below). Sketch some of curves of flux of the field.



4.3 Integral Representations of Harmonic Functions

A function f has an **integral representation formula** if $f(z)$ is equal to an integral of an expression containing z as a parameter. A perfect example is the Cauchy Integral Formula:

$$f(z) = \frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

Integral representation formulas are important tools in solving boundary-value problems. In this section, we explore such formulas in two special but important cases: the disc $\{z: |z| < 1\}$ and the upper half-plane $\{z: \operatorname{Im} z > 0\}$.

Suppose first that u is a real-valued function that is harmonic on some open set that includes the closed disc $\{z: |z| \leq 1\}$. We shall derive a formula for the values of $u(x, y)$ when $x^2 + y^2 < 1$, which makes use only of the values of $u(x, y)$ when $x^2 + y^2 = 1$. Since u is harmonic on some disc centered at the origin and of radius

larger than 1, there is an analytic function f on this same disc with $\operatorname{Re} f = u$. Thus, for a particular z with $|z| < 1$, we have two formulas:

$$f(z) = \frac{1}{2\pi i} \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta \quad (1)$$

and

$$0 = \frac{1}{2\pi i} \int_{\Gamma} \frac{f(\zeta)\bar{z}}{1 - \bar{z}\zeta} d\zeta. \quad (2)$$

The first, (1), is the Cauchy Integral Formula with Γ the circle $|\zeta| = 1$ oriented positively. The second, (2), is just Cauchy's Theorem applied to the function

$$G(\zeta) = \frac{f(\zeta)\bar{z}}{1 - \bar{z}\zeta},$$

which is analytic on an open set containing $|\zeta| \leq 1$. We set $\zeta = e^{it}$, $0 \leq t \leq 2\pi$, as usual, and find that

$$f(z) = \frac{1}{2\pi} \int_0^{2\pi} \frac{f(e^{it})}{1 - ze^{-it}} dt \quad (3)$$

and

$$0 = \frac{1}{2\pi} \int_0^{2\pi} \frac{f(e^{it})\bar{z}e^{it}}{1 - \bar{z}e^{it}} dt. \quad (4)$$

However,

$$\frac{1}{1 - ze^{-it}} + \frac{\bar{z}e^{it}}{1 - \bar{z}e^{it}} = \frac{1 - |z|^2}{|1 - \bar{z}e^{it}|^2}.$$

When we add (3) and (4), we therefore obtain

$$f(z) = \frac{1}{2\pi} \int_0^{2\pi} f(e^{it}) \frac{1 - |z|^2}{|1 - \bar{z}e^{it}|^2} dt.$$

The function appearing after $f(e^{it})$ under the integral sign has a special notation. Set

$$P_r(t) = \frac{1 - r^2}{1 - 2r \cos t + r^2} \quad (5)$$

$$= \operatorname{Re} \left\{ \frac{1 + z}{1 - z} \right\} \quad \text{if } z = re^{it}.$$

$P_r(t)$ is called the **Poisson* kernel** for r and t . Thus, we have shown that

$$f(re^{i\theta}) = \frac{1}{2\pi} \int_0^{2\pi} f(e^{it}) P_r(\theta - t) dt. \quad (6)$$

Since $P_r(\theta - t)$ is real-valued, we may take the real part of both sides in this formula and conclude that

$$u(re^{i\theta}) = \frac{1}{2\pi} \int_0^{2\pi} u(e^{it}) P_r(\theta - t) dt. \quad (7)$$

This is the **Poisson Integral Formula**; it gives the value of the function u in the disc $|z| < 1$ in terms of the values of $u(e^{it})$, $0 \leq t \leq 2\pi$.

Several extensions of this formula are immediate. First, if the disc is not $|z| \leq 1$ but rather $|w - w_0| \leq R$, then the change of variables

$$z = \frac{w - w_0}{R}$$

produces the formula

$$u(w) = \frac{1}{2\pi} \int_0^{2\pi} u(w_0 + Re^{it}) \frac{R^2 - |w - w_0|^2}{|R - (\bar{w} - \bar{w}_0)e^{it}|^2} dt. \quad (8)$$

The formula is valid for $|w - w_0| < R$.

Substantially more important, we need not assume that u is harmonic on an open set containing the disc $|z| \leq 1$. Suppose, in fact, only that u is a continuous function on the circle $|z| = 1$. Set

$$U(re^{i\theta}) = \frac{1}{2\pi} \int_0^{2\pi} u(e^{it}) P_r(\theta - t) dt. \quad (9)$$

It is then true that $U(z)$, $z = re^{i\theta}$, is a harmonic function on the disc $|z| < 1$ and at all points λ on the circle $|\lambda| = 1$,

$$\lim_{z \rightarrow \lambda} U(z) = u(\lambda). \quad (10)$$

That is, the function defined by

$$\begin{cases} U(z) & \text{if } |z| < 1 \\ u(z) & \text{if } |z| = 1 \end{cases}$$

is continuous on the closed set $\{z: |z| \leq 1\}$ and harmonic on the open set $\{z: |z| < 1\}$. Part of this assertion is easy: If u has only real values, then

* Siméon Denis Poisson, 1781–1840.

$$\begin{aligned} U(z) &= \frac{1}{2\pi} \int_0^{2\pi} u(e^{it}) \operatorname{Re} \left(\frac{e^{it} + z}{e^{it} - z} \right) dt \\ &= \operatorname{Re} \left\{ \frac{1}{2\pi} \int_0^{2\pi} u(e^{it}) \frac{e^{it} + z}{e^{it} - z} dt \right\}. \end{aligned}$$

The expression within the braces is an analytic function of z , so U is harmonic, being the real part of an analytic function. In general, $u = u_1 + iu_2$, where u_1 and u_2 are real-valued. Hence, $U = U_1 + iU_2$, where both U_1 and U_2 are harmonic; thus, U is also harmonic. The proof that (10) holds is more difficult; it is outlined in the exercises at the end of this section.

There is even a further extension of (7). A function $u(e^{it})$ is **piecewise continuous** if the circle $\{z: |z| = 1\}$ can be broken up into closed arcs I_1, \dots, I_N that are disjoint except for their endpoints (Fig. 4.27) and, furthermore, on each of the arcs I_j , $u(e^{it})$ is continuous. (This gives two values to u at each of the N endpoints of the arcs; this situation will not bother us, because such a set “does not count” in integration.) If $u(e^{it})$ is piecewise continuous and U is defined by (9), then U is harmonic in the disc $|z| < 1$, and, except for the N points that are the endpoints of the arcs I_1, \dots, I_N ,

$$\lim_{z \rightarrow \lambda} U(z) = u(\lambda), \quad \lambda \text{ is not an endpoint of } I_1, \dots, I_N.$$




Figure 4.27

Example 1 A long cylinder with circular cross section is split lengthwise down the middle, and the two halves are insulated from one another. One half is given a charge of $+1$ and the other half is given the charge of -1 . Find the resulting electrostatic potential inside the cylinder.

Solution We may assume that a cross section of the cylinder is the circle $x^2 + y^2 = 1$. Hence, we must find a function $u(x, y)$ that is harmonic for $x^2 + y^2 < 1$ and that has the value $+1$ on the semicircle $x^2 + y^2 = 1$, $y > 0$, and the value -1 on the semicircle $x^2 + y^2 = 1$, $y < 0$. The answer is given to us immediately by the Poisson Integral Formula:

$$u(re^{i\theta}) = \frac{1}{2\pi} \int_0^\pi \frac{1-r^2}{1-2r \cos(\theta-t) + r^2} dt - \frac{1}{2\pi} \int_{-\pi}^0 \frac{1-r^2}{1-2r \cos(\theta-t) + r^2} dt.$$

We evaluate the integrals in the following way. Since

$$\operatorname{Re}\left(\frac{1+z}{1-z}\right) = \frac{1-r^2}{1-2r \cos \theta + r^2}, \quad z = re^{i\theta},$$

and since

$$\frac{1+z}{1-z} = 1 + \frac{2z}{1-z},$$

we have

$$\frac{1+ze^{-it}}{1-ze^{-it}} = 1 + \frac{2ze^{-it}}{1-ze^{-it}}.$$

Thus,

$$\begin{aligned} \int_0^\pi \frac{1+ze^{-it}}{1-ze^{-it}} dt &= \pi + 2z \left(\overline{\int_0^\pi \frac{e^{it}}{1-\bar{z}e^{it}} dt} \right) \\ &= \pi + 2z \left(\overline{\frac{1}{i} \int_\gamma \frac{dw}{1-\bar{z}w}} \right), \end{aligned}$$

where γ is the semicircle $\{e^{it}: 0 \leq t \leq \pi\}$. The antiderivative of $\bar{z}(1-\bar{z}w)^{-1}$ with respect to w is $\log(1-\bar{z}w)$, so

$$\int_0^\pi \frac{1-r^2}{1-2r \cos(\theta-t) + r^2} dt = \pi + 2\operatorname{Arg}\left(\frac{1+z}{1-z}\right).$$

Likewise,

$$\int_{-\pi}^0 \frac{1-r^2}{1-2r \cos(\theta-t) + r^2} dt = \pi - 2\operatorname{Arg}\left(\frac{1+z}{1-z}\right).$$

Putting these two together yields

$$u(z) = \frac{2}{\pi} \operatorname{Arg}\left(\frac{1+z}{1-z}\right). \quad \square$$

The Poisson Integral Formula in the Upper Half-Plane

Suppose that $w(t)$ is a bounded, piecewise continuous function on the real axis, $-\infty < t < \infty$. There is an integral formula, much like the Poisson Integral Formula, that yields a function $W(\zeta)$ that is bounded and harmonic in the upper half-plane $\{\zeta = \sigma + it: t > 0\}$ and satisfies

$$\lim_{\zeta \rightarrow s} W(\zeta) = w(s)$$

for all points s on the real axis at which $w(t)$ is continuous (that is, at all but a finite number of points). The formula is this:

$$W(\sigma + i\tau) = \frac{1}{\pi} \int_{-\infty}^{\infty} w(t) \frac{\tau}{(\sigma - t)^2 + \tau^2} dt, \quad \tau > 0. \quad (11)$$

Its derivation is simple. Let

$$\zeta = \psi(z) = i \frac{1+z}{1-z}.$$

Then ψ is a linear fractional transformation that maps the disc $\{z : |z| < 1\}$ onto the upper half-plane, and ψ maps the circle $|z| = 1$ onto the real axis. Hence,

$$u(e^{i\theta}) = w\left(i \frac{1+e^{i\theta}}{1-e^{i\theta}}\right)$$

is bounded and piecewise continuous on the circle $\{e^{i\theta} : 0 \leq \theta \leq 2\pi\}$. Consequently, the function

$$U(z) = \frac{1}{2\pi} \int_0^{2\pi} u(e^{i\theta}) \frac{1-r^2}{1-2r \cos(s-\theta)+r^2} d\theta, \quad z = re^{is}$$

is bounded and harmonic in the disc $|z| < 1$, and $U(z) \rightarrow u(e^{i\theta})$ as $z \rightarrow e^{i\theta}$. Suppose that $w(t)$ is real-valued; then u is real-valued and

$$U(z) = \operatorname{Re} \left\{ \frac{1}{2\pi} \int_0^{2\pi} u(e^{i\theta}) \frac{e^{i\theta} + z}{e^{i\theta} - z} d\theta \right\}.$$

Now $z = (\zeta - i)/(\zeta + i)$, so

$$W(\zeta) = U\left(\frac{\zeta - i}{\zeta + i}\right)$$

is harmonic on the upper half-plane. But $e^{i\theta} = (t - i)/(t + i)$ and $e^{i\theta} d\theta = 2(t + i)^{-2} dt$. With these changes of variables,

$$\begin{aligned} W(\zeta) &= \operatorname{Re} \left[\frac{1}{2\pi} \int_0^{2\pi} u(e^{i\theta}) \frac{e^{i\theta} + z}{e^{i\theta} - z} d\theta \right] \\ &= \operatorname{Re} \left[\frac{1}{2\pi} \int_{-\infty}^{\infty} w(t) \frac{t\zeta + 1}{i(t - \zeta)} \frac{2}{1+t^2} dt \right] \\ &= \frac{1}{\pi} \int_{-\infty}^{\infty} w(t) \frac{\tau}{(t - \sigma)^2 + \tau^2} dt. \end{aligned}$$

This is the desired formula; it is called the **Poisson Integral Formula for the upper half-plane**.

Example 2 Find the electrostatic potential in the region $y > 0$ if the segment $(-\sigma, \sigma)$ of the real axis is kept at potential $V_0 > 0$ and the remainder of the real axis is kept at potential 0.

Solution The electrostatic potential $V(x, y)$ is a harmonic function for $y > 0$ and must satisfy

To find such a harmonic function, we use the Poisson Integral Formula for the upper half-plane.

$$\begin{aligned} V(x, y) &= \frac{1}{\pi} \int_{-\infty}^{\infty} V(t, 0) \frac{y}{(x-t)^2 + y^2} dt \\ &= \frac{V_0}{\pi} \int_{-\sigma}^{\sigma} \frac{y}{(t-x)^2 + y^2} dt \\ &= \frac{V_0}{\pi} \left\{ \arctan\left(\frac{\sigma-x}{y}\right) - \arctan\left(\frac{-\sigma-x}{y}\right) \right\} \end{aligned}$$

A function $U(x, y)$ that is a harmonic conjugate of $V(x, y)$ in the upper half-plane is given by




Figure 4.28

$$U(x, y) = \frac{V_0}{\pi} \log \left| \frac{z - \sigma}{z + \sigma} \right|.$$

The level curves of U represent the curves of flux of the electric field; these curves are just the Circles of Apollonius,

$$|z - \sigma| = \rho |z + \sigma|, \quad 0 < \rho < \infty,$$

which are centered at the real axis, either in the interval $(-\infty, -\sigma)$ or in the interval (σ, ∞) (Fig. 4.28); see Section 2, Chapter 1. \square

The Reflection Principle

The Poisson integral formula in a disc can be used to give a proof of the following theorem, which is called, for reasons that will be clear from its statement, the reflection principle.

THEOREM 1 The Reflection Principle Suppose D is a domain in the upper half-plane $\{x + iy: y > 0\}$, which includes in its boundary a segment (a, b) of the real axis. Suppose that $f = u + iv$ is analytic in D and, moreover, is continuous at each point of (a, b) . If $v(x) = 0$ for all x in the interval (a, b) , then f extends to be analytic in the domain $\Omega = D \cup (a, b) \cup D^*$, where D^* is the reflection of D over the real axis:

$$D^* = \{z: \bar{z} \in D\}.$$

Furthermore, f satisfies the relation

$$f(z) = \overline{f(\bar{z})}, \quad z \in \Omega.$$

Proof The sets D , D^* , and Ω are illustrated in Figure 4.29. We begin the proof by defining a function f^* in D^* by




Figure 4.29

$$f^*(z) = \overline{f(\bar{z})}, \quad z \in D^*.$$

We shall show that f^* is analytic in D^* by this simple argument. If $z_0 \in D^*$, then $\bar{z}_0 \in D$. Now f is analytic in some disc $|z - \bar{z}_0| < \delta$ centered at \bar{z}_0 ; thus, f has a power series valid in this disc:

$$f(z) = \sum_{k=0}^{\infty} A_k(z - \bar{z}_0)^k, \quad |z - \bar{z}_0| < \delta.$$

Hence, in the disc $|z - z_0| < \delta$,

$$\begin{aligned} f^*(z) &= \overline{f(\bar{z})} = \overline{\sum_{k=0}^{\infty} A_k(\bar{z} - \bar{z}_0)^k} \\ &= \sum_{k=0}^{\infty} \bar{A}_k(z - z_0)^k. \end{aligned}$$

Consequently, f^* is analytic for $|z - z_0| < \delta$, since it has a representation there as the sum of a convergent power series.

Next, we define a function F on Ω by

$$F(z) = \begin{cases} f(z), & z \in D \cup (a, b) \\ f^*(z), & z \in D^*. \end{cases}$$

The assumption that $v(x) = 0$ for $a < x < b$ implies that F is continuous on Ω . The functions u and v are extended by D^* by the rules $u(z) = u(\bar{z})$, $v(z) = -v(\bar{z})$, $z \in D^*$. The tricky part of the proof is to show that $F = u + iv$ is analytic across the segment (a, b) . To see this, let $x_0 \in (a, b)$, and let r be a positive number so small that the disc $|z - x_0| < 2r$ lies in Ω . Let γ be the positively oriented circle $|z - x_0| = r$. On this circle, the function

$$w(\theta) = \begin{cases} v(x_0 + re^{i\theta}), & \text{if } 0 \leq \theta \leq \pi \\ -v(x_0 + re^{-i\theta}), & \text{if } -\pi \leq \theta \leq 0 \end{cases}$$

is continuous and odd: $w(\theta) = -w(-\theta)$. Let v_1 be the harmonic function on the disc $|z - x_0| < r$, whose values on γ are w ; v_1 is found by using the Poisson integral, appropriately scaled. Then v_1 satisfies $v_1(z) = -v_1(\bar{z})$ if $|z - x_0| < r$ (also, see Exercise 8). Thus, v_1 vanishes on the real segment $x_0 - r \leq x \leq x_0 + r$. The function $v - v_1$ is then harmonic on the half-disc $\mathcal{U} = \{z: |z - x_0| < r \text{ and } \operatorname{Im} z > 0\}$ and vanishes on the boundary of \mathcal{U} , since $v = v_1 = w$ on the semicircle $z = x_0 + re^{i\theta}$, $0 \leq \theta \leq \pi$, and $v = v_1 = 0$ on the segment $x_0 - r \leq x \leq x_0 + r$. Hence, the maximum principle for harmonic functions [(5) in Section 1] implies that $v = v_1$ on all of \mathcal{U} . Likewise, $v = v_1$ on all of $\mathcal{U}^* = \{z: \bar{z} \in \mathcal{U}\}$. Thus, $v = v_1$ on the disc $|z - x_0| < r$. But v_1 is harmonic on all this disc; hence, so must be v . The function u is a harmonic conjugate of v on both \mathcal{U} and \mathcal{U}^* . Since v is harmonic on all the disc

$|z - x_0| < r$, it has a harmonic conjugate on all this disc that must differ from u by a constant on $\mathcal{U} \cup \mathcal{U}^*$. Hence, u is also harmonic on the disc $|z - x_0| < r$, so F is actually analytic on this disc. ■

REMARK: The reflection principle is frequently proved by use of Morera's Theorem.

Example 3 Show that an entire function that is real-valued on some interval (a, b) in the real axis is real-valued on all the real axis.

Solution Let f be an entire function that is real-valued on the interval (a, b) . Set

$$F(z) = \begin{cases} f(z), & \text{if } \operatorname{Im} z \geq 0 \\ \overline{f(\bar{z})}, & \text{if } \operatorname{Im} z < 0. \end{cases}$$

The reflection principle implies immediately that F is analytic on the domain Ω , which consists of the union of the upper and lower half-planes and the interval (a, b) . Furthermore, f itself is of course analytic on Ω and equals F if $\operatorname{Im} z > 0$. Thus, $f = F$ throughout Ω ; in particular,

$$f(z) = \overline{f(\bar{z})}, \quad \operatorname{Im} z \neq 0.$$

Since f is actually continuous on the whole plane, it follows that for any real number x ,

$$f(x) = \lim_{y \rightarrow 0} f(x + iy) = \lim_{y \rightarrow 0} \overline{f(x - iy)} = \overline{f(x)},$$

which says exactly that f is real on the real axis. □

EXERCISES FOR SECTION 4.3

- An arc of θ_0 radians in a circle is kept at temperature T_1 , while the remainder of the circle is kept at temperature T_2 . Find the temperature distribution inside the circle. Sketch some of the equitherms.
- The segment $[-\sigma, \sigma]$ is kept at temperature T_1 , while the remainder of the real axis is kept at temperature T_2 . Find the temperature distribution in the upper half-plane; sketch some of the equitherms.
- Let $w(t)$ be a bounded piecewise continuous function on the real axis with $\int_{-\infty}^{\infty} |w(t)| dt < \infty$ and $w(t) \rightarrow 0$ as $|t| \rightarrow \infty$. Let $W(\zeta)$ be the harmonic function on the upper half-plane $\operatorname{Im} \zeta > 0$, with $W(\sigma, 0) = w(\sigma)$, given by formula (11).
 - Show that $\lim_{|\zeta| \rightarrow \infty} W(\zeta) = 0$.
 - Show that if $W_1(\zeta)$ is (another) harmonic function on $\operatorname{Im} \zeta > 0$ with $W_1(\sigma, 0) = w(\sigma)$, and that if W_1 also satisfies (a), then $W_1 = W$.
 - Show that $\int_{-\infty}^{\infty} |W(x + iy)| dx \leq \int_{-\infty}^{\infty} |w(t)| dt$ for each $y > 0$.
- Let $P_r(t)$ be the Poisson kernel, defined in formula (5). Show that
 - $P_r(t) > 0$ for $0 \leq r < 1$, $-\pi \leq t \leq \pi$,

$$(ii) \frac{1}{2\pi} \int_{-\pi}^{\pi} P_r(t) dt = 1, \quad 0 \leq r < 1,$$

(iii) if $\delta > 0$, then $\lim_{r \rightarrow 1^-} \{\max_{\pi \geq |t| \geq \delta} P_r(t)\} = 0$.

5. We show here that (10) holds. Fix a point λ with $|\lambda| = 1$ and write $\lambda = e^{i\theta_0}$. Fill in the reasons for each of the following steps:

(i) Given $\varepsilon > 0$, there is a $\delta > 0$ such that

$$|u(e^{it}) - u(e^{i\theta_0})| < \frac{\varepsilon}{2} \quad \text{if } |t - \theta_0| < \delta.$$

$$(ii) U(re^{i\theta}) - u(e^{i\theta_0}) = \frac{1}{2\pi} \int_{-\pi}^{\pi} \{u(e^{it}) - u(e^{i\theta_0})\} P_r(\theta - t) dt.$$

Now consider θ with $|\theta - \theta_0| < \delta/4$. Write the integral as the sum of two integrals: The first, I_1 , is over those t with $|t - \theta_0| < \delta$; the second, I_2 , is over all the remaining t .

$$(iii) |I_1| < \frac{\varepsilon}{2} \frac{1}{2\pi} \int_{|t-\theta_0|<\delta} P_r(\theta - t) dt < \frac{\varepsilon}{2}.$$

$$(iv) |I_2| < 2M \left(\max \left\{ P_r(\theta - t) : |\theta - t| \geq \frac{3\delta}{4} \right\} \right),$$

where M is an upper bound on $|u(e^{it})|$, $-\pi \leq t \leq \pi$.

$$(v) |I_2| < \frac{\varepsilon}{2} \text{ if } r \text{ is close enough to 1.}$$

$$(vi) |U(re^{i\theta}) - u(e^{i\theta_0})| < \varepsilon \text{ if } |\theta - \theta_0| < \frac{\delta}{4} \text{ and } r \text{ is close enough to 1.}$$

6. Show that the function

$$u(x, y) = \frac{1}{\pi} \int_0^\infty \left\{ \frac{y}{(x-t)^2 + y^2} - \frac{y}{(x+t)^2 + y^2} \right\} f(t) dt$$

is harmonic in the upper half-plane and satisfies

$$u(0, y) = 0, \quad 0 < y < \infty$$

$$u(x, 0) = f(x), \quad 0 < x < \infty.$$

7. Use Exercise 6 to find an integral representation for a harmonic function $u(x, y)$ in the first quadrant $\{x + iy : x > 0, y > 0\}$ with

$$u(x, 0) = f_1(x), \quad 0 < x < \infty$$

$$u(0, y) = f_2(y), \quad 0 < y < \infty.$$

8. Show that the function

$$u(re^{i\theta}) = \frac{1}{2\pi} \int_0^\pi \{P_r(t - \theta) - P_r(t + \theta)\} f(e^{it}) dt$$

is harmonic in the half-disc $\{re^{i\theta}: 0 < r < 1, 0 < \theta < \pi\}$ and satisfies the boundary conditions

$$u(re^{i\theta}) = f(e^{i\theta}), \quad 0 < \theta < \pi$$

$$u(x) = 0, \quad -1 < x < 1.$$

9. Solve the Dirichlet problem for the exterior of the disc of radius 1 centered at the origin. That is, give an integral formula for a harmonic function $v(z)$ on the region $1 < |z| \leq \infty$ (including ∞) that satisfies

$$\lim_{z \rightarrow \lambda} v(z) = f(\lambda), \quad |\lambda| = 1.$$

10. Use the result of Example 2 to find the electrostatic potential in the region

$$D = \{x + iy: x^2 + y^2 > \sigma^2 \text{ and } y > 0\}$$

when the potential is 0 on both the segments $(-\infty, -\sigma)$ and (σ, ∞) and 1 on the semicircle $x^2 + y^2 = \sigma^2, y > 0$.

11. The function

$$u(re^{i\theta}) = \frac{1 - r^2}{1 - 2r \cos \theta + r^2}$$

is harmonic on the disc $\{z: |z| < 1\}$. Show that

$$\lim_{r \rightarrow 1^-} u(re^{i\theta}) = 0 \quad \text{if} \quad \pi \geq |\theta| > 0.$$

Show, moreover, that $u(re^{i\theta})$ represents the temperature distribution in the disc $0 \leq r < 1$ due to a “hot spot” at the point $r = 1, \theta = 0$ (that is, the temperature there is “ ∞ ”). (**Hint:** Find the temperature distribution u_n in the disc due to the boundary temperature

$$u_n(e^{i\theta}) = \begin{cases} n, & |\theta| < \frac{\pi}{2n} \\ 0, & \frac{\pi}{2n} < |\theta| \leq \pi. \end{cases}$$

Then let $n \rightarrow \infty$.)

12. Extend the result in Exercise 11 to the case when there are N “hot spots” at points $e^{i\theta_1}, \dots, e^{i\theta_N}$ of the circle of radius 1 centered at the origin. Discuss what happens if some of the “hot spots” are instead “cold spots” (that is, the temperature there is $-\infty$).
13. Give the temperature distribution in the upper half-plane $y > 0$ that is produced

by a single “hot spot” at the point x_0 on the real axis. Do the same for several “hot/cold spots” at the points x_1, \dots, x_N on the real axis.

14. Argue on physical grounds that if $v(re^{i\theta})$ is a nonnegative harmonic function on the disc $\{z: |z| < 1\}$ with

$$v(re^{i\theta}) \leq \frac{1 - r^2}{1 - 2r \cos \theta + r^2} = P_r(\theta), \quad 0 \leq r < 1, \quad 0 \leq \theta \leq 2\pi,$$

then $v(re^{i\theta}) = \lambda P_r(\theta)$ for some $\lambda, 0 < \lambda \leq 1$. (Hint: What temperature distribution on $|z| = 1$ does v represent?)

15. Suppose $u(z)$ is a real-valued harmonic function on the disc $\{z: |z| < 1 + \delta\}$ for some $\delta > 0$; let v be the harmonic conjugate of u on this disc, which is zero at the origin. Show that

$$v(re^{i\theta}) = \frac{1}{2\pi} \int_0^{2\pi} u(e^{it}) \frac{2r \sin(\theta - t)}{1 - 2r \cos(\theta - t) + r^2} dt.$$

(Hint: The function

$$f(z) = \frac{1}{2\pi} \int_0^{2\pi} u(e^{it}) \frac{e^{it} + z}{e^{it} - z} dt$$

is analytic on the disc $\{z: |z| < 1\}$, and its real part is $u(z)$, by (7).)

Fourier Series and Harmonic Functions*

16. Let $u(e^{i\theta})$ be a piecewise continuous function on the circle $\{e^{i\theta}: 0 \leq \theta \leq 2\pi\}$. Set

$$\hat{u}(n) = \frac{1}{2\pi} \int_0^{2\pi} u(e^{i\theta}) e^{-in\theta} d\theta, \quad n = 0, \pm 1, \pm 2, \dots;$$

$\hat{u}(n)$ is the n th Fourier coefficient of u . Show that

$$|\hat{u}(n)| \leq \max_{0 \leq \theta \leq 2\pi} |u(e^{i\theta})|.$$

17. Let k be an integer. Show that

$$\frac{1}{2\pi} \int_0^{2\pi} e^{ik\theta} d\theta = \begin{cases} 1, & k = 0 \\ 0, & k \neq 0. \end{cases}$$

18. If $u(e^{i\theta}) = \sum_{-\infty}^{\infty} a_n e^{in\theta}$, where $\sum_{-\infty}^{\infty} |a_n| < \infty$, then $a_n = \hat{u}(n)$.

19. Let $P_r(\theta)$ be the Poisson kernel:

$$P_r(\theta) = \frac{1 - r^2}{1 - 2r \cos \theta + r^2} = \operatorname{Re} \left(\frac{1 + re^{i\theta}}{1 - re^{i\theta}} \right).$$

Show that

$$P_r(\theta) = \sum_{-\infty}^{\infty} r^{|n|} e^{in\theta}.$$

(Hint: $1/(1-z) = 1 + z + z^2 + \dots$, if $0 \leq |z| < 1$.)

20. Let $U(re^{i\theta})$ be the harmonic function on the disc $\{z: |z| < 1\}$ with $\lim_{r \rightarrow 1^-} U(re^{i\theta}) = u(e^{i\theta})$, except at the finitely many points of discontinuity of $u(e^{i\theta})$. Show that

$$U(re^{i\theta}) = \sum_{-\infty}^{\infty} \hat{u}(n) r^{|n|} e^{in\theta}, \quad 0 \leq r < 1, \quad 0 \leq \theta \leq 2\pi.$$

21. Suppose $v(e^{i\theta})$ is another piecewise continuous function on the circle $\{e^{i\theta}: 0 \leq \theta \leq 2\pi\}$ and $V(re^{i\theta})$ is its harmonic extension to the disc $\{z: |z| < 1\}$. Show that for each r , $0 < r < 1$,

$$\frac{1}{2\pi} \int_0^{2\pi} U(re^{i\theta}) \overline{V(re^{i\theta})} d\theta = \sum_{-\infty}^{\infty} r^{2n} \hat{u}(n) \overline{\hat{v}(n)}. \quad (12)$$

(Hint: By Exercise 18, with $U(re^{i\theta})$ in place of $u(e^{i\theta})$, we know that the n th Fourier coefficient of $U(re^{i\theta})$ is $r^{|n|} \hat{u}(n)$.)

22. Let r increase to 1 in (12); conclude that

$$\frac{1}{2\pi} \int_0^{2\pi} u(e^{i\theta}) \overline{v(e^{i\theta})} d\theta = \sum_{-\infty}^{\infty} \hat{u}(n) \overline{\hat{v}(n)}. \quad (13)$$

(This is a correct formula, but this passage from (12) to (13) needs more mathematical justification than just letting $r \rightarrow 1$.)

23. In Exercise 22, take $v = u$ and conclude that

$$\frac{1}{2\pi} \int_0^{2\pi} |u(e^{i\theta})|^2 d\theta = \sum_{-\infty}^{\infty} |\hat{u}(n)|^2. \quad (14)$$

This equation is known as **Parseval's equality**.

The Reflection Principle

24. Suppose that g is analytic in a domain D in the right half-plane, $\operatorname{Re} z > 0$, which includes a segment $(i\alpha, i\beta)$ of the imaginary axis in its boundary. Show that if g is real-valued on this segment, then g extends to be analytic in the domain formed by the union of D , the reflection of D over the imaginary axis ($=\{-\bar{z}: z \in D\}$), and the interval $(i\alpha, i\beta)$.
25. Suppose that f is an entire function that is real-valued on some segment of the real axis and also real-valued on some segment of the imaginary axis. Show that $f(-z) = f(z)$ for all z . (Hint: Reflect f over the imaginary axis.)
26. Use the reflection principle to prove this result. Let D be a domain in the upper

half-plane with the segment (a, b) of the real axis in its boundary. Suppose that f is analytic on D , continuous on $D \cup (a, b)$, and $f(z) = 0$ if z is in (a, b) . Then $f(z) = 0$ for all $z \in D$.

27. Find the form of a rational function M that is real on the circle $\{z: |z| = 1\}$; assume that M has no poles on this circle.
28. Show that an entire function has only real coefficients in its power series expansion about the origin if and only if it is real-valued on the real axis.
29. Suppose that f is analytic on the open disc $\{z: |z| < 1\}$ and continuous on the closed disc $\{z: |z| \leq 1\}$. Show that if $|f(z)| = 1$ whenever $|z| = 1$, then f can be extended by reflection to be analytic, except for finitely many poles, in the whole complex plane by the rule

$$f^*(z) = \frac{1}{f\left(\frac{1}{\bar{z}}\right)}, \quad |z| > 1.$$

Show further that f is a rational function and, in particular, has the form

$$f(z) = \lambda \frac{z - a_1}{1 - \bar{a}_1 z} \cdots \frac{z - a_M}{1 - \bar{a}_M z}, \quad |a_j| < 1, \quad |\lambda| = 1.$$

30. Suppose f is analytic on the disc $\{z: |z| < 1\}$, continuous on the set $\{z: |z| \leq 1\}$, and real-valued on the circle $\{z: |z| = 1\}$. Use the reflection principle to show that f is constant. (Hint: Extend f to the set $\{z: |z| > 1\}$ by $f^*(z) = \overline{f(1/\bar{z})}$)

4.4 Boundary-Value Problems

In a **boundary-value problem**, we are given information about an unknown function on the boundary of a domain and an equation (usually a differential equation) that the function satisfies on the domain; we are challenged to find the function itself on the domain. In complex variables, the boundary-value problems typically have one of two different forms. In the first, we are to find a harmonic function u on the domain D from the knowledge of the function itself on the boundary of the domain; this is the **Dirichlet*** problem for D . In the second, we are again to find a harmonic function on D , but this time we know only the normal derivative of the function on the boundary of D ; this is the **Neumann†** problem for D . Of course, there is also a mixture of these two cases, where we know the function on part of the boundary and its normal derivative elsewhere on the boundary.

There are two basic techniques in solving boundary-value problems. The first is to use an explicit integral representation formula such as the Poisson Integral Formula. This technique was illustrated in Section 3, where we examined the Poisson Integral Formula both on the disc and on the upper half-plane. The second

* Peter Gustav Lejeune Dirichlet, 1805–1859.

† Carl Neumann, 1832–1925.

basic technique is to transform the given region by a conformal mapping to a new region on which the boundary-value problem is easily solved. This solution can then be carried back to the original region by means of the conformal mapping, thus yielding an explicit solution to the problem. Care must be taken to be sure that the boundaries of the two regions correspond to each other in the correct way, but this happens automatically in most cases.

We illustrate this second technique of conformal mapping by several examples.




Figure 4.30

Example 1 Find the electrostatic potential between two long, parallel, charged, hollow, circular cylinders if one is inside the other.

Solution A cross section of the cylinders perpendicular to their axes gives two circles, one inside the other. By a simple translation, rotation, and change of scale we may assume that the outer circle is the circle $\Gamma_0 = \{z: |z| = 1\}$ and the inner circle is the circle $\Gamma_1 = \{z: |z - x_0| = r_0\}$, where $0 \leq x_0 < 1$ and $x_0 + r_0 < 1$ (Fig. 4.30a). Let the charge on the inner and outer cylinders be C_1 and C_0 , respectively. Thus, we are searching for a harmonic function on the region between the two circles Γ_0 and Γ_1 , which has the value C_0 on the outer circle Γ_0 and the value C_1 on the inner circle Γ_1 . If the smaller circle is centered at the origin (that is, if the two cylinders are coaxial), then the solution is immediate*; namely,

$$U(re^{i\theta}) = C_0 \frac{\log r_0 - \log r}{\log r_0} + C_1 \frac{\log r}{\log r_0}, \quad r_0 \leq r \leq 1,$$

because we know that $w(re^{i\theta}) = \log r$ is a harmonic function (see Exercise 3 of

* The domain and the boundary data are circularly symmetric, so the solution $U(re^{i\theta})$ should be as well. However, the only harmonic function that is independent of θ is $A + B \log r$ (see Exercise 5(b) in Section 1).

Section 4.1). However, it may not be the case that the circles are concentric, so we must first transform the region between Γ_0 and Γ_1 into the region between concentric circles. This can be accomplished by use of an appropriate linear fractional transformation. A computation (or see Exercise 16, Section 5, Chapter 3) shows that there is a real number b such that the function

$$\phi(z) = \frac{z - b}{1 - bz}$$

maps the inner circle Γ_1 onto a circle of radius r_1 centered at the origin; of course, ϕ also maps the outer circle Γ_0 onto itself. Consequently, from the solution to the problem in the coaxial case, we obtain the desired solution in this case:

$$u(z) = C_0 \frac{\log r_1 - \log |\phi(z)|}{\log r_1} + C_1 \frac{\log |\phi(z)|}{\log r_1}. \quad \square$$

Example 2 A magnetic field is established by a north and south pole, each of large rectangular cross section (Fig. 4.31a). Find the magnetic field in the surrounding space.




Figure 4.31

Solution We arbitrarily assign the value -1 to the field on the boundary of the north pole and $+1$ to the field on the boundary of the south pole. Our task is then to find a harmonic function in the region D outside the poles with these boundary values. (In practice, of course, the poles do not have infinite extent; this model assumes that the gap between the poles is small in comparison to the width of the poles. The model is accurate for the magnetic field in the region and close to the gap, where it is of the most interest in any event.)

In Section 5, Chapter 3 we found the Schwarz–Christoffel transformation that maps the upper half-plane U onto the domain D :

$$\phi(z) = \frac{2a}{\pi} \left\{ \sqrt{z^2 - 1} + \arcsin \left(\frac{1}{z} \right) \right\};$$

ϕ carries the ray $(-\infty, 0)$ onto the edge of the north pole and the ray $(0, \infty)$ to the edge of the south pole (Fig. 4.31b). The function $1 - (2/\pi) \operatorname{Arg} z = U(z)$ is harmonic on the upper half-plane, and its boundary values are 1 on $(0, \infty)$ and -1 on $(-\infty, 0)$. Therefore, the function

$$u(w) = 1 - \frac{2}{\pi} \operatorname{Arg}(\phi^{-1}(w))$$

is the desired function on D . The curves of magnetic flux are the images under ϕ of the level lines of V , a harmonic conjugate of U : $V(z) = (2/\pi) \log|z|$. These level lines are circles centered at the origin, so the curves of magnetic flux of the field are the curves

$$\Gamma_r = \frac{2a}{\pi} \left\{ \sqrt{r^2 e^{2it} - 1} + \arcsin\left(\frac{e^{-it}}{r}\right) : 0 \leq t \leq \pi \right\}$$

for $0 < r < \infty$. Some of these curves are shown in Figure 4.32. \square




Figure 4.32 The curves of magnetic flux and equipotentials.

The Neumann Problem

In this boundary-value problem, we are given a function g on the boundary of a domain D , and we are to find a harmonic function u on D with normal derivative equal to g on the boundary of D : $\partial u / \partial n = g$ on ∂D . This is physically realizable as specifying the flux over the boundary of the region of a field or the permeability of the boundary for diffusion. This problem, like the Dirichlet problem, can be explicitly

solved in the case of a disc or a half-plane. We work out the case for the disc $|z| < 1$ here and leave the other setting (which is handled similarly) for the exercises.

Let g be the value of the normal derivative at the unit circle $|z| = 1$. We seek a function $v(re^{i\theta})$ that is harmonic on the disc $\{re^{i\theta}: 0 \leq r < 1\}$ and satisfies

$$\frac{\partial v}{\partial r}(e^{it}) = g(t), \quad 0 \leq t \leq 2\pi.$$

Consider the function

$$v(re^{i\theta}) = \frac{-1}{2\pi} \int_0^{2\pi} g(t) \operatorname{Log}(1 - 2r \cos(\theta - t) + r^2) dt.$$

This function is harmonic, since it is the real part of the analytic function

$$\frac{-1}{\pi} \int_0^{2\pi} g(t) \operatorname{Log}(1 - ze^{-it}) dt, \quad z = re^{i\theta},$$

assuming $g(t)$ is real. Furthermore,

$$\begin{aligned} \frac{\partial v}{\partial r}(se^{i\theta}) &= -\frac{1}{2\pi} \int_0^{2\pi} g(t) \frac{\partial}{\partial r} \operatorname{Log}(1 - 2r \cos(\theta - t) + r^2) \Big|_{r=s} dt \\ &= \frac{1}{2\pi} \int_0^{2\pi} g(t) \frac{2 \cos(\theta - t) - 2s}{1 - 2s \cos(\theta - t) + s^2} dt \\ &= \frac{1}{2\pi s} \int_0^{2\pi} g(t) \left[\frac{1 - s^2}{1 - 2s \cos(\theta - t) + s^2} - 1 \right] dt \\ &= \frac{1}{2\pi s} \int_0^{2\pi} g(t) P_s(\theta - t) dt - \frac{1}{2\pi s} \int_0^{2\pi} g(t) dt, \end{aligned}$$

where $P_s(\theta - t)$ is the Poisson kernel. However, we know from Section 3 and the properties of the Poisson kernel that

$$\lim_{s \rightarrow 1} \frac{1}{2\pi} \int_0^{2\pi} g(t) P_s(\theta - t) dt = g(\theta).$$

Hence,

$$\lim_{s \rightarrow 1} \frac{\partial v}{\partial r}(se^{i\theta}) = g(\theta) - \frac{1}{2\pi} \int_0^{2\pi} g(t) dt.$$

However, the integral of g must vanish, since there are no sources or sinks within the disc $\{z: |z| < 1\}$. Therefore, v is the desired solution. (Another way to see that

$\int_0^{2\pi} g(t) dt = 0$ is to use Green's Theorem:

$$\int_0^{2\pi} g(t) dt = \int_{\Gamma} \frac{\partial v}{\partial n} ds = \iint_{\Omega} \Delta v \, dx \, dy = 0, \quad \text{since } \Delta v \equiv 0.$$

Example 3 A long cylinder with a circular cross section (for example, a blood vessel) contains a fluid with a certain chemical A dissolved within it. The walls of the cylinder allow the chemical A to permeate through with a permeability that depends only on the angular location; moreover, there is a “line” source of the chemical A within the cylinder of strength exactly sufficient to keep the whole system in steady state. If the permeability function is given, find the strength of the source and the lines of flow of the chemical within a cross section of the cylinder (Fig. 4.33).




Figure 4.33

Solution By taking a cross section, we work within a circle, which we can take to be the circle $x^2 + y^2 = 1$. The total flux of the chemical A through the circle is

$$Q = \int_0^{2\pi} g(t) dt,$$

where $g(t)$ is the coefficient of permeability at the point e^{it} , $0 \leq t \leq 2\pi$. Since the system is in steady state, Q is also the strength of the source. If the source is located at the point z_0 , it is given by

$$u_1(z) = \frac{Q}{2\pi} \log \left| \frac{1 - \bar{z}_0 z}{z - z_0} \right|, \quad |z| < 1.$$

We wish, then, to find a function $u(z)$ that is harmonic on the disc $|z| < 1$ except at the point z_0 and that satisfies

$$\frac{\partial u}{\partial r}(e^{it}) = g(t).$$

The function $u_1(z)$ satisfies

$$-\frac{\partial u_1}{\partial r}(e^{it}) = \frac{Q}{2\pi} \left(\frac{1 - r_0^2}{1 - 2r_0 \cos(\theta_0 - t) + r_0^2} \right), \quad z_0 = r_0 e^{i\theta_0}.$$

(See Exercise 13 at the end of this section.) Furthermore, the function defined by

$$u_2(re^{i\theta}) = -\frac{1}{2\pi} \int_0^{2\pi} \left\{ g(t) - \frac{\partial u_1}{\partial r}(e^{it}) \right\} \operatorname{Log}(1 - 2r \cos(\theta - t) + r^2) dt$$

is harmonic on the disc $|z| < 1$ and satisfies

$$\frac{\partial u_2}{\partial r}(e^{it}) = g(t) - \frac{\partial u_1}{\partial r}(e^{it}).$$

Consequently, $u(z) = u_1(z) + u_2(z)$ is the desired solution to the problem. In particular, in the very special case when the permeability is constant and the source is at the origin, the solution is

$$u(z) = -\frac{Q}{2\pi} \operatorname{Log}|z|. \quad \square$$

Example 4 A thin layer of a chemical solution occupies the lower half-plane $\{x + iy: y < 0\}$. The segments $(-\infty, -1]$ and $[1, \infty)$ are insulated so that there is no diffusion across these portions of the boundary, but across the segment $(-1, 1)$ there is a semiporous membrane that allows diffusion at a constant rate into the upper half-plane. Find the steady-state concentration of the substance in the upper half-plane, and the lines of flow of the chemical.

Solution The concentration function $c(x, y)$ for which we are searching must satisfy $\lim\{c(z): z \rightarrow \infty, y > 0\} = 0$, and

$$\frac{\partial c}{\partial y}(x, 0) = \begin{cases} 0, & \text{if } |x| > 1 \\ c_0, & \text{if } |x| < 1. \end{cases} \quad (1)$$

As a first guess, we try

$$b(x, y) = \frac{c_0}{2\pi} \int_{-1}^1 \log[(x - t)^2 + y^2] dt.$$

The function b has the correct normal derivative on the real axis, since

$$\frac{\partial b}{\partial y} = \frac{c_0}{\pi} \int_{-1}^1 \frac{y}{(x - t)^2 + y^2} dt,$$

and by the Poisson Integral Formula for the upper-half-plane, we know that

$$\lim_{y \rightarrow 0} \frac{\partial b}{\partial y}(x, y) = \begin{cases} c_0, & |x| < 1 \\ 0, & |x| > 1 \end{cases}$$

(also see Exercise 1 in this section). However, this function is not the solution we wish, because it is clearly unbounded for large values of $x^2 + y^2$. Instead, we consider

$$c(x, y) = \frac{c_0}{2\pi} \left[\int_{-1}^1 \log[(x-t)^2 + y^2] dt - 2 \log(x^2 + y^2) \right].$$

This function is still harmonic for $y > 0$ and certainly satisfies (1). Furthermore, if $x^2 + y^2$ is large, $c(x, y)$ is virtually zero. Thus, $c(x, y)$ given above is the desired solution. \square

Example 5 Groundwater flows out of an underground aquifer bounded on the bottom by a layer of impervious rock and on the top by a semi-infinite layer of impervious rock; the remainder of the top is fully permeable (Fig. 4.34). Find the flow and its streamlines if the discharge per unit width of the aquifer is q .




Figure 4.34

Solution By the introduction of a coordinate system with the x -axis along the bottom face of the aquifer and the y -axis passing through the left end of the top face, we have the situation shown in Figure 4.35a. The flow has a source “at ∞ ” (that is, far to the right) of total strength $Q = qh$. We now consider the domain D , which is the horizontal strip $\{x + iy : -\infty < x < \infty, 0 < y < h\}$. See Figure 4.35b. On D we seek a harmonic function $v(x, y)$ (the potential) with these boundary conditions:

$$\begin{aligned} \frac{\partial v}{\partial n}(x, y) &= 0 & \text{if} & \begin{cases} y = 0 \text{ and } -\infty < x < \infty \\ y = h \text{ and } 0 < x < \infty \end{cases} \\ v(x, y) &= 0 & \text{if} & y = h \text{ and } -\infty < x < 0. \end{aligned}$$

The first of these conditions says that the boundary is impermeable, and the second




Figure 4.35

says that the groundwater is free to go out of the aquifer from that portion of the top that is not blocked by the impervious rock. The flow must, of course, have a source of strength Q far to the right. Therefore, we are faced with a mixed Dirichlet–Neumann problem; we solve it by finding a conformal mapping of the strip D onto a domain in which the corresponding problem is solved by inspection.

The function

$$\zeta = \frac{-1 + \cos w}{2}$$

maps the vertical half-strip $S = \{w: \operatorname{Im} w > 0 \text{ and } -\pi < \operatorname{Re} w < 0\}$ onto the upper half-plane, $\operatorname{Im} \zeta > 0$. In turn, the function $z = (h/\pi) \operatorname{Log} \zeta$ maps the upper half-plane onto D . Thus,

$$z = f(w) = \frac{h}{\pi} \operatorname{Log} \left(\frac{-1 + \cos w}{2} \right)$$

is a one-to-one analytic mapping of the strip S onto D . Furthermore, the mapping has been carefully arranged so that the left vertical edge of the strip S is mapped to the segment $\{x + ih: 0 < x < \infty\}$, the right vertical edge of S is sent to the real axis, and the horizontal bottom edge of S is sent to the segment $\{x + ih: -\infty < x < 0\}$.

On S , the function $(Q/\pi) \operatorname{Im} w$ has all the required properties: it is zero on the bottom edge; its normal derivative (that is, its partial derivative with respect to $\operatorname{Re} w$) is zero on both the vertical edges; and the flow given by the gradient of $(Q/\pi) \operatorname{Im} w$ has a source at ∞ at strength Q . Since

$$w = \arccos \left[1 + 2 \exp \left(\frac{z\pi}{h} \right) \right],$$

we see that

$$v(x, y) = \frac{Q}{\pi} \operatorname{Im} \left\{ \arccos \left[1 + 2 \exp \left(\frac{z\pi}{h} \right) \right] \right\}$$

is the required potential. The complex potential is, then,

$$G(z) = -i \frac{Q}{\pi} \arccos \left[1 + 2 \exp \left(\frac{z\pi}{h} \right) \right]. \quad \square$$

EXERCISES FOR SECTION 4.4

1. Derive this formula for a solution to the Neumann problem in the upper half-plane $y > 0$:

$$u(x, y) = \frac{1}{2\pi} \int_{-\infty}^{\infty} g(t) \log[(x-t)^2 + y^2] dt + c,$$

where we assume that $|g(t)| \leq Ct^{-v}$ for all large t and for some $v > 1$, to make the integral converge.

2. Suppose that $g(-t) = -g(t)$ for all t ; show that

$$u(x, y) = \frac{1}{2\pi} \int_0^{\infty} g(t) \log \frac{(t-x)^2 + y^2}{(t+x)^2 + y^2} dt + c$$

is a harmonic function in the upper half-plane $y > 0$ with $u(0, y) = 0$ and $(\partial u / \partial y) = g$ on the real line.

3. Suppose that $g(t) = 0$ if $|t| > M$; let

$$u(x, y) = \frac{1}{2\pi} \int_{-M}^M g(t) \log[(x-t)^2 + y^2] dt - A \log(x^2 + y^2),$$

where $A = (1/2\pi) \int_{-M}^M g(t) dt$. Show that $u(x, y)$ is harmonic in the upper half-plane, $u(x, y) \rightarrow 0$ as $x^2 + y^2 \rightarrow \infty$ and

$$\lim_{\delta \downarrow 0} \frac{\partial u}{\partial y}(x, \delta) = g(x), \quad -\infty < x < \infty.$$

4. Show that the function

$$u(x, y) = \frac{1}{\pi} \int_0^{\infty} g(t) \left\{ \frac{y}{(x-t)^2 + y^2} + \frac{y}{(t+x)^2 + y^2} \right\} dt$$

is harmonic in the first quadrant and satisfies

$$u(x, 0) = g(x), \quad 0 < x < \infty$$

$$\frac{\partial u}{\partial x}(0, y) = 0.$$

5. Modify the integral formula in Exercise 4 to find an integral formula for a function $u(x, y)$ harmonic in the first quadrant $x > 0, y > 0$, which satisfies

$$u(0, y) = f(y)$$

$$\frac{\partial u}{\partial y}(x, 0) = 0.$$

6. Find the temperature distribution $T(x, y)$ in the upper half-plane if $T(x, 0) = T_1$ for $x < -\sigma$, $T(x, 0) = T_2$ for $x > \sigma$, and the segment $-\sigma < x < \sigma$ is insulated (that is, $(\partial T/\partial y)(x, 0) = 0$ for $-\sigma < x < \sigma$).
 7. Find the temperature distribution in the first quadrant if $T(0, y) = T_1, 0 < y < \infty$, $T(x, 0) = T_2$ for $\sigma < x < \infty$, and the segment $(0, \sigma)$ on the real axis is insulated.
 8. Find the temperature distribution in an infinite straight rod if the boundary temperatures are as shown in Figure 4.36. (**Hint:** Assume that the width is π and use $w = e^z$ to map the strip onto the upper half-plane.)




Figure 4.36

9. Find the electrostatic potential between two long parallel cylinders of circular cross section if one is inside the other and they touch along one line. (The line of contact is assumed to be insulated.)
 10. Find the electrostatic potential in the region exterior to two cylinders (neither inside the other). (See Fig. 4.37 for a cross section.) (**Hint:** Invert over a point inside one of the cylinders; you may also need to use Exercise 16, Section 5, Chapter 3.)




Figure 4.37

11. Find the temperature distribution inside the half-disc $\{x^2 + y^2 < 1, y > 0\}$, if the temperature is T_1 along the bottom and T_2 along the semicircle. (Hint: Use a conformal mapping onto the first quadrant.)
12. Find the temperature distribution in the lens-shaped region within both the circles $(x - 1)^2 + y^2 = 1$ and $x^2 + (y + 1)^2 = 1$ if one edge of the lens is held at temperature T_1 and the other at temperature T_2 .
13. Let $|a| < 1$; show that the function

$$g(z) = \log \left| \frac{1 - \bar{a}z}{z - a} \right|$$

satisfies

$$-\frac{\partial g}{\partial r}(e^{it}) = \frac{1 - |a|^2}{1 - 2|a|\cos t + |a|^2}.$$

14. Justify the use of conformal mapping to solve the Neumann problem by showing that a Neumann condition on a segment of the boundary of one domain is transformed to a Neumann condition on the corresponding segment in the boundary of the second domain—provided that the mapping is actually conformal, not only inside the domain, but on these segments in the boundaries as well.
15. Let D be the region obtained from the strip $\{x + iy: 0 < y < \pi\}$ by deleting the half-line $\{x + (i\pi/2): 0 \leq x < \infty\}$ (Fig. 4.38). Exercise 14 in Section 5, Chapter 3 gives the Schwarz–Christoffel transformation of the upper half-plane U onto D . Use this to determine the potential in D if the top and bottom surfaces are held at potential zero and the middle (semi-infinite) surface is held at potential $V_0 \neq 0$.




Figure 4.38

4.5 Impulse Functions and the Green's Function of a Domain

It is often very useful in the analysis of physical systems to imagine an “impulse function” with these two properties: (1) The impulse function is zero everywhere except at one point, where it has the value ∞ , and (2) the total area (or volume) under the graph of the function is 1. Such “functions” represent an idealization of an impact of short duration or a short, strong, driving pulse, for instance.* We can

* In mechanics, this type of function is also referred to as a “concentrated force.” In electrostatics, a unit point charge is like an impulse function.

understand an impulse function (say, on a domain D in the plane) by the following limit argument. Fix a point p in D , the point where the impulse will be felt, and let r be a small positive number. Set

$$h_r(z) = \begin{cases} 0, & \text{if } |z - p| > r, z \in D \\ (\pi r^2)^{-1}, & \text{if } |z - p| \leq r. \end{cases}$$

Then the total volume under the graph of h_r is 1, and for each continuous function f on D ,

$$\lim_{r \rightarrow 0} \iint_D f(z) h_r(z) dx dy = f(p). \quad (1)$$

Equation (1) is valid by the following argument. Since the volume under the graph of h_r is 1,

$$\iint_D f(z) h_r(z) dx dy - f(p) = \frac{1}{\pi r^2} \iint_{|z-p| \leq r} \{f(z) - f(p)\} dx dy.$$

The function f is continuous at p , so if $\varepsilon > 0$ is given, then r can be chosen with

$$|f(z) - f(p)| < \varepsilon \quad \text{whenever } |z - p| < r.$$

Hence,

$$\left| \iint_D f(z) h_r(z) dx dy - f(p) \right| \leq \varepsilon \quad \text{if } r \text{ is small,}$$

which is what we wished to show. Thus, in some sense, the impulse function δ_p equals $\lim_{r \rightarrow 0} h_r$.

In complex variables (particularly in the applications of complex variables), we are interested in harmonic functions u —that is, in solutions of the equation $\Delta u = 0$. Often, as well, the values of u are prescribed on the boundary of the domain D . Let us consider here the function G , which is the solution of the boundary-value problem

$$\begin{cases} \Delta G = \delta_p & \text{on } D \\ G = 0 & \text{on the boundary of } D, \end{cases}$$

where δ_p is the impulse function for $p \in D$. Thus, G is harmonic on the open set $D \setminus \{p\}$ and vanishes on the boundary of D .^{*} Green's Formula (formula (9) of Section 6, Chapter 1) gives us the relation

* That is, G is the **response** to the impulse at p for a system governed by the Laplacian, with zero boundary data.

$$\int_{\Gamma} \left(\frac{\partial G}{\partial n} f - G \frac{\partial f}{\partial n} \right) ds = \iint_D \{(\Delta G)f - G(\Delta f)\} dx dy,$$

where Γ is the boundary of D . If we assume that f is continuous on $D \cup \Gamma$ and harmonic on D , we find that

$$\int_{\Gamma} f \frac{\partial G}{\partial n} ds = \iint_D (\Delta G)f dx dy = \iint_D \delta_p f dx dy = f(p).$$

Consequently, the normal derivative of G is precisely the function that takes the boundary values f and produces the value at p of the harmonic function with these boundary values; that is, $\partial G/\partial n$ solves the Dirichlet problem for the domain D . Of course, the derivation carried out above is only heuristic, but it can be made mathematically sound (see the exercises).

We now give the “official” definition of the Green’s function and then some examples that illustrate the Green’s function and its use.

The Definition of the Green’s Function of a Domain

Suppose that D is a bounded domain whose boundary is a finite number of disjoint smooth simple closed curves (Fig. 4.39). Let p be a point in D . The **Green’s function for D with singularity at p** is the function $G(z; p)$ with these three properties:




Figure 4.39 A domain bounded by three disjoint, piecewise smooth simple closed curves.

- (a) $G(z; p)$ is harmonic for z in D , $z \neq p$,
- (b) $G(z; p) + \log|z - p|$ is harmonic near p , (2)
- (c) $G(z; p) = 0$ if z is in the boundary of D .

Such a function $G(z; p)$ can be found in the following way. Let v be the harmonic function on D with $v(z) = \log|z - p|$ if z is in the boundary of D ; then set

$$G(z; p) = v(z) - \log|z - p|.$$

Exercise 1 at the end of this section shows that $G(z; p)$ is unique.

The Green's function for an unbounded domain is defined exactly as in (a), (b), and (c) above if p is not the point at ∞ . If ∞ is a point in the domain D , then the Green's function for D with singularity at ∞ is the function $G(z; \infty)$ satisfying

- (a)' $G(z; \infty)$ is harmonic on $D \setminus \{\infty\}$,
- (b)' $G(z; \infty) - \log|z|$ is harmonic for $|z|$ large, (3)
- (c)' $G(z; \infty) = 0$ if z is in the boundary of D .

Domains that include the point at ∞ , and their Green's functions, are important in the subject of potential theory.

Example 1 Let D be the disc $\{z: |z| < 1\}$ and fix $p \in D$. Set

$$G(z; p) = \log \left| \frac{1 - \bar{p}z}{z - p} \right|, \quad |z| < 1.$$

Then G is harmonic except at p , $G = 0$ on the boundary of D , since if $|z| = 1$, then

$$\left| \frac{1 - \bar{p}z}{z - p} \right| = 1.$$

On the disc D ,

$$G(z; p) + \log|z - p| = \log|1 - \bar{p}z|.$$

Note that $\log|1 - \bar{p}z|$ is harmonic on D . Furthermore, the (inward) normal derivative of G at the boundary is just minus the partial derivative of G with respect to r , which is

$$-\frac{\partial}{\partial r} \left\{ \log \left| \frac{1 - \bar{p}re^{i\theta}}{re^{i\theta} - p} \right| \right\} \Big|_{r=1} = \frac{1 - |p|^2}{|e^{i\theta} - p|^2}.$$

You should recognize this function immediately as the Poisson kernel for the point p , $|p| < 1$. \square

Example 2 Let D be a simply-connected domain, bounded or unbounded, with at least two points in its boundary, and fix a point $p \in D$. Let ϕ be the Riemann mapping of D onto the disc $\Delta = \{w: |w| < 1\}$ with $\phi(p) = 0$, $\phi'(p) > 0$; such a mapping exists and is unique, by Theorem 1, Section 5, Chapter 3. Show that

$$G(z; p) = -\log|\phi(z)|, \quad z \in D$$

is the Green's function for D with singularity at p .

Solution Let us suppose first that p is not the point at ∞ (if this point is even in D). On $D \setminus \{p\}$, the function ϕ is analytic and nonzero, so $\log|\phi|$ is harmonic, by Exercise 2 of Section 4.1. Furthermore, ϕ has a zero of order one at p , so

$$\phi(z) = (z - p)g(z),$$

where $g(z)$ is never zero on D . Thus,

$$\log|\phi(z)| = \log|z - p| + \log|g(z)|, \quad z \in D,$$

and $\log|g|$ is harmonic on all of D . This shows that $\log|\phi|$ satisfies (a) and (b) of (2); to see that $\log|\phi|$ satisfies (c), we need to use this fact:

if $\{z_n\}$ is a sequence of points in D with $\lim_{n \rightarrow \infty} z_n = \lambda$

for some λ in the boundary of D , then $\lim_{n \rightarrow \infty} |\phi(z_n)| = 1$.

(See Exercise 19, Section 5, Chapter 3.) From this, it is evident that (c) holds, so $-\log|\phi(z)|$ is, in fact, the Green's function for D with singularity at p .

The case when p is the point at ∞ is left for the exercises, but it is done in much the same way. \square

Example 3 Let D be the complement of the segment $[a, b]$, including the point at ∞ . Find the Green's function for D with singularity at ∞ and evaluate

$$\kappa = \lim_{|z| \rightarrow \infty} \{G(z; \infty) - \log|z|\}.$$

Solution Let

$$\phi_1(z) = \frac{2z}{b-a} + \frac{a+b}{a-b}.$$

$\phi_1(z)$ is a one-to-one analytic function that maps the segment $[a, b]$ onto $[-1, 1]$ and D onto the domain D_1 that is the complement of $[-1, 1]$. Let $\phi_2(w) = w - \sqrt{w^2 - 1}$; $\phi_2(w)$ is an analytic function that maps the domain D_1 one-to-one onto the disc $\Delta = \{\zeta : |\zeta| < 1\}$; ϕ_2 is the inverse of the function $\psi(\zeta) = \frac{1}{2}(\zeta + \zeta^{-1})$, which maps the unit disc Δ onto D_1 (see Example 5, Section 4, Chapter 3). The composition $\phi(z) = \phi_2(\phi_1(z))$ is the Riemann mapping of D onto Δ with $\phi(\infty) = 0$. According to Example 2, the Green's function for D with singularity at ∞ is

$$\begin{aligned} G(z; \infty) &= -\log|\phi(z)| \\ &= -\log|(Az + B) - \sqrt{(Az + B)^2 - 1}|, \end{aligned}$$

where $A = 2/(b - a)$ and $B = (a + b)/(a - b)$. Next, for $|w|$ large,

$$\sqrt{w^2 - 1} = w\sqrt{1 - w^{-2}} = w \left\{ 1 - \frac{1}{2w^2} - \frac{c_2}{w^4} - \frac{c_3}{w^6} \dots \right\},$$

where c_2, c_3, \dots are constants (see Exercises 25 to 27 of Section 2, Chapter 2). Hence,

$$w - \sqrt{w^2 - 1} = \frac{1}{2w} + \frac{c_2}{w^3} + \frac{c_3}{w^5} + \dots,$$

and consequently,

$$\begin{aligned} -\log|\phi_2(w)| &= -\log \left| \frac{1}{2w} + \frac{c_2}{w^3} + \dots \right| \\ &= \log|2w| - \log \left| 1 + \frac{2c_2}{w^2} + \dots \right|. \end{aligned}$$

Replacing w with $Az + B$ gives

$$G(z; \infty) - \log|z| = \log \left| 2 \left(A + \frac{B}{z} \right) \right| - \log \left| 1 + \frac{2c_2}{(Az + B)^2} + \dots \right|,$$

so finally,

$$\begin{aligned} \kappa &= \lim_{|z| \rightarrow \infty} \{G(z; \infty) - \log|z|\} = \log(2A) \\ &= \log \left(\frac{4}{b - a} \right). \end{aligned}$$
□

EXERCISES FOR SECTION 4.5

- Let $p \in D$. Show that the Green's function of the domain D with singularity at p is unique. (Hint: If both $G(z; p)$ and $G_1(z; p)$ satisfy the definition of the Green's function, show that the difference, $G - G_1$, is harmonic and bounded in D and vanishes on the boundary of D .)
- Let D_1 and D_2 be two domains with Green's functions G_1 and G_2 , respectively. Suppose that ϕ is a one-to-one analytic function mapping D_1 onto D_2 . Show that $G_1(z; p) = G_2(\phi(z); \phi(p))$.

In Exercises 3 to 6, use Exercise 2 or the definition of the Green's function to explicitly find the Green's function for the domain D with singularity at $p \in D$.

3. $D = \{z = x + iy: y > 0\}$
4. $D = \{z: |z - z_0| < R\}, \quad z_0, R \text{ fixed}, R > 0$
5. $D = \{z = x + iy: x > 0 \text{ and } y > 0\}$
6. $D = \{z = x + iy: y > 0 \text{ and } |z| < 1\}$

In Exercises 7 to 9, use Example 2 to find the Green's function for the domains indicated.

7. $D = \{z = x + iy: 0 < y < \pi\}$
8. $D = \{z = x + iy: -\pi/2 < x < \pi/2 \text{ and } y > 0\}$
9. $D = \{z = x + iy: x \notin (-\infty, 0]\}$
10. Show that the Green's function for a domain D with singularity at p is always positive on $D \setminus \{p\}$. (Hint: Show that $G(z; p)$ is positive on the circle $|z - p| = \varepsilon$ for all small ε ; then use the maximum/minimum principle on the domain

$$D \setminus \{z: |z - p| \leq \varepsilon\}.$$

11. Argue on physical grounds that the Green's function is symmetric: $G(q; p) = G(p; q)$ for all $p \neq q$ in D .
12. Let D be a bounded domain whose boundary, Γ , consists of a finite number of disjoint piecewise smooth simple closed curves. It is a fact that the normal derivative of the Green's function exists and is continuous on Γ . Use this to show that

$$\frac{1}{2\pi} \int_{\Gamma} u(z) \frac{\partial}{\partial n} \{G(z; p)\} ds = u(p)$$

for any function u that is continuous on $D \cup \Gamma$ and harmonic on D . (Hint: Apply Green's Formula on the domain $D_\varepsilon = \{z \in D: |z - p| \geq \varepsilon\}$; you'll have to show that the term

$$\frac{1}{2\pi} \int_{|z-p|=\varepsilon} \left\{ \frac{\partial G}{\partial n} u - \frac{\partial u}{\partial n} G \right\} ds$$

approaches $u(p)$ as ε decreases to 0. This is accomplished by noting that

$$\frac{\partial G}{\partial n} = \frac{1}{|z - p|} + \text{bounded term}$$

on this circle. Next, estimate $|G(z; p)|$ by $\log(1/|z - p|)$ and recall that $\varepsilon \log \varepsilon$ goes to zero as $\varepsilon \rightarrow 0$.)

13. Let D be an unbounded domain including the point at ∞ , and let E be the complement of D ; let $G(z; \infty)$ be the Green's function for D with singularity at

∞ . The number $c(E)$ defined by

$$-\log c(E) = \lim_{|z| \rightarrow \infty} \{G(z; \infty) - \log|z|\}$$

is called the **logarithmic capacity** of E . Find the logarithmic capacity of E if (a) $E = \{z: |z| \leq 1\}$; (b) $E = \{z: |z - z_0| \leq r_0\}$; (c) $E = [a, b]$, $a < b$ (see Example 3); (d) $E = \{x + iy: (x^2/a^2) + (y^2/b^2) \leq 1\}$.

14. Complete Example 2 by showing that if D is a simply-connected domain containing the point at ∞ , then

$$G(z; \infty) = -\log|\phi(z)|,$$

where $\phi(z)$ is the Riemann mapping of D onto the disc $\{w: |w| < 1\}$ with $\phi(\infty) = 0$.

Impulse Functions

15. Show that the Poisson kernel

$$P_r(\theta) = \frac{1 - r^2}{1 - 2r \cos \theta + r^2}$$

converges to the impulse function at $\theta = 0$ as $r \rightarrow 1$ in the sense that $\lim_{r \rightarrow 1} (1/2\pi) \int_{-\pi}^{\pi} h(e^{i\theta}) P_r(e^{i\theta}) d\theta = h(1)$ for each function that is continuous on the circle $\{e^{i\theta}: -\pi \leq \theta \leq \pi\}$.

16. Let $[a, b]$ be an interval in the line, and let $a < p < b$. Show that the “impulse function” δ_p for $[a, b]$ has the property that

$$\int_a^b f(x) \delta_p(x) dx = f(p)$$

for every continuous function $f(x)$ on $[a, b]$. (Hint: Let $u_n(x)$ be zero for $|x - p| > 1/(2n)$ and $u_n(x)$ be n for $|x - p| < 1/(2n)$. Show that

$$\lim_{n \rightarrow \infty} \int_a^b f(x) u_n(x) dx = f(p).$$

17. Let $H(x) = \begin{cases} 0, & a \leq x < p \\ 1, & p < x \leq b. \end{cases}$ Use integration by parts to “show” that $H' = \delta_p$. (Hint: If $f(b) = 0$, then

$$\int_a^b f'(x) H(x) dx = - \int_a^b f(x) H'(x) dx.$$

But $\int_a^b f'(x) H(x) dx = \int_p^b f'(x) dx = f(b) - f(p) = -f(p)$.

Further Reading

The following two books present a substantial number of applications of complex variables to engineering and science problems.

Henrici, P. *Applied and computational analysis*. Vol. 1. New York: Wiley, 1974.
Kyrala, A. *Applied functions of a complex variable*. New York: Wiley, 1972.

More specialized books that I have found useful in dealing with the topics in this chapter include the following:

- Chorin, L., and Marsden, J. *A mathematical introduction to fluid mechanics*.
New York: Springer-Verlag, 1979.
- Crank, J. *The mathematics of diffusion*. New York: Oxford University Press,
1956.
- Kellogg, O. D. *Foundations of potential theory*. New York: Dover, 1953.
- Milne-Thompson, L. M. *Theoretical aerodynamics*. 4th ed. New York: Macmillan,
1966.
- Milne-Thompson, L. M. *Theoretical hydrodynamics*. 5th ed. New York: Macmillan,
1968.
- Muskhelishvili, N. I. *Some basic problems of the mathematical theory of elasticity*.
Groningen, Netherlands: P. Noordhoff, 1963.
- Verruijt, A. *Theory of groundwater flow*. New York: Macmillan, 1970.

5

Transform Methods

5.1 The Fourier* Transform: Basic Properties

The Fourier transform is a powerful tool in both applied and theoretical mathematics. Its utility is due in no small part to the simple relationships between a function and its Fourier transform. However, a careful mathematical justification of these relationships is not elementary. For that reason, we postpone our discussion of these relationships until Section 2. In this section, we define the Fourier transform, demonstrate some of its basic properties, and show some of its applications. The relationships between the function and its transform are stated in Section 2, and several examples illustrate their uses. At the end of Section 2, we sketch a portion of the mathematics establishing these relationships.

We begin with the notion of a piecewise smooth function on the real line. Suppose t_0 is a real number and u is a function defined for t near t_0 ; then we say that u has a **limit at t_0 from the right** if

$$\lim_{h \downarrow 0} u(t_0 + h) = u^+(t_0), \quad h > 0$$

exists; we say that u has a **limit at t_0 from the left** if

$$\lim_{h \downarrow 0} u(t_0 - h) = u^-(t_0), \quad h > 0$$

exists. We do not make any requirement about the possible equality of $u^+(t_0)$ and $u^-(t_0)$. If it happens that these two limits exist and are both equal to $u(t_0)$, then (and only then) u is continuous at t_0 .

Example 1 The function

$$u(t) = \begin{cases} 1, & |t| < \sigma \\ 0, & |t| > \sigma \end{cases}$$

* Jean Baptiste Joseph Fourier, 1768–1830.

has both left and right limits at all points t_0 . The left and right limits coincide except at $t_0 = \pm\sigma$; at $t_0 = -\sigma$, the limit from the left is zero and the limit from the right is 1; at $t_0 = \sigma$, exactly the reverse situation occurs. \square

A function u is **piecewise smooth** if the following conditions are met: there are a finite number of points $t_1 < t_2 < \dots < t_N$ in the real line such that

- both u and u' are continuous on all the intervals $(-\infty, t_1), (t_1, t_2), \dots, (t_N, \infty)$.
- both u and u' have limits from the right *and* left at each of the points t_1, \dots, t_N .

We do not require that the right and left limits of either u or u' coincide at any of the points t_1, \dots, t_N , although this may happen.

Example 2 The function

$$u(t) = \begin{cases} 1, & |t| < \sigma \\ 0, & |t| > \sigma \end{cases}$$

from Example 1 is piecewise smooth. The graph of u is shown in Figure 5.3a. \square

Example 3 The function

$$U(t) = \begin{cases} 0, & -\infty < t < -\sigma \\ t + \sigma, & -\sigma \leq t \leq \sigma \\ 2\sigma, & \sigma < t < \infty \end{cases}$$

is piecewise smooth. U is continuous and differentiable (except at $\pm\sigma$). However, $U' = u$, the function from Example 2, so U' is discontinuous at $\pm\sigma$. \square

Example 4 If u is piecewise smooth and U is any function with $U'(t) = u(t)$ for all $t \neq t_j, j = 1, \dots, N$, then U is piecewise smooth. The elementary demonstration of this is left to the exercises. \square

Example 5 The functions

$$u_1(t) = \begin{cases} \frac{1}{t}, & t > 0 \\ 0, & t < 0 \end{cases}$$

and

$$u_2(t) = \begin{cases} t \log t, & t > 0 \\ 0, & t \leq 0 \end{cases}$$

are not piecewise smooth. u_1 does not have a (finite) limit from the right at $t = 0$. u_2 is continuous on the real line; however, u'_2 fails to have a (finite) limit from the right at $t = 0$. \square

The Fourier Transform

Let u be a piecewise smooth function on the real line, and assume that

$$\int_{-\infty}^{\infty} |u(t)| dt < \infty. \quad (1)$$

Condition (1) will certainly hold if $u(t) = 0$ for $|t| > M$ or if $|u(t)| \leq C/t^2$ for all large t . We define the **Fourier transform** of u by the rule

$$\hat{u}(x) = \int_{-\infty}^{\infty} u(t) e^{-itx} dt, \quad -\infty < x < \infty. \quad (2)$$

The condition (1), together with the fact that $|e^{-itx}| = 1$ if both t and x are real, assures us that the integral defining $\hat{u}(x)$ makes good sense (see Exercises 16 to 19 in Section 2). As we shall see, the Residue Theorem will be an indispensable tool in the computation of Fourier transforms.

Example 6 Find the Fourier transform of the function

$$u(t) = \begin{cases} 1, & |t| < \sigma \\ 0, & |t| > \sigma. \end{cases}$$

Solution By the definition of the Fourier transform,

$$\begin{aligned} \hat{u}(x) &= \int_{-\sigma}^{\sigma} e^{-itx} dt = \frac{-1}{ix} \left[e^{-itx} \right]_{t=-\sigma}^{t=\sigma} \\ &= \frac{2}{x} \left[\frac{e^{i\sigma x} - e^{-i\sigma x}}{2i} \right] \\ &= 2 \frac{\sin \sigma x}{x}. \end{aligned}$$
□

Example 7 Find the Fourier transform of $u(t) = e^{-t^2}$.

Solution The Fourier transform of u is given by

$$\begin{aligned} \hat{u}(x) &= \int_{-\infty}^{\infty} e^{-t^2} e^{-itx} dt = \int_{-\infty}^{\infty} e^{-(t+ix/2)^2} e^{-x^2/4} dt \\ &= \{\exp(-x^2/4)\} \int_{-\infty}^{\infty} \exp\left[-\left(t + \frac{ix}{2}\right)^2\right] dt. \end{aligned}$$

To evaluate this last integral, we make use of some complex analysis. Let R be a large positive number; the integral

$$\int_{-R}^R \exp\left\{-\left(t + \frac{ix}{2}\right)^2\right\} dt$$

converges to

$$\int_{-\infty}^{\infty} \exp\left\{-\left(t + \frac{ix}{2}\right)^2\right\} dt$$

as $R \rightarrow \infty$. However, the integral from $-R$ to R is nothing but the integral

$$\int_{\gamma_R} \exp(-z^2) dz, \quad z = t + \frac{ix}{2}, \quad -R \leq t \leq R,$$

where γ_R is the line segment joining $-R + (ix/2)$ to $R + (ix/2)$. Consider now the rectangle Γ_R with vertices at $-R$, R , $R + (ix/2)$, and $-R + (ix/2)$ (Fig. 5.1). By Cauchy's Theorem, the line integral of $f(z) = \exp(-z^2)$ about Γ_R is zero. Furthermore, on the vertical edges of Γ_R , we may make the estimate

$$|f(z)| = |\exp(-z^2)| \\ = \exp\left[-R^2 + s^2\right], \quad 0 \leq s \leq \frac{x}{2},$$

so the integral of f over the two vertical edges of Γ_R is very small if R is very large. The integral of $f(z)$ as z runs from $-R$ to R cannot be computed exactly, but as $R \rightarrow \infty$ it converges to

$$\int_{-\infty}^{\infty} e^{-t^2} dt = \sqrt{\pi}.$$

(The value of this integral should be familiar from calculus; also see Exercise 20, Section 3, Chapter 2.) Hence, we have shown that

$$\int_{-\infty}^{\infty} \exp(-t^2 - ixt) dt = \sqrt{\pi} \exp\left[-\frac{x^2}{4}\right],$$

so




Figure 5.1

$$\hat{u}(x) = \sqrt{\pi} \exp\left[-\frac{x^2}{4}\right].$$

That is, the Fourier transform of e^{-t^2} is $\sqrt{\pi}e^{-x^2/4}$. \square

Example 8 Find the Fourier transform of

$$u(t) = \frac{1}{1+t^2}.$$

Solution The Fourier transform of u is

$$\hat{u}(x) = \int_{-\infty}^{\infty} \frac{e^{-ixt}}{1+t^2} dt.$$

To evaluate this integral, we use the Residue Theorem. Let

$$f(z) = \frac{e^{-izx}}{1+z^2},$$

and set $z = \sigma + i\tau$. Thus,

$$\begin{aligned} |f(z)| &\leq \frac{e^{x\tau}}{|z|^2 - 1}, \quad |z| \text{ large} \\ &\leq \frac{1}{|z|^2 - 1} \quad \text{if } x\tau \leq 0. \end{aligned}$$

The necessity that $x\tau$ be less than or equal to zero requires us to consider two cases. If x is negative, we integrate f over the contour shown in Figure 5.2a; if x is positive, we integrate f over the contour in Figure 5.2b. We can apply the Residue Theorem




Figure 5.2

(or just use Cauchy's Formula). For the case where $x \leq 0$, the pole is at i , and we find that

$$\int_{\Gamma_R} f(z) dz = 2\pi i \frac{e^{-ix(i)}}{2i} = \pi e^x, \quad x \leq 0.$$

For the case where $x \geq 0$, the pole is at $-i$. As before,

$$\int_{\Gamma_R} f(z) dz = 2\pi i \frac{e^{-ix(-i)}}{-2i} = -\pi e^{-x}, \quad x \geq 0.$$

Since the integral of f over the semicircular part of Γ_R goes to zero as $R \rightarrow \infty$, we find that

$$\hat{u}(x) = \int_{-\infty}^{\infty} \frac{e^{-ixt}}{1+t^2} dt = \begin{cases} \pi e^x, & x \leq 0 \\ \pi e^{-x}, & x \geq 0. \end{cases}$$

Consequently,

$$\hat{u}(x) = \pi e^{-|x|}, \quad x \text{ real.} \quad \square$$

Basic Properties of the Fourier Transform

The operation of taking the Fourier transform of a function has certain convenient and useful properties, which we list here. (In all cases, we assume that (1) holds.)

Property 1: Linearity

If u_1 and u_2 are functions and λ_1 and λ_2 are complex numbers, then

$$(\lambda_1 u_1 + \lambda_2 u_2) \hat{ }(x) = \lambda_1 \hat{u}_1(x) + \lambda_2 \hat{u}_2(x). \quad (3)$$

Property 2: Translation

If a is a real number and v is the function defined by $v(t) = u(a+t)$, $-\infty < t < \infty$, then

$$\hat{v}(x) = e^{iax} \hat{u}(x), \quad -\infty < x < \infty. \quad (4)$$

Property 3: Scaling

If b is a real number, $b \neq 0$, and w is the function defined by $w(t) = u(bt)$, $-\infty < t < \infty$, then

$$\hat{w}(x) = \frac{1}{|b|} \hat{u}\left(\frac{x}{b}\right). \quad (5)$$

Property 4: Transform of the Derivative

If u is differentiable everywhere on the real axis, and if u and u' both satisfy (1), then

$$\widehat{u'}(x) = ix\hat{u}(x), \quad -\infty < x < \infty. \quad (6)$$

Property 5: Derivative of the Transform

$$\frac{d}{dx} \hat{u}(x) = -i(tu(t))^\wedge(x), \quad (7)$$

assuming that

$$\int_{-\infty}^{\infty} |tu(t)| dt < \infty.$$

The demonstrations of (3) through (7) are simple and left for the exercises; (4) and (5) are done by a change of variables and (6) by an integration by parts. (See Exercise 20, Section 2 of this chapter, to conclude that $\lim_{|t| \rightarrow \infty} u(t) = 0$.)

Example 9 Find the Fourier transform of

$$(i) \quad u_1(t) = (t^2 + 4t + 5)^{-1}.$$

$$(ii) \quad u_2(t) = \exp\left[-\frac{t^2}{2}\right].$$

$$(iii) \quad u_3(t) = \begin{cases} 1+t, & -1 \leq t \leq 0 \\ 1-t, & 0 \leq t \leq 1 \\ 0, & |t| \geq 1. \end{cases}$$

Solution (i) u_1 is given by

$$u_1(t) = \frac{1}{(t+2)^2 + 1} = u(t+2),$$

where u is the function in Example 8. Hence, by Property 2,

$$\begin{aligned} \hat{u}_1(x) &= e^{2ix} \hat{u}(x) \\ &= \pi e^{2ix} e^{-|x|}, \quad -\infty < x < \infty. \end{aligned}$$

(ii) Next, $u_2(t)$ is exactly $u(t/\sqrt{2})$, where u is the function from Example 7. Consequently, with $b = 1/\sqrt{2}$, Property 3 gives

$$\begin{aligned} \hat{u}_2(x) &= \sqrt{2} \hat{u}(\sqrt{2}x) \\ &= \sqrt{2\pi} \exp\left[-\frac{2x^2}{4}\right] \\ &= \sqrt{2\pi} u_2(x). \end{aligned}$$

Thus, u_2 is a constant multiple of its Fourier transform.

(iii) Finally, the function $u_3(t)$ has a very simple derivative:

$$u'_3(t) = \begin{cases} 1, & -1 < t < 0 \\ -1, & 0 < t < 1 \\ 0, & |t| > 1. \end{cases}$$

The Fourier transform of u'_3 is easily computed to be

$$\widehat{u'_3}(x) = \frac{2(\cos x - 1)}{ix}.$$

Thus

$$\widehat{u}_3(x) = \frac{1}{ix} \widehat{u'_3}(x) = 2 \frac{1 - \cos x}{x^2}. \quad \square$$

There is one more very important property of the Fourier transform, but we need a definition in order to introduce it. Suppose u and v are two functions satisfying (1); the **convolution of u and v** is the function $u * v$ defined by

$$(u * v)(t) = \int_{-\infty}^{\infty} u(s)v(t-s) ds, \quad -\infty < t < \infty. \quad (8)$$

Example 10 Let

$$u(t) = v(t) = \begin{cases} 1, & |t| < \sigma \\ 0, & |t| > \sigma. \end{cases}$$

Find $u * v$.

Solution The convolution of u and v at t is given by

$$(u * v)(t) = \int_{-\infty}^{\infty} u(s)v(t-s) ds.$$

Now $v(t-s) = 0$ unless $|t-s| < \sigma$; since $u(s) = 1$ only when $|s| < \sigma$, we know that

$$u(s)v(t-s) = \begin{cases} 1, & \text{if } |t-s| < \sigma \text{ and } |s| < \sigma \\ 0, & \text{otherwise.} \end{cases}$$

Thus, $u(s)v(t-s) = 0$ for all s if $|t| > 2\sigma$, so $(u * v)(t) = 0$ if $|t| > 2\sigma$. Moreover, if $0 \leq t < 2\sigma$, then

$$u(s)v(t-s) = \begin{cases} 1, & \text{if } t - \sigma < s < \sigma \\ 0, & \text{otherwise,} \end{cases}$$

and, for $-2\sigma < t \leq 0$,

$$u(s)v(t-s) = \begin{cases} 1, & \text{if } -\sigma < s < t + \sigma \\ 0, & \text{otherwise.} \end{cases}$$

Thus,

$$(u * v)(t) = \begin{cases} 0, & t \leq -2\sigma \\ t + 2\sigma, & -2\sigma \leq t \leq 0 \\ 2\sigma - t, & 0 \leq t \leq 2\sigma \\ 0, & t \geq 2\sigma. \end{cases}$$

A sketch of the graphs of u , v , and $u * v$ is shown in Figure 5.3a and 5.3b. \square




Figure 5.3

Example 11 Let u be a bounded continuous function on $(-\infty, \infty)$, satisfying (1), and, for $y > 0$, let

$$K_y(t) = \frac{1}{\pi} \frac{y}{t^2 + y^2}.$$

Then the function

$$U(x, y) = (u * K_y)(x) = \frac{1}{\pi} \int_{-\infty}^{\infty} u(t) \frac{y}{(x-t)^2 + y^2} dt$$

is exactly the bounded harmonic function of $x + iy$ defined in the upper half-plane $\{x + iy : y > 0\}$, which satisfies

$$\lim_{x+iy \rightarrow s} U(x, y) = u(s), \quad -\infty < s < \infty;$$

see Section 3, Chapter 4. \square

In the exercises, the reader is asked to show that $u * v$ is the same as $v * u$; that is, the order of the convolution is not important. It is also true that convolution “distributes” across addition just as multiplication does:

$$u * (v + w) = u * v + u * w.$$

The critical connection between the Fourier transform and convolution is that the Fourier transform of the convolution of u and v is just the *product* of the Fourier transforms of u and v , respectively. In symbols,

$$(u * v)^{\hat{}}(x) = \hat{u}(x)\hat{v}(x), \quad -\infty < x < \infty. \quad (9)$$

This is not too hard to show (if one takes on faith the reversibility of the order of integration in an iterated integral):

$$\begin{aligned} (u * v)^{\hat{}}(x) &= \int_{-\infty}^{\infty} (u * v)(t)e^{-itx} dt \\ &= \int_{-\infty}^{\infty} \left\{ \int_{-\infty}^{\infty} u(s)v(t-s) ds \right\} e^{-itx} dt \\ &= \int_{-\infty}^{\infty} u(s) \left\{ \int_{-\infty}^{\infty} v(t-s)e^{-i(t-s)x} dt \right\} e^{-isx} ds \\ &= \left(\int_{-\infty}^{\infty} u(s)e^{-isx} ds \right) \hat{v}(x) \\ &= \hat{u}(x)\hat{v}(x). \end{aligned}$$

The Fourier transform has powerful applications in the area of differential equations; we illustrate with two examples. Both examples make use of the fact that if u_1 and u_2 are two functions with $\hat{u}_1 = \hat{u}_2$, then $u_1 = u_2$; see Exercise 12, Section 2.

Example 12 At a particular time, a very long thin rod has a given temperature distribution $f(s)$, $-\infty < s < \infty$. Find the temperature at all future times.

Solution The rod is one-dimensional (it just has length), so the temperature is a function of the position s on the rod at the time t ; we let $u(t, s)$ be the temperature at time t and position s . The function $u(t, s)$ satisfies the **heat equation**

$$\frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial s^2},$$

with **initial condition**

$$u(0, s) = f(s), \quad -\infty < s < \infty.$$

We now use the Fourier transform (on the s variable) to solve this equation. We begin by setting

$$\phi(t, x) = \int_{-\infty}^{\infty} u(t, s) e^{-isx} ds, \quad -\infty < x < \infty.$$

Then

$$\begin{aligned}\frac{\partial \phi}{\partial t} &= \int_{-\infty}^{\infty} \frac{\partial u}{\partial t}(t, s) e^{-isx} ds \\ &= \int_{-\infty}^{\infty} \frac{\partial^2 u}{\partial s^2}(t, s) e^{-isx} ds.\end{aligned}$$

In this second integral, we use (6) (or integrate twice by parts); we also make the natural assumption that $u(t, s)$ and $(\partial u / \partial s)(t, s)$ both tend to zero as $s \rightarrow \pm\infty$ for each t . The net result, then, is that

$$\begin{aligned}\frac{\partial \phi}{\partial t}(t, x) &= -x^2 \int_{-\infty}^{\infty} u(t, s) e^{-isx} ds \\ &= -x^2 \phi(t, x).\end{aligned}$$

Furthermore,

$$\begin{aligned}\phi(0, x) &= \int_{-\infty}^{\infty} u(0, s) e^{-isx} ds \\ &= \int_{-\infty}^{\infty} f(s) e^{-isx} ds \\ &= \hat{f}(x).\end{aligned}$$

Consequently, for each fixed x , $\phi(x, t)$ satisfies the ordinary differential equation

$$\frac{d\phi}{dt} = -x^2 \phi, \quad \phi(0, x) = \hat{f}(x).$$

Hence, we must have

$$\phi(t, x) = \hat{f}(x) \exp[-tx^2].$$

However, we recognize (and this is critical!) that $\exp[-x^2 t]$ is the Fourier transform of

$$K(s) = \frac{1}{2\sqrt{\pi t}} e^{-s^2/4t}, \quad t > 0.$$

Since $\phi(t, x)$ is the Fourier transform of $u(t, s)$, we use the convolution theorem to discover that $u(t, s)$ must be the convolution of f and K :

$$\begin{aligned} u(t, s) &= \int_{-\infty}^{\infty} f(y) \frac{1}{2\sqrt{\pi t}} e^{-(s-y)^2/4t} dy \\ &= \frac{1}{2\sqrt{\pi t}} \int_{-\infty}^{\infty} f(y) e^{-(s-y)^2/4t} dy. \end{aligned}$$

This explicitly displays the dependence of the solution $u(t, s)$ on the initial temperature distribution f . The function K given above is the **heat kernel**. \square

Example 13 (Biharmonic Functions in the Upper Half-Plane) In this example, we show how the Fourier transform can be used to give an integral representation formula for a function $u(x, y)$ that is biharmonic on the upper half-plane, $U = \{x + iy: y > 0\}$, and that has given values on the real axis. The formula will be in many ways analogous to the Poisson Integral Formula for the upper half-plane. Refer to Section 2, Chapter 4, for a physical problem whose solution is biharmonic.

We wish to find a function $u(x, y)$ satisfying

$$\begin{aligned} \text{(i)} \quad \Delta(\Delta u) &= \frac{\partial^4 u}{\partial x^4} + 2 \frac{\partial^4 u}{\partial x^2 \partial y^2} + \frac{\partial^4 u}{\partial y^4} = 0 \quad \text{on } U \\ \text{(ii)} \quad u(x, 0) &= f(x), \quad -\infty < x < \infty. \end{aligned}$$

Let $U(s, y)$ be the Fourier transform of $u(x, y)$ on the x -variable:

$$U(s, y) = \int_{-\infty}^{\infty} u(x, y) e^{-isx} dx.$$

We hold s fixed and treat U as a function of y . Differentiation yields

$$\begin{aligned} \frac{d^4 U}{dy^4} &= \int_{-\infty}^{\infty} \frac{\partial^4 u}{\partial y^4} e^{-isx} dx \\ &= \int_{-\infty}^{\infty} \left\{ -\frac{2\partial^4 u}{\partial y^2 \partial x^2} - \frac{\partial^4 u}{\partial x^4} \right\} e^{-isx} dx. \end{aligned}$$

Using (6), or integrating by parts, we find that

$$\begin{aligned}\frac{d^4U}{dy^4} &= 2s^2 \int_{-\infty}^{\infty} \frac{\partial^2 u}{\partial y^2} e^{-isx} dx - s^4 \int_{-\infty}^{\infty} u(x, y) e^{-isx} dx \\ &= 2s^2 \frac{d^2 U}{dy^2} - s^4 U.\end{aligned}$$

Hence, U satisfies the fourth-order linear ordinary differential equation

$$\frac{d^4U}{dy^4} - 2s^2 \frac{d^2 U}{dy^2} + s^4 U = 0,$$

with

$$\begin{aligned}U(s, 0) &= \int_{-\infty}^{\infty} u(x, 0) e^{-isx} dx \\ &= \int_{-\infty}^{\infty} f(x) e^{-isx} dx \\ &= \hat{f}(s).\end{aligned}$$

The linearly independent solutions of $v^{(4)} - 2s^2 v'' + s^4 v = 0$ are e^{sy} , ye^{sy} , e^{-sy} , and ye^{-sy} . In our problem, y is positive, and the solution U must be a Fourier transform. This leads to the conclusion that

$$U(s, y) = A(s)e^{-y|s|} + B(s)ye^{-y|s|}, \quad -\infty < s < \infty,$$

where $A(s)$ and $B(s)$ are functions of s (see Exercise 21 at the end of this section). The condition $U(s, 0) = \hat{f}(s)$ yields $A(s) = \hat{f}(s)$. Furthermore, $e^{-y|s|}$ is the Fourier transform of

$$u_1(t) = \frac{1}{\pi} \frac{1}{t^2 + y^2}.$$

(See Example 8 and formula (5).) Likewise, $ye^{-y|s|}$ is the Fourier transform of

$$u_2(t) = \frac{1}{\pi} \frac{y^2}{t^2 + y^2}.$$

The convolution result, equation (9), then gives

$$u(x, y) = \frac{1}{\pi} \int_{-\infty}^{\infty} f(t) \frac{y}{(x-t)^2 + y^2} dt + \frac{1}{\pi} \int_{-\infty}^{\infty} g(t) \frac{y^2}{(x-t)^2 + y^2} dt.$$

The Poisson Integral Formula shows that the first term on the right is a harmonic function on the upper half-plane U whose boundary values on the real axis are f . The second term is just $yG(x, y)$, where G is a harmonic function on U whose boundary values on the real axis are g . Thus, as y decreases to zero, $yG(x, y)$ converges to $0 \cdot g(x) = 0$. Since G is harmonic, $yG(x, y)$ is biharmonic. The function g is arbitrary, within the limitations that the integral defining $G(x, y)$ makes sense; for instance, g could be any piecewise continuous, bounded function. \square

EXERCISES FOR SECTION 5.1

In Exercises 1 to 12, find \hat{u} from the given u .

1. $u(t) = \begin{cases} t, & |t| < b \\ 0, & |t| > b \end{cases}; \quad b \text{ a positive constant}$
2. $u(t) = \begin{cases} -1, & -\sigma < t < 0 \\ 1, & 0 < t < \sigma; \quad \sigma \text{ a positive constant} \\ 0, & |t| > \sigma \end{cases}$
3. $u(t) = \begin{cases} (1+t)^2, & -1 \leq t \leq 0 \\ (1-t)^2, & 0 \leq t \leq 1 \\ 0, & |t| \geq 1 \end{cases}$
4. $u(t) = 16e^{-4t^2}$
5. $u(t) = (20 + 8t + t^2)^{-1}$
6. $u(t) = (a^2 + b^2 t^2)^{-1}; \quad a \text{ and } b \text{ positive constants}$
7. $u(t) = \int_{-\sigma}^{\sigma} \frac{1}{1+(t-s)^2} ds; \quad \sigma \text{ a positive constant}$
8. $u(t) = \int_{-\infty}^{\infty} \frac{1}{1+s^2} \frac{1}{1+(t-s)^2} ds$
9. $u(t) = \begin{cases} e^{-t^2}, & |t| > \sigma \\ e^{-t^2} - e^{-\sigma^2}, & |t| < \sigma; \quad \sigma \text{ a positive constant} \end{cases}$
- (Hint: Let $u_1(t) = e^{-t^2}$ and $u_2 = u - u_1$. Find \hat{u}_1 and \hat{u}_2 .)
10. u has the graph shown in Figure 5.4.




Figure 5.4

11. u has the graph shown in Figure 5.5.




Figure 5.5

$$12. u(t) = \begin{cases} A, & |t| < a \\ B, & a < |t| < b; \\ 0, & |t| > b \end{cases} \quad a, b, A, \text{ and } B \text{ constants, and } 0 < a < b$$

(Hint: Write $u = u_1 + u_2$, where each of u_1 and u_2 has just one step.)

13. Give the details to show that (i) (3) is valid; (ii) (4) is valid; (iii) (5) is valid; and (iv) (6) is valid.
14. Let u be piecewise smooth and $U'(t) = u(t)$. Show that U is also piecewise smooth.
15. Use the Residue Theorem in the manner of Example 8 to find $\hat{u}(x)$ if $u(t) = (1 + t^4)^{-1}$.
16. Let $u(t) = 2^{-n}$ if $n \leq |t| < n + 1$, $n = 0, 1, 2, \dots$. Find $\hat{u}(x)$.
17. Let $*$ be the operation of convolution defined in (8). Show that $u * v = v * u$ and that $u * (v_1 + v_2) = u * v_1 + u * v_2$.
18. Let $\delta(t)$ be the “impulse function” at 0 discussed in Section 5, Chapter 4, Exercise 16. Show that $\delta * \delta = \delta$ and that $\hat{\delta}(x) = 1$ for all x .
19. Suppose that f is piecewise continuous on $(-\infty, \infty)$ and satisfies

$$\int_{-\infty}^{\infty} |f(t)| dt = M.$$

Let

$$u(x, y) = \frac{1}{\pi} \int_{-\infty}^{\infty} f(t) \frac{y}{(x-t)^2 + y^2} dt, \quad y > 0.$$

Show that

$$\int_{-\infty}^{\infty} |u(x, y)| dx \leq M, \quad \text{for each } y > 0.$$

20. Suppose that $u(x, y)$ satisfies

$$\int_{-\infty}^{\infty} |u(x, y)| dx \leq M,$$

for each $y, y \geq 0$. Let

$$U(s, y) = \int_{-\infty}^{\infty} u(x, y) e^{-isx} dx.$$

Show that $|U(s, y)| \leq M$ for all $y \geq 0$ and all $s, -\infty < s < \infty$.

21. Let $U(s, y)$ be a bounded function on the region $-\infty < s < \infty, y > 0$. Suppose that U can be written as

$$U(s, y) = [A(s) + yB(s)]e^{sy} + [C(s) + yD(s)]e^{-sy}.$$

Show that

$$A(s) = B(s) = 0 \quad \text{if } s > 0$$

and

$$C(s) = D(s) = 0 \quad \text{if } s < 0.$$

(Hint: e^{sy} is unboundedly large as $y \uparrow \infty$ if $s > 0$, and e^{-sy} and ye^{-sy} both go to zero as $y \uparrow \infty$ if $s > 0$.)

22. Let u and v satisfy (1), and let w be their convolution:

$$w(x) = \int_{-\infty}^{\infty} u(x-t)v(t) dt.$$

Show that

$$\int_{-\infty}^{\infty} |w(x)| dx \leq \left(\int_{-\infty}^{\infty} |u(t)| dt \right) \left(\int_{-\infty}^{\infty} |v(t)| dt \right).$$

In Exercises 23 to 26, use the Fourier transform in the manner of Examples 12 and 13 to solve the given partial differential equations.

23. Solve the wave equation

$$u_{tt} = u_{xx}, \quad t \geq 0, \quad -\infty < x < \infty,$$

subject to

$$u(x, 0) = 0$$

$$\frac{\partial u}{\partial t}(x, 0) = g(x), \quad -\infty < x < \infty.$$

24. Solve the Dirichlet problem in the upper half-plane:

$$u_{xx} + u_{yy} = 0, \quad u(x, 0) = f(x), \quad -\infty < x < \infty.$$

(At one point, you will need to make use of the conclusion of Exercise 21.) Compare your answer to the solution given in Section 3, Chapter 4.

25. Solve the Neumann problem in the upper half-plane:

$$u_{xx} + u_{yy} = 0, \quad \lim_{t \downarrow 0} \frac{\partial u}{\partial y}(x, t) = g(x), \quad -\infty < x < \infty.$$

Compare your answer to the solution given in Section 4, Chapter 4. (As in Exercise 24, at one point you will need to use the conclusion of Exercise 21.)

26. Solve the **modified heat equation**,

$$\frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2} + g''(x), \quad -\infty < x < \infty, \quad t \geq 0,$$

with

$$u(0; x) = f(x), \quad -\infty < x < \infty.$$

Here g is a function of x alone; we assume that both g and g' satisfy (1).

27. (**The telegraph equation***) Current flows along a long cable according to the equation

$$\frac{\partial^2 u}{\partial x^2} = CL \frac{\partial^2 u}{\partial t^2} + (RC + SL) \frac{\partial u}{\partial t} + RSu,$$

where $u(x; t)$ is the current at point x , $-\infty < x < \infty$, and time t , $t \geq 0$. The constants C , L , R , and S are, respectively, the capacitance, inductance, resistance, and leakage, all per unit length. Suppose that the initial conditions are $u(x; 0) = f(x)$ and $(\partial u / \partial t)(x; 0) = g(x)$, $-\infty < x < \infty$. We shall take steps toward solving this equation. Set $a = RS/CL$, $b = \frac{1}{2}[(R/L) + (S/C)]$, and $c = (CL)^{-1/2}$. The equation becomes

$$c^2 \frac{\partial^2 u}{\partial x^2} = \frac{\partial^2 u}{\partial t^2} + 2b \frac{\partial u}{\partial t} + au.$$

Define

$$U(t) = \int_{-\infty}^{\infty} u(x; t) e^{-ixs} dx, \quad s \text{ fixed.}$$

* The telegraph equation was discovered by Oliver Heaviside; see Section 3.

- (i) Show that U satisfies the ordinary differential equation

$$U'' + 2bU' + (a + s^2c^2)U = 0,$$

and solve for U .

- (ii) In the special case where $RC = SL$, complete the work by solving for $u(x; t)$.

5.2 Formulas Relating u and \hat{u}

The relationships between a function u and its Fourier transform \hat{u} are set forth in the following two formulas. The first, the **Fourier Inversion Formula**, is

$$u(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \hat{u}(x)e^{ixt} dx, \quad (1)$$

if the function u is *continuous* at t . Since u could possibly be discontinuous at a finite number of points, a more general version of (1) is

$$\frac{1}{2}(u^+(t) + u^-(t)) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \hat{u}(x)e^{ixt} dx. \quad (1')$$

Either of these formulas shows that the function u can be recovered from its Fourier transform \hat{u} by means of a formula almost identical to the formula that defines \hat{u} in terms of u .

The second formula states that 2π times the integral over the real axis of $|u|^2$ is exactly the same as the integral of $|\hat{u}|^2$:

$$2\pi \int_{-\infty}^{\infty} |u(t)|^2 dt = \int_{-\infty}^{\infty} |\hat{u}(x)|^2 dx. \quad (2)$$

These two formulas are at the heart of the theory of the Fourier transform. As mentioned in the opening paragraph of Section 1, a thorough mathematical justification of formulas (1) and (2) is not elementary.

The equality in (2) is called **Parseval's equality**. In physical terms, it states that the total energy of a system is preserved by the Fourier transform—the transformed system has exactly as much energy as the original system. Furthermore, equation (2) is equivalent to the seemingly more general equality

$$2\pi \int_{-\infty}^{\infty} u(t)\overline{v(t)} dt = \int_{-\infty}^{\infty} \hat{u}(x)\overline{\hat{v}(x)} dx. \quad (3)$$

Indeed, (3) follows from (2) by replacing u with $u + v$ and using the known equality (2) for both $|u|^2$ and $|v|^2$. (See the exercises at the end of this section.)

Example 1 Find the Fourier transform of $f(t) = (\sin t)/t$.

Solution From Example 6, Section 1, we know that $2(\sin x)/x$ is the Fourier transform of the function

$$u(t) = \begin{cases} 1, & |t| < 1 \\ 0, & |t| > 1. \end{cases}$$

Thus, by the Fourier Inversion Formula,

$$\hat{f}(x) = \begin{cases} \pi, & |x| < 1 \\ 0, & |x| > 1. \end{cases}$$
□

You might try to compute $\hat{f}(x)$ directly, to appreciate better the strength of the Fourier Inversion Formula we are using here.

Example 2 Compute

$$\int_{-\infty}^{\infty} \frac{(1 - \cos x)^2}{x^4} dx.$$

Solution The function

$$f(x) = 2 \frac{1 - \cos x}{x^2}$$

is the Fourier transform of the function

$$u(t) = \begin{cases} 1 + t, & -1 \leq t \leq 0 \\ 1 - t, & 0 \leq t \leq 1 \\ 0, & |t| > 1. \end{cases}$$

(See the function u_3 in Example 9, Section 1.) The integral we want is one-quarter the integral of the square of $f = \hat{u}$; by (2), this is equal to

$$\left(\frac{1}{4}\right) 2\pi \int_{-\infty}^{\infty} (u(t))^2 dt = \frac{\pi}{3}.$$

Hence,

$$\int_{-\infty}^{\infty} \frac{(1 - \cos x)^2}{x^4} dx = \frac{\pi}{3}.$$
□

The Complex Fourier Transform

Let us look back at formula (2) in Section 1, which defines the Fourier transform of a function u , but now replace the real variable x with the complex variable

$z = x + iy$:

$$\hat{u}(z) = \int_{-\infty}^{\infty} u(t)e^{-izt} dt. \quad (4)$$

We need to concern ourselves first with the question of whether the integral makes sense and, if so, for which z . We note that

$$|u(t)e^{-izt}| = |u(t)|e^{yt}.$$

First take $-\infty < t \leq 0$; in order that the integral on $(-\infty, 0)$ converges, we assume that

$$|u(t)| \leq M_1 e^{-\tau_1 t}, \quad t \leq 0 \quad (5)$$

for some constant M_1 and some number τ_1 . It follows from this that

$$|u(t)e^{-izt}| \leq M_1 e^{t(y-\tau_1)}, \quad t \leq 0,$$

so

$$\int_{-\infty}^0 u(t)e^{-izt} dt$$

converges for all z with $y = \operatorname{Im} z > \tau_1$ (see Exercises 16 to 19 at the end of this section). Likewise, if u satisfies the growth condition

$$|u(t)| \leq M_2 e^{-\tau_2 t} \quad \text{for } t \geq 0 \quad (6)$$

for some number τ_2 , then the integral

$$\int_0^{\infty} u(t)e^{-izt} dt$$

is absolutely convergent for all z with $-\tau_2 + y < 0$. Thus, if both (5) and (6) hold and if $\tau_1 < \tau_2$, then the integral defining $\hat{u}(z)$ is convergent for all z with

$$\tau_1 < \operatorname{Im} z < \tau_2. \quad (7)$$

In this way, we see that the complex Fourier transform $\hat{u}(z)$ is defined for all z in the horizontal strip defined by (7). Moreover, in this strip, $\hat{u}(z)$ is an analytic function of z with derivative

$$[\hat{u}]'(z) = -i \int_{-\infty}^{\infty} u(t)te^{-izt} dt. \quad (8)$$

The analyticity of $\hat{u}(z)$ can be validated more fully by use of Morera's Theorem (see the exercises at the end of this section). Thus, for example, if $u(t) = 0$ for all t with $|t| \geq A > 0$, then surely both (5) and (6) hold for every τ_1 and τ_2 , so $\hat{u}(z)$ is analytic for all z ; that is, \hat{u} is an entire function.

Example 3 Let $u(t) = e^{-|t|}$. Then $\tau_1 = -1$ and $\tau_2 = 1$, so $\hat{u}(z)$ is analytic within the strip $|\operatorname{Im} z| < 1$. In fact,

$$\hat{u}(z) = 2 \left(\frac{1}{1 + z^2} \right),$$

as Example 8, Section 1, and the Fourier Inversion Formula show. Thus, the strip of analyticity of \hat{u} is as wide as possible: \hat{u} is not analytic in any wider (open) strip. \square

Example 4 Let us look at Example 3 the other way around. Take $u(t) = 2(1 + t^2)^{-1}$; then $\hat{u}(x) = 2\pi e^{-|x|}$, and this function has no analytic extension to a strip of the form $\tau_1 < \operatorname{Im} z < \tau_2$, no matter how τ_1 and τ_2 are chosen, $\tau_1 \leq 0 \leq \tau_2$. Thus the bounds on the growth of $|u(t)|$ given by (5) and (6) are not at all superfluous. \square

To invert the complex Fourier transform, we use the usual formula, but in a new location. Let y_0 be any number with $\tau_1 < y_0 < \tau_2$; then

$$\begin{aligned} |u(t)e^{-it(x+iy_0)}| &= |u(t)|e^{ty_0} \\ &\leq \begin{cases} M_1 e^{-\varepsilon_1 t} & \text{if } t \geq 0 \\ M_2 e^{+\varepsilon_2 t} & \text{if } t \leq 0, \end{cases} \end{aligned}$$

where M_1 , M_2 , ε_1 , and ε_2 are positive constants. (See (5) and (6).) Let

$$u_1(t) = u(t)e^{ty_0}.$$

Then for $-\infty < x < \infty$,

$$\hat{u}_1(x) = \hat{u}(x + iy_0).$$

Consequently, by the Fourier Inversion Formula,

$$\begin{aligned} u(t) &= e^{-ty_0} u_1(t) \\ &= e^{-ty_0} \frac{1}{2\pi} \int_{-\infty}^{\infty} \hat{u}_1(x) e^{ix} dx \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} \hat{u}(x + iy_0) e^{it(x+iy_0)} dx \\ &= \frac{1}{2\pi} \int_{-\infty+iy_0}^{\infty+iy_0} \hat{u}(z) e^{itz} dz. \end{aligned}$$

The lower and upper limits on this last integral mean that z traverses the entire horizontal line $y = y_0$ from left to right. Since y_0 is any number in the interval (τ_1, τ_2) , we can write

$$u(t) = \frac{1}{2\pi} \int_{-\infty+iy_0}^{\infty+iy_0} \hat{u}(z) e^{izt} dz, \quad \tau_1 < y < \tau_2.$$

$\hat{u}(z)$ is analytic; obviously so is e^{izt} . Thus, it is frequently possible to use the Residue Theorem, or some other similar device, to evaluate this integral.

Example 5 Find u if $\hat{u}(z) = 1/z^2$.

Solution $\hat{u}(z)$ is analytic except at the origin, so we may use the Residue Theorem to evaluate the integral

$$\int_{-\infty+iy_0}^{\infty+iy_0} \frac{e^{izt}}{z^2} dz.$$

We choose y_0 to be a positive number. If t is positive, we integrate over the semicircle shown in Figure 5.6a; if t is negative, we integrate over the semicircle in Figure 5.6b. In both cases, the integral over the curved portion goes to zero as $R \rightarrow \infty$. Cauchy's Theorem and the Residue Theorem then give, respectively,

$$u(t) = \begin{cases} 0, & t > 0 \\ t, & t < 0. \end{cases}$$
□




Figure 5.6

A Mathematical Justification of Formulas (1) and (2)

The correctness of what is done as follows depends on the validity of two steps that, although clearly believable, are themselves in need of further mathematical

justification. We will point these out and comment further on them after the derivation of the formulas.

Let u and v be two piecewise smooth functions that satisfy the conditions

$$\int_{-\infty}^{\infty} |u(t)| dt < \infty; \quad \int_{-\infty}^{\infty} |u(t)|^2 dt < \infty; \quad \lim_{|t| \rightarrow \infty} u(t) = \lim_{|t| \rightarrow \infty} u'(t) = 0,$$

and the same with v in place of u . Then

$$\begin{aligned} \int_{-\infty}^{\infty} \hat{u}(x)v(x)e^{isx} dx &= \int_{-\infty}^{\infty} v(x) \left\{ \int_{-\infty}^{\infty} u(t)e^{-ixt} dt \right\} e^{isx} dx \\ &= \int_{-\infty}^{\infty} u(t) \left\{ \int_{-\infty}^{\infty} v(x)e^{-ix(t-s)} dx \right\} dt \\ &= \int_{-\infty}^{\infty} u(t)\hat{v}(t-s) dt = \int_{-\infty}^{\infty} u(t+s)\hat{v}(t) dt. \end{aligned}$$

Now take $v(x) = e^{-\epsilon x^2}$, where ϵ is a small positive number. We know, from Example 7 in Section 1 and (5) in Section 1, that

$$\hat{v}(t) = \sqrt{\pi} \frac{e^{-t^2/4\epsilon^2}}{\epsilon}.$$

Hence,

$$\begin{aligned} \int_{-\infty}^{\infty} e^{-\epsilon x^2} \hat{u}(x)e^{isx} dx &= \sqrt{\pi} \int_{-\infty}^{\infty} u(t+s) \frac{e^{-t^2/4\epsilon^2}}{\epsilon} dt \\ &= \sqrt{\pi} \int_{-\infty}^{\infty} u(\epsilon y + s) e^{-y^2/4} dy. \end{aligned}$$

Now let ϵ decrease to zero; the left-hand side converges to

$$\int_{-\infty}^{\infty} \hat{u}(x)e^{isx} dx.$$

The right-hand side converges to

$$\sqrt{\pi} \int_{-\infty}^{\infty} u(s) e^{-y^2/4} dy = u(s)2\pi$$

if u is continuous at s or to $(2\pi)^{\frac{1}{2}}(u^+(s) + u^-(s))$ if s happens to be one of the finite

number of points of discontinuity of u . This gives (1) if u is continuous at s or (1') if u is discontinuous at s .*

To prove (2), let $\varepsilon > 0$ and set

$$v_\varepsilon(t) = \overline{\hat{u}(t)} e^{-\varepsilon^2 t^2},$$

$$w_\varepsilon(t) = \frac{1}{2\sqrt{\pi\varepsilon}} e^{-t^2/4\varepsilon^2},$$

and

$$p(t) = \overline{u(-t)}.$$

A simple computation gives $\overline{\hat{u}(t)} = \hat{p}(t)$, so

$$v_\varepsilon(t) = \hat{p}(t) \hat{w}_\varepsilon(t) = (p * w_\varepsilon)^\wedge(t).$$

Using (1), we find that

$$\begin{aligned} \frac{1}{2\pi} \hat{v}_\varepsilon(x) &= (p * w_\varepsilon)(-x) \\ &= \int_{-\infty}^{\infty} p(-x - s) w_\varepsilon(s) ds \\ &= \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} p(-x - 2\varepsilon t) e^{-t^2} dt \\ &= \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} \overline{u(x + 2\varepsilon t)} e^{-t^2} dt. \end{aligned}$$

In the formula

$$\int_{-\infty}^{\infty} \hat{u}(x) v(x) e^{isx} dx = \int_{-\infty}^{\infty} u(x + s) \hat{v}(x) dx,$$

derived above, we take $s = 0$ and $v = v_\varepsilon$. This produces the relation

$$\begin{aligned} \int_{-\infty}^{\infty} |\hat{u}(x)|^2 e^{-\varepsilon^2 x^2} dx &= \int_{-\infty}^{\infty} u(x) \left\{ \int_{-\infty}^{\infty} (2\sqrt{\pi}) \overline{u(x + 2\varepsilon t)} e^{-t^2} dt \right\} dx \\ &= 2\sqrt{\pi} \int_{-\infty}^{\infty} e^{-t^2} \left\{ \int_{-\infty}^{\infty} u(x) \overline{u(x + 2\varepsilon t)} dx \right\} dt. \end{aligned}$$

Upon letting ε decrease to zero, we obtain (2).

* An alternative justification of (1) is given in Exercises 22 to 25.

A few comments now on the preceding mathematics. To prove (1), we wrote down the integral of $\hat{u}(x)v(x)$, then replaced $\hat{u}(x)$ with its integral formulation, and then interchanged the order of integration in the resulting iterated integral. This is mathematically justifiable here, but only with some effort. We then made the choice $v(x) = \exp(-ex^2)$; after a change of variables, we then asserted that

$$\lim_{\varepsilon \rightarrow 0} \int_{-\infty}^{\infty} (\quad) dx = \int_{-\infty}^{\infty} \lim_{\varepsilon \rightarrow 0} (\quad) dx,$$

where the expression (\quad) depends on ε as well as the variable of integration. Again, this is reasonable but needs mathematical justification.

EXERCISES FOR SECTION 5.2

1. Use (3) to show that

$$\int_{-\infty}^{\infty} \frac{\sin \alpha x \sin \beta x}{x^2} dx = \pi \min\{\alpha, \beta\}, \quad \alpha \text{ and } \beta \text{ positive.}$$

2. Show that

$$\int_0^{\infty} e^{-t} \frac{\sin \alpha t}{t} dt = \arctan \alpha, \quad \alpha \text{ positive.}$$

(Hint: Rewrite the integral as

$$\frac{1}{2} \int_{-\infty}^{\infty} e^{-|t|} \frac{\sin \alpha t}{t} dt$$

and use (3).)

3. Compute the value of

$$\int_{-\infty}^{\infty} \left(\frac{1 - \cos \sigma x}{x} \right)^2 dx$$

using (2) and Exercise 2, Section 1.

4. Compute the value of

$$\int_{-\infty}^{\infty} \frac{(x - \sin x)^2}{x^6} dx$$

using (2) and Exercise 3, Section 1.

5. Using (2), compute the value of

$$\int_{-\infty}^{\infty} \frac{dx}{(1 + x^2)^2}.$$

6. Find \hat{u} if

$$u(t) = \int_{-\infty}^{\infty} e^{-|t-s|} \frac{\sin \sigma s}{s} ds.$$

7. Find u if

$$\hat{u}(x) = e^{-|x|} \left(\frac{\sin 4x}{x} \right).$$

8. Suppose that $u(t) = 0$ if $|t| > A$. Use Morera's Theorem to show that $\hat{u}(z)$ is an entire function of the complex variable z ; moreover, \hat{u} is bounded on the real axis and, in general, satisfies $|\hat{u}(z)| \leq 2AMe^{4|y|}$ if $z = x + iy$, $M = \max\{|u(t)| : |t| \leq A\}$.
9. Use (3) to find the value of

$$\int_{-\infty}^{\infty} \frac{(x - \sin x)(\sin \sigma x)}{x^4} dx.$$

10. Use (3) to show that

$$\int_{-\infty}^{\infty} \frac{(\sin^2 \alpha x)(\sin^2 \beta x)}{x^2} dx = \frac{\pi}{2} \min\{\alpha, \beta\}.$$

(Hint: Use $2 \sin^2 \theta = 1 - \cos 2\theta$ and Exercise 2, Section 5.1.)

11. Suppose that $u(t) = 0$ for $|t| \geq A$ and that $\hat{u}(x) = 0$ for $|x| \geq B$, A and $B > 0$. Show that u is identically zero. (Hint: Refer to Exercise 8.)
12. Suppose that u_1 and u_2 are two functions satisfying (1) of Section 5.1 and u_1 and u_2 have the same Fourier transform. Show that $u_1 = u_2$ at all points of continuity of u_1 and u_2 .
13. Give the details of how (2) implies (3).
14. Suppose that $\int_{-\infty}^{\infty} |u(t)| dt < \infty$; show that $\hat{u}(x)$ is a continuous function of x . (Hint: We can write

$$\hat{u}(x_2) - \hat{u}(x_1) = \int_{-\infty}^{\infty} u(t) \{e^{-ix_2 t} - e^{-ix_1 t}\} dt = \int_{-M}^{M} + \int_M^{\infty} + \int_{-\infty}^{-M}.$$

The second and third integrals are small for all x_1 and x_2 if only M is big, while for M fixed the first integral is small if x_1 is near x_2 .)

15. Suppose (1) of Section 1 holds. Fill in the details of the following steps to show that the **Riemann–Lebesgue Lemma** holds:

$$\lim_{x \rightarrow \infty} \hat{u}(x) = 0.$$

(a) $\hat{u}(x) = - \int_{-\infty}^{\infty} e^{-ixt} u(t + (\pi/x)) dt$

(b) $\hat{u}(x) = \int_{-\infty}^{\infty} e^{-ixt} \frac{1}{2} \{u(t) - u(t + \pi/x)\} dt$
 (c) $|\hat{u}(x)| \leq \left\{ \int_{-M}^M + \int_M^{\infty} + \int_{-\infty}^{-M} \right\} w(t) dt,$

where $w(t) = |u(t) - u(t + \pi/x)|$. Show that the second and third integrals are small for any x if M is big; show that the first integral is small for M fixed if x is big and u is continuous.

- (d) Modify step (c) above to include the case when u is merely piecewise continuous.

Improper Integrals on $(-\infty, \infty)$

A piecewise continuous function u on the real axis has an **improper integral** on $(-\infty, \infty)$ if

$$\lim_{\substack{M_1 \rightarrow \infty \\ M_2 \rightarrow \infty}} \int_{-M_2}^{M_1} u(t) dt$$

exists, no matter how M_1 and M_2 go to ∞ . $u(t)$ is **absolutely integrable** if $|u(t)|$ has an improper integral. We denote the improper integral of u on $(-\infty, \infty)$ by $\int_{-\infty}^{\infty} u(t) dt$.

16. Show that u is absolutely integrable if and only if there is a constant C such that

$$\int_{-M}^M |u(t)| dt \leq C$$

for all choices of M . Show that the numbers $\int_{-M}^M |u(t)| dt$ increase to $\int_{-\infty}^{\infty} |u(t)| dt$ as $M \uparrow \infty$.

17. Let u be absolutely integrable. Show that

$$\lim_{B \rightarrow \infty} \int_{-\infty}^{-B} |u(t)| dt = \lim_{A \rightarrow \infty} \int_A^{\infty} |u(t)| dt = 0.$$

18. Let u be absolutely integrable. Show that u has an improper integral on $(-\infty, \infty)$ and that

$$\left| \int_{-\infty}^{\infty} u(t) dt \right| \leq \int_{-\infty}^{\infty} |u(t)| dt.$$

(Hint: For any $A < B$,

$$\left| \int_A^B u(t) dt \right| \leq \int_A^B |u(t)| dt.$$

Next show that for M_1 and M_2 sufficiently large and $M'_1 > M_1$, $M'_2 > M_2$, we have

$$\left| \int_{M_1}^{M'_1} u(t) dt \right| \leq \int_{M_1}^{M'_1} |u(t)| dt < \varepsilon$$

$$\left| \int_{-M'_2}^{-M_2} u(t) dt \right| \leq \int_{-M'_2}^{-M_2} |u(t)| dt < \varepsilon.)$$

19. Let $|u(t)| \leq |v(t)|$ for $-\infty < t < \infty$, and suppose that v is absolutely integrable on $(-\infty, \infty)$. Show that u is also absolutely integrable on $(-\infty, \infty)$.
20. Suppose that $\int_{-\infty}^{\infty} |u'(t)| dt < \infty$ and $\int_{-\infty}^{\infty} |u(t)| dt < \infty$. Show that $\lim_{x \rightarrow \infty} u(x) = 0$. (Hint: Given a small $\varepsilon > 0$, choose M so large that $\int_M^{\infty} |u'(t)| dt < \varepsilon$ (see Exercise 17). For any $s > 0$,

$$|u(s+M) - u(M)| = \left| \int_M^{M+s} u'(t) dt \right| \leq \int_M^{M+s} |u'(t)| dt < \varepsilon.$$

Now let $s \rightarrow \infty$. This shows that $\lim_{x \rightarrow \infty} u(x)$ exists. Now show that this limit must be zero.)

21. Use Morera's Theorem to prove that $\hat{u}(z)$ is analytic for $\tau_1 < \operatorname{Im} z < \tau_2$ if u satisfies (5) and (6).

An Alternate Proof of the Fourier Inversion Formula (1)*

We shall show that, for x fixed,

$$\lim_{M \rightarrow \infty} \frac{1}{2\pi} \int_{-M}^M \hat{u}(s) e^{ixs} ds = \frac{1}{2}(u^+(x) + u^-(x)).$$

22. Show that

$$\begin{aligned} \int_{-M}^M \hat{u}(s) e^{ixs} ds &= 2 \int_{-\infty}^{\infty} u(t) \frac{\sin M(x-t)}{x-t} dt \\ &= 2 \int_{-\infty}^{\infty} u(x+t) \frac{\sin Mt}{t} dt. \end{aligned}$$

23. Use Exercise 15 to show that for each x and each $\delta > 0$,

$$(i) \quad \lim_{M \rightarrow \infty} \int_{\delta}^{\infty} u(x+t) \frac{\sin Mt}{t} dt = 0$$

and

$$(ii) \quad \lim_{M \rightarrow \infty} \int_{-\infty}^{-\delta} u(x+t) \frac{\sin Mt}{t} dt = 0.$$

24. Choose δ so small that $u(x+t) - u^+(x)$ is continuous on $[0, \delta]$ and $u(x+t) - u^-(x)$ is continuous on $[-\delta, 0]$. Show that

$$(iii) \quad \lim_{M \rightarrow \infty} \int_0^\delta (u(x+t) - u^+(x)) \frac{\sin Mt}{t} dt = 0,$$

$$(iv) \quad \lim_{M \rightarrow \infty} \int_{-\delta}^0 (u(x+t) - u^-(x)) \frac{\sin Mt}{t} dt = 0.$$

25. Conclude from Exercises 23 and 24 that

$$\begin{aligned} \lim_{M \rightarrow \infty} \int_{-\infty}^{\infty} u(x+t) \frac{\sin Mt}{t} dt &= (u^+(x) + u^-(x)) \int_0^{\infty} \frac{\sin x}{x} dx \\ &= \frac{\pi}{2}(u^+(x) + u^-(x)), \end{aligned}$$

and hence that

$$\lim_{M \rightarrow \infty} \frac{1}{2\pi} \int_{-M}^M \hat{u}(s) e^{ixs} ds = \frac{1}{2}(u^+(x) + u^-(x)).$$

5.3 The Laplace Transform

Roughly speaking, the Laplace transform is just the Fourier transform rotated by 90° and applied to a function that vanishes on the ray $(-\infty, 0)$. Specifically, suppose that u is a piecewise continuous function on $[0, \infty)$ such that both u and u' have limits from the right at $t = 0$. The **Laplace transform** of u is

$$(\mathcal{L}u)(s) = \int_0^{\infty} u(t) e^{-st} dt. \quad (1)$$

Initially, we can think of s as a real variable, but with the knowledge we already have about the complex Fourier transform, it is natural to investigate immediately for which complex numbers s the integral in (1) makes sense. We shall henceforth suppose that the function u has **exponential growth** a ; that is, there is a constant M such that

$$|u(t)| \leq M e^{at}, \quad t > 0. \quad (2)$$

Let the complex number s be $s = \sigma + it$; then

$$\begin{aligned} |u(t)e^{-st}| &\leq M e^{at} e^{-\sigma t} \\ &= M e^{t(a-\sigma)}. \end{aligned}$$

Hence, if $a - \sigma < 0$, then the integral in (1) is absolutely convergent. We see,

therefore, that the Laplace transform of u is defined at least for all s with $\operatorname{Re} s > a$, where a is the exponential order of growth of u from (2). Furthermore, within the half-plane $\operatorname{Re} s > a$, the Laplace transform of u is an analytic function of the complex variable s . One way to see that $\mathcal{L}u(s)$ is analytic is to write down the elementary relationship between the Laplace and Fourier transforms of the function u :

$$(\mathcal{L}u)(s) = \hat{u}(-is), \quad \operatorname{Re} s > a. \quad (3)$$

Since $\hat{u}(z)$ is analytic in the half-plane $\operatorname{Im} z < -a$, it follows that $\mathcal{L}u(s)$ is analytic for $\operatorname{Re} s > a$.*

Why should we bother with the Laplace transform when it is so much like the Fourier transform? There are several reasons. One is that in a physical system, there is typically no output until there is input. That is, the function that represents the output must vanish on the ray $(-\infty, 0)$ (assuming that the input begins at $t = 0$); such functions are termed **causal**. Another reason is that by beginning the integration at $t = 0$, when we integrate by parts the boundary terms will incorporate the numbers $u(0)$, $u'(0)$, etc., so the Laplace transform is an effective tool for handling differential equations on the ray $(0, \infty)$ that include **initial conditions**.

Finally, the Laplace transform is a better tool than the Fourier transform for dealing with the impulse function $\delta(t)$ and the **Heaviside[†] function** H , defined by

$$H(t) = \begin{cases} 0, & t < 0 \\ 1, & t > 0. \end{cases}$$

As was pointed out in Exercise 17, Section 5, Chapter 4, H is in some sense an indefinite integral of $\delta(t)$. Furthermore, the Fourier transform of $H(t_0 - t)$ makes no sense for any t_0 , whereas the Laplace transform is defined for all t_0 :

$$\begin{aligned} \int_0^\infty H(t_0 - t)e^{-st} dt &= \int_0^{t_0} e^{-st} dt \\ &= \frac{1}{s}[1 - e^{-st_0}] \quad \text{if } t_0 > 0, \end{aligned}$$

and

$$\int_0^\infty H(t_0 - t)e^{-st} dt = 0, \quad \text{if } t_0 \leq 0.$$

Example 1 Find the Laplace transform of each of the following functions (all the functions vanish for $t < 0$):

* It is possible to prove that there is a number s_0 such that the Laplace transform of u converges if $\operatorname{Re} z > s_0$ and diverges if $\operatorname{Re} z < s_0$; s_0 is much like the radius of convergence of a power series.

† Oliver Heaviside, 1850–1925.

- a. $u_1(t) = 1$;
- b. $u_2(t) = t^k$, k a positive integer;
- c. $u_3(t) = \exp(at)$, a a complex number;
- d. $u_4(t)$ is the impulse function at the point t_0 , $t_0 > 0$.

Solution

- a. $\mathcal{L}u_1(s) = \int_0^\infty e^{-st} dt = 1/s$.
- b. $\mathcal{L}u_2(s) = \int_0^\infty t^k e^{-st} dt = k!/s^{k+1}$; integrate by parts k times.
- c. $\mathcal{L}u_3(s) = \int_0^\infty e^{at} e^{-st} dt = 1/(s-a)$ for $\operatorname{Re} s > \operatorname{Re} a$.
- d. $\mathcal{L}u_4(s) = \int_0^\infty \delta(t - t_0) e^{-st} dt = e^{-st_0}$. \square

Example 2 Find the Laplace transform of u' , u'' , ..., in terms of the Laplace transform of u and the initial data $u(0)$, $u'(0)$,

Solution We assume that u and all its derivatives have exponential growth a . For $\operatorname{Re} s > a$,

$$\begin{aligned} (\mathcal{L}u')(s) &= \int_0^\infty u'(t) e^{-st} dt = u(t) e^{-st} \Big|_0^\infty + s \int_0^\infty u(t) e^{-st} dt \\ &= -u(0) + s\mathcal{L}u(s). \end{aligned}$$

Likewise,

$$\begin{aligned} (\mathcal{L}u'')(s) &= \int_0^\infty u''(t) e^{-st} dt = u'(t) e^{-st} \Big|_0^\infty + s \int_0^\infty u'(t) e^{-st} dt \\ &= -u'(0) + s(\mathcal{L}u')(s) = -u'(0) - su(0) + s^2\mathcal{L}u(s). \end{aligned}$$

Continuing in this way, we obtain the formula

$$(\mathcal{L}u^{(n)})(s) = s^n \mathcal{L}u(s) - s^{n-1}u(0) - s^{n-2}u'(0) - \cdots - u^{(n-1)}(0), \quad (4)$$

which is valid for all complex numbers s with $\operatorname{Re} s > a$. \square

Further Properties of the Laplace Transform

It is clear that the Laplace transform is linear:

$$\mathcal{L}(\lambda_1 u_1 + \lambda_2 u_2)(s) = \lambda_1 \mathcal{L}u_1(s) + \lambda_2 \mathcal{L}u_2(s), \quad \operatorname{Re} s > a,$$

for any two functions u_1 , u_2 , both of exponential growth a , and complex numbers λ_1 , λ_2 . In addition, like the Fourier transform, the Laplace transform of the convolution of two functions is just the product of their Laplace transforms. Since the functions vanish on $(-\infty, 0)$, the convolution takes the form

$$(u * v)(t) = \int_0^t u(x)v(t-x) dx, \quad t > 0,$$

and the transform formula is

$$[\mathcal{L}(u * v)](s) = (\mathcal{L}u(s))(\mathcal{L}v(s)). \quad (5)$$

Another useful formula relates the derivative of the Laplace transform of u to the transform of $tu(t)$. The formula is

$$\frac{d}{ds}(\mathcal{L}u)(s) = -\mathcal{L}(tu(t))(s). \quad (6)$$

The derivation of (6) is immediate (see Exercise 18).

Next, if the Laplace transform of u is identically zero, then u is also identically zero. Consequently, if u_1 and u_2 are two functions with the same Laplace transform, then u_1 and u_2 are the same function.

Furthermore, because of the close connection between the Fourier and Laplace transforms, we can quickly and easily obtain the inversion formula for the Laplace transform. Suppose that $u(t)$ vanishes for $t < 0$ and is of exponential growth a ; let σ be any number larger than a . The Laplace and Fourier transforms are related by

$$\hat{u}(-is) = (\mathcal{L}u)(s), \quad \operatorname{Re} s > a.$$

By the Fourier Inversion Formula, we know that

$$u(t) = \frac{1}{2\pi} \int_{-\infty+i\sigma}^{\infty+i\sigma} \hat{u}(z)e^{izt} dz.$$

In this integral, set $z = -is$; this produces the formula

$$u(t) = \frac{1}{2\pi i} \int_{\sigma-i\infty}^{\sigma+i\infty} (\mathcal{L}u)(s)e^{st} ds, \quad (7)$$

where σ is any number larger than a , and the integration is over the entire vertical line $x = \sigma$. Formula (7) is the **inversion formula for the Laplace transform**. Typically, the integral is evaluated by means of the Residue Theorem, as the following examples show.

Example 3 Find u if

$$\mathcal{L}u(s) = \frac{1}{s^2 - 2s + 3}.$$

Solution The function u is found from the Laplace inversion formula (7). With $\mathcal{L}u$ as given, we obtain

$$u(t) = \frac{1}{2\pi i} \int_{\sigma-i\infty}^{\sigma+i\infty} \frac{e^{st}}{s^2 - 2s + 3} ds.$$




Figure 5.7

The function $(s^2 - 2s + 3)^{-1}$ has poles at $1 \pm i\sqrt{2}$ and so is analytic for $\operatorname{Re} s > 1$; hence, we choose $\sigma > 1$ for the vertical line of integration. We distinguish two cases: $t \geq 0$ and $t < 0$. For $t \geq 0$, we integrate

$$f(z) = \frac{e^{zt}}{z^2 - 2z + 3}$$

over the curve Γ shown in Figure 5.7. On the semicircular part of Γ , $z = \sigma + Re^{it}$, $\pi/2 \leq \theta \leq 3\pi/2$; this gives the estimate (for $t \geq 0$)

$$\begin{aligned} |f(z)| &\leq e^{\sigma t} \frac{e^{tR \cos \theta}}{R^2 - 2R - 3}, \quad \text{if } R \text{ is large,} \\ &\leq \frac{4e^{\sigma t}}{R^2 - 8R - 12}. \end{aligned}$$

Thus, the integral of f over the semicircular part of Γ is no larger than

$$\frac{4e^{\sigma t}}{R^2 - 8R - 12} \pi R \rightarrow 0 \quad \text{as } R \rightarrow \infty$$

(see formula (3), Section 6, Chapter 1). The function f has simple poles at $1 \pm i\sqrt{2}$ with residues

$$\operatorname{Res}(f; 1 + i\sqrt{2}) = \frac{e^t e^{it\sqrt{2}}}{2i\sqrt{2}}$$

and

$$\operatorname{Res}(f; 1 - i\sqrt{2}) = \frac{e^t e^{-it\sqrt{2}}}{-2i\sqrt{2}}.$$




Figure 5.8

An application of the Residue Theorem and passage to the limit as $R \rightarrow \infty$ then yields

$$\begin{aligned} u(t) &= \operatorname{Res}(f; 1 + i\sqrt{2}) + \operatorname{Res}(f; 1 - i\sqrt{2}) \\ &= \frac{e^t}{\sqrt{2}} \sin(\sqrt{2}t), \quad t \geq 0. \end{aligned}$$

On the other hand, if $t < 0$, the integration is taken over the curve Γ shown in Figure 5.8.

Estimates such as those just completed show that the integral of f over the semicircular part of Γ goes to zero as $R \rightarrow \infty$. However, because f is analytic on and within Γ , Cauchy's Theorem (and passage to the limit as $R \rightarrow \infty$) yields $u(t) = 0$ for $t < 0$. Hence,

$$u(t) = \begin{cases} e^t \sin(\sqrt{2}t), & t \geq 0 \\ 0, & t < 0. \end{cases}$$
□

Example 4 Find u if

$$\mathcal{L}u(s) = \frac{s}{(s^2 + A^2)^2}.$$

Solution We recall that

$$\mathcal{L}(\sin At)(s) = \frac{A}{s^2 + A^2}.$$

Hence,

$$\frac{d}{ds} \mathcal{L}(\sin At)(s) = \frac{-2sA}{(s^2 + A^2)^2}.$$

However, by formula (6), we also have

$$-\mathcal{L}(t \sin At) = \frac{d}{ds} \mathcal{L}(\sin At)(s) = \frac{-2sA}{(s^2 + A^2)^2}.$$

Consequently,

$$u(t) = \frac{1}{2A} t \sin At.$$

Another technique to find u is to use the Residue Theorem (see the exercises). \square

Example 5 Let u be a piecewise smooth function that satisfies the equation

$$u(t) = u(t + T), \quad t \geq 0$$

for some fixed positive number T . Find the Laplace transform of u .

Solution For $\operatorname{Re} s > 0$,

$$\begin{aligned} (\mathcal{L}u)(s) &= \int_0^\infty u(t)e^{-st} dt \\ &= \int_0^T u(t)e^{-st} dt + \int_T^{2T} u(t)e^{-st} dt + \int_{2T}^{3T} u(t)e^{-st} dt + \cdots \\ &= \int_0^T u(t)e^{-st} dt + \int_0^T u(t)e^{-s(t+T)} dt + \int_0^T u(t)e^{s(t+2T)} dt + \cdots \\ &= \int_0^T u(t)e^{-st} \{1 + e^{-sT} + e^{-2sT} + \cdots\} dt \\ &= \frac{\int_0^T u(t)e^{-st} dt}{1 - e^{-sT}}. \end{aligned}$$

The number T is called a **period** of u , and the smallest such T is called the period of u . \square

EXERCISES FOR SECTION 5.3

In Exercises 1 to 10, find the Laplace transform of the given function u . In all cases, $u(t) = 0$ if $t < 0$.

1. $u(t) = \cos At$
2. $u(t) = \sinh At$
3. $u(t) = e^{-Bt} \sin At$

4. $u(t) = e^{-Bt} \cos At$

5. $u(t) = \begin{cases} 1, & a < t < b, a > 0 \\ 0, & \text{otherwise} \end{cases}$

6. $u(t) = \begin{cases} 1, & 0 < t < \sigma \\ -1, & \sigma < t < 2\sigma \\ 0, & t > 2\sigma \end{cases}$

7. $u(t) = \begin{cases} t, & 0 < t < \sigma \\ 0, & t > \sigma \end{cases}$

8. $u(t) = \begin{cases} 1, & 0 < t < \sigma \\ 0, & \sigma < t < 2\sigma \end{cases}$ and $u(t) = u(t + 2\sigma)$, $t > 0$

9. $u(t) = \int_0^t H(s)H(t-s) ds$, H = Heaviside function

10. $u(t) = \int_0^t \cos(t-u) \sin u du$

11. Use term-by-term integration to show that the Laplace transform of

$$u(t) = \begin{cases} 0, & t < 0 \\ \frac{\sin t}{t}, & t > 0 \end{cases}$$

is $\arctan(1/s)$, $s > 1$. This can also be done by using formula (6).

12. Use term-by-term integration to show that the Laplace transform of the 0th Bessel function J_0 is $1/\sqrt{s^2 + 1}$, $s > 1$. (Hint: Compute $\mathcal{L}J_0$ and then square it.)

In Exercises 13 to 17, find u from $\mathcal{L}u$ by means of the Residue Theorem or another technique.

13. $\mathcal{L}u(s) = \frac{s}{(s^2 + A^2)^2}$

14. $\mathcal{L}u(s) = \frac{1}{(s-1)^2}$

15. $\mathcal{L}u(s) = \frac{1}{(s-1)^4}$

16. $\mathcal{L}u(s) = \frac{(s-2)e^{-s}}{(s^2 - 4s + 3)}$

17. $\mathcal{L}u(s) = \frac{1}{(1-is)}$

18. Establish the formula (6):

$$\frac{d}{ds}(\mathcal{L}u)(s) = -\mathcal{L}(tu(t))(s).$$

(Hint: The differentiation on the s variable can be brought inside the integral; this can be proved by techniques from real analysis.)

19. Show that if $u(t) = 0$ for $t \geq A$, $A > 0$, then $\mathcal{L}u(s)$ is an entire function of the complex variable s , and

$$(\mathcal{L}u)(s) = \sum_{n=0}^{\infty} s^n \left\{ \frac{(-1)^n}{n!} \int_0^A u(t) t^n dt \right\}$$

is its power series expansion about $s = 0$.

20. Find the Laplace transform of the function

$$u(t) = \begin{cases} 0, & t \leq 0 \\ \frac{1}{\sqrt{t}} e^{-c/t}, & t > 0, \end{cases}$$

where c is a positive constant.

$$\left(\text{Answer: } \mathcal{L}u(s) = \sqrt{\frac{\pi}{s}} e^{-2\sqrt{cs}}, s > 0. \right)$$

Further Integral Transforms*: The Gamma and Beta Functions

The **Gamma function** is defined by

$$\Gamma(z) = \int_0^\infty e^{-t} t^{z-1} dt, \quad \operatorname{Re} z > 0. \quad (8)$$

For $\operatorname{Re} z = x > 0$,

$$|e^{-t} t^{z-1}| = e^{-t} t^{x-1},$$

so the integral is absolutely convergent.

21. Show that $t^{z-1} = e^{(z-1)\log t}$ is analytic on the half-plane $\{z: \operatorname{Re} z > 0\}$.
22. Use Morera's Theorem and an interchange of the order of integration to show that $\Gamma(z)$ is analytic on the half-plane $\{z: \operatorname{Re} z > 0\}$.
23. Let $z = x$ be real and bigger than 1; use integration by parts to show that

$$\Gamma(x) = (x - 1)\Gamma(x - 1).$$

24. Conclude from Exercise 23 that $\Gamma(z) = (z - 1)\Gamma(z - 1)$ if $\operatorname{Re} z > 1$.
25. Show that $\Gamma(n) = (n - 1)!$ if n is a positive integer.
26. Derive the formula

$$\frac{\Gamma(x)\Gamma(y)}{\Gamma(x + y)} = \int_0^\infty \frac{t^{y-1}}{(1+t)^{x+y}} dt. \quad (9)$$

(Hint: Write

$$\Gamma(x)\Gamma(y) = \int_0^\infty t^{x-1} e^{-t} dt \int_0^\infty u^{y-1} e^{-u} du.$$

Now let $u = tv$ in the second integral; we obtain

$$\Gamma(x)\Gamma(y) = \int_0^\infty t^{x+y-1} \int_0^\infty e^{-t(v+1)} v^{y-1} dv dt.$$

Then set $s = (1 + v)t$ to obtain

$$\Gamma(x)\Gamma(y) = \Gamma(x + y) \int_0^\infty \frac{v^{y-1}}{(1+v)^{x+y}} dv.$$

The **Beta function** is defined by

$$\beta(x, y) = \frac{\Gamma(x)\Gamma(y)}{\Gamma(x + y)},$$

for x and y positive.

27. In the integral in (9), substitute $u = 1/(1 + t)$ and conclude that

$$\begin{aligned} \beta(x, y) &= \frac{\Gamma(x)\Gamma(y)}{\Gamma(x + y)} = \int_0^1 u^{x-1}(1-u)^{y-1} du \\ &= 2 \int_0^{\pi/2} (\sin \theta)^{2x-1} (\cos \theta)^{2y-1} d\theta. \end{aligned}$$

28. Set $x = y = \frac{1}{2}$ and use $\Gamma(1) = 1$ to get $\Gamma(\frac{1}{2}) = \sqrt{\pi}$.

29. Put $y = 1 - x$, $0 < x < 1$; this yields

$$\Gamma(x)\Gamma(1-x) = \int_0^\infty \frac{u^{x-1}}{1+u} du.$$

Now employ formula (8), Section 6, Chapter 2, to obtain

$$\Gamma(x)\Gamma(1-x) = \frac{\pi}{\sin(\pi x)}, \quad 0 < x < 1.$$

30. Use the conclusion of Exercise 29 to deduce that

$$\Gamma(z)\Gamma(1-z) = \frac{\pi}{\sin(\pi z)} \quad \text{for } 0 < \operatorname{Re} z < 1.$$

Now use the fact that $\Gamma(z)$ is analytic for $\operatorname{Re} z > 0$ to define $\Gamma(w)$ for those w with $\operatorname{Re} w \leq 0$:

$$\Gamma(w) = \frac{\pi}{\sin(\pi(1-w))} \cdot \frac{1}{\Gamma(1-w)}, \quad \operatorname{Re} w \leq 0.$$

More on Bessel Functions*

31. Let v be a real number, $v \geq 0$; show that one solution of the differential equation

$$z^2 f''(z) + zf'(z) + (z^2 - v^2)f(z) = 0$$

is

$$J_v(z) = \sum_{n=1}^{\infty} (-1)^n \frac{z^{2n+v}}{2^{2n+v} n! \Gamma(v+n+1)}, \quad \operatorname{Re} z > 0,$$

where Γ is the Gamma function. Note how this extends the results of Exercise 29, Section 4, Chapter 2.

5.4 Applications of the Laplace Transform to Differential Equations

The Laplace transform finds a major application in the solution of linear ordinary differential equations; it is frequently used to solve certain partial differential equations as well. Its advantages as a tool for solving such equations are twofold. First, the initial conditions and the nonhomogeneous term are incorporated from the start, so the solution is generated without any “constants of integration” that need further evaluation. Second, an ordinary differential equation is converted into an algebraic equation. The technique is illustrated by means of several examples.

Example 1 Solve the equation

$$u'(t) + u(t) = \begin{cases} 1, & \text{if } 0 < t < 1 \\ 0, & \text{if } 1 < t < \infty \end{cases}$$

with initial condition $u(0) = 1$.

Solution We take the Laplace transform of both sides of the equation; this yields

$$\mathcal{L}u'(s) + \mathcal{L}u(s) = \frac{1}{s}(1 - e^{-s}).$$

However, $\mathcal{L}u'(s) = -u(0) + s\mathcal{L}u(s)$, by Example 2, Section 3, so we obtain the equation

$$-1 + s\mathcal{L}u(s) + \mathcal{L}u(s) = \frac{1}{s}(1 - e^{-s}),$$

and hence

$$-1 + (s + 1)(\mathcal{L}u)(s) = \frac{1 - e^{-s}}{s}$$

or

$$\mathcal{L}u(s) = \frac{1}{s} - e^{-s} \left\{ \frac{1}{s} - \frac{1}{s+1} \right\}.$$

The function e^{-s} is the Laplace transform of the impulse function at $t_0 = 1$; the functions $1/s$ and $1/(s+1)$ are the Laplace transforms of 1 and e^{-t} , respectively. Hence, by the convolution result,

$$\frac{1}{s}e^{-s} \quad \text{and} \quad \frac{1}{s+1}e^{-s}$$

are the Laplace transforms of the convolution of the impulse function at $t_0 = 1$ with 1 and e^{-t} , respectively. These are, respectively,

$$\int_0^t 1 \delta(1-x) dx = \begin{cases} 0, & \text{if } 0 < t < 1 \\ 1, & \text{if } 1 \leq t < \infty \end{cases}$$

and

$$\int_0^t e^{-(t-x)} \delta(1-x) dx = \begin{cases} 0, & \text{if } 0 < t < 1 \\ e^{-(t-1)}, & \text{if } 1 \leq t < \infty. \end{cases}$$

Consequently,

$$u(t) = \begin{cases} 1, & \text{if } 0 < t < 1 \\ e^{-(t-1)}, & \text{if } 1 \leq t < \infty. \end{cases}$$
□

Note that the solution $u(t)$, although continuous on $[0, \infty)$, fails to be differentiable at $t_0 = 1$, exactly because the function on the right-hand side of the original equation is not continuous at $t_0 = 1$.

Example 2 Solve each of the following differential equations by means of the Laplace transform

(a) $u'' + u = f$; $u(0) = 1$, $u'(0) = 0$, and

$$f(t) = \begin{cases} 1, & 0 < t < \pi \\ 0, & \pi < t < 2\pi, \end{cases} \quad f(t) = f(t + 2\pi).$$

(b) $u'' + 2u' + 5u = \delta(t - \pi)$; $u(0) = u'(0) = 0$.

Solutions

(a) The Laplace transform of the 2π -periodic function $f(t)$ is

$$\mathcal{L}f(s) = \frac{\int_0^{2\pi} f(t)e^{-st} dt}{1 - e^{-2\pi s}} = \frac{1}{s} \frac{1 - e^{-\pi s}}{1 - e^{-2\pi s}} = \frac{1}{s(1 + e^{-\pi s})}.$$

Hence, the Laplace transform of the equation gives us

$$s^2 \mathcal{L}u(s) - s + \mathcal{L}u(s) = \frac{1}{s(1 + e^{-\pi s})}$$

or

$$\mathcal{L}u(s) = \frac{s}{1 + s^2} + \frac{1}{1 + s^2} \cdot \frac{1}{s(1 + e^{-\pi s})}.$$

The function $s/(1 + s^2)$ is the Laplace transform of $\cos t$, and $1/(1 + s^2)$ is the transform of $\sin t$. The Convolution Theorem, (8) of Section 1, then gives

$$\begin{aligned} u(t) &= \cos t + (f * \sin)(t) \\ &= \cos t + \int_0^t f(x) \sin(t-x) dx \\ &= \begin{cases} 1, & 0 < t \leq \pi \\ -\cos t, & \pi \leq t \leq 2\pi \\ 1 - 2\cos t, & 2\pi \leq t \leq 3\pi \\ -3\cos t, & 3\pi \leq t \leq 4\pi. \\ \text{etc.} & \end{cases} \end{aligned}$$

(b) Here, after taking Laplace transforms, we obtain

$$s^2 \mathcal{L}u(s) + 2s \mathcal{L}u(s) + 5 \mathcal{L}u(s) = e^{-\pi s}.$$

Thus,

$$\mathcal{L}u(s) = \frac{e^{-\pi s}}{s^2 + 2s + 5}.$$

The function $1/(s^2 + 2s + 5)$ is the Laplace transform of $\frac{1}{2}e^{-t} \sin 2t$, so the Convolution Theorem gives us

$$\begin{aligned} u(t) &= \int_0^t \delta(x - \pi) \frac{1}{2} e^{-(t-x)} \sin 2(t-x) dx \\ &= \begin{cases} 0, & 0 < t \leq \pi \\ \frac{1}{2} e^{-(t-\pi)} \sin 2t, & \pi \leq t < \infty. \end{cases} \end{aligned}$$
□

Example 3 Use the Laplace transform to solve the wave equation $u_{xx} = u_{tt}$, subject to

$$\begin{cases} u(0, t) = u(\pi, t) = 0, & t > 0 \\ u(x, 0) = \sin x, & 0 < x < \pi \\ u_t(x, 0) = 0. \end{cases}$$

Solution Let $U(x, s)$ be the Laplace transform of $u(x, t)$ on the t variable:

$$U(x, s) = \int_0^\infty u(x, t)e^{-st} dt.$$

Then $U(0, s) = U(\pi, s) = \int_0^\infty 0 \cdot e^{-st} dt = 0$ for each s . Treating U as a function of x alone (holding s fixed), we find that

$$\begin{aligned} \frac{d^2 U}{dx^2} &= \int_0^\infty u_{xx}(x, t)e^{-st} dt \\ &= \int_0^\infty u_{tt}(x, t)e^{-st} dt. \end{aligned}$$

We now integrate twice by parts and use the conditions $u_t(x, 0) = 0, u(x, 0) = \sin x$. The end result is the equation

$$\frac{d^2 U}{dx^2} = s^2 U - s \sin x, \quad 0 < x < \pi,$$

with

$$U(0) = U(\pi) = 0.$$

This is an elementary linear second-order equation whose solution is easily found to be

$$U(x, s) = \frac{s}{s^2 + 1} \sin x.$$

The term $\sin x$ is constant with respect to s , and $s/(s^2 + 1)$ is the Laplace transform of $\cos t$. Hence, the solution is

$$u(x, t) = \cos t \sin x. \quad \square$$

You may be aware that in the previous example, the original equation is easily solved by separation of variables. However, the example is designed to illustrate,

with a minimum of extraneous details, the use of the Laplace transform in solving partial differential equations. The following two examples are further illustrations of this technique—this time, of a more complicated nature.

Example 4 A string of length 1 is clamped at its left end and initially is straight and at rest. At time $t = 0$, the right end begins to move up and down, with height $\sin t$ at time t . Find the resulting displacement at position x , $0 \leq x \leq 1$, and time t , $t > 0$.

Solution Let $u(x, t)$ be the displacement at position x and time t , $0 \leq x \leq 1$, $t \geq 0$. The equation that determines u is the wave equation $u_{xx} = u_{tt}$, subject to

$$\begin{cases} u(x, 0) = u_t(x, 0) = 0 & 0 < x < 1 \\ u(0, t) = 0, & t > 0 \\ u(1, t) = \sin t, & t > 0. \end{cases}$$

Let $U(x, s)$ be the Laplace transform of $u(x, t)$ on the t variable:

$$U(x, s) = \int_0^\infty u(x, t)e^{-st} dt.$$

We treat U as a function of x alone and differentiate twice. This gives

$$\begin{aligned} \frac{d^2 U}{dx^2} &= \int_0^\infty u_{xx}(x, t)e^{-st} dt \\ &= \int_0^\infty u_{tt}(x, t)e^{-st} dt \\ &= s^2 \int_0^\infty u(x, t)e^{-st} dt, \end{aligned}$$

after two integrations by parts (or the use of the formula in Example 2, Section 3) and the use of the initial conditions $u(x, 0) = u_t(x, 0) = 0$. The boundary conditions on u yield

$$U(0) = \int_0^\infty u(0, t)e^{-st} dt = \int_0^\infty 0 \cdot e^{-st} dt = 0$$

and

$$U(1) = \int_0^\infty u(1, t)e^{-st} dt = \int_0^\infty (\sin t)e^{-st} dt = \frac{1}{1 + s^2}.$$

Hence, U satisfies the elementary second-order equation

$$\frac{d^2U}{dx^2} = s^2 U \quad \text{with } U(0) = 0, \quad U(1) = \frac{1}{1+s^2}.$$

This equation has solution

$$U(x, s) = \frac{1}{1+s^2} \frac{\sinh xs}{\sinh s}.$$

Our final task, then, is to invert U and recover u :

$$u(x, t) = \frac{1}{2\pi i} \int_{\sigma_0-i\infty}^{\sigma_0+i\infty} \frac{1}{1+s^2} \frac{\sinh xs}{\sinh s} e^{st} ds.$$

The function U has simple poles at $\pm i$ and $\pm in\pi$, $n = 1, 2, \dots$. The residues of the integrand $U(s, x)e^{st}$ are

$$\text{Res}(Ue^{st}; \pm i) = \pm \frac{1}{2i} \frac{\sin x}{\sin 1} e^{\pm it}$$

and

$$\text{Res}(Ue^{st}; in\pi) = \frac{i}{1 - \pi^2 n^2} \frac{\sin(\pi n x)}{(-1)^n} e^{in\pi t}, \quad n = \pm 1, \pm 2, \dots$$

Hence, the solution is

$$u(x, t) = \frac{\sin x}{\sin 1} \sin t + \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{\pi^2 n^2 - 1} \{\sin(\pi n x)\} \{\sin(\pi n t)\}. \quad \square$$

Example 5 A steel wire of length 1 is clamped at both ends. The wire is straight and at rest when a sinusoidal magnetic field is activated. Find the resulting displacement of the wire at each future time.

Solution Let $u(x, t)$, $0 \leq x \leq 1$, $0 \leq t$, be the displacement at the position x and the time t from the rest position. The partial differential equation that determines the function $u(x, t)$ is

$$u_{xx} = u_{tt} - \sin \omega t, \quad 0 < x < 1, \quad t > 0,$$

with boundary and initial conditions

$$u(0, t) = u(1, t) = 0, \quad t > 0 \quad (\text{the ends are clamped})$$

$$u(x, 0) = u_t(x, 0) = 0, \quad 0 < x < 1 \quad (\text{the wire is initially straight and at rest}).$$

Let $U(x, s)$ be the Laplace transform of $u(x, t)$ on the time variable t :

$$U(x, s) = \int_0^\infty u(x, t) e^{-st} dt.$$

We hold s fixed and treat U as a function of x . Then

$$\begin{aligned}\frac{d^2 U}{dx^2} &= \int_0^\infty u_{xx}(x, t) e^{-st} dt \\ &= \int_0^\infty u_{tt}(x, t) e^{-st} dt - \int_0^\infty (\sin \omega t) e^{-st} dt \\ &= s^2 U - \frac{\omega}{s^2 + \omega^2}.\end{aligned}$$

Furthermore, we have the boundary conditions $U(0) = U(1) = 0$. This is an ordinary differential equation for U whose solution is

$$\begin{aligned}U(x, s) &= \frac{\omega}{s^2(s^2 + \omega^2)} \frac{(e^s - e^{sx})(1 - e^{-sx})}{1 + e^s} \\ &= \frac{2\omega}{s^2(s^2 + \omega^2)} \frac{\sinh(sx/2) \sinh((s/2)(1-x))}{\cosh(s/2)}.\end{aligned}$$

The frequency ω of the magnetic field plays a critical role in the nature of the solution u . We shall first analyze the case when ω is not an odd integer multiple of π . We must find $u(x, t)$ from $U(x, s)$; that is, we must invert the Laplace transform. The Inversion Formula, (7) of Section 3, gives

$$u(x, t) = \frac{1}{2\pi i} \int_{\sigma-i\infty}^{\sigma+i\infty} \frac{\omega}{s^2(s^2 + \omega^2)} \frac{(e^s - e^{sx})(1 - e^{-sx})}{1 + e^s} e^{st} ds.$$

The integrand has simple poles at $s = \pm i\omega$ and at $s = \pm i\pi, \pm 3i\pi, \dots$; there is no pole at $s = 0$, since $(e^s - e^{sx})(1 - e^{-sx})$ has a double zero at $s = 0$. The residue at $s = ik\pi$, $k = \pm 1, \pm 3, \dots$, is

$$A_k = \frac{2i\omega}{\pi^4} \frac{\sin k\pi x}{k^2(k^2 - (\omega/\pi)^2)} e^{ik\pi t},$$

so

$$A_k + A_{-k} = -\frac{4\omega}{\pi^4} \frac{\sin k\pi x \sin k\pi t}{k^2(k^2 - (\omega/\pi)^2)}, \quad k = 1, 3, 5, \dots$$

The residues at $s = i\omega$ and $s = -i\omega$ sum to

$$B(x, t) = \frac{\sin \omega t}{\omega^2} \left\{ \frac{(e^{i\omega} - e^{i\omega x})(1 - e^{-i\omega x})}{1 + e^{i\omega}} \right\}.$$

This gives the solution

$$u(x, t) = B(x, t) - \frac{4\omega}{\pi^4} \sum_{k \text{ odd}} \frac{(\sin k\pi x)(\sin k\pi t)}{k^2 \left(k^2 - \left(\frac{\omega}{\pi} \right)^2 \right)}.$$

When ω is an odd integer multiple of π , there is a new difficulty; we shall examine the case when $\omega = \pi$. We are still faced with the job of inverting U :

$$u(x, t) = \frac{1}{2\pi i} \int_{\sigma-i\infty}^{\sigma+i\infty} \frac{\pi}{s^2(s^2 + \pi^2)} \frac{(e^s - e^{sx})(1 - e^{-sx})}{1 + e^s} e^{st} ds.$$

The integrand again has simple poles at $s = \pm 3i\pi, \pm 5i\pi, \dots$, and again there is no pole at $s = 0$. However, there are poles of order two at $s = \pm i\pi$. The residue at $s = ik\pi, k = \pm 3, \pm 5, \dots$, is

$$A_k = \frac{2i}{\pi^3} \frac{\sin k\pi x}{k^2(k^2 - 1)} e^{ik\pi t}, \quad k = \pm 3, \pm 5, \dots,$$

and, as before,

$$A_k + A_{-k} = -\frac{4}{\pi^3} \frac{\sin k\pi x \sin k\pi t}{k^2(k^2 - 1)}, \quad k = 3, 5, \dots.$$

The residue at both $i\pi$ and $-i\pi$ is more complicated, because these points are poles of order two. Diligent computation shows these residues to be

$$A_1 = te^{i\pi t} \frac{\sin \pi x}{\pi^2} + i \left[\frac{2\pi + (\cos \pi x)(4\pi x - 2\pi) - 10 \sin \pi x}{4\pi^4} \right]$$

and

$$A_{-1} = te^{-i\pi t} \frac{\sin \pi x}{\pi^2} - i \left[\frac{2\pi + (\cos \pi x)(4\pi x - 2\pi) - 10 \sin \pi x}{4\pi^4} \right].$$

Their sum is

$$A_1 + A_{-1} = \frac{2t \cos \pi t \sin \pi x}{\pi^2},$$

so the solution is given by

$$u(x, t) = \frac{2t \cos \pi t \sin \pi x}{\pi^2} - \frac{4}{\pi^3} \sum_{n=1}^{\infty} \frac{(\sin(2n+1)\pi x)(\sin(2n+1)\pi t)}{(2n+1)^2[(2n+1)^2 - 1]}.$$

Note that the second term of the solution remains bounded for all t , but the first term can become unboundedly large for large t . This is the phenomenon of **resonance**. \square

EXERCISES FOR SECTION 5.4

In Exercises 1 to 5, use the Laplace transform to solve the given differential equation.

1. $u'' - 2u' + 2u = 6e^{-t}; \quad u(0) = 0, \quad u'(0) = 1$
2. $u'' + 9u = \begin{cases} 1, & 0 < t < \pi \\ 0, & \pi < t < \infty \end{cases}; \quad u(0) = 1, \quad u'(0) = 0$
3. $u''(t) + tu'(t) + u(t) = 0; \quad u(0) = 1, \quad u'(0) = 0$
4. $u'' + 2u' + 2u = \delta(t - 2\pi); \quad u(0) = 1, \quad u'(0) = 0$
5. $u'' + u = \begin{cases} 0, & 0 \leq t \leq \pi \\ \cos t, & \pi < t < \infty \end{cases}; \quad u(0) = 0, \quad u'(0) = 1$

6. Solve the **Volterra*** integral equation

$$\phi(t) + \int_0^t k(t-s)\phi(s) ds = f(t)$$

for the unknown function ϕ by taking Laplace transforms. Express the answer in terms of the Laplace transform of ϕ ; f and k are assumed known.

In Exercises 7 to 11, use the Laplace transform to solve the given partial differential equation.

7. (The traveling wave) Solve the wave equation

$$u_{tt} = u_{xx}, \quad x > 0, t > 0,$$

subject to

$$\begin{cases} u(x, 0) = u_t(x, 0) = 0, & x > 0 \\ u(0, t) = f(t), & t \geq 0, f(0) = 0 \\ \lim_{x \rightarrow \infty} u(x, t) = 0, & t \geq 0. \end{cases}$$

8. (The falling string) Solve the modified wave equation

$$u_{tt} = u_{xx} - g, \quad x > 0, t > 0,$$

(g is a positive constant) subject to

$$\begin{cases} u(x, 0) = u_t(x, 0) = 0, & x > 0 \\ u(0, t) = 0, & t \geq 0 \\ \lim_{x \rightarrow \infty} u_x(x, t) = 0, & t \geq 0. \end{cases}$$

* Vito Volterra, 1860–1940.

9. Solve the wave equation $u_{tt} = u_{xx}$, subject to

$$\begin{cases} u(x, 0) = u_t(x, 0) = 0, & 0 < x < 1 \\ u(0, t) = 0, & t \geq 0 \\ u(1, t) = \sin \omega t, & t \geq 0. \end{cases}$$

The value of ω will affect the nature of the solution.

10. Solve the wave equation $u_{tt} = u_{xx}$, subject to

$$\begin{cases} u(x, 0) = u_t(x, 0) = 0, & 0 < x < 1 \\ u(0, t) = 0, & t \geq 0 \\ u_x(1, t) = c_0, & t \geq 0. \end{cases}$$

Here c_0 is a constant.

11. Solve the heat equation

$$\frac{\partial u}{\partial t} = k \frac{\partial^2 u}{\partial x^2}, \quad x > 0, t > 0,$$

subject to the conditions

$$u(x, 0) = 0, \quad x > 0$$

$$-k \frac{\partial u}{\partial x}(0, t) = \phi_0, \quad \phi_0 = \text{constant}, \quad t > 0$$

$$\lim_{x \rightarrow \infty} u(x, t) = 0, \quad t > 0.$$

To invert $U(x, s)$, you will need to use the conclusion of Exercise 20, Section 5.3.
 k is a constant.

5.5 The Z-Transform

The Z-transform provides a technique to deal with sequences (and solve difference equations involving sequences) in much the same way that the Laplace transform provides a technique to solve differential equations involving functions. The Z-transform is a valuable tool in areas such as numerical analysis, where a problem may involve solving a difference equation of discrete data.

Let $\{a_j\}$ be a sequence of complex numbers satisfying the growth condition

$$|a_j| \leq M r_0^j, \quad j = 0, 1, 2, \dots \quad (1)$$

for some positive numbers M and r_0 . The **Z-transform** of $\{a_j\}$ is the function

$$Z(\{a_j\}) = \sum_{j=0}^{\infty} \frac{a_j}{z^j}.$$

The series converges absolutely for all z with $|z| = r > r_0$, because

$$\left| \frac{a_j}{z^j} \right| = \frac{|a_j|}{r^j} \leq M \left(\frac{r_0}{r} \right)^j,$$

and the series $\sum (r_0/r)^j$ converges, since the ratio r_0/r is less than 1. Consequently, the Z-transform of $\{a_j\}$ is an analytic function on the exterior of the circle $|z| = r_0$, including at ∞ .

Example 1 Find the Z-transform of the sequence $\{a_j\}$ if $a_j = 3$ for all j .

Solution If $a_j = 3$ for all j , then, directly from the definition,

$$Z(\{3\}) = \sum_{j=0}^{\infty} \frac{3}{z^j} = \frac{3}{1 - \frac{1}{z}} = \frac{3z}{z - 1}.$$

This function is analytic outside the circle $|z| = 1$. □

Example 2 Find the Z-transform of the sequence $a_j = 1/j!$.

Solution Using the definition

$$Z\left(\left\{\frac{1}{j!}\right\}\right) = \sum_{j=0}^{\infty} \frac{1}{j! z^j} = e^{1/z}, \quad |z| > 0,$$

this function is analytic for all z with $|z| > 0$. □

Basic Properties of the Z-Transform

The Z-transform is a discrete version of the Laplace transform. To see this, let δ be a positive number and set $x_n = n\delta$, $n = 1, 2, 3, \dots$. The Laplace transform of a function u is given by the integral $\int_0^\infty u(t)e^{-st} dt$. This integral is approximated by the sum $\sum_{n=0}^{\infty} u(x_n) e^{-sx_n} \Delta x_n = \delta \sum_{n=0}^{\infty} u(n\delta) e^{-s\delta n}$. The function $z = F(s) = e^{s\delta}$ maps the half-plane $\operatorname{Re}(s) > a$ onto the exterior of the circle $|z| = e^{a\delta} = 1/r_0$. Set $a_n = \delta u(n\delta)$; the Laplace transform of u ,

$$(\mathcal{L}u)(s) = \int_0^\infty u(t)e^{-st} dt,$$

is then approximated by

$$\sum_0^{\infty} \frac{a_n}{z^n} = Z(\{a_n\}), \quad z = e^{s\delta}.$$

Note that the condition $|u(t)| \leq M e^{-at}$ is translated into the condition $|a_n| \leq \delta M r_0^n$,

$n = 0, 1, 2, \dots$. This is a reason for saying that the Z-transform is a “discrete Laplace transform.”

The Z-transform shares many of the general properties of the Laplace transform. First, it is linear:

If $\{a_j\}$ and $\{b_j\}$ are two sequences both satisfying (1) and if λ and μ are two complex numbers, then $Z(\{\lambda a_j + \mu b_j\}) = \lambda Z(\{a_j\}) + \mu Z(\{b_j\})$. (2)

Second, like the Laplace transform, the Z-transform turns a convolution into a product. Convolution is defined in this way. If $\{a_j\}$ and $\{b_j\}$ are two sequences of complex numbers, then the **convolution of $\{a_j\}$ and $\{b_j\}$** is the sequence $\{c_n\}$ given by

$$c_n = \sum_{k=0}^n a_k b_{n-k}, \quad n = 0, 1, 2, \dots . \quad (3)$$

We make note of the fact that if $\{a_k\}$ and $\{b_k\}$ both satisfy (1), then $|c_n| \leq MM'(n+1)r_0^n < M''r^n$ for each $r > r_0$. Hence, the Z-transform of $\{c_n\}$ is well-defined on the domain $|z| > r_0$.

The Z-transform of $\{c_n\}$ is given in the following:

If $\{c_n\}$ is the convolution of $\{a_j\}$ and $\{b_j\}$, then
 $Z(\{c_n\}) = Z(\{a_j\})Z(\{b_j\})$. (4)

The demonstration of (4) is easy:

$$\begin{aligned} Z(\{a_j\})Z(\{b_j\}) &= \left(\sum_{j=0}^{\infty} \frac{a_j}{z^j} \right) \left(\sum_{k=0}^{\infty} \frac{b_k}{z^k} \right) \\ &= \sum_{n=0}^{\infty} \frac{1}{z^n} \left(\sum_{k=0}^n a_k b_{n-k} \right) \\ &= \sum_{n=0}^{\infty} \frac{c_n}{z^n} = Z(\{c_n\}). \end{aligned}$$

The passage from the first line to the second occurs simply by multiplying out the two series and collecting equal powers of z (see Section 2, Chapter 2).

The convolution result in (4) is enormously useful; we give one elementary example here to illustrate this fact.

Example 3 Consider the (infinite) system of linear equations

$$b_0 = a_0$$

$$b_1 = a_1 - 2a_0$$

$$b_2 = a_2 - 2a_1 + a_0$$

$$b_3 = a_3 - 2a_2 + a_1$$

....

In general, $b_n = a_n - 2a_{n-1} + a_{n-2}$, $n \geq 2$. Solve this system for the sequence $\{a_j\}$ in terms of the sequence $\{b_j\}$.

Solution The system may be concisely expressed as

$$\{b_j\} = \{a_j\} * \{f_j\},$$

where

$$f_0 = 1, \quad f_1 = -2, \quad f_2 = 1, \quad \text{and } f_j = 0 \quad \text{for } j \geq 3.$$

Note that $\{b_n\}$ satisfies (1) with constant $(1 + (1/r_0))^2 M$. After taking Z-transforms,

$$\begin{aligned} Z(\{b_j\}) &= Z(\{a_j\})Z(\{f_j\}) = Z(\{a_j\}) \left(1 + \frac{-2}{z} + \frac{1}{z^2} \right) \\ &= \left(\frac{z-1}{z} \right)^2 Z(\{a_j\}). \end{aligned}$$

Division then yields this equation for the Z-transform of $\{a_j\}$:

$$Z(\{a_j\}) = \left(\frac{z}{z-1} \right)^2 Z(\{b_j\}).$$

The convolution result implies that

$$\{a_j\} = \{g_j\} * \{b_j\},$$

where $\{g_j\}$ is the sequence whose Z-transform is $(z/(z-1))^2$. To find $\{g_j\}$, we must expand $(z/(z-1))^2$ in powers of $1/z$:

$$\begin{aligned} \frac{z^2}{(z-1)^2} &= \left(1 + \frac{1}{z-1} \right)^2 = 1 + \frac{2}{z-1} + \frac{1}{(z-1)^2} \\ &= 1 + \frac{2}{z-1} - \left(\frac{1}{z-1} \right)' \\ &= 1 + 2 \left\{ \frac{1}{z} + \frac{1}{z^2} + \dots \right\} - \left\{ \frac{1}{z} + \frac{1}{z^2} + \dots \right\}' \\ &= 1 + \frac{2}{z} + \frac{2}{z^2} + \dots + \frac{1}{z^2} + \frac{2}{z^3} + \frac{3}{z^4} + \dots \\ &= 1 + \frac{2}{z} + \frac{3}{z^2} + \frac{4}{z^3} + \frac{5}{z^4} + \dots \end{aligned}$$

Hence, $g_j = j + 1$, $j = 0, 1, 2, \dots$, so

$$a_j = \sum_{n=0}^j (n+1)b_{j-n}, \quad j = 0, 1, 2, \dots$$

In particular, $a_0 = b_0$, $a_1 = b_1 + 2b_0$, $a_2 = b_2 + 2b_1 + 3b_0$, etc. \square

Shifting

If the sequence $\{a_j\}$ is shifted by one unit to form the new sequence $\{b_j\}$, $b_j = a_{j+1}$, $j = 0, 1, 2, \dots$, then

$$Z(\{b_j\}) = z[Z(\{a_j\}) - a_0]. \quad (5)$$

This follows because

$$\begin{aligned} Z(\{b_j\}) &= b_0 + \frac{b_1}{z} + \frac{b_2}{z^2} + \cdots = a_1 + \frac{a_2}{z} + \frac{a_3}{z^2} + \cdots \\ &= z \left[a_0 + \frac{a_1}{z} + \frac{a_2}{z^2} + \cdots - a_0 \right] \\ &= z[Z(\{a_j\}) - a_0]. \end{aligned}$$

More generally, if N is a fixed positive integer and

$$b_j = a_{j+N}, \quad j = 0, 1, 2, \dots,$$

then

$$Z(\{b_j\}) = z^N \left[Z(\{a_j\}) - \sum_{i=0}^{N-1} \frac{a_i}{z^i} \right]. \quad (6)$$

The result in (5) or (6) is referred to as the **Theorem on Shifting**.

The Z-transform is a particularly effective tool for studying the solution of a **linear difference equation with constant coefficients**. Such an equation has the form

$$\sum_{j=0}^P A_j y_{j+n} = \sum_{j=0}^Q B_j x_{j+n}, \quad n = 0, 1, 2, \dots. \quad (7)$$

Here P and Q are fixed nonnegative integers; A_0, \dots, A_P and B_0, \dots, B_Q are known real or complex numbers, the coefficients of the system. The numbers $\{x_j\}$ (which satisfy (1) for some r_0), called the **forcing function** or the **input**, are known. We wish to solve for the numbers $\{y_j\}$, the **response** or the **output of the system** to the forcing function. The numbers y_0, \dots, y_{P-1} are **initial conditions** and are assumed to be known as well. (If $P = 0$, then there are no initial conditions.)

We begin by taking the Z -transform of both sides of this equation. This yields

$$\sum_{j=0}^P A_j z^j \left[Y(z) - \sum_{k=0}^{j-1} y_k z^{-k} \right] = \sum_{j=0}^Q B_j z^j \left[X(z) - \sum_{k=0}^{j-1} x_k z^{-k} \right],$$

where

$$Y(z) = \sum_{j=0}^{\infty} \frac{y_j}{z^j}, \quad X(z) = \sum_{j=0}^{\infty} \frac{x_j}{z^j}$$

are the Z -transforms of $\{y_j\}$ and $\{x_j\}$, respectively. We now solve for $Y(z)$:

$$Y(z) = \frac{\sum_{j=0}^Q B_j z^j [X(z) - \sum_{k=0}^{j-1} x_k z^{-k}] + \sum_{j=0}^P A_j z^j \sum_{k=0}^{j-1} y_k z^{-k}}{\sum_{j=0}^P A_j z^j}. \quad (8)$$

The solution is completed by applying the inverse Z -transform or by expanding the right-hand side in powers of $1/z$ (which is the same thing).* One of the advantages of this form of the answer is that it prominently displays the denominator, which is critical in the determination of the stability of the system. This is discussed in detail in Section 5.5.1.

Example 4 Solve the linear difference equation

$$2y_n + y_{n+1} = x_n - x_{n+2}, \quad n = 0, 1, \dots, \quad y_0 = 1.$$

Solution Here $P = 1$ with $A_0 = 2$, and $A_1 = 1$; $Q = 2$ with $B_0 = 1$, $B_1 = 0$, and $B_2 = -1$. The formula for $Y(z)$ yields

$$\begin{aligned} Y(z) &= \frac{X(z) - z^2 \left[X(z) - \left(x_0 + \frac{x_1}{z} \right) \right] + 2 + z}{2 + z} \\ &= \frac{\left(x_0 + \frac{x_1}{z} + \frac{x_2}{z^2} + \dots \right) - \left(x_2 + \frac{x_3}{z} + \frac{x_4}{z^2} + \dots \right) + 2 + z}{2 + z}. \end{aligned}$$

* Here the material at the end of Section 5, Chapter 2, can be useful.

The expansion $1/(2 + z)$ in powers of $1/z$, valid for $|z| > 2$, is

$$\begin{aligned}\frac{1}{2+z} &= \left(\frac{1}{z}\right) \frac{1}{1+\frac{2}{z}} \\ &= \left(\frac{1}{z}\right) \left(1 - \frac{2}{z} + \left(\frac{2}{z}\right)^2 - \left(\frac{2}{z}\right)^3 + \dots\right) \\ &= \frac{1}{z} - \frac{2}{z^2} + \frac{4}{z^3} - \frac{8}{z^4} + \dots.\end{aligned}$$

This yields

$$\begin{aligned}Y(z) &= 1 + \left(\frac{1}{2+z}\right) \left(x_0 - x_2 + \frac{x_1 - x_3}{z} + \frac{x_2 - x_4}{z^2} + \dots\right) \\ &= 1 + \left(\frac{1}{z} - \frac{2}{z^2} + \frac{4}{z^3} - \frac{8}{z^4} + \dots\right) \left(x_0 - x_2 + \frac{x_1 - x_3}{z} + \frac{x_2 - x_4}{z^2} + \dots\right) \\ &= 1 + \frac{x_0 - x_2}{z} + \frac{-2(x_0 - x_2) + x_1 - x_3}{z^2} \\ &\quad + \frac{x_2 - x_4 - 2(x_1 - x_3) + 4(x_0 - x_2)}{z^3} + \dots.\end{aligned}$$

Hence, $y_1 = x_0 - x_2$, $y_2 = -2x_0 + x_1 + 2x_2 - x_3$, and in general,

$$y_n = \sum_{j=1}^n (-2)^{j-1} (x_{n-j} - x_{n+2-j}), \quad n = 1, 2, 3, \dots \quad \square$$

Example 5 Find the current i_k in each step of the ladder network shown in Figure 5.9.




Figure 5.9

Solution In the first loop

$$3Ri_0 - Ri_1 = V$$

and in the other loops

$$-Ri_k + 4Ri_{k+1} - Ri_{k+2} = 0.$$

Let $I(z) = \sum_{k=0}^{\infty} i_k/z^k$ be the Z-transform of $\{i_k\}$. Using the difference equations previously given and (6), we obtain

$$i_1 = 3i_0 - V/R$$

and

$$I(z) - 4z(I(z) - i_0) + z^2(I(z) - i_0 - i_1/z) = 0.$$

Upon substituting for i_1 and solving, we find that

$$I(z) = i_0 \frac{z^2 - z \left(1 + \frac{V}{i_0 R} \right)}{z^2 - 4z + 1}.$$

The numbers i_1, i_2, \dots are then obtained by expanding this rational function in powers of $1/z$. One way to accomplish this is by using equations (12) and (13) of Section 5, Chapter 2. We first write

$$I(z) = i_0 \left[1 + \frac{-1 + z \left(3 - \frac{V}{i_0 R} \right)}{z^2 - 4z + 1} \right].$$

The rational function

$$f(z) = \frac{-1 + z \left(3 - \frac{V}{i_0 R} \right)}{z^2 - 4z + 1}$$

has poles of order one at $z_1 = 2 - \sqrt{3}$ and $z_2 = 2 + \sqrt{3}$ with residues

$$r_1 = \text{Res}(f; z_1) = \frac{-1 + (2 - \sqrt{3}) \left(3 - \frac{V}{i_0 R} \right)}{-2\sqrt{3}}$$

and

$$r_2 = \text{Res}(f; z_2) = \frac{-1 + (2 + \sqrt{3}) \left(3 - \frac{V}{i_0 R} \right)}{2\sqrt{3}}.$$

Hence, for $|z| > z_2$, we obtain the expansion

$$I(z) = i_0 \left[1 + \sum_{k=1}^{\infty} \frac{i_k}{z_k} \right],$$

where

$$i_k = r_1 z_1^{k-1} + r_2 z_2^{k-1}, \quad k = 1, 2, 3, \dots. \quad \square$$

EXERCISES FOR SECTION 5.5

In Exercises 1 to 7, find the Z-transform of the given sequence $\{a_j\}$ and the region in which $Z(\{a_j\})$ is analytic.

$$1. a_j = \begin{cases} 1, & j \text{ even} \\ 0, & j \text{ odd} \end{cases}$$

$$2. a_j = \begin{cases} 0, & j \text{ even} \\ (-1)^k \frac{1}{(2k+1)!}, & j = 2k+1, k = 0, 1, 2, \dots \end{cases}$$

$$3. a_j = \frac{2^j}{j!}$$

$$4. a_j = \begin{cases} 0, & j \text{ odd} \\ \frac{1}{j!}, & j \text{ even} \end{cases}$$

$$5. a_j = \frac{1}{j+1}$$

$$6. \{a_j\} = \{b_j\} * \{b_j\}, \quad \text{where } b_j = 1 \text{ for all } j$$

$$7. a_j = \begin{cases} 4^j, & j \text{ even} \\ 3^{-j}, & j \text{ odd} \end{cases}$$

In Exercises 8 to 14, find the sequence $\{a_j\}$ given its Z-transform $Z(\{a_j\})$.

$$8. Z(\{a_j\}) = \cos \frac{1}{z}$$

$$9. Z(\{a_j\}) = \sinh \frac{2}{z}$$

$$10. Z(\{a_j\}) = \text{Log} \left(1 + \frac{1}{z} \right), \quad |z| > 1$$

$$11. Z(\{a_j\}) = z(e^{1/z} - 1)$$

$$12. Z(\{a_j\}) = (z^2 + z + 1) \left(\cos \left(\frac{1}{z} \right) - 1 \right)$$

$$13. Z(\{a_j\}) = \left(\frac{1}{z-1} \right)^2$$

$$14. Z(\{a_j\}) = \frac{1}{z^2 + z - 2}$$

15. Show that if $\{a_j\} * \{b_j\} = \{0\}$, then either $a_j = 0$ for all j or $b_j = 0$ for all j .
(Hint: The hypothesis gives

$$\left(\sum_{j=0}^{\infty} \frac{a_j}{z^j} \right) \left(\sum_{j=0}^{\infty} \frac{b_j}{z^j} \right) = 0 \quad \text{for all } z \text{ with } |z| > r_0^{-1}.$$

Hence, one or the other of the Z -transforms must be identically zero.)

16. Let $P(z) = A_0 + A_1 z + \cdots + A_N z^N$ be a polynomial of degree one or more, all of whose zeros lie within the disc $|z| < 1$. Let $\{y_j\}$ be any solution of the system.

$$\sum_{j=0}^N A_j y_{j+n} = x_n, \quad n = 0, 1, 2, \dots$$

Show that if the forcing term $\{x_n\}$ is **bounded** (that is, $|x_n| \leq M$ for all n), then $Y(z)$, the Z -transform of $\{y_j\}$, is analytic in the region $\{z : |z| > 1\}$.

Solve the following linear systems.

$$17. y_n - y_{n+1} = x_n + x_{n+1} + x_{n+2}; \quad y_0 = 1$$

$$18. y_n + 2y_{n+1} + y_{n+2} = x_n - x_{n+1}; \quad y_0 = 1, y_1 = 0$$

$$19. y_n + y_{n+2} = x_n; \quad y_0 = 0, y_1 = 1$$

$$20. y_n - y_{n+1} - 4y_{n+2} + 4y_{n+3} = x_n + x_{n+1}; \quad y_0 = y_1 = y_2 = 0$$

$$21. y_n - 2y_{n+1} = x_n - 2x_{n+1} + x_{n+2}; \quad y_0 = 1$$

5.5.1 The Stability of a Discrete Linear System*

Let us look back at the linear difference equation in (7) and make three simplifying assumptions: first, that the initial conditions are all zero, $y_0 = \cdots = y_{p-1} = 0$; second, that in the forcing term (or input), we have $x_0 = \cdots = x_{q-1} = 0$; and third, that $P \geq Q$. Then we obtain a simple explicit representation for the Z -transform, $Y(z)$, of the solution given by (8):

$$Y(z) = F(z)X(z),$$

where

$$F(z) = \frac{\sum_0^Q B_j z^j}{\sum_0^P A_j z^j} = \sum_{j=0}^{\infty} \frac{f_j}{z^j}. \quad (9)$$

Equivalently,

$$y_n = \sum_{k=0}^n f_k x_{n-k}, \quad n = 0, 1, 2, \dots \quad (10)$$

That is, the output $\{y_n\}$ of the system is given in terms of the input $\{x_j\}$ by the convolution equation (10). The rational function F given in (9) is called the **system transfer function**. The linear system is **stable** if bounded input always produces bounded output. The stability of the system obviously depends on the system transfer function F , and we shall discover a bit later in this section exactly what properties of F correspond to stability. But first we shall work out a few examples to get an understanding of some of the mechanics of stability.

Example 1 $x_0 = y_0, x_1 - x_0 = y_1, x_2 - x_1 = y_2, \dots$, and, in general,

$$x_{n+1} - x_n = y_{n+1}, \quad n = 0, 1, 2, \dots$$

This is the system (7) with $B_0 = -1, B_1 = 1$, and $A_0 = 0, A_1 = 1$. Hence, the system transfer function is

$$F(z) = \frac{-1 + z}{z} = \frac{z - 1}{z} = 1 - \frac{1}{z}.$$

The system is clearly stable, since $|x_n| \leq M$ for all n gives $|y_n| \leq 2M$ for all n . \square

Example 2 $x_0 = y_0, y_1 - y_0 = x_1, y_2 - y_1 = x_2$, and, in general,

$$y_{n+1} - y_n = x_{n+1}, \quad n = 0, 1, 2, \dots$$

This is the system (7) with $B_0 = 0, B_1 = 1$, and $A_0 = -1, A_1 = 1$. The system transfer function is

$$F(z) = \frac{z}{-1 + z} = 1 + \frac{1}{z} + \frac{1}{z^2} + \dots,$$

and the solution is

$$y_n = x_0 + x_1 + \dots + x_n, \quad n = 0, 1, 2, \dots$$

This system is clearly *not* stable, since the bounded input $x_n = 1$ for all n produce the unbounded output $y_n = n + 1$ for all n . \square

Example 3 $x_0 = y_0, y_1 - \frac{1}{2}y_0 = x_1, y_2 - \frac{1}{2}y_1 = x_2$, and in general

$$y_{n+1} - \frac{1}{2}y_n = x_{n+1}, \quad n = 0, 1, 2, \dots$$

Once again comparing with (7), we find that $A_1 = 1, A_0 = -\frac{1}{2}$, and $B_0 = 0, B_1 = 1$.

The system transfer function here is

$$F(z) = \frac{z}{-\frac{1}{2} + z} = \frac{2z}{2z - 1} = 1 + \frac{1}{2z} + \frac{1}{4z^2} + \dots$$

The solution is $y_0 = x_0$ and

$$y_n = x_n + \frac{1}{2}x_{n-1} + \frac{1}{4}x_{n-2} + \dots + \frac{1}{2^n}x_0, \quad n = 0, 1, 2, \dots$$

The system is stable, since $|x_n| \leq M$ for all n yields

$$|y_n| \leq M + \frac{1}{2}M + \frac{1}{4}M + \dots + \frac{1}{2^n}M < 2M \quad \text{for all } n. \quad \square$$

Example 4 $x_0 = y_0, y_1 + y_0 = x_1, y_2 + y_1 = x_2$; generally,

$$y_{n+1} + y_n = x_{n+1}, \quad n = 0, 1, 2, \dots$$

The system corresponds to (7) with $A_0 = A_1 = 1$ and $B_0 = 0, B_1 = 1$. The system transfer function is

$$F(z) = \frac{z}{1+z} = 1 - \frac{1}{z} + \frac{1}{z^2} - \frac{1}{z^3} + \dots$$

and the solution is

$$y_n = x_n - x_{n-1} + x_{n-2} - \dots + (-1)^n x_0, \quad n = 0, 1, 2, \dots$$

This particular system is not stable, since the bounded input $x_j = (-1)^{j+1}, j = 0, 1, 2, \dots$ produces the unbounded output $y_n = (-1)^{n+1}(n+1)$. \square

Conditions for Stability

The fundamental question on stability is completely answered in the following theorem.

THEOREM 1 The linear difference equation (10) with system transfer function F given in (9) is stable if and only if all the poles of the rational function F lie in the open disc $\{z: |z| < 1\}$.

Proof Suppose first that all the poles of F lie in the open disc $\{z: |z| < 1\}$. Then there is a number $r_0 < 1$ such that all the poles of F lie in the disc $\{z: |z| \leq r_0\}$; this is because F has only a finite number of poles. Hence, F is analytic on the set $\{z: |z| > r_0\}$, including at ∞ , and so has a power series valid there, centered

at ∞ :

$$F(z) = A_0 + \frac{A_1}{z} + \frac{A_2}{z^2} + \cdots, \quad |z| > r_0.$$

However, F has the power series (in $1/z$) representation (9), so $A_j = f_j$ for $j = 0, 1, 2, \dots$. Thus, the series $\sum f_j/z^j$ converges exterior to the circle $|z| = r_0$ and, by the basic properties of power series, is absolutely convergent on any circle of radius larger than r_0 . In particular, this holds on the circle $|z| = 1$. Consequently,

$$S = \sum_{n=0}^{\infty} |f_n|$$

is finite. Suppose now that $|x_k| \leq M$ for all k . Then

$$\begin{aligned} |y_n| &= \left| \sum_{k=0}^n x_k f_{n-k} \right| \leq \sum_{k=0}^n |x_k| |f_{n-k}| \\ &\leq M \sum_{k=0}^n |f_{n-k}| \leq MS, \end{aligned}$$

for each $n = 0, 1, 2, \dots$. Therefore, the linear system is stable.

Conversely, suppose that the linear system (10) is stable. We note first that if $\{y_n\}$ is a bounded sequence (say, $|y_n| \leq M$ for all n), then the Z -transform $Y(z)$ of $\{y_n\}$ satisfies

$$\begin{aligned} |Y(z)| &= \left| \sum_{n=0}^{\infty} \frac{y_n}{z^n} \right| \leq \sum_{n=0}^{\infty} \frac{|y_n|}{|z|^n} \\ &\leq M \sum_{n=0}^{\infty} \frac{1}{|z|^n} = M \frac{|z|}{|z| - 1}, \quad |z| > 1. \end{aligned}$$

In particular, the function $Y(z)$ is analytic on the region $\{z: |z| > 1\}$ because its power series in $1/z$ is absolutely convergent there. Since

$$Y(z) = F(z)X(z),$$

we see immediately that F has no poles in the region $\{z: |z| > 1\}$: just choose $x_0 = 1$ and $x_j = 0$, $j = 1, 2, \dots$. Moreover, if F had a pole at a point λ , $|\lambda| = 1$, then

$$F(z) = \frac{G(z)}{(z - \lambda)^m},$$

where G is analytic near λ , $G(\lambda) \neq 0$, and $m \geq 1$ (see Section 5, Chapter 2). Let

$x_k = \lambda^k, k = 0, 1, 2, \dots$; then

$$X(z) = 1 + \frac{\lambda}{z} + \frac{\lambda^2}{z^2} + \cdots$$

$$= \frac{z}{z - \lambda}.$$

The rule for the Z-transform of the output $\{y_n\}$ corresponding to this bounded input yields

$$Y(z) = F(z)X(z), \quad |z| > 1$$

$$= G(z)z \frac{1}{(z - \lambda)^{m+1}}, \quad |z| > 1, z \text{ near } \lambda.$$

Hence,

$$|Y(z)| \geq \frac{\frac{1}{2}|G(\lambda)|}{|z - \lambda|^{m+1}}, \quad z \text{ near } \lambda, |z| > 1.$$

By setting $z = \lambda t, t > 1$, we get

$$|Y(\lambda t)| \geq \frac{\frac{1}{2}|G(\lambda)|}{(t - 1)^{m+1}}.$$

This contradicts the estimate

$$|Y(\lambda t)| \leq \frac{Mt}{t - 1},$$

obtained above, if t is near 1. We conclude that F has no poles on $|z| = 1$, so all the poles of F lie within the disc $\{z: |z| < 1\}$. ■

Examine Examples 1 to 4 again in light of Theorem 1 and note the role that the location of the pole played.

EXERCISES FOR SECTION 5.5.1

- Let p be a polynomial of degree N . Show that all the zeros of p lie within the disc $|z| < 1$ if and only if all the zeros of rational function

$$R(w) = p\left(\frac{w + 1}{w - 1}\right)$$

lie in the left half-plane; that is, satisfy $\operatorname{Re} w < 0$.

2. Let p be a polynomial of degree N ; show that

$$q(w) = (w - 1)^N p\left(\frac{w + 1}{w - 1}\right)$$

is a polynomial of degree N or less in w .

3. Let p and q be related as in Exercise 2. Find $q(w)$ if $p(z)$ is
- (a) $p(z) = Az + B$; (b) $p(z) = 27z^4 - 24z^3 + 18z^2 - 8z + 3$;
 - (c) $p(z) = z^2 - z$; (d) $p(z) = z^3 - 3z^2 + 3z - 2$.
4. Let q be given as in Exercise 2, and suppose that p and q have the same degree. Show that all the zeros of q lie in the left half-plane if and only if all the zeros of p lie in the disc $\{z: |z| < 1\}$. (Hint: Combine Exercises 1 and 2.) What can happen to the degree of q ? Show that degree q is less than degree p if and only if p has a zero at $z = 1$.
5. Use Exercise 4 and, if necessary, the Routh–Hurwitz criterion (Theorem 1, Section 1.1, Chapter 3) to determine whether all the zeros of the following polynomials lie within the disc $\{z: |z| < 1\}$.
- (a) $p(z) = 5z^3 + 3z^2 - z + 1$; (b) $p(z) = 10z^3 - 4z^2 + 2z$;
 - (c) $p(z) = 27z^4 - 24z^3 + 18z^2 - 8z + 3$;
 - (d) $p(z) = 59z^3 - 89z^2 + 37z - 7$;
 - (e) $p(z) = 10z^3 + 6z^2 - 20z + 6$.
6. Test the following linear systems for stability:
- (a) $y_n - y_{n+1} - 4y_{n+2} + 4y_{n+3} = x_n - x_{n+1}$
 - (b) $y_n - y_{n+1} + 3y_{n+2} + 5y_{n+3} = x_n + 2x_{n+1} - x_{n+3}$
 - (c) $y_n - y_{n+1} - 4y_{n+2} - 4y_{n+3} = x_n$
 - (d) $3y_n - 8y_{n+1} + 18y_{n+2} - 24y_{n+3} + 27y_{n+4} = x_n + x_{n+1} + x_{n+2}$
 - (e) $y_n - 2y_{n+1} = x_n + x_{n+1} - x_{n+2}$
 - (f) $2y_n + y_{n+1} = x_n + 2x_{n+1}$
 - (g) $y_n - 4y_{n+2} = 3x_n + 2x_{n+1}$
 - (h) $6y_n - 20y_{n+1} + 6y_{n+2} + 10y_{n+3} = x_n + x_{n+1}$

Further Reading

More on the Fourier transform can be found in Titchmarsh, E. C. *The Fourier transform*. New York: Oxford University Press, 1937; and Carslaw, H. S. *Theory of Fourier series and integrals*. New York: Dover, 1950.

The Laplace (and Fourier) transform is covered in Sneddon, I. N. *Fourier transforms*. New York: McGraw-Hill, 1951; and LePage, W. R. *Complex variables and the Laplace transform for engineers*. New York: Dover, 1980.

Several of the applications in the text come from Churchill, R. V. *Operational mathematics*. New York: McGraw-Hill, 1958; and Powers, D. L. *Boundary-value problems*. New York: Academic Press, 1972.

A good reference for the Z-transform, including many applications, is Jury, E. J. *Theory and applications of the Z-transform*. New York: John Wiley, 1964.

This page is intentionally left blank

Locating the Zeros of a Polynomial

A polynomial of degree d has precisely d zeros, counting multiplicities, in the complex plane. This appendix contains several theorems that help to locate these zeros. The first result is the “grandfather” of the others.

THEOREM 1 If every zero of the polynomial p lies in the half-plane $\operatorname{Re}(Az + B) > 0$, then so does every zero of its derivative p' .

Proof This result is called the **Gauss^{*}–Lucas[†] Theorem**; the proof is not at all difficult. Write

$$p(z) = a \prod_1^N (z - z_j)^{m_j}, \quad a \neq 0,$$

where z_1, \dots, z_N are distinct points and m_1, \dots, m_N are positive integers. A simple computation yields

$$\begin{aligned} \frac{p'(z)}{p(z)} &= \sum_1^N \frac{m_j}{z - z_j} \\ &= \sum_1^N \frac{m_j(\bar{z} - \bar{z}_j)}{|z - z_j|^2}. \end{aligned}$$

Clearly, we need only look at those points b that are zeros of p' but not zeros of p . If b is such a point, then

$$0 = \sum_1^N \frac{m_j}{|b - z_j|^2} (\bar{b} - \bar{z}_j) = \bar{b}c_0 + c_1,$$

where $c_0 = \sum_1^N m_j |b - z_j|^{-2}$ is positive and $c_1 = -\sum_1^N m_j \bar{z}_j |b - z_j|^{-2}$. Hence,

* Karl Friedrich Gauss, 1777–1855.

† Francois Édouard Anatole Lucas, 1842–1891.

$$\begin{aligned}
 \operatorname{Re}(Ab + B) &= \operatorname{Re}\left(-A\frac{\bar{c}_1}{c_0} + B\right) = \frac{1}{c_0} \operatorname{Re}(-A\bar{c}_1 + Bc_0) \\
 &= \frac{1}{c_0} \operatorname{Re} \sum_1^N m_j \left(\frac{Az_j + B}{|b - z_j|^2} \right) \\
 &= \frac{1}{c_0} \sum_1^N \frac{m_j}{|b - z_j|^2} \operatorname{Re}(Az_j + B) > 0. \quad \blacksquare
 \end{aligned}$$

There is another classical result on the location of the zeros of a polynomial, this time ensuring that they lie in a disc. This theorem is known as the **Eneström–Kakeya Theorem**.

THEOREM 2 Suppose that $0 < a_0 < a_1 < \cdots < a_n$. Then all the zeros of the polynomial $p(z) = a_0 + a_1z + \cdots + a_nz^n$ lie in the disc $\{z : |z| < 1\}$.

Proof Set $q(z) = z^n p(1/z) = a_n + a_{n-1}z + \cdots + a_0z^n$; we shall show that all the zeros of q lie in the set $|z| > 1$. We begin by noting that

$$\begin{aligned}
 (1 - z)q(z) &= a_n + z(a_{n-1} - a_n) + z^2(a_{n-2} - a_{n-1}) \\
 &\quad + \cdots + z^n(a_0 - a_1) - a_0z^{n+1}.
 \end{aligned}$$

Thus, for $|z| \leq 1$,

$$\begin{aligned}
 |1 - z||q(z)| &\geq a_n - |(a_n - a_{n-1})z + \cdots + (a_1 - a_0)z^n + a_0z^{n+1}| \\
 &\geq a_n - (a_n - a_{n-1}) - \cdots - (a_1 - a_0) - a_0 = 0.
 \end{aligned}$$

Furthermore, equality holds in the second inequality only if $|z| = 1$ and *all* of the terms $(a_j - a_{j-1})z^{n-j}$ have the same argument, $j = 0, 1, \dots, n$; this can only happen when $z = 1$. But at $z = 1$, $p(1) = a_0 + a_1 + \cdots + a_n > 0$. ■

The Zeros of Cubic and Quartic Polynomials

The familiar quadratic formula for finding the zeros of a second-degree polynomial has an extension (albeit not so simple), which yields the zeros of a third- or fourth-degree polynomial. These formulas (or better, *techniques*) date back to the sixteenth century.

Cubic Polynomials

We wish to find the roots of the equation

$$z^3 + Az^2 + Bz + C = 0. \quad (1)$$

We begin by making the substitution $z = w + d$ and then choosing d such that the

resulting quadratic term vanishes. Specifically, if $z = w + d$, we obtain

$$w^3 + (A + 3d)w^2 + (3d^2 + 2Ad + B)w + (d^3 + Ad^2 + Bd + C) = 0.$$

The choice $d = -A/3$ yields

$$w^3 + aw + b = 0 \quad (2)$$

with

$$a = B - \frac{1}{3}A^2 \quad \text{and} \quad b = \frac{2}{27}A^3 - \frac{1}{3}AB + C.$$

If it happens that $a = 0$, then the equation becomes

$$w^3 + b = 0,$$

and the roots are obtained easily. If $a \neq 0$, then let λ be a square root of $-a/3$, and set $w = \lambda q$. This substitution changes (2) to the equation

$$q^3 - 3q + \beta = 0, \quad (3)$$

where $\beta = b\lambda^{-3}$. Long ago, someone noticed that the substitution $q = p + 1/p$ changes (3) into the very simple equation

$$p^6 + \beta p^3 + 1 = 0. \quad (4)$$

Indeed, this was the whole point of putting (2) into the form of (3). Equation (4) is nothing but a quadratic in the variable p^3 . Its zeros produce the q 's that solve (3) and so in turn produce the zeros of (2) and finally the zeros of (1). This three-step process yields the zeros of the general cubic polynomial (1).

Example 1 The cubic $z^3 + 6z^2 + 12z + 35 = 0$ becomes

$$w^3 + 27 = 0 \quad (5)$$

after the substitution $z = w - 2$. The roots of (5) are

$$w = -3 \quad \text{and} \quad \frac{1}{2}(3 \pm i\sqrt{27}),$$

so the roots of the original equation are

$$z = -5 \quad \text{and} \quad \frac{1}{2}(-1 \pm i\sqrt{27}). \quad \square$$

Example 2 The cubic $z^3 + 6z^2 + 4 = 0$ is changed to the cubic

$$w^3 - 12w + 20 = 0$$

by the substitution $z = w - 2$. With $a = -12$ and $b = 20$, we find that $\lambda = 2$ and $\beta = 5/2$, so we must solve

$$p^6 - \left(\frac{5}{2}\right)p^3 + 1 = 0.$$

This gives $p^3 = 2$ or $p^3 = 1/2$; we work with the cube roots of 2, which are

$$\gamma_1 = (2)^{1/3}, \quad \gamma_2 = (2)^{-2/3}(-1 + i\sqrt{3}), \quad \gamma_3 = (2)^{-2/3}(-1 - i\sqrt{3}).$$

Hence, the roots of $w^3 - 12w + 20 = 0$ are

$$w_1 = \gamma_1 + \frac{1}{\gamma_1}, \quad w_2 = \gamma_2 + \frac{1}{\gamma_2}, \quad w_3 = \gamma_3 + \frac{1}{\gamma_3},$$

so the roots of the original equation are $z_j = w_j - 2$, $j = 1, 2, 3$. □

Quartic Polynomials

We wish to find the roots of $z^4 + Az^3 + Bz^2 + Cz + D = 0$. We rewrite this as

$$\left(z^2 + \frac{A}{2}z\right)^2 = \left(\frac{A^2}{4} - B\right)z^2 - Cz - D. \quad (6)$$

If the right-hand side of (6) is a perfect square (say, $(Ez + F)^2$), then we obtain two quadratic equations:

$$z^2 + \frac{A}{2}z = Ez + F$$

and

$$z^2 + \frac{A}{2}z = -(Ez + F).$$

Each of these equations has two roots, and these four numbers are the roots we seek. If the right-hand side of (6) is not a perfect square, we add $w^2 + 2(z^2 + (A/2)z)w$ to both sides of (6). This yields

$$\left(z^2 + \frac{A}{2}z + w\right)^2 = \left(\frac{A^2}{4} - B + 2w\right)z^2 + (Aw - C)z + (w^2 - D). \quad (7)$$

We now choose w such that the right-hand side of (7) is a perfect square. This will

be so if the discriminant of the right-hand side vanishes—that is, if

$$(Aw - C)^2 - 4\left(\frac{A^2}{4} - B + 2w\right)(w^2 - D) = 0. \quad (8)$$

This is a cubic in w and can be solved as outlined above. We can then take any root w^* of this cubic, substitute it into (7), and then solve the resulting equation:

$$\left(z^2 + \frac{A}{2}z + w^*\right)^2 = \left(\frac{A^2}{4} - B + 2w^*\right)\left(z + \frac{Aw^* - C}{2\left(\frac{A^2}{4}\right) - B + 2w^*}\right)^2. \quad (9)$$

This will yield the four zeros of $z^4 + Az^3 + Bz^2 + Cz + D$.

Example 3 Find the zeros of $z^4 + z^2 - 4iz - 4$.

Solution Here $A = 0$, so we obtain

$$\begin{aligned}(z^2)^2 &= z^4 = -z^2 + 4iz + 4 \\ &= (iz + 2)^2.\end{aligned}$$

Hence, we must solve the two quadratic equations

$$z^2 = iz + 2 \quad \text{and} \quad z^2 = -iz - 2.$$

The solutions of these are

$$\frac{i \pm \sqrt{7}}{2} \quad \text{and} \quad \frac{-i \pm 3i}{2},$$

respectively. Hence,

$$i, \quad -2i, \quad \frac{i + \sqrt{7}}{2}, \quad \text{and} \quad \frac{i - \sqrt{7}}{2}$$

are the four zeros of $z^4 + z^2 - 4iz - 4$. □

Example 4 Find the zeros of $z^4 + 2z^3 + 4z^2 + 2z + 3$.

Solution Here $((A^2/4) - B)z^2 - Cz - D = -3z^2 - 2z - 3$ is not a perfect square, so we must find a zero of the cubic $(2w - 2)^2 - 4(-3 + 2w)(w^2 - 3)$. One root of this is $w^* = 2$. Putting this into (9), we obtain

$$(z^2 + z + 2)^2 = (1)\left(z + \frac{2}{2}\right)^2.$$

This yields the two equations

$$z^2 + z + 2 = z + 1 \quad \text{and} \quad z^2 + z + 2 = -z - 1.$$

The first equation has roots $\pm i$ and the second has roots $-1 \pm i\sqrt{2}$. These four numbers are the zeros of the original polynomial. \square

A final remark The reader may well draw the conclusion that the zeros of any polynomial may eventually be found by a succession of the steps that work for the quadratic, cubic, and quartic cases—that is, by a sequence of substitutions and extractions of roots. Quite surprisingly this is not the case if the degree is five or more. In fact, one of the more profound mathematical results of the early nineteenth century was that there is no formula of this type that will solve every fifth- (or higher) degree polynomial.

Sturm Sequences

Let p be a polynomial with real coefficients, and let $a < b$ be two points on the real line with $p(a) \neq 0$ and $p(b) \neq 0$. We shall describe a simple technique that counts the number of distinct zeros of p in the interval (a, b) ; the zeros are counted *without multiplicity*.

We begin in a somewhat more general context. Let p_1, p_2, \dots, p_m be polynomials with real coefficients satisfying these three conditions:

- (1) if $p_k(x_0) = 0$ for some $x_0 \in (a, b)$ and some k , $2 \leq k \leq m - 1$, then $p_{k-1}(x_0)p_{k+1}(x_0) < 0$.
- (2) $p_m(x) \neq 0$ for $a < x < b$.
- (3) $p_1(a) \neq 0, p_1(b) \neq 0$; if $p_1(x_0) = 0$ for some $x_0 \in (a, b)$, then the sign of $p_1(x)p_2(x)$ changes from minus to plus as x passes through x_0 from left to right.

Let $S(x)$ be the number of changes of sign in the sequence $\{p_1(x), p_2(x), \dots, p_m(x)\}$ for $a \leq x \leq b$. A zero of some $p_k(x)$, $1 \leq k \leq m - 1$, can be counted either as a plus or a minus without affecting $S(x)$, because of conditions (1) and (3).

THEOREM 3 $S(a) - S(b)$ is the number of zeros of p_1 on the interval (a, b) . The zeros are counted without multiplicity.

Proof Suppose that x is a point at which no p_k is zero. Then by the continuity of p_j , there is an interval $(x - \delta, x + \delta)$ in which no p_j is zero, so $S(x)$ does not change in $(x - \delta, x + \delta)$. Hence, any change in $S(x)$ must occur at a zero of some p_k .

If $p_k(x_0) = 0$ for $k = 2, \dots, m - 1$, then condition (1) shows that the number of sign changes in the triple $p_{k-1}(x), p_k(x), p_{k+1}(x)$ remains exactly one for all x in some small interval about x_0 . Consequently, the only changes in $S(x)$ can occur, if at all, at points x_0 , where $p_1(x_0) = 0$. By condition (3), however, we see that $S(x)$ decreases by exactly 1 as x passes through a zero of p_1 from left to right. Thus, the theorem is proved. ■

To fit the theorem to our context, we take $p_1 = p$ and $p_2 = p'$, the derivative of p . Simple calculus shows that condition (3) holds; that is, that the sign of pp' changes from minus to plus as x passes through a zero of p from left to right. The remainder of the sequence p_3, \dots, p_m are found by long division. Write $p_1 = q_1 p_2 - p_3$, where q_1 is a polynomial and $\text{degree}(p_3) < \text{degree}(p_2)$. Continuing $p_2 = q_2 p_3 - p_4$, where $\text{degree}(p_4) < \text{degree}(p_3)$, and so on until the remainder is a polynomial of degree zero—that is, a constant. If the remainder is nonzero, then p_m is this remainder, and (2) holds. If the remainder is zero, then p_m is the previous remainder, which, therefore, divides p_{m-1} exactly. There is just one point that deserves mention: It may happen that p_m does not have degree zero; indeed, p_m may have zeros in $[a, b]$. If this is the case, it is relatively easy to check that p_m actually divides all the preceding polynomials $p_{m-1}, p_{m-2}, \dots, p_2, p_1$. Thus, p_m appears as a factor in all the terms of the sequence $\{p_1, p_2, \dots, p_m\}$; then the value of $S(a) - S(b)$ is just the same as it is for the sequence

$$\left\{ \frac{p_1}{p_m}, \frac{p_2}{p_m}, \dots, \frac{p_{m-1}}{p_m}, 1 \right\}.$$

Further Reading

An enormous amount of material on the location of zeros of polynomials is in Marden, M. *The geometry of zeros of polynomials*. Providence, R.I.: American Mathematical Society, 1966.









A Table of Conformal Mappings

$z = x + iy$, $w = u + iv$. Lowercase letters are mapped to uppercase letters, cross-hatched regions onto shaded regions.

z -plane	Mapping	w -plane
	$w = e^z$	

The vertical segment $x = x_0$, $0 \leq y \leq \pi$ is mapped to the semicircle $u^2 + v^2 = e^{2x_0}$, $v \geq 0$.











	$w = z^\alpha$ $0 < \alpha$	
	$w = \sin\left(\frac{\pi z}{2\alpha}\right)$	

z -plane	Mapping	w -plane
	$w = \sin z$	
The horizontal line $y = y_0$ goes to the ellipse $\frac{u^2}{\cosh^2 y_0} + \frac{v^2}{\sinh^2 y_0} = 1$.		
	$w = \log z$	
	$w = z^2$	
The vertical line $x = x_0$ is mapped to the parabola $u = x_0^2 - \frac{v^2}{4x_0^2}$, $x_0 \neq 0$.		
	$w = z^2$	

The hyperbola $xy = \alpha$ is mapped to the horizontal line $v = \frac{\alpha}{2}$. The hyperbola $x^2 - y^2 = \beta$, $\beta > 0$, is mapped to the vertical line $u = \beta$.

<i>z</i> -plane	Mapping	<i>w</i> -plane
	$w = \frac{1}{z}$	
	$w = \frac{1}{z}$	
	$w = i \left(\frac{1+z}{1-z} \right)$ $z = \frac{i-w}{i+w}$ is the inverse mapping.	
	$w = i \left(\frac{1-z}{1+z} \right)$	
	$w = -i \frac{z^2 + 2iz + 1}{z^2 - 2iz + 1}$	

z -plane	Mapping	w -plane
A diagram of the complex plane showing the unit disk $ z < 1$. The origin is labeled 0. The real axis is labeled x , and the imaginary axis is labeled y . Points i and $-i$ are marked on the y -axis. Point a is marked in the first quadrant. A radius vector from the origin to point a is labeled a .	$w = \frac{a-z}{1-\bar{a}z}$ $z = \frac{a-w}{1-\bar{a}w}$ is the inverse mapping. 0 and a are interchanged.	A diagram of the complex plane showing the unit disk $ w < 1$. The origin is labeled 0. The real axis is labeled u , and the imaginary axis is labeled v . Points i and $-i$ are marked on the v -axis. Point a is marked in the second quadrant. A radius vector from the origin to point a is labeled a .
A diagram of the complex plane showing a horizontal strip $b < y < a$ between two parallel lines at $y=b$ and $y=a$. The real axis is labeled x , and the imaginary axis is labeled y . Points c and d are marked on the x -axis. The width of the strip is labeled h .	$w = \cosh\left(\frac{\pi z}{h}\right)$	A diagram of the complex plane showing a horizontal strip $-1 < u < +1$ between two parallel lines at $u=-1$ and $u=+1$. The real axis is labeled u , and the imaginary axis is labeled v . Points A , B , C , and D are marked on the u -axis. The width of the strip is labeled $+1 - (-1) = 2$.
A diagram of the complex plane showing a quarter-disk in the first quadrant bounded by the x -axis, y -axis, and a semicircle from $(1,0)$ to $(0,i)$. The origin is labeled 0. The real axis is labeled x , and the imaginary axis is labeled y . Points c , d , and e are marked on the x -axis. Points i and $-i$ are marked on the y -axis. Points a and b are marked on the x -axis to the left of the origin.	$w = \frac{\alpha}{2} \left(z + \frac{1}{z} \right)$	A diagram of the complex plane showing a quarter-disk in the first quadrant bounded by the u -axis, v -axis, and a semicircle from $(\alpha,0)$ to $(0,\alpha)$. The origin is labeled 0. The real axis is labeled u , and the imaginary axis is labeled v . Points A , B , C , D , and E are marked on the u -axis. Points $-\alpha$ and α are marked on the v -axis.
A diagram of the complex plane showing a sector $\alpha < \theta < \pi$ in the second quadrant. The origin is labeled 0. The real axis is labeled x , and the imaginary axis is labeled y . Points c , d , and e are marked on the x -axis. Points f and i are marked on the y -axis. Points a and b are marked on the x -axis to the left of the origin. A small circle of radius r is shown near the origin.	$w = z + \frac{1}{z}$	A diagram of the complex plane showing a sector $\alpha < \theta < \pi$ in the second quadrant. The origin is labeled 0. The real axis is labeled u , and the imaginary axis is labeled v . Points A , B , C , D , and E are marked on the u -axis. Points F and i are marked on the v -axis.
The circle $ z = \alpha$ is mapped onto the ellipse $\frac{u^2}{\left(\alpha + \frac{1}{\alpha}\right)^2} + \frac{v^2}{\left(\alpha - \frac{1}{\alpha}\right)^2} = 1$.		
A diagram of the complex plane showing the upper half-plane $y > 0$. The real axis is labeled x , and the imaginary axis is labeled y . Points $i\pi$ and $-i\pi$ are marked on the y -axis.	$w = z + e^z$	A diagram of the complex plane showing the upper half-plane $v > 0$. The real axis is labeled u , and the imaginary axis is labeled v . Points $-1 + i\pi$ and $-1 - i\pi$ are marked on the v -axis.

z -plane	Mapping	w -plane
	$w = \frac{\alpha}{\pi} [(z^2 - 1)^{1/2} + \operatorname{Log}(z + (z^2 - 1)^{1/2})]$	
	$w = \frac{2\alpha i}{\pi} [z(1 - z^2)^{1/2} + \arcsin z]$	
	$w = \frac{2\alpha}{\pi} \left[(z^2 - 1)^{1/2} + \arcsin \frac{1}{z} \right]$	
	$w = \alpha(z^2 - 1)^{1/2}$	
	$w = \frac{i\sigma_0}{\pi} \left\{ \operatorname{Log} \left(\frac{1 + \zeta}{1 - \zeta} \right) - i\alpha \operatorname{Log} \left(\frac{1 + i\alpha\zeta}{1 - i\alpha\zeta} \right) \right\}$ $\alpha = \frac{\tau_0}{\sigma_0}; \quad \zeta^2 = \frac{z + 1}{z - \alpha^2}$	

z-plane	Mapping	w-plane
	$w = 2\sqrt{z+1} + \operatorname{Log}\left(\frac{\sqrt{z+1}-1}{\sqrt{z+1}+1}\right)$	
 $-1 < a < b < 1, \quad a + b \neq 0$	$w = \frac{z - \lambda}{1 - \bar{\lambda}z}, \quad \text{where}$ $\lambda = \frac{1 + ab - \sqrt{(1 - a^2)(1 - b^2)}}{a + b}$ and $R = \frac{1 - ab - \sqrt{(1 - a^2)(1 - b^2)}}{b - a}.$	

A Table of Laplace Transforms

$u(x); \quad u(x) = 0 \text{ if } x < 0$	$(\mathcal{L}u)(s)$
1. e^{ax}	$\frac{1}{s - a}$
2. x^m	$\frac{m!}{s^{m+1}}, \quad m = 0, 1, 2, \dots$
3. $\sin bx$	$\frac{b}{b^2 + s^2}$
4. $\cos bx$	$\frac{s}{b^2 + s^2}$
5. $\sinh bx$	$\frac{b}{s^2 - b^2}$
6. $\cosh bx$	$\frac{s}{s^2 - b^2}$
7. $e^{ax} \sin bx$	$\frac{b}{(s - a)^2 + b^2}$
8. $e^{ax} \cos bx$	$\frac{s - a}{(s - a)^2 + b^2}$
9. $x \sin bx$	$\frac{2sb}{(s^2 + b^2)^2}$
10. $x \cos bx$	$\frac{s^2 - b^2}{(s^2 + b^2)^2}$
11. $\begin{cases} 1, & 0 \leq x \leq \sigma \\ 0, & \sigma < x \end{cases}$	$\frac{1 - e^{-s\sigma}}{s}$
12. Impulse function for σ	$e^{-s\sigma}$
13. x^p	$\frac{\Gamma(p + 1)}{s^{p+1}}, \quad p > -1$
14. $\begin{cases} 1, & 0 < x < \sigma \\ -1, & \sigma < x < 2\sigma \\ 0, & 2\sigma < x \end{cases}$	$\frac{(1 - e^{-s\sigma})^2}{s}$
15. $\frac{\cos(b\sqrt{x})}{\sqrt{x}}$	$\sqrt{\frac{\pi}{s}} \exp\left(-\frac{b^2}{4s}\right)$
16. $\frac{e^{-b^2/x}}{\sqrt{x}}$	$\sqrt{\frac{\pi}{s}} e^{-2b\sqrt{s}}, \quad b \geq 0$
17. $\sin(b\sqrt{x})$	$\frac{1}{2}b\sqrt{\pi} s^{-3/2} e^{-b^2/4s}$

This page is intentionally left blank

Solutions to Odd-Numbered Exercises

Chapter 1

Section 1.1

1. (a) $7 - i$ (b) $-i$ (c) $-3 + 4i$ (d) $4 - 12i$ (e) $4/25$
(f) $-i$ (g) $18 + 18i$
3. (a) circle of radius 3 centered at 0 (b) circle of radius 1 centered at 2
(c) circle of radius 2 centered at -2
(d) perpendicular bisector of the line segment joining 2 to -2
(e) the two points $1 \pm \sqrt{2}$ (f) the line $y = x$ (g) the line $y = -x$
5. (a) $\sqrt{2} \left(\cos \frac{3\pi}{4} + i \sin \frac{3\pi}{4} \right)$ (b) $2 \left(\cos \frac{\pi}{3} + i \sin \frac{\pi}{3} \right)$
(c) $1 \left(\cos \left(-\frac{\pi}{2} \right) + i \sin \left(-\frac{\pi}{2} \right) \right)$ (d) $5(\cos \theta + i \sin \theta)$ where $\tan \theta = -4/3$
(e) 5 (f) $\sqrt{6}(\cos \theta_0 + i \sin \theta_0)$, where $\theta_0 = \arctan(-1/\sqrt{5})$
(g) $2\sqrt{2} \left(\cos \frac{5\pi}{4} + i \sin \frac{5\pi}{4} \right)$ (h) $1 \left(\cos \left(\frac{-\pi}{4} \right) + i \sin \left(\frac{-\pi}{4} \right) \right)$
(i) $1(\cos \pi + i \sin \pi)$
7. The roots are $z_1 = (-b + \sqrt{b^2 - 4ac})/2a$ and $z_2 = (-b - \sqrt{b^2 - 4ac})/2a$. These are complex conjugates because $b^2 < 4ac$.
9. $|z| = 1$ iff $|z|^2 = 1$ iff $z\bar{z} = 1$ iff $\bar{z} = 1/z$
11. $|z + w|^2 - |z - w|^2 = |z|^2 + 2 \operatorname{Re}(z\bar{w}) + |w|^2 - (|z|^2 - 2 \operatorname{Re}(z\bar{w}) + |w|^2)$
 $= 4 \operatorname{Re}(z\bar{w})$
13. (a) Yes (b) No (c) Yes
15. The lengths squared of the legs are $|z|^2, |w|^2, |z - w|^2$. These are equal iff $\{|z|^2 = |w|^2 = |z - w|^2\}$ iff $\{|z|^2 = |w|^2 = |z|^2 - 2 \operatorname{Re}(z\bar{w}) + |w|^2\}$ iff $\{|z|^2 = |w|^2 = 2 \operatorname{Re}(z\bar{w})\}$.

$$\begin{aligned}
17. \quad \left| \frac{z}{w} \right|^2 &= \left| \frac{(xs + yt) + i(ys - xt)}{s^2 + t^2} \right|^2 = \frac{(xs + yt)^2 + (ys - xt)^2}{(s^2 + t^2)^2} \\
&= \frac{x^2 s^2 + y^2 t^2 + y^2 s^2 + x^2 t^2}{(s^2 + t^2)^2} = \frac{(x^2 + y^2)(s^2 + t^2)}{(s^2 + t^2)^2} = \frac{x^2 + y^2}{s^2 + t^2} \\
&= \frac{|z|^2}{|w|^2}.
\end{aligned}$$

$$19. \quad \cos n\theta + i \sin n\theta = (\cos \theta + i \sin \theta)^n = \sum_{k=0}^n \binom{n}{k} (\cos \theta)^{n-k} (i \sin \theta)^k$$

For k even, there is no factor of i ; for k odd, there is a factor of i .

21. For all complex λ ,

$$\begin{aligned}
0 &\leq \sum_{j=1}^n |a_j - \lambda b_j|^2 = \sum_1^n |a_j|^2 - 2 \operatorname{Re} \bar{\lambda} \sum_1^n a_j \bar{b}_j + |\lambda|^2 \sum_1^n |b_j|^2 \\
&= B - 2 \operatorname{Re} \bar{\lambda} A + |\lambda|^2 C.
\end{aligned}$$

Apply the results of Exercise 20.

23. Equality iff $|a_j - \lambda b_j|^2 = 0$ for some λ and for all j iff $a_j = \lambda b_j$, $j = 1, \dots, n$.

Section 1.1.1

1. $(-z)^2 = ((-x)^2 - (-y)^2, 2(-x)(-y)) = (x^2 - y^2, 2xy) = z^2$.
3. $(0, 1) = z^2 = (x^2 - y^2, 2xy)$, so $x^2 = y^2$ and $2xy = 1$. If $x = -y$, then $-2x^2 = 1$, which is impossible. If $x = y$, then $2x^2 = 1$. Hence, $x = y = \pm 1/\sqrt{2}$.
7. $z_1 z_2 = (x_1 x_2 - y_1 y_2, x_1 y_2 + x_2 y_1)$. Hence, $|z_1 z_2|^2 = (x_1 x_2 - y_1 y_2)^2 + (x_1 y_2 + x_2 y_1)^2 = x_1^2 x_2^2 + y_1^2 y_2^2 + x_1^2 y_2^2 + x_2^2 y_1^2 = (x_1^2 + y_1^2)(x_2^2 + y_2^2) = |z_1|^2 |z_2|^2$.
9. If $z_1 z_2 = 0$, then $0 = |z_1 z_2| = |z_1| |z_2|$. Hence, $|z_1| = 0$ or $|z_2| = 0$.

Section 1.2

- | | |
|---|--|
| 1. The y -axis | 3. The line $y = 4x + 6$ |
| 5. The ellipse $1 = \frac{4x^2}{25} + \frac{4y^2}{9}$ | 7. The hyperbola $4 = x^2 - y^2$ |
| 9. The points $z = \pm 1$ | 11. $ z - (4 + i) = 2$ |
| 13. $\operatorname{Re}(z + 3) = 0$ | 15. $\left z - \left(\frac{1+i}{2} \right) \right = \frac{1}{\sqrt{2}}$ |
| 17. $\operatorname{Re}((2 - i)z - 1) = 0$ | |
| 19. $ z - p = cx$ iff $x^2 + y^2 - 2px + p^2 = c^2 x^2$ | |
| (a) $0 < c < 1$ gives $\left(x - \frac{p}{1 - c^2} \right)^2 + \frac{y^2}{1 - c^2} = \frac{p^2 c^2}{(1 - c^2)^2}$, an ellipse. | |
| (b) $c = 1$ gives $2px = y^2 + p^2$, a parabola. | |
| (c) $1 < c < \infty$ gives $\left(x + \frac{p}{c^2 - 1} \right)^2 = p^2 \left[1 + \frac{1}{(c^2 - 1)^2} \right] + \frac{1}{c^2 - 1} y^2$, a hyperbola. | |

21. (a) $|z - \alpha| < |1 - \bar{\alpha}z|$ iff $|z - \alpha|^2 < |1 - \bar{\alpha}z|^2$ iff $|z|^2 + |\alpha|^2 < 1 + |\alpha|^2|z|^2$ iff $|\alpha|^2(1 - |z|^2) < 1 - |z|^2$ iff $1 - |z|^2 > 0$, since $0 < |\alpha| < 1$. The others are similar.

23. $z = \cos\left(\frac{\pi}{10} + \frac{2}{5}\pi k\right) + i \sin\left(\frac{\pi}{10} + \frac{2}{5}\pi k\right), \quad k = 0, 1, 2, 3, 4$

25. $z = \cos\left(\frac{\pi}{8} + \frac{\pi}{4}k\right) + i \sin\left(\frac{\pi}{8} + \frac{\pi}{4}k\right), \quad k = 0, 1, \dots, 7$

27. Let $n = 2m + 1$ and let r_0 be the positive number with $r_0^{2m+1} = |w|$. The roots of $z^n = w$ are


$$z = r_0 \left\{ \cos\left(\frac{\pi}{2m+1} + \frac{2\pi k}{2m+1}\right) + i \sin\left(\frac{\pi}{2m+1} + \frac{2\pi k}{2m+1}\right) \right\}, \quad k = 0, 1, \dots, 2m.$$

The choice $k = m$ gives a negative real root.

29. The roots of $z^2 + bz + c = 0$ are

$$z_1 = \frac{-b + \sqrt{b^2 - 4c}}{2} \quad \text{and} \quad z_2 = \frac{-b - \sqrt{b^2 - 4c}}{2}.$$

z_1 and z_2 are complex conjugates iff $0 = \operatorname{Im}(z_1 + z_2) = -\operatorname{Im} b$ and $0 = \operatorname{Re}(z_1 - z_2) = \operatorname{Re}(\sqrt{b^2 - 4c})$. These give the conclusions.



$$\begin{aligned} x &= \pi - 2\theta + \pi - 2\psi = 2\pi - 2(\theta + \psi). \\ \text{Hence, } \theta + \psi &= \pi - x/2, \text{ a quantity independent of } P. \end{aligned}$$

33. The translation by β of C is the circle centered at $z_0 + \beta$ with the same radius. If the line \mathcal{L} is $\{z: \operatorname{Re}(Az + B) = 0\}$, then the translation of \mathcal{L} by β is the line $\{w: \operatorname{Re}(Aw + B - A\beta) = 0\}$.

35. $L = \{x + ia: -\infty < x < \infty\}; \left| \frac{1}{x + ia} + \frac{i}{2a} \right| = \left| \frac{2a + ix - a}{(2a)(x + ia)} \right| = \frac{1}{2a}$

37. $C = \{z: |z - c| = r\} = \{c + re^{i\theta}: 0 \leq \theta < 2\pi\}.$

$$\left| \frac{1}{z} - \frac{c}{c^2 - r^2} \right| = \left| \frac{1}{c + re^{i\theta}} - \frac{c}{c^2 - r^2} \right| = \left| \frac{c^2 - r^2 - c(c + re^{i\theta})}{(c^2 - r^2)(c + re^{i\theta})} \right| = \frac{r}{c^2 - r^2}$$

Section 1.3

- (a) interior = $\{x > 2 \text{ and } y < 4\}$;
boundary = $\{x + 4i: 2 \leq x < \infty\} \cup \{2 + iy: -\infty < y \leq 4\}$
(b) closed (c) interior is connected
- (a) interior = $\{x^2 < y\}$; boundary = $\{x^2 = y\}$
(b) open (c) interior is connected

5. (a) interior = $\{x^2 + y^2 > 2\}$; boundary = $\{x^2 + y^2 = 2\}$
 (b) closed (c) interior is connected
7. (a) interior = $\{x < 0 \text{ and } |z+1| > 1\}$;
 boundary = $\{iy: -\infty < y < \infty\} \cup \{|z+1| = 1\}$
 (b) neither open nor closed (c) interior is connected
9. Let $\alpha = |\alpha|e^{i\theta_0}$
- (a) $\{\alpha z + \beta: z \text{ in 1st quadrant}\} = \left\{ \beta + se^{i\psi}: 0 < s < \infty, \theta_0 < \psi < \theta_0 + \frac{\pi}{2} \right\}$
 (b) $\{\beta + se^{i\psi}: 0 < s < \infty, \theta_0 < \psi < \theta_0 + \pi\}$
 (c) $\{\beta + se^{i\psi}: 0 \leq s < R|\alpha|; 0 \leq \psi \leq 2\pi\}$
11. A, C, D, E, G , and H are unbounded.
13. Suppose that D_1, D_2 are open. Let $p \in D_1 \cap D_2$. There is an $r_1 > 0$ such that $\{|z-p| < r_1\} \subset D_1$; likewise, there is an $r_2 > 0$ such that $\{|z-p| < r_2\} \subset D_2$. Let $r = \min(r_1, r_2)$. Then $\{|z-p| < r\}$ lies inside both D_1 and D_2 . Hence, $D_1 \cap D_2$ is open. Next, let C_1, C_2 be closed. Then $\mathbb{C} \setminus (C_1 \cup C_2) = (\mathbb{C} \setminus C_1) \cap (\mathbb{C} \setminus C_2)$ and this is open, being the intersection of two open sets. Hence, $C_1 \cup C_2$ is closed. Likewise, $\mathbb{C} \setminus (C_1 \cap C_2) = (\mathbb{C} \setminus C_1) \cup (\mathbb{C} \setminus C_2)$, and this is open, being the union of two open sets. Hence, $C_1 \cap C_2$ is closed.
17. The ray $\{tz_0: 0 \leq t < \infty\}$ is closed, so D is open. Any two points p and q in D can be joined by the union of one, two, or three line segments, in turn parallel or perpendicular to the ray.
19. (a) Star-shaped with respect to 0 (b) not star-shaped (c) not star-shaped
 (d) star-shaped with respect to any x in $(1, \underline{\infty})$
21. If $p' \in C$ and $|z_0 - p'| \leq r$, then the segment pp' lies in C and also in the set $\{|z_0 - z| < r\}$. Thus, there is a point of C closer to z_0 than p , a contradiction.
25. Use the definition of the boundary B of D to show that the complement of B is open.
27. Use 25 and the definition of the boundary.
29. Use the definition of "interior point" of a set.

Section 1.4

1. 0 3. 0 5. No limit exists.
 7. 0 9. 2
 11. 1 at $z_0 = 8$; no limit exists at $8 + i$.
 13. 0 15. Continuous everywhere
 17. Discontinuous only at all points of the line $y = x$
 19. Discontinuous only at all points of the circle $|z| = 1$, except $z = 1$
 21. 0 23. 4
 25. f, g continuous at z_0 means $\lim_{z \rightarrow z_0} f(z) = f(z_0), \lim_{z \rightarrow z_0} g(z) = g(z_0)$. Hence,
 $\lim_{z \rightarrow z_0} (f(z) + g(z)) = f(z_0) + g(z_0)$, since the limit of the sum is the sum of the limits.
 Likewise, fg and $1/g$ are continuous at z_0 .
 27. Let $w = f(z)$; then $\lim_{z \rightarrow z_0} f(z) = f(z_0) = w_0$, so $\lim_{z \rightarrow z_0} g(f(z)) = \lim_{w \rightarrow w_0} g(w) = g(w_0) = g(f(z_0))$.
 29. Outline. Fix any point z_0 of D and let $c = u(z_0)$. Let z_1 be any point of D that may be joined to z_0 by a line segment

$$L = \{z(t) = (1-t)z_0 + tz_1, \quad 0 \leq t \leq 1\};$$

see Exercise 20, Section 2. Let

$$A = \{t \in [0, 1], \quad u(z(t)) = c\}.$$

Since u is equal to c on some disc centered at z_0 , the interval $[0, \delta)$ lies in A for some $\delta > 0$; δ is the radius of the disc. Since u is continuous at the point $z(\delta)$, we also have $\delta \in A$. However, u is constant on some disc centered at $z(\delta)$, so u is equal to c on this (second) disc. Hence, there is a $\delta_1 > \delta$, so $[0, \delta_1)$ lies in A . Continuing in this fashion, it becomes clear that A contains all points of $[0, 1]$, and so $u(z_1) = c$, also. If z_2 may be reached from z_1 by a single line segment, the same argument shows that $u(z_2) = c$. Since a finite number of line segments may be used to join z_0 to any point of D , it follows that $u(z) = c$ for all z in D .

31. Converges, since $\left| \frac{1+2i}{6} \right| = \frac{\sqrt{5}}{6} < 1$.
33. Diverges, since $\left| \frac{2+i}{\sqrt{5}} \right| = 1$, so terms do not go to 0.
35. Converges (ratio test).
37. Converges; use the technique of Example 29.
39. Converges (ratio test).
41. Use the hint and the fact that $|z| < 1$.

Section 1.5

1. $\frac{1+i}{\sqrt{2}}$
3. $\log 2 + i\left(\frac{\pi}{3} + 2k\pi\right), \quad k = 0, \pm 1, \dots$
5. $\exp\left(-\frac{\pi}{4} + 2k\pi\right)[\cos(\log \sqrt{2}) + i \sin(\log \sqrt{2})], \quad k = 0, \pm 1, \pm 2, \dots$
7. $\frac{1}{2} - i\frac{\sqrt{3}}{2}$
9. $\log 4\sqrt{2} - \frac{i\pi}{4}$
11. $\cos\left(\pi\sqrt{3}\left[\frac{1}{2} + 2k\right]\right) + i \sin\left(\pi\sqrt{3}\left[\frac{1}{2} + 2k\right]\right), \quad k = 0, \pm 1, \dots$
13. $4\left\{\log \sqrt{2} + i\left(-\frac{\pi}{4} + 2k\pi\right)\right\}, \quad k = 0, \pm 1, \dots$
15. $\log(e^z) = w$ or $e^z = e^w$. So all solutions are $z = w + 2\pi ik, \quad k = 0, \pm 1, \dots$
17. $0 = \cos z = \cos x \cosh y - i \sin x \sinh y$ iff $[0 = \cos x \cosh y \text{ and } 0 = \sin x \sinh y]$
iff $[0 = \cos x \text{ and } 0 = \sin x \sinh y]$ iff $[x = \pi/2 + \pi n, n = 0, \pm 1, \dots, \text{and } \sinh y = 0]$
iff $[x = \pi/2 + \pi n, n = 0, \pm 1, \dots, \text{and } y = 0]$. Likewise, $\sin z = 0$ only if $z = n\pi, n = 0, \pm 1, \dots$
19. $\cos(iy) = \frac{1}{2}(e^{iy} + e^{-iy}) = \frac{1}{2}(e^{-y} + e^y) \rightarrow \infty$ as $y \rightarrow \infty$. Furthermore, $|\cos(x+iy)| = |\frac{1}{2}(e^{i(x+iy)} + e^{-i(x+iy)})| \leq \frac{1}{2}(e^{-y} + e^y) \leq e^y$ if $y \geq 0$. Likewise, $|\sin(x+iy)| \leq e^y$ if $y \geq 0$.
23. On the line $\operatorname{Im} z = \pi/2$, the function e^z has the values $e^{x+iy/2} = e^x e^{iy/2} = ie^x$ and e^z assumes all values in $(0, \infty)$. Likewise, e^z maps to line $\operatorname{Im} z = -\pi/2$ onto the negative imaginary axis.
25. Follow the hint.
27. Use the principal branch of $\log z$.
29. Let $\zeta = -i \log(iz + \sqrt{1 - z^2})$; then

$$\begin{aligned}\sin \zeta &= \frac{1}{2i}(e^{i\zeta} - e^{-i\zeta}) = \frac{1}{2i}(\exp(\log[iz + \sqrt{1-z^2}]) - \exp(-\log[iz + \sqrt{1-z^2}])) \\&= \frac{1}{2i} \left(iz + \sqrt{1-z^2} - \frac{1}{iz + \sqrt{1-z^2}} \right) = \frac{1}{2i} \left[\frac{-z^2 + 2iz\sqrt{1-z^2} + 1-z^2 - 1}{iz + \sqrt{1-z^2}} \right] \\&= \frac{1}{2i} \left[\frac{2iz\sqrt{1-z^2} - 2z^2}{iz + \sqrt{1-z^2}} \right] = \frac{z(\sqrt{1-z^2} + iz)}{iz + \sqrt{1-z^2}} = z.\end{aligned}$$

Section 1.6

1. 0 3. $\frac{20}{3}(1+i)$ 5. $\frac{1}{2}(i-1)$

7. $\int_{\gamma_1} z^2 dz = -\frac{2}{3}$; $\int_{\gamma_2} z^2 dz = -\frac{2}{3}$. They are equal because $\gamma = \gamma_1 - \gamma_2$ is a closed curve and $\int_{\gamma} z^2 dz = 0$, by Example 10. However, $\int_{\gamma_1} \bar{z} dz = i\pi$, while $\int_{\gamma_2} \bar{z} dz = -i\pi$.

9. (a) For $k \neq 0$, $\int_0^{2\pi} e^{ik\theta} d\theta = \frac{1}{ik} e^{ik\theta} \Big|_{\theta=0}^{\theta=2\pi} = \frac{1}{ik} [1 - 1] = 0$.

11. $\int_{\gamma} f(x) dz = \int_{\gamma} (u + iv)(dx + i dy) = \int_{\gamma} u dx - v dy + i \int_{\gamma} v dx + u dy$. Also,

$$\begin{aligned}\iint_{\Omega} \left(\frac{\partial f}{\partial x} + i \frac{\partial f}{\partial y} \right) dx dy &= \iint_{\Omega} \left(\frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} + i \left[\frac{\partial u}{\partial y} + i \frac{\partial v}{\partial y} \right] \right) dx dy \\&= \iint_{\Omega} \left(\frac{\partial u}{\partial x} - \frac{\partial v}{\partial y} \right) dx dy + i \iint_{\Omega} \left(\frac{\partial v}{\partial x} + \frac{\partial u}{\partial y} \right) dx dy.\end{aligned}$$

13. In (8), replace u with $-v \frac{\partial u}{\partial y}$ and replace v with $v \frac{\partial u}{\partial x}$.

15. Establish the estimate: $\left| \int_{\gamma_R} \frac{u(z)}{(z - z_0)^2} dz \right| \leq \frac{C}{(R - |z_0|)^2} 2\pi R$, by using (3).

17. $\int_{\gamma} P dx + Q dy = \int_{\gamma} \frac{\partial g}{\partial x} dx + \frac{\partial g}{\partial y} dy = \int_a^b \frac{d}{dt} [g(\gamma(t))] dt = g(\gamma(b)) - g(\gamma(a)) = 0$.

21. $F(z) = \frac{1}{m+1} z^{m+1}$, $m = 0, 1, 2, \dots$.

Chapter 2

Section 2.1

1. (a)-(d) follow from (6) and the definition of these functions; (e) follows from (a), (b), and (4); (f) and (g) follow from the chain rule, (5).
3. $(3z^2 - 1) \exp(z^3 - z)$ 5. $-12z^2(z^3 + 100)^{-5}$
7. $e^z \cosh(e^z)$ 9. $-\frac{1}{z} + \frac{1}{3}z^3$

11. $\frac{1}{2} \sinh(2z)$

13. $f(z_0 + h) = f(z_0) + hf'(z_0) + hE_1, E_1 \rightarrow 0$ as $|h| \rightarrow 0$
 $g(w_0 + p) = g(w_0) + pg'(w_0) + pE_2, E_2 \rightarrow 0$ as $|p| \rightarrow 0$; $w_0 = f(z_0)$.
Hence,

$$\begin{aligned} g(f(z_0 + h)) &= g(f(z_0) + h(f'(z_0) + E_1)) = g(f(z_0)) + h(f'(z_0) + E_1)[g'(w_0) + E_2] \\ &= g(f(z_0)) + hf'(z_0)g'(w_0) + hE_1g'(w_0) + hf'(z_0)E_2 + hE_1E_2 \\ &= g(f(z_0)) + hf'(z_0)g'(w_0) + hE_3, \quad E_3 \rightarrow 0 \text{ as } h \rightarrow 0. \end{aligned}$$

15. $0 = f'$ on D so $\frac{\partial u}{\partial x} = \frac{\partial u}{\partial y} = \frac{\partial v}{\partial x} = \frac{\partial v}{\partial y}$ at all points in D .

As in Theorem 2, f is constant on D .

17. Follow the hint.

21. $\int_{\gamma} F'(z) dz = \int_a^b F'(\gamma(t))\gamma'(t) dt = \int_a^b \frac{d}{dt} F(\gamma(t)) dt = F(\gamma(b)) - F(\gamma(a)) = 0.$

23. Suppose the range of f lies in the straight line $\operatorname{Re}(Aw + B) = 0$. Then $g(z) = Af(z) + B$ is analytic, and $\operatorname{Re} g = 0$. Hence, g is constant, and therefore so is f . A similar argument may be applied if the range of f lies in a circle.

Section 2.1.1

1. Globally sourceless and irrotational; $G(z) = \frac{1}{4}z^4$
3. Globally sourceless and irrotational; $G(z) = \sin z$
5. Globally sourceless and irrotational; $G(z) = \frac{1}{3}z^3$
7. Neither sourceless nor irrotational
9. Use Exercise 21 in Section 1.

Section 2.2

1. $R = 1$

3. $R = 2^{1/3}$

5. $R = 1$

7. $\sum_0^{\infty} \frac{(-1)^n}{n!} z^n$

9. $\sum_{n=1}^{\infty} z^{3n}$

11. $\sum_{n=1}^{\infty} \frac{n}{4^{n+1}} z^{n+1}$

13. $z^2 \sum_{n=0}^{\infty} (-1)^n z^{2n} \left(\sum_{j=0}^n \frac{1}{(2j+1)! (2n+1-2j)!} \right)$

15. $\frac{z^3}{1-z^3}$

17. e^{-z}

19. Let $|z - z_0| = s < R$; then $\sum_0^{\infty} \left| \frac{a_n}{n+1} \right| |z - z_0|^{n+1} \leq s \sum_0^{\infty} |a_n| s^n$, which is finite by hypothesis.

21. (a) Follow the hint.

- (b) $(n+1)\rho^{n+1}/n\rho = ((n+1)/n)\rho < \frac{1}{2}(1+\rho)$ when $n > 2\rho/(1-\rho)$. Hence, $\{n\rho^n\}$ decreases geometrically fast once $n > 2\rho/(1-\rho)$.

- (c) If $1 \leq M$, then $1 \leq M^{1/n} \leq n^{1/n}$ once $n \geq M$. But $n^{1/n} \rightarrow 1$ by (a). If $0 < M < 1$, then $1 \leq (1/M)^{1/n} \leq n^{1/n}$, n large.

23. Use the ratio or root test on the given series.

Section 2.3

1. 0

3. $2\pi i$

5. $\frac{2\pi}{\sqrt{3}}$

7. $\frac{\pi}{\sqrt{a^2 - b^2}}$

9. i

11. $e - \frac{1}{e}$

13. By Cauchy's Theorem,

$$0 = \int_{\gamma} e^{iz^2} dz = \int_0^R e^{ix^2} dx + \int_0^{\pi/4} e^{i(Re^{i\theta})^2} i \operatorname{Re}^{i\theta} d\theta + \int_R^0 e^{i(t(1+i/\sqrt{2}))^2} \frac{1+i}{\sqrt{2}} dt.$$

The middle integral is

$$\int_0^{\pi/4} \exp[iR^2(\cos 2\theta + i \sin 2\theta)] ie^{i\theta} R d\theta.$$

This is estimated above by

$$R \int_0^{\pi/4} \exp[-R^2 \sin 2\theta] d\theta \leq R \int_0^{\pi/4} \exp[-2R^2\theta/\pi] d\theta < \frac{\pi}{2R}.$$

Let $R \rightarrow \infty$; we obtain

$$0 = \int_0^{\infty} [\cos(x^2) + i \sin(x^2)] dx - \frac{1+i}{\sqrt{2}} \int_0^{\infty} e^{-r^2} dt.$$

Use Exercise 20 and separate real and imaginary parts.

19. Follow the hint.
21. (c), (d) Cauchy's Theorem and Formula

Section 2.4

1. Zeros of order one at $n\pi$, $n = \pm 1, \pm 2, \dots$
3. Zeros of order three at -2 and 1
5. A zero of order four at 0 ; zeros of order two at $2\pi n$, $n = \pm 1, \pm 2, \dots$
7. Zeros of order one at $\pi i(1+2n)$, $n = 0, \pm 1, \dots$, and at $\log 4 + 2\pi im$, $m = 0, \pm 1, \dots$

9. $\sum_1^{\infty} \frac{z^{k+1}}{k!}$; valid everywhere.

11. $11(z-1) + 9(z-1)^2 + (z-1)^3$

13. $\frac{1}{2} + \sum_1^{\infty} (-1)^{k+1} \frac{(z+1)^k}{2^{k+1}}$

15. $\sum_1^{\infty} (-1)^j \pi^{2j} [(2j)!]^{-1} (z - \frac{1}{2})^{2j}$
17. $f(z) = (z - z_0)^m g(z)$; $g(z_0) \neq 0$. Thus, $f'(z) = (z - z_0)^{m-1} [mg(z) + (z - z_0)g'(z)] = (z - z_0)^{m-1} h(z)$; $h(z_0) = mg(z_0) \neq 0$. Likewise, $f''(z) = (z - z_0)^{m-2} g''(z)$; $g''(z_0) \neq 0$.
19. $|f'(z_0)| \leq 1! \frac{M}{s}$ if $|f(z)| \leq M$ for all z ; let $s \rightarrow \infty$.

21. For $n > m$, $|f^{(n)}(z_0)| \leq \frac{n!}{s^n} As^m$ if $|z| = s \geq R_0$; let $s \rightarrow \infty$.

23. Write $G(z) = \sum_0^{\infty} c_k z^k$ for $|z| < 1/R$. Set $w = 1/z$; then

$$F(w) = G(z) = \sum_0^{\infty} c_k z^k = \sum_0^{\infty} c_k \frac{1}{w^k}, \quad |w| > R.$$

Furthermore, if $0 < r < 1/R$, then

$$\begin{aligned} c_k &= \frac{1}{2\pi i} \int_{|\zeta|=r} \frac{G(\zeta)}{\zeta^{k+1}} d\zeta. \text{ Let } \zeta = \frac{1}{\xi}, |\xi| = s = \frac{1}{r} > R; \quad \text{therefore,} \\ &= \frac{1}{2\pi i} \int_{|\xi|=s} F(\xi) \xi^{k-1} d\xi, \quad k = 0, 1, 2, \dots \end{aligned}$$

25. Write $f(z) = \sum_0^{\infty} a_k z^k$; substitute into the differential equation and collect equal powers of z . This gives equations for a_2, a_3, \dots , in terms of a_0 and a_1 :

$$a_{2k} = \frac{(-1)^k}{(2k)!} a_0 \beta^{2k} \quad \text{and} \quad a_{2k+1} = \frac{(-1)^k}{(2k+1)!} a_1 \beta^{2k}.$$

27. Set $g(z) = \exp[-\int_{z_0}^z A(w) dw]$. Then

$$(fg)'(z) = f'(z)g(z) + f(z)g'(z) = A(z)f(z)g(z) - A(z)g(z)f(z) = 0.$$

Hence, $f = C/g = C \exp[\int_{z_0}^z A(w) dw]$.

29. Compute J'_v, J''_v , substitute into the given equation, and compute.

Section 2.5

1. A removable singularity at 0; the value at 0 is 1.
3. A removable singularity at 1; the value at 1 is 4.
5. A pole of order one at -2

7. $\frac{1}{z} + \frac{1}{2} + \frac{1}{6} z + \sum_2^{\infty} \frac{1}{(k+2)!} z^k; \quad \text{Res}(f; 0) = 1$

9. $\frac{-1}{z-\pi} + \sum_0^{\infty} (-1)^{k+1} \frac{(z-\pi)^{2k+1}}{(2k+1)!}; \quad \text{Res}(f; \pi) = -1$

11. $\frac{bc-ad}{c^2} \frac{1}{z+d/c} + \frac{a}{c}; \quad \text{Res}\left(f; -\frac{d}{c}\right) = \frac{bc-ad}{c^2} \quad (\text{all if } c \neq 0).$

13. $\frac{2}{z^2} + \frac{1}{6} + \frac{1}{120}z^2 + \dots; \quad \text{Res}(f; 0) = 0.$

15. $f(z) = (z - z_0)^{-l}g(z)$, $l \geq 1$. Thus, $f'(z) = (z - z_0)^{-l}[g'(z) - l(z - z_0)^{-1}g(z)]$, and

$$\frac{f'(z)}{f(z)} = \frac{g'(z)}{g(z)} + \frac{-l}{z - z_0}.$$

17. Follow the hint: If g is bounded near z_0 , then z_0 is a removable singularity for g ; let $A = g(z_0)$. Next, $f(z) = w + 1/g(z)$. If g has a zero at z_0 ($A = 0$), then f has a pole at z_0 . If $A \neq 0$, then f is bounded near z_0 .

19. Let γ be the circle $|z - z_0| = s$, $0 < s < r$. Then for any integer m ,

$$\begin{aligned} 0 &= \frac{1}{2\pi i} \int_{\gamma} (z - z_0)^{-m} \left(\sum_{-\infty}^{\infty} a_n (z - z_0)^n \right) dz \\ &= \frac{1}{2\pi} \int_0^{2\pi} s^{-m} e^{-im\theta} \left(\sum_{-\infty}^{\infty} a_n s^n e^{in\theta} \right) s e^{i\theta} d\theta \\ &= \sum_{-\infty}^{\infty} a_n s^{-m} s^{n+1} \frac{1}{2\pi} \int_0^{2\pi} e^{i(n+1-m)\theta} d\theta = a_{m-1}. \end{aligned}$$

21. By (5),

$$\begin{aligned} \text{Res}(F; z_0) &= \frac{1}{2\pi i} \int_{|z-z_0|=s} F(z) dz = \frac{1}{2\pi i} \int_{|z-z_0|=s} G'(z) dz \\ &= \frac{1}{2\pi} \int_0^{2\pi} G'(z_0 + se^{i\theta}) s e^{i\theta} d\theta \\ &= \frac{1}{2\pi} [G(z_0 + se^{2\pi i}) - G(z_0 + se^{i0})] = 0. \end{aligned}$$

23. In $1 < |z| < 2$

(a) $\sum_{-\infty}^{\infty} a_n z^n, \quad a_n = \begin{cases} \frac{(-1)^n}{3}, & n = -1, -2, \dots \\ \frac{-4}{3^{n+2}}, & n = 0, 1, 2, \dots \end{cases}$

$$\text{In } 2 < |z| < \infty, \quad \sum_1^{\infty} \frac{a_k}{z^k}, \quad a_k = \frac{(-1)^k}{3} + \frac{4}{3^{k+2}}, \quad k = 1, 2, \dots$$

(b) In $0 \leq |z| < 1$, $\sum_0^{\infty} a_k z^k, \quad a_k = \frac{1}{24} [15(-1)^k - 3^{-k+1}]$

$$\text{In } 1 < |z| < 3, \quad \sum_{-\infty}^{\infty} a_k z^k, \quad a_k = \begin{cases} \left(\frac{5}{8}\right)(-1)^{k+1}, & k = -1, -2, \dots \\ \left(\frac{3}{8}\right)(3^{-k-1})((-1)^k - 1), & k = 0, 1, 2, \dots \end{cases}$$

(c) In $2 < |z| < 3$, $\sum_{-\infty}^{\infty} a_k z^k$, $a_k = \begin{cases} 2^{-k-1}, & k = -1, -2, \dots \\ \frac{4}{3^{k+1}} - \frac{1}{5^{k+1}}, & k = 0, 1, 2, \dots \end{cases}$
 $\text{In } 5 < |z| < \infty, \sum_1^{\infty} \frac{a_n}{z^n}, a_n = 2^{n-1} - 4(3)^{n-1} + 5^{n-1}, n = 1, 2, 3, \dots$

25. ∞ is removable for f iff 0 is removable for $g(z) = f(1/z)$ iff g is bounded near 0 iff f is bounded near ∞ .

27. (a) pole of order two at ∞ ; $3z^2 + 4 - 1/z$
(b) pole of order two at ∞ ; $-z^2 + 5z - 4$

(c) pole of order one at ∞ ; $z + 4 + \frac{4^2}{z} + \frac{4^3}{z^2} + \dots$

(d) zero of order two at ∞ ; $-\frac{1}{z^2} + \frac{2}{z^3} - \frac{7}{z^4} + \dots$

(e) removable singularity at ∞ ; value is 1 ; $1 + z^{-1} + \frac{z^{-2}}{2!} + \frac{z^{-3}}{3!} + \dots$

(f) essential singularity at ∞ ; $1 - z^2 + \frac{z^4}{2!} - \frac{z^6}{3!} + \dots$

(g) pole of order three at ∞ ; $z^3 + 3z + \frac{3}{z} + \frac{1}{z^3}$

(h) removable singularity at ∞ ; value is 0 ; $z^{-1} - \frac{z^{-3}}{3!} + \frac{z^{-5}}{5!} - \dots$

(i) removable singularity at ∞ ; value is 1 ; $1 - \frac{z^{-2}}{2!} - \frac{z^{-4}}{4!} - \dots = \cos\left(\frac{1}{z}\right)$

29. First, $\bar{G}(\bar{z}; u) = \exp[(u/2)(\bar{z} - 1/\bar{z})] = \exp[(u/2)(z - 1/z)] = G(z; u)$ if u is real. Hence, $J_n(u)$ is real if u is real. Apply (19) with $s = 1$. This yields

$$J_n(u) = \operatorname{Re}(J_n(u)) = \operatorname{Re}\left(\frac{1}{2\pi} \int_0^{2\pi} e^{iu \sin \theta - n\theta} d\theta\right)$$

$$= \frac{1}{2\pi} \int_0^{2\pi} \cos(u \sin \theta - n\theta) d\theta.$$

31. $\sum_{-\infty}^{\infty} J_n(u) z^n = G(z; u) = \exp\left[\frac{u}{2}\left(z - \frac{1}{z}\right)\right] = G\left(-\frac{1}{z}; u\right) = \sum_{-\infty}^{\infty} J_n(u) \frac{(-1)^n}{z^n}$

$$= \sum_{-\infty}^{\infty} J_{-n}(u) (-1)^n z^n. \quad \text{So } J_{-n}(u) = (-1)^n J_n(u).$$

Section 2.6

1. $\frac{\pi}{4} \left[\sin\left(\frac{3\pi}{8}\right) \right]^{-1}$

3. $\frac{\pi}{ab(a+b)}$

5. $\pi \exp\left(-\frac{\sqrt{2}}{2}\right) \sin\left(\frac{\sqrt{2}}{2}\right)$

7. $\frac{\pi}{\beta} e^{-\beta} \cos \alpha$

9. $\frac{4\pi}{3\sqrt{3}}$

11. $\frac{2\pi a^2}{1-a^2}$

13. $\frac{\pi[2^\alpha - 1]}{\sin \pi\alpha}$

15. $\frac{\pi}{\sin \omega} \frac{\sin \omega\lambda}{\sin \pi\lambda}$

17. 0

19. $\left(\frac{5}{32}\right)\pi^5$

21. Follow the hint.

23. Use the technique of Examples 1 and 2; the poles in U have order two (for (a)) or three (for (b)).25. Use (11) to write $g = h + \sum_1^N P_j$, where P_j is a polynomial and h is analytic in D .27. $C(z) = \pi \cot \pi z = \pi \frac{\cos \pi z}{\sin \pi z}$ has a pole of order one at each integer. Furthermore,

$$\text{Res}(C; k) = \lim_{z \rightarrow k} (z - k)\pi \frac{\cos \pi z}{\sin \pi z} = \lim_{z \rightarrow k} (\cos \pi z) \frac{\pi}{\frac{\sin \pi z}{z - k}} = \pi \frac{\cos \pi k}{(\sin \pi z)'(k)} = 1.$$

29. $\left| \int_{\gamma_N} C(z)f(z) dz \right| \leq \left(\frac{A}{N^2} \right) \text{length } (\gamma_N) \leq \frac{B}{N} \rightarrow 0 \quad \text{as } N \rightarrow \infty.$ Now apply the Residue Theorem.

31. (a) $\frac{1}{2} \left(\frac{\pi \cosh \pi a}{a \sinh \pi a} - \frac{1}{a^2} \right)$ (b) $\left(\frac{\pi}{\sin \pi a} \right)^2$ (c) $\frac{\pi^2}{8}$ (d) $\frac{\pi^4}{90}$

33. In 32, the left-hand side goes to zero as $N \rightarrow \infty$; write the sum over $k = 1, \dots, N$.

Chapter 3

Section 3.1

1. One zero in first quadrant
3. One zero in first quadrant
5. Two zeros in first quadrant
7. Two zeros in the upper half-plane
9. On the imaginary axis, $f(z) = z^4 - 5z^2 + 3 - e^{-z}$ has no solution, since $|e^{-iy}| = 1$ while $y^4 + 5y^2 + 3 \geq 3$. Let $g(z) = z^4 - 5z^2 + 3$. Then $|f - g| < |g|$ on the contour in Figure 3.4. g has two zeros in the right half-plane.
11. $\sin z$ has zeros of order one at $z = k\pi$, $k = 0, \pm 1, \dots$. The square of an analytic function can only have zeros of even order.
13. $z^4 - 2z - 2$ has four zeros in $\frac{1}{2} < |z| < \frac{3}{2}$.
15. $4z^3 - 12z^2 + 2z + 10$ has two zeros in $\frac{1}{2} < |z - 1| < 2$.
17. (a) No solutions (b) No solutions (c) One solution
19. Follow the hints.
21. $p - q$ is a polynomial of degree n with $n + 1$ zeros.
23. The hint shows how.
25. g has no poles, so it is entire.
27. The hint has it.

29. $\frac{P(z)}{Q(z)} z^{M-N} = \frac{a_0/z^N + \dots + a_N}{b_0/z^M + \dots + b_M} \rightarrow \frac{a_N}{b_M} \neq 0 \quad \text{as } |z| \rightarrow \infty.$

31. $R(z) - \alpha = \frac{P(z) - \alpha Q(z)}{Q(z)}.$

Now $\deg(P - \alpha Q) = \max(N, M)$ except if $M = N$ and $\alpha b_N = a_N$. In this case, $\deg(P - \alpha Q) < N = \deg Q = \max(N, M)$.

33. Use Cauchy's Formula: Let γ be a contour surrounding t . Then

$$\frac{1}{2^n} \frac{1}{2\pi i} \int_{\gamma} \frac{(w^2 - 1)^n}{(w - z)^{n+1}} dw = \frac{1}{2^n} \frac{1}{n!} \left. \frac{d^n}{dw^n} (w^2 - 1)^n \right|_{w=z} = P_n(z).$$

$$\begin{aligned} 35. \quad \sum P_n(z) t^n &= \sum t^n \frac{1}{2\pi} \int_0^{2\pi} (z + \sqrt{z^2 - 1} \cos \theta)^n d\theta \\ &= \frac{1}{2\pi} \int_0^{2\pi} \sum t^n (z + \sqrt{z^2 - 1} \cos \theta)^n d\theta \\ &= \frac{1}{2\pi} \int_0^{2\pi} \frac{1}{1 - t(z + \sqrt{z^2 - 1} \cos \theta)} d\theta \\ &= \frac{1}{2\pi} \int_0^{2\pi} \frac{1}{a - b \cos \theta} d\theta, \end{aligned}$$

with $a = 1 - tz$, $b = t\sqrt{z^2 - 1}$. Now use Exercise 7, Section 2.3, to evaluate the integral.

Section 3.1.1

1. (a) Not stable (b) Stable (c) Not stable
- (d) Not stable (e) Not stable (f) Stable
- (g) Not stable (h) Not stable (i) Stable
- (j) Not stable (k) Not stable (l) Stable
3. (a) Stable (b) Stable (c) Not stable
- (d) Not stable (e) Stable (f) Not stable
7. Check Theorem 2, Section 3.1, the argument principle.

Section 3.2

1. 1 3. $\frac{1}{2}$
5. S is not continuous at $z = 1$.

7. $0 = \operatorname{Re} F(z_0) = \frac{1}{2\pi} \int_0^{2\pi} \operatorname{Re} F(z_0 + re^{it}) dt.$

Hence, $\operatorname{Re} F$ must be both positive and negative somewhere on the circle $|z - z_0| = r$.

9. Use minimum modulus.
11. Follow the hint.
13. The function $g(z) = f(z)/M$ is bounded by 1 in $|z| < 1$. Let $\alpha_0 = g(z_0)$ and $\psi(z) = (z - \alpha_0)(1 - \bar{\alpha}_0 z)^{-1}$. Then $h = \psi \circ g$ is also bounded by 1 and is zero at z_0 . Use Exercise 12 with z_0 in place of α .

15. Follow the hint.
17. Let $h = g/f$ and use the maximum-modulus principle for h on γ . An example of failure: $f(z) = z$, $g(z) = 1$, and $\gamma = \{w: |w| = 1\}$.
19. $\tilde{p}(z) = \overline{(p(-\bar{z}))} = p(z)$. Let $Q(z) = \sum_0^n a_k z^k$; then $Q = \tilde{Q}$ if and only if a_k is real for k even and pure imaginary if k is odd.
21. p stable implies that $\tilde{p}(z)/p(z)$ is analytic on $\operatorname{Re} z > 0$ and continuous on $\operatorname{Re} z \geq 0$. Apply maximum modulus to \tilde{p}/p on the contour $\{\operatorname{Re}^{it}: -\pi/2 \leq t \leq \pi/2\} \cup \{iy: R \geq y \geq -R\}$; let $R \rightarrow \infty$. Conversely, if $|\tilde{p}(z)| \leq |p(z)|$ for $\operatorname{Re} z \geq 0$, and \tilde{p} and p have no common zero, then p has no zero in $\{\operatorname{Re} z \geq 0\}$.
23. p stable implies $r = \tilde{p}/p$ maps $\{\operatorname{Re} z > 0\}$ into $\{w: |w| < 1\}$. This implies that $R = (1 - r)/(1 + r)$ maps $\{\operatorname{Re} z > 0\}$ into $\{\operatorname{Re} \zeta > 0\}$, so R is positive. The converse is similar; use the results of Exercise 21.

Section 3.3

1. Compute

3. Compute

5. (a) $\lambda(1-i)\frac{z+1}{z-1}$, λ any real number (b) $z+1$ (c) $iz+\frac{1}{2}$
- (d) $\frac{z-z_0}{r}$ (e) $4i-4+\frac{4}{z}$
7. (a) $\frac{(1-i)z}{z-i}$ (b) $\frac{1}{z+1}$ (c) $\frac{z+4}{2z+2}$ (d) $\frac{z-1}{z+1}$
9. (a) Fixed points at 0, -1 ; sink at 0; source at -1
 (b) Fixed points at ± 2 ; sink at 2; source at -2
 (c) Fixed points at 1, ∞ ; source at 1; sink at ∞ ; straight-line motion outward from $z = 1$
 (d) Fixed points at 0, ∞ ; source at ∞ ; sink at 0; inward clockwise spiral to origin
 (e) Translation to the right; dipole at ∞
11. Solve $T(1) = \frac{a+b}{c+d} = 1$ and $T(-1) = \frac{-a+b}{-c+d} = -1$.
13. Use the results of Exercise 11.
15. Write $S(z) = \frac{az+b}{cz+d}$ and solve $z = S(S(z))$ for a , b , c , and d :
- $$S(z) = \frac{az+b}{cz-a}, \quad \text{where } a^2 + bc \neq 0.$$
17. Follow the extensive hints.
19. (a) and (c) are clear from the definition. (b) If L is the real axis, then $d = |\operatorname{Im} z|$.
23. A reflection over a line: $z^* = \lambda\bar{z} + b$, $|\lambda| = 1$; a reflection over a circle: $w^* = a + R^2/(\bar{w} - \bar{a})$. Putting two such transforms together produces $Az + B$ or $(Az + B)/(Cz + D)$.
25. Given α , find a, b with $(a\alpha + b)/(-b\alpha + \bar{a}) = 0$ and $|a|^2 + |b|^2 = 1$. Take $a = (1 + |\alpha|^2)^{-1/2}$ and $b = -a\alpha$.
27. If S is analytic and preserves chordal distance and $S(0) = 0$, then

$$x(z, 0) = x(S(z), S(0)) = x(S(z), 0),$$

$$\frac{|z|^2}{1+|z|^2} = \frac{|S(z)|^2}{1+|S(z)|^2}.$$

This gives $|z|^2 = |S(z)|^2$. Schwarz's Lemma (Theorem 2, Section 3.2) implies that $S(z) = \lambda z$, $|\lambda| = 1$.

Section 3.4

1. $z^2 = w^2$ iff $0 = z^2 - w^2 = (z+w)(z-w)$. If $z+w=0$, then $\operatorname{Re} z = -\operatorname{Re} w$, which cannot happen. If U is open and strictly bigger than $\{\operatorname{Re} z > 0\}$, there is a $z_0 \in U$ with $\operatorname{Re} z_0 < 0$. Hence, $-z_0 \in \{\operatorname{Re} z > 0\}$, and $(-z_0)^2 = z_0^2$.
3. (a) Let $\alpha_0 = f(0)$; by Theorem 1, Section 3.2, there is a $\delta > 0$ such that the disc $\{w: |w - \alpha_0| < \delta\}$ is a subset of $\{f(z): |z| < 1\}$. Hence, $|f(z) - \alpha_0| \geq \delta$ if $|z| \geq 1$. (Why?) Show that $(f - \alpha_0)^{-1}$ has a removable singularity at ∞ . Conclude that f has a pole at ∞ of order m , $m \geq 1$, and so is a polynomial of degree m . Then deduce that $m = 1$.
- (b) Compose with a linear fractional transformation so the pole is at ∞ . Then apply the result of 3(a).
5. The quadratic $z^2 - 2wz + R^2 = 0$ is left unchanged by the substitution of R^2/z for z . Hence, its two roots z_1, z_2 are related by $z_1 z_2 = R^2$. Therefore, these roots lie on opposite sides of the circle $|z| = R$, or both lie on the circle $|z| = R$ and are complex conjugates of each other. The latter holds if and only if $w \in [-R, R]$.
7. (a) See Figure 3.16 with $\frac{1}{2}, 2$ in place of $-1, 1$, respectively.
 (b) See Figure 3.19 translated over one unit.
 (c) This is a dipole at 1 with circles centered on the real axis and on the line $\operatorname{Re} z = 1$.
 (d) See Figure 3.15 moved over and up one unit: $z^2 - 2z = (z-1)^2 - 1$.
9. Write $p(z) = \prod_1^n (z - z_k)^{m_k}$; $\gamma_\varepsilon = \{z: |\prod_1^n |z - z_k|^{m_k}| = \varepsilon\}$. For the product to be small, one factor must be small, so z is near some z_j (and so the other factors are bounded). For the product to be large, one factor is large (and so all the factors are large). Let $n = \deg p$. Then

$$\frac{|p(z)|}{|z|^n} \rightarrow 1 \quad \text{as } R = |z| \rightarrow \infty,$$

so γ_ε is "almost" the circle $|z| = R$ if R is big.

11. Follow the suggested route.
13. $g(z) = z + e^z$ so $g'(z) = 1 + e^z$ and $\operatorname{Im}[g'(z)] = e^x \sin y > 0$ if $0 < y < \pi$. Now apply the results of Exercise 12 with $f = -ig$ to see that g is $1 - 1$.
15. Follow the suggestions.

Section 3.4.1

1. $G(iy) = -\frac{1}{2}(y^2 + (1/y^2)) \in (-\infty, -1]$ if $1 \leq y < \infty$; $G(e^{i\theta}) = \cos 2\theta \in [-1, 1]$ if $\pi/2 \geq \theta \geq 0$; $G(x) = \frac{1}{2}(x^2 + (1/x^2)) \in [1, \infty)$ if $1 \leq x < \infty$. Hence, G maps the boundary of D onto the real axis. G is also $1 - 1$, since $G(z) = G(w)$ iff $z^2 - w^2 = z^{-2}w^{-2}(z^2 - w^2)$ iff $z^2 = w^2$ (since $|z| > 1$, $|w| > 1$) iff $z = w$ (since $\operatorname{Re} z > 0$, $\operatorname{Re} w > 0$). The streamlines are $\Gamma_c = \{z: z^2 = t + ic + i\sqrt{1 - (t + ic)^2}, -\infty < t < \infty\}, c > 0$.
3. Refer to Exercise 5, Section 3.4, for the properties of K . The streamlines are

$$\Gamma_c = \left\{ z: z = \left(-\frac{t^2}{4} - i\frac{tc}{4} \right) + \frac{c^2 - 1}{4}, \quad -\infty < t < \infty \right\}, \quad c > 0.$$

5. $G(z) = G(w)$ iff $(1 - z^2)^2 = (1 - w^2)^2$ iff $(z^2 - w^2)(z^2 + w^2) = 2(z^2 - w^2)$ iff either $z^2 = w^2$ or

$z^2 + w^2 = 2$. But z and w are in the first quadrant, so $0 < \operatorname{Im}(z^2 + w^2)$. Hence, $z^2 = w^2$, so $z = w$, since $\operatorname{Re} z > 0$, $\operatorname{Re} w > 0$. Next, $G(iy) = -(1 + y^2)^2 \in (-\infty, -1]$ if $0 \leq y < \infty$; $G(x) = -(1 - x^2)^2 \in [-1, 0]$ if $0 \leq x \leq 1$; $G(x + i\sqrt{x^2 - 1}) = 4x\sqrt{x^2 - 1} \in [0, \infty)$ if $1 \leq x < \infty$. Hence, G maps the boundary of D onto the real axis.

7. $G'(z) = \frac{2}{z^2 - 1} = \frac{1}{z-1} - \frac{1}{z+1}$.

(See Fig. 3.16.) The streamlines are the circles $\arg[(z-1)/(z+1)] = c$.

9. See Figure 3.16 with p, q in place of $-1, 1$, respectively.
 11. Set $\gamma(t) = H(u(t))$; then $G(\gamma(t)) = u(t) \in L$, so $\operatorname{Im} G(\gamma(t)) = c_0$. Moreover, $\gamma'(t) = H'(u(t))u'(t) = f'(\gamma(t))$, so γ parametrizes the curve.

Section 3.5

1. $\phi(z) = z^2$

3. $\phi(z) = i\frac{2-i+z}{2+i-z}$

5. $\phi(z) = \frac{1}{2R}\left(z + \frac{R^2}{z}\right)$

7. $\phi(z) = \frac{\pi}{2} + \arcsin(e^z)$

9. $\phi(z) = \frac{4}{\pi}(\operatorname{Log} z) - 1$

11. $f(z) = \arcsin z$

13. $f(z) = -iS(z^2 - 1)^{3/4}$

15. $f(z) = \frac{-a}{\pi} \left[2\sqrt{z+1} + \operatorname{Log} \left(\frac{\sqrt{z+1} - 1}{\sqrt{z+1} + 1} \right) \right]$

17. Write $f = u + iv$; the change-of-variables formula gives

$$\begin{aligned} \text{area}(D) &= \iint_D 1 \, du \, dv = \iint_{|z|<1} \det \begin{pmatrix} u_x & u_y \\ v_x & v_y \end{pmatrix} dx \, dy \quad (\text{apply Cauchy-Riemann}) \\ &= \iint_{|z|<1} (u_x^2 + u_y^2) \, dx \, dy = \iint_{|z|<1} |f'(z)|^2 \, dx \, dy. \end{aligned}$$

19. If $\{|\phi(z_k)|\}$ does not converge to 1, then there is a subsequence $\{z_{k_i}\}$ with $\phi(z_{k_i}) \rightarrow q$, $|q| < 1$. This contradicts Exercise 14, Section 4.

Chapter 4

Section 4.1

1. (a) u is harmonic; $v = 4x^3y - 4xy^3$.
 (b) u is harmonic; $v = [\sin(2xy)][e^{x^2-y^2}]$.
 (c) u is not harmonic.
 (d) u is harmonic; $v = [\cos(x^2 - y^2)][\sinh(2xy)]$.
 (e) u is not harmonic.
3. Take $f(z) = z$ on $0 < |z| < R$ and apply the conclusion of Exercise 2.
5. (a) A function u of θ alone is harmonic iff $u = A + B\theta$.
 (b) A function u of r alone is harmonic iff $u = C + D \log r$, $0 < r < \infty$.

7. This is a computation.


9. If $z = re^{i\theta}$, then $\frac{1 - r^2}{1 - 2r \cos \theta + r^2} = \operatorname{Re}\left(\frac{1+z}{1-z}\right)$.

11. Compute Δu and set it equal to 0.

13. Given $\delta > 0$, choose $r > 0$ such that $|u(z)| < \delta$ if $|z| < r$ and $\operatorname{Im} z \geq 0$; this is possible by continuity. Let

$$V_\varepsilon(x, y) = u(x, y) - \varepsilon \log \frac{x^2 + y^2}{r^2};$$

consider V_ε on the contour shown. For large R , V_ε is negative on the large semicircle; V_ε is also negative on the segments $(-R, -r)$ and (r, R) . On the small semicircle, V_ε equals u and so is less than δ . Hence, $V_\varepsilon < \delta$ on D by the maximum principle. Let $R \rightarrow \infty$; then $\varepsilon \downarrow 0$. This gives $u \leq \delta$ on $\{z: \operatorname{Im} z > 0, |z| > r\}$. Let $r \downarrow 0$ and $\delta \downarrow 0$. Then repeat the argument with $-u$ in place of u .



15. Let f be analytic on $|z - z_0| < \delta$ with $u = \operatorname{Re} f$. Then $g(z) = \overline{f(\bar{z})}$ is also analytic on $|z - z_0| < \delta$ (why?), and $f(x) = g(x)$, $x_0 - \delta < x < x_0 + \delta$. Hence, $g = f$, so $\operatorname{Re} g = \operatorname{Re} f = u$.
17. Follow the numerous hints.

Section 4.2

- $\Gamma_0 = \{z: |z - i\alpha_0| = \sqrt{R^2 + \alpha_0^2}\}$. Thus, if \sim is one of $<$, $=$, $>$, then $|z - i\alpha_0|^2 \sim R^2 + \alpha_0^2$ iff $|z|^2 - 2 \operatorname{Re}(i\alpha_0 z) + \alpha_0^2 \sim R^2 + \alpha_0^2$ iff $|z|^2 - 2y\alpha_0 - R^2 \sim 0$.
- Let h be another analytic function outside Γ with $h(z) = a + (c/2\pi iz) + \dots$, $\operatorname{Im} H = 0$ on Γ , and $H' = h$. Then $g - h$ is analytic outside Γ ; near ∞ , $g(z) - h(z) = a_2/z^2 + \dots$, and $g - h$ has a single-valued indefinite integral J . Therefore, $\operatorname{Im} J = 0$ on Γ , so J is a real constant. This gives $0 = J' = g - h$.
- The flow has a source at -1 and a sink at λ , $|\lambda| = 1$, $\lambda \neq -1$. Thus,

$$F(z) = \frac{Q}{\pi} \log \left(\frac{1+z}{1-\bar{\lambda}z} \right).$$

- Let the sources be at $\pm\pi + ip$; then $F(z) = \frac{Q}{\pi} \left[\operatorname{Log} \left\{ \sin^2 \left(\frac{z}{2} \right) - \sin^2 \left(\frac{p}{2} \right) \right\} \right]$.

- Let the source be at x_0 , $x_0 \in (-\pi, \pi)$; then

$$F(z) = \frac{Q}{\pi} \operatorname{Log} \left\{ \sin \left(\frac{z}{2} \right) - \sin \left(\frac{x_0}{2} \right) \right\}.$$

- (a) $z^2 + 1$ is a conformal mapping of the first quadrant onto the upper half-plane U . $\operatorname{Log} w$ is a conformal mapping of U onto the strip S .

- (b) There is a source of strength π at $z_0 = i$.
(c) If p is a point in the boundary of the first quadrant, then $\operatorname{Log}(z^2 - p^2)$ gives the potential of a flow with a source at p .

$$\begin{aligned} 13. \text{ potential } &= \phi(z) = c_0 \int_{-1}^1 \log \left| \frac{1}{z-t} \right| dt \\ &= (2+y)c_0 - c_0 \log|1-z^2| - c_0 \operatorname{Re} \left[z \operatorname{Log} \left(\frac{1+z}{1-z} \right) \right]. \end{aligned}$$

15. $\phi(z) = (a^2 - b^2)^{-1}[az - b\sqrt{z^2 - (a^2 - b^2)}]$ maps the region in the upper half-plane and outside the ellipse onto the upper half-plane.

17. Use the conclusion of Exercise 16 for each term. Write

$$g(z) = \frac{K_2}{2\pi i} \frac{\bar{z}_1 - \bar{z}_2}{(\bar{z} - \bar{z}_1)(\bar{z} - \bar{z}_2)} + \frac{K_1 - K_2}{2\pi i} \frac{1}{\bar{z} - \bar{z}_1}.$$

For $|z|$ large, the first term is very small in comparison to the second, so the direction of the flow given by g is determined by the sign of $K_1 - K_2$: clockwise if $K_1 > K_2$, counterclockwise if $K_1 < K_2$.

19. Assuming that the origin is midway between the edges of the plates, the potential is

$$G(z) = \frac{V_1 + V_0}{\pi} + \frac{V_1 - V_0}{\pi} \operatorname{Re} \left(\arcsin \frac{z}{\sigma} \right).$$

Section 4.3

1. Let the arc for temperature T_1 be $\{e^{i\theta} : 0 \leq \theta < \theta_0\}$. Then

$$T(z) = T_2 + \frac{T_1 - T_2}{\pi} \operatorname{Arg} \left\{ i \frac{1+z}{1-z} - i \frac{1+e^{i\theta_0}}{1-e^{i\theta_0}} \right\}.$$

3. (a) Choose M so that $|w(t)| < \varepsilon/4$ if $|t| > M$. Then

$$\frac{1}{\pi} \int_{-\infty}^{-M} \frac{\tau |w(t)|}{(\sigma - t)^2 + \tau^2} dt + \frac{1}{\pi} \int_M^{\infty} \frac{\tau |w(t)|}{(\sigma - t)^2 + \tau^2} dt < \frac{\varepsilon}{4} \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{\tau}{(\sigma - t)^2 + \tau^2} dt = \frac{\varepsilon}{4}.$$

Moreover,

$$\begin{aligned} \frac{1}{\pi} \int_{-M}^M \frac{\tau |w(t)|}{(\sigma - t)^2 + \tau^2} dt &\leq \left(\frac{1}{\pi} \int_{-M}^M |w(t)| dt \right) \left(\max_{|t| \leq M} \frac{\tau}{(\sigma - t)^2 + \tau^2} \right) \\ &\leq \left(\frac{1}{\pi} \int_{-\infty}^{\infty} |w(t)| dt \right) \frac{|\zeta|}{(|\zeta| - M)^2} \\ &< \frac{\varepsilon}{2} \quad \text{if } |\zeta| \text{ is very large.} \end{aligned}$$

- (b) The difference, $h = W - W_1$, is harmonic on $\{x + iy: y > 0\}$, is 0 on $\{y = 0\}$, and is bounded, since both W and W_1 go to 0 at ∞ ; now apply Exercise 13, Section 4.1.

$$\begin{aligned} \text{(c)} \quad \int_{-\infty}^{\infty} |W(\sigma + i\tau)| d\sigma &\leq \int_{-\infty}^{\infty} \left(\frac{1}{\pi} \int_{-\infty}^{\infty} |w(t)| \frac{\tau}{(\sigma - t)^2 + \tau^2} dt \right) d\sigma \\ &= \int_{-\infty}^{\infty} |w(t)| \left(\frac{1}{\pi} \int_{-\infty}^{\infty} \frac{\tau}{(\sigma - t)^2 + \tau^2} d\sigma \right) dt \\ &= \int_{-\infty}^{\infty} |w(t)| dt. \end{aligned}$$

5. (i) u is continuous at $e^{i\theta_0}$.
(ii) Apply (ii) from Exercise 4.
(iii) $|u(e^{it}) - u(e^{i\theta_0})| < \varepsilon/2$, since $|t - \theta_0| < \delta/4$, and (ii) from Exercise 4 holds.
(iv) The integral of a function is less than the maximum of the function times the length of the interval.
(v) Apply (iii) of Exercise 4 and (iv).
(vi) Combine the two estimates on I_1 and I_2 .
7. Let $u = u_1 + u_2$, where

$$u_1(x, y) = \int_0^{\infty} f_1(t) \left\{ \frac{y}{(x-t)^2 + y^2} - \frac{y}{(x+t)^2 + y^2} \right\} dt$$

and

$$u_2(x, y) = \int_0^{\infty} f_2(t) \left\{ \frac{x}{(y-t)^2 + x^2} - \frac{x}{(y+t)^2 + x^2} \right\} dt.$$

$$9. \quad V(z) = \frac{1}{2\pi} \int_0^{2\pi} f(e^{-i\theta}) \frac{|z|^2 - 1}{|ze^{i\theta} - 1|^2} d\theta, \quad |z| > 1.$$

11. $u(re^{i\theta}) = \operatorname{Re}((1+z)/(1-z))$, $z = re^{i\theta}$, if $\theta \neq 0$, then $\cos \theta \neq 1$, so $\lim_{r \uparrow 1} u(re^{i\theta}) = 0$. Follow the hint for the “hot spot.”

$$13. \quad T(x+iy) = \frac{1}{\pi} \sum_{j=1}^N A_j \frac{y}{(x-x_j)^2 + y^2}, \quad \text{where } A_j = \begin{cases} 1 & \text{if } x_j \text{ is “hot”} \\ -1 & \text{if } x_j \text{ is “cold”.} \end{cases}$$

15. Follow the hint.

$$17. \quad \int_0^{2\pi} e^{ik\theta} d\theta = \frac{1}{ik} e^{ik\theta} \Big|_0^{2\pi} = \frac{1}{ik} [e^{ik2\pi} - e^0] = 0 \quad \text{if } k \neq 0.$$

$$\begin{aligned} 19. \quad P_r(\theta) &= \operatorname{Re} \left[\frac{1+re^{i\theta}}{1-re^{i\theta}} \right] = \operatorname{Re} [(1+re^{i\theta})(1+re^{i\theta} + r^2 e^{2i\theta} + \cdots)] \\ &= \operatorname{Re}(1+2re^{i\theta} + 2r^2 e^{2i\theta} + \cdots) \\ &= 1 + r(e^{i\theta} + e^{-i\theta}) + r^2(e^{2i\theta} + e^{-2i\theta}) + \cdots \\ &= \sum_{n=0}^{\infty} r^{|n|} e^{in\theta}. \end{aligned}$$

$$\begin{aligned}
 21. \quad & \int_0^{2\pi} U(re^{i\theta}) \overline{V(re^{i\theta})} d\theta = \int_0^{2\pi} \left(\sum_{n=-\infty}^{\infty} r^{|n|} \hat{u}(n) e^{in\theta} \right) \left(\sum_{m=-\infty}^{\infty} r^{|m|} \overline{\hat{v}(m)} e^{-im\theta} \right) d\theta \\
 &= \sum_{n=-\infty}^{\infty} \sum_{m=-\infty}^{\infty} r^{|n|} r^{|m|} \hat{u}(n) \overline{\hat{v}(m)} \int_0^{2\pi} e^{i(n-m)\theta} d\theta \\
 &= 2\pi \sum_{n=-\infty}^{\infty} r^{2|n|} \hat{u}(n) \overline{\hat{v}(n)}, \quad \text{by Exercise 17.}
 \end{aligned}$$

23. Do what is suggested.
 25. Use the reflection principle to show that $\overline{f(\bar{z})} = f(z)$ and then that $\overline{f(-\bar{z})} = f(z)$. Hence, $f(-w) = f(w)$, as well.
 27. Let

$$z = \phi(w) = \frac{iw + 1}{iw - 1};$$

ϕ maps the upper half-plane onto the disc $\{|z| < 1\}$. Set $R = \phi \circ M$; then R is rational, and $|R(e^{i\theta})| = 1$ for all θ . Let a_1, \dots, a_p be the zeros of R in $\{|z| < 1\}$ and b_1, \dots, b_q be the poles of R in $\{|z| < 1\}$. Show that

$$R(z) = \lambda \prod_1^p \frac{z - a_j}{1 - \bar{a}_j z} \prod_1^q \frac{1 - \bar{b}_k z}{z - b_k}, \quad |\lambda| = 1.$$

- Then recover M : $M = \phi^{-1} \circ R$.
 29. One method is outlined; here is another. Let

$$\phi(w) = \frac{iw + 1}{iw - 1}.$$

Then $g(w) = \phi^{-1}(f(\phi(w)))$ is analytic on the upper half-plane, real on the real axis (except for those points x where $f(\phi(x)) = 1$). By the reflection principle, g extends to the lower half-plane. Hence, f extends analytically across the unit circle, except possibly at a finite number of points where f might have a pole. But f is continuous at all points of the unit circle. Hence, f extends analytically over the unit circle. Now apply the argument used in the solution to Exercise 27.

Section 4.4

1. Note that $\log[(x-t)^2 + y^2]$ is a harmonic function of (x, y) for each t , so

$$\Delta u = \frac{1}{2\pi} \int_{-\infty}^{\infty} g(t) \Delta(\log[(x-t)^2 + y^2]) dt = 0.$$

Furthermore,

$$\frac{\partial u}{\partial y} = \frac{1}{\pi} \int_{-\infty}^{\infty} g(t) \frac{y}{(x-t)^2 + y^2} dt,$$

the Poisson integral of g , so

$$\lim_{y \downarrow 0} \frac{\partial u}{\partial y}(x, y) = g(x).$$

3. As in Exercise 1; $\Delta u = 0$, and $\frac{\partial u}{\partial y}$ = Poisson integral of $g - \frac{2Ay}{x^2 + y^2}$.


Finally,

$$\begin{aligned}|u(x, y)| &= \left| \frac{1}{2\pi} \int_{-M}^M g(t) [\log\{(x-t)^2 + y^2\} - \log(x^2 + y^2)] dt \right| \\&\leq \left[\frac{1}{2\pi} \int_{-M}^M |g(t)| dt \right] \max_{|t| \leq M} \left| \log \frac{(x-t)^2 + y^2}{x^2 + y^2} \right| \\&\leq C \max_{|t| \leq M} \left| \log \left(1 - \frac{2xt - t^2}{x^2 + y^2} \right) \right|; \quad \text{but } |\log(1-\delta)| \leq 2\delta \text{ for small } \delta \\&\leq C \frac{2|z|M + M^2}{|z|^2} \rightarrow 0 \quad \text{as } |z| \rightarrow \infty.\end{aligned}$$

5. $u(x, y) = \frac{1}{\pi} \int_0^\infty f(t) \left[\frac{x}{(y-t)^2 + x^2} + \frac{x}{(y+t)^2 + x^2} \right] dt$

7. $T(x, y) = \frac{T_1 + T_2}{2} + \frac{T_2 - T_1}{\pi} \operatorname{Re} \left[\arcsin \left(\frac{z^2 - \frac{1}{2}\sigma^2}{\frac{1}{2}\sigma^2} \right) \right].$

9. Let the circles be centered on the real axis and internally tangent at $z_0 = 1$ (see diagram).



Then the potential is

$$\phi(z) = \frac{(1-a)V_0 - (1-b)V_1}{b-a} + \frac{(1-a)(1-b)}{b-a}(V_0 - V_1) \operatorname{Re} \left(\frac{1}{z-1} \right).$$

11. $T(z) = T_1 + \frac{2}{\pi} (T_2 - T_1) \operatorname{Arg} \left(\frac{1+z}{1-z} \right)$

13. Write $z = re^{it}$ and $a = |a|e^{i\theta}$; then

$$2g(re^{it}) = \log(1 - 2r|a| \cos(\theta - t) + |a|^2 r^2) - \log(r^2 - 2|a|r \cos(\theta - t) + |a|^2).$$

Now compute.

15. $v(z) = \frac{V_0}{\pi} [\operatorname{Arg}(\sqrt{1 + e^{-2z}} - 1) - \operatorname{Arg}(\sqrt{1 + e^{-2z}} + 1)]$

Section 4.5

1. $G - G_1$ has no singularity, is harmonic and bounded on D , and vanishes on the boundary of D .

3. $G(z; p) = \log \left| \frac{z - \bar{p}}{z - p} \right|$

5. $G(z; p) = \log \left| \frac{z^2 - \bar{p}^2}{z^2 - p^2} \right|$

7. $G(z; p) = \log \left| \frac{e^z - e^p}{e^z - e^p} \right|$

9. $G(z; p) = \log \left| \frac{\sqrt{z} + \sqrt{p}}{\sqrt{z} - \sqrt{p}} \right|$

11. The response at q to an impulse at p equals the response at p to an impulse at q .

13. (a) 1 (b) r_0 (c) $\frac{b-a}{4}$ (d) $\frac{a+b}{2}$

15. Let h be continuous on the unit circle; then $H(re^{i\theta}) = (1/2\pi) \int_0^{2\pi} h(e^{it}) P_r(\theta - t) dt$ is harmonic in $|z| < 1$, and by (10) of Section 3, $\lim_{r \uparrow 1} H(re^{i\theta}) = h(e^{i\theta})$, all θ . Take $\theta = 0$.

17. Follow the hint.

Chapter 5

Section 5.1

1. Note: $\hat{u}(0) = 0$; $\hat{u}(x) = \frac{2i}{x} \left[b \cos bx - \frac{\sin xb}{x} \right]$, $x \neq 0$

3. $\hat{u}(x) = \frac{4}{x^2} \left[1 - \frac{\sin x}{x} \right]$

5. $\hat{u}(x) = \frac{\pi}{2} \exp[2(2ix - |x|)]$

7. $\hat{u}(x) = 2\pi \left(\frac{\sin \sigma x}{x} \right) e^{-|x|}$

9. $\hat{u}(x) = \sqrt{\pi} e^{-x^2/4} - 2 \frac{\sin \sigma x}{x} e^{-\sigma^2}$

11. $\hat{u}(x) = \frac{\lambda}{v} \left[\frac{2 \sin(vx/2)}{x} \right]^2$

13. (ii) Make the change of variables $a + t = s$.

- (iii) Make the change of variables $s = tb$.

- (iv) Use the conclusion of Exercise 16, Section 5.2, that $u(x) \rightarrow 0$ as $|x| \rightarrow \infty$. Then integrate by parts.

15. Set $f(z) = e^{-izx}/(1 + z^4)$ and integrate f over the contour in Figure 5.2a if $x < 0$ and the contour in Figure 5.2b if $x > 0$. When $x < 0$, f has poles of order one at $\gamma = \frac{1+i}{\sqrt{2}}$ and $i\gamma$ and the Residue Theorem gives

$$\begin{aligned}\hat{u}(x) &= 2\pi i(\text{Res}(f; \gamma) + \text{Res}(f; i\gamma)) \\ &= \frac{-i\pi}{\sqrt{2}} e^{x/\sqrt{2}} \{ \cos(x/\sqrt{2}) + \sin(x/\sqrt{2}) \}, \quad x < 0\end{aligned}$$

Likewise, if $x > 0$

$$\hat{u}(x) = \frac{i\pi}{\sqrt{2}} e^{-x/\sqrt{2}} \{ \cos(x/\sqrt{2}) - \sin(x/\sqrt{2}) \}.$$

17. Just compute.

$$\begin{aligned}19. \int_{-\infty}^{\infty} |u(x, y)| dx &\leq \frac{1}{\pi} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |f(t)| \frac{y}{(x-t)^2 + y^2} dt dx \\ &= \int_{-\infty}^{\infty} |f(t)| \left\{ \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{y}{(x-t)^2 + y^2} dx \right\} dt \\ &= \int_{-\infty}^{\infty} |f(t)| dt = M\end{aligned}$$

21. Follow the hints.

$$23. u(x, t) = \frac{1}{2} \int_{-t}^t g(x - \zeta) d\zeta$$

$$25. u(x, y) = \frac{1}{2\pi} \int_{-\infty}^{\infty} g(t) \log((x-t)^2 + y^2) dt$$

27. Following the hints, the equation simplifies to

$$c^2 \frac{\partial^2 u}{\partial x^2} = \frac{\partial^2 u}{\partial t^2} + 2b \frac{\partial u}{\partial t} + au.$$

Let $U(t) = \int_{-\infty}^{\infty} u(x, t) e^{-ixs} dx$, s fixed. Then $u'' + 2bu' + (a + c^2 s^2)u = 0$, so $U(t) = A \exp(\gamma_1 t) + B \exp(\gamma_2 t)$, where $\gamma_1 = -b + \sqrt{b^2 - a - c^2 s^2}$, $\gamma_2 = -b - \sqrt{b^2 - a - c^2 s^2}$ are the two roots of $r^2 + 2rb + (a + c^2 s^2) = 0$, and A and B are found from initial conditions to be

$$A = \frac{\gamma_2 \hat{f}(s) - \hat{g}(s)}{\gamma_2 - \gamma_1}, \quad B = \frac{\hat{g}(s) - \gamma_1 \hat{f}(s)}{\gamma_2 - \gamma_1}.$$

- (ii) In the case that $RC = SL$, we find that $b^2 = a$; then $U = \hat{u}$ simplifies to

$$\begin{aligned}\hat{u}(s) &= (\gamma_2 - \gamma_1)^{-1} e^{-bt} [(\gamma_2 \hat{f}(s) - \hat{g}(s)) e^{its} + (\hat{g}(s) - \gamma_1 \hat{f}(s)) e^{-its}] \\ &= (\gamma_2 - \gamma_1)^{-1} e^{-bt} \{ [(\gamma_2 f - g) * \delta_{-ic}] \hat{(s)} + [(g - \gamma_1 f) * \delta_{ic}] \hat{(s)} \},\end{aligned}$$

and consequently,

$$u(x; t) = \frac{e^{-bt}}{\gamma_2 - \gamma_1} [\gamma_2 f(x + tc) - \gamma_1 f(x - tc) + g(x - tc) - g(x + tc)].$$

Section 5.2

1. Let $u(t) = \begin{cases} 1, & |t| < \alpha \\ 0, & |t| > \alpha \end{cases}$; $v(t) = \begin{cases} 1, & |t| < \beta \\ 0, & |t| > \beta \end{cases}$.

Then by (3),

$$4\pi \min(\alpha, \beta) = 2\pi \int_{-\infty}^{\infty} u\bar{v} = \int_{-\infty}^{\infty} \hat{u}\bar{v} = 4 \int_{-\infty}^{\infty} \frac{\sin \alpha x \sin \beta x}{x^2} dx.$$

3. $\int_{-\infty}^{\infty} \left[\frac{1 - \cos \sigma x}{x} \right]^2 dx = \pi\sigma$

5. Take $u(t) = \frac{1}{1+t^2}$; $\hat{u}(x) = \pi e^{-|x|}$.

Hence, $2\pi \int_{-\infty}^{\infty} \frac{dt}{(1+t^2)^2} = \int_{-\infty}^{\infty} \pi^2 e^{-2|x|} dx = \pi^2$.

The value of the desired integral is $\pi/2$.

7. $u(t) = \frac{1}{2\pi} [\arctan(4+t) + \arctan(4-t)]$

9. $\frac{\pi}{12}(3\sigma - 3\sigma^2 + \sigma^3)$

11. By Exercise 8, $\hat{u}(z)$ is entire. Hence, $\hat{u} \equiv 0$, so $u \equiv 0$, by the Inversion Formula.

13. In (2), replace u with $u + v$; expand and use the known equality for u and v .

15. Follow the given steps.

17. Let $J = \lim_{M \rightarrow \infty} \int_{-M}^M |u(t)| dt$.

Then for $A, B \geq M_0$, $\varepsilon > J - \int_{-M_0}^{M_0} |u(t)| dt \geq \int_{-\infty}^{-B} + \int_A^{\infty}$.

19. Since $|u(t)| \leq |v(t)|$ for $-\infty < t < \infty$,

$$\int_{-M}^M |u(t)| dt \leq \int_{-M}^M |v(t)| dt \leq C \quad \text{for all } M.$$

21. Let γ be a triangle in the strip $\tau_1 < \operatorname{Im} z < \tau_2$. Then

$$\begin{aligned} \int_{\gamma} \hat{u}(z) dz &= \int_{\gamma} \left[\int_{-\infty}^{\infty} u(t) e^{-itz} dt \right] dz \\ &= \int_{-\infty}^{\infty} u(t) \left[\int_{\gamma} e^{-itz} dz \right] dt = \int_{-\infty}^{\infty} u(t) \cdot 0 dt = 0. \end{aligned}$$

23. Use the Riemann–Lebesgue Lemma (Exercise 15) with

$$w(t) = \begin{cases} 0, & -\infty < t < \delta \\ \frac{u(x+t)}{t}, & \delta \leq t < \infty. \end{cases}$$

This gives

$$0 = \lim_{M \rightarrow \infty} \operatorname{Re} \hat{w}(M) = \lim_{M \rightarrow \infty} \int_{\delta}^{\infty} \frac{u(x+t)}{t} \sin Mt dt.$$

Similarly,

$$\lim_{M \rightarrow \infty} \int_{-\infty}^{-\delta} \frac{u(x+t)}{t} \sin Mt dt = 0.$$

25. Combine 23 and 24:

$$\begin{aligned} \int_{-\infty}^{\infty} u(x+t) \frac{\sin Mt}{t} dt &= \int_{-\infty}^{-\delta} \frac{u(x+t)}{t} \sin Mt dt + \int_{\delta}^{\infty} \frac{u(x+t)}{t} \sin Mt dt \\ &\quad + \int_0^{\delta} \frac{u(x+t) - u^+(x)}{t} \sin Mt dt \\ &\quad + \int_{-\delta}^0 \frac{u(x+t) - u^-(x)}{t} \sin Mt dt \\ &\quad + [u^+(x) + u^-(x)] \int_0^{\delta} \frac{\sin Mt}{t} dt. \end{aligned}$$

The first four terms are small if δ is small and M is large. The fifth integral converges to

$$\int_0^{\infty} \frac{\sin x}{x} dx = \frac{\pi}{2}$$

as $M \rightarrow \infty$, δ fixed. (Make the change of variables $s = Mt$.)

Section 5.3

1. $(\mathcal{L}u)(s) = \frac{s}{A^2 + s^2}$

3. $(\mathcal{L}u)(s) = \frac{A}{(B+s)^2 + A^2}$

5. $(\mathcal{L}u)(s) = \frac{1}{s} [e^{-as} - e^{-bs}]$

7. $(\mathcal{L}u)(s) = \frac{1}{s^2} [1 - s\sigma e^{-s\sigma} - e^{-s\sigma}]$

9. $(\mathcal{L}u)(s) = \frac{1}{s^2}$

11. $\frac{\sin t}{t} = \sum_{k=0}^{\infty} (-1)^k \frac{t^{2k}}{(2k+1)!}; \quad \text{next, } \int_0^{\infty} t^{2k} e^{-st} dt = \frac{(2k)!}{s^{2k+1}}.$

Hence,

$$\mathcal{L}\left(\frac{\sin t}{t}\right)(s) = \sum_{k=0}^{\infty} (-1)^k \frac{1}{(2k+1)!} \left(\int_0^{\infty} t^{2k} e^{-st} dt \right) = \sum_{k=0}^{\infty} \frac{(-1)^k}{2k+1} s^{-(2k+1)}.$$

Furthermore

$$\begin{aligned} \frac{d}{ds} \left[\mathcal{L}\left(\frac{\sin t}{t}\right)(s) \right] &= \sum_{k=0}^{\infty} (-1)^{k+1} s^{-2k-2} = \frac{-1}{s^2 + 1}, \quad 1 < s < \infty \\ &= \frac{d}{ds} \arctan \frac{1}{s}, \quad 1 < s < \infty. \end{aligned}$$

13. $u(t) = \frac{t(\sin At)}{2A}$

15. $u(t) = \frac{1}{6}t^3 e^t$

17. $u(t) = ie^{-it}$

19. $\mathcal{L}u$ is entire by Morera's Theorem. Furthermore,

$$\begin{aligned} (\mathcal{L}u)(s) &= \int_0^A u(t) e^{-st} dt = \int_0^A u(t) \sum_{k=0}^{\infty} (-1)^k \frac{s^k t^k}{k!} dt \\ &= \sum_{k=0}^{\infty} (-1)^k \frac{s^k}{k!} \int_0^A u(t) t^k dt. \end{aligned}$$

21. $e^{(z-1)\log t}$ is an entire function of z ; $\Gamma(z)$ is defined for $\operatorname{Re} z > 0$ and analytic there, by Morera's Theorem.

23. Integrate by parts.

25. For $n = 1$, $\Gamma(1) = \int_0^{\infty} e^{-t} dt = 1$. Use induction and Exercise 23 for $n \geq 2$.

27. Follow the suggestion.

29. Again, follow the suggestions.

Section 5.4

1. $u(t) = \frac{6}{5}e^{-t} + 2e^t \left\{ -\frac{3}{5}\cos t + \frac{17}{10}\sin t \right\}$

3. $u(t) = e^{-t^2/2}$

$$5. \quad u(t) = \begin{cases} \sin t, & 0 \leq t \leq \pi \\ \left(1 + \frac{t-\pi}{2}\right) \sin t, & \pi < t < \infty \end{cases}$$

$$7. \quad u(x; t) = \begin{cases} 0, & \text{if } x > t \\ f(t-x), & \text{if } x \leq t \end{cases}$$

9. Case 1: ω is not an integer multiple of π . Then

$$u(x, t) = \frac{\sin \omega x \sin \omega t}{\sin \omega} + 2\omega \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{\omega^2 - n^2\pi^2} \sin n\pi x \sin n\pi t.$$

Case 2: $\omega = \pi$ (representative of $\omega = m\pi$).

$$u(x, t) = -2t \sin \pi x \cos \pi t - x \cos \pi x \sin \pi t - \frac{1}{2\pi} \sin \pi x \sin \pi t$$

$$+ \frac{2}{\pi} \sum_{n=2}^{\infty} \frac{(-1)^n \sin \pi nx \sin \pi nt}{n^2 - 1}.$$

$$11. \quad u(x, t) = \frac{\phi_0}{\sqrt{\pi k}} \int_0^t \frac{e^{-x^2/4ks}}{\sqrt{s}} ds$$

Section 5.5

$$1. \quad Z(\{a_j\}) = \frac{z^2}{z^2 - 1}, \quad |z| > 1$$

$$3. \quad Z(\{a_j\}) = e^{2/z}, \quad z \neq 0$$

$$5. \quad Z(\{a_j\}) = z \operatorname{Log} \left(1 - \frac{1}{z}\right), \quad |z| > 1$$

$$7. \quad Z(\{a_j\}) = \frac{z^2}{z^2 - 16} + \frac{3z}{9z^2 - 1}, \quad |z| > 4$$

$$9. \quad \begin{cases} 0, & j \text{ even} \\ \frac{2^j}{j!}, & j \text{ odd} \end{cases}$$

$$11. \quad a_j = \frac{1}{(j+1)!}$$

$$13. \quad a_0 = a_1 = 0; \quad a_j = j-1, \quad j = 2, 3, 4, \dots$$

15. $\{a_j\} * \{b_j\} = \{0\}$, so $0 = Z(\{0\}) = Z(\{a_j\} * \{b_j\}) = Z(\{a_j\})Z(\{b_j\})$. Since $Z(\{a_j\})$ and $Z(\{b_j\})$ are both analytic, one or the other must vanish identically; for that one, all its coefficients must be zero.

$$17. \quad y_0 = 1; \quad y_{r+1} = -\sum_{k=0}^r (x_k + x_{k+1} + x_{k+2}), \quad r = 0, 1, 2, \dots$$

19. $y_0 = 0; \quad y_1 = 1; \quad y_2 = x_0 + 1; \quad y_3 = -1 + x_1; \quad$ generally,

$$y_r = \begin{cases} (-1)^{m-1}(x_0 + 1) + \sum_{k=0}^{m-2} (-1)^k x_{r-2k-2}, & r = 2m, m = 2, 3, \dots \\ 1 + \sum_{k=0}^m (-1)^k x_{r-2k-2}, & r = 2m + 1 \end{cases}$$

21. $y_0 = 1; \quad y_{r+1} = -\sum_{k=0}^r \frac{x_k - 2x_{k+1} + x_{k+2}}{2^{r-k+1}}, \quad r = 0, 1, 2, \dots$

Section 5.5.1

1. $0 = R(w) = p\left(\frac{w+1}{w-1}\right)$ if and only if $\left|\frac{w+1}{w-1}\right| < 1$ if and only if $\operatorname{Re} w < 0$
3. (a) $q(w) = (A + B)w + (A - B)$
 (b) $q(w) = 16[w^4 + 4w^3 + 9w^2 + 8w + 5]$
 (c) $q(w) = 2w + 2$
 (d) $q(w) = -w^3 + 3w^2 - 3w + 9$
5. (a) p has a root at -1 .
 (b) All roots within $\{z: |z| < 1\}$
 (c) All roots within $\{z: |z| < 1\}$
 (d) One root outside $\{z: |z| < 1\}$
 (e) Some root outside $\{z: |z| < 1\}$

Index

absolutely convergent series, 38
absolutely integrable, 344
absolute value, 1, 3
additive identity, 11
additive inverse, 11
analytic function, definition of, 77
angle between curves, 209
aquifer, 305
arc-length, 72, 259
area formula, 244
argument, 7
argument principle, 175
associative, 11

Bernoulli's law, 270–271
Bessel functions, 135, 153, 353, 356
beta function, 355
biharmonic function, 280, 329–330
binomial formula, 98, 103, 105
boundary, 24
boundary point, 24
boundary-value problem, 298–307
bounded set, 28, 192
branch of $\log z$, 46

Cauchy estimates, 133
Cauchy–Goursat Theorem, 119
Cauchy–Riemann equations, 80, 82
Cauchy–Riemann equations in polar coordinates, 86
Cauchy's Formula, 111, 123
Cauchy's Theorem, 107
causal functions, 347
center of a power series, 93
chain rule, 78
charge density, 260
chordal distance, 207
circle, 16–20, 21, 199, 243
Circles of Appolonius, 20
circulation, 271–274
circulation of a linear fractional transformation, 201

closed curve, 56
closed half-plane, 27
closed set, 25, 191
coaxial, 299
commutative, 11
complex conjugate, 2
complex Fourier transform, 336–339
complex number, 1
complex plane, 9
complex potential, 132, 219
concentrated force, 309
concentration function, 278, 304
conformally equivalent domains, 225
conformal mapping, 209, 210–211, 221, 224–241
table of, 389–394
connected set, 25–26
conservative field, 88
continuous at a point, 36, 42, 79
continuous function, 36–37
continuous on a set, 36
convergent sequence, 33
convergent series, 37–38
convex set, 27
convolution, 325, 367
 $\cos z$, 48, 100
 $\cosh z$, 54, 126
Coulomb's Law, 255
cross-ratio, 206
cubic polynomial, 253, 382–383
curve, 56–59
 closed, 56
 orientation of, 59
 piecewise smooth, 57
 simple, 56
 smooth, 57

definite integrals evaluated by the Residue Theorem, 155–156
degree of a rational function, 181
de Moivre's Theorem, 6, 10

derivative of a power series, 97
differentiable at a point, 77
differential equation in the complex plane, 134
differential equations solved by the Laplace transform, 356–364
diffusion, 278, 303–305
dipole, 203
directional derivative, 71
Dirichlet problem, 298–301, 334
discrete Laplace transform, 366
divergent sequence, 33
divergent series, 38
domain, 27
domain of definition, 30

electrostatic field, 255–260
electrostatic potential, 258, 260, 308
energy integral, 249
Eneström–Kakaya Theorem, 382
entire function, 77, 130
equation
 Cauchy–Riemann, 80, 82
 heat, 327, 334, 365
 Laplace, 81, 245
 linear difference, 369, 376
 telegraph, 334
 Volterra integral, 364
equilibrium point, 258
essential singularity, 137, 150, 152
exact differential, 75
exponential function, 43–45, 78, 100
exponential growth, 346

falling string, 364
feedback control system, 190
field, 11
fixed point, 181, 198, 205
flow, 87, 131, 261–278
flux, 88
fluxless field, 88

- forcing function, 369
formula
 area, 244
 binomial, 98, 103, 105
 Cauchy, 111, 123
 Fourier inversion, 335, 345–346
 Laplace inversion, 349
 Poisson integral, 286–291
 Routh–Hurwitz condition, 188
 Fourier coefficient, 296
 Fourier inversion formula, 335, 345–346
 Fourier series, 296
 Fourier transform, 320–342
 Fresnel integrals, 117
function
 analytic, 77
 Bessel, 135, 153, 353, 356
 beta, 355
 biharmonic, 280, 329–330
 causal, 347
 concentration, 278, 304
 continuous, 36
 definition of, 30
 entire, 77
 exponential, 43
 forcing, 369
 gamma, 354
 Green's, 311–314
 harmonic, 73, 81, 245–252, 284–293, 296–297
 Heaviside, 347
 impulse, 309, 316
 inverse trigonometric, 52–53, 56
 logarithm, 45
 periodic, 49, 352
 piecewise continuous, 287
 positive rational, 196
 potential, 260–261
 rational, 37
 response, 369
 stream, 222, 261
 stress, 279
 trigonometric, 48
Fundamental Theorem of Algebra, 179
Gamma function, 354
Gauss–Lucas Theorem, 381
generating formula for Legendre polynomials, 182
geometric series, 39
gradient, 219, 258
graph, 32, 249–252
Green's Formula, 71–72, 310
Green's function, 311–314
Green's Identity, 74, 281
Green's Theorem, 65–70, 90
groundwater flow, 305
half-plane, 27
harmonic conjugate, 81, 247, 296
harmonic function, 73, 81, 245–252, 284–293, 296–297
heat equation, 327, 334, 365
heat kernel, 329
Heaviside function, 347
ideal fluid, 261
imaginary axis, 9
imaginary part, 1, 9
improper integral, 344–345
impulse function, 309, 316
infinite series, 37–40
initial conditions, 184, 347, 356, 369
input, 369, 374
inside of a curve, 57
integral representation of harmonic functions, 284–291
interior point, 22
inverse trigonometric function, 52–53, 56
inversion, 22, 200
irrotational flow, 88, 90
isolated singularity, 135
isolated zero, 172
Jordan Curve Theorem, 56
Jordan's Lemma, 159
Joukowski airfoil, 275–278
Kelvin's Minimum-Energy Theorem, 281
kernel
 heat, 329
 Poisson, 114, 118, 286
keyhole contour, 58, 165
kinetic energy, 281
Kutta–Joukowski Theorem, 274
ladder network, 371
Laplace inversion formula, 349
Laplace's equation, 81, 245
Laplace transform, 346–352
 table of, 395
Laplacian, 72, 310
Laplacian in polar coordinates, 252
Laurent series, 141–149
law
 Bernoulli's, 270–271
 Coulomb's, 255
Legendre polynomial, 182
Legendre's differential equation, 182
lemniscate, 218
level curve, 212
lifting force, 274
limit, 33–36, 318
linear difference equation, 369, 376
linear fractional transformation, 31, 85, 196–204, 211, 246
line integral, 60–65
Liouville's Theorem, 130, 133
locally irrotational, 90
locally sourceless, 90
local maximum, 191–192
logarithmic capacity, 316
logarithm of an analytic function, 131
logarithm of z , 45–48, 83
maximum modulus principle, 191, 192, 193, 248
Mean-Value Theorem, 193, 248
modulus, 1
Morera's Theorem, 129
multiplicative identity, 11
multiplicative inverse, 11
multiplicity of a zero, 171
Neumann problem, 301–307, 309
normal derivative, 71–72, 301
 n th root, 14
 n th partial sum, 37
Nyquist stability criterion, 190
one-to-one, 45, 191, 196, 210, 225
open disc, 22
open half-plane, 27
open mapping, 191
open set, 22, 30
order of a pole, 137
order of a zero, 128, 171
orientation of a curve, 59
output, 369, 374
outside of a curve, 57
parametrization of a curve, 57
Parseval's equality, 297, 335
period, 352
periodic function, 49, 352
permeability, 301–304, 305
piecewise continuous, 287
piecewise smooth curve, 57
piecewise smooth function, 319
point at infinity, 28
Poisson integral formula, 286–291
Poisson kernel, 114, 118, 286
polar representation, 4–5
pole, 137
polygon, 225, 227–230
polygonal curve, 25, 29
polynomial, 37, 78, 155, 179, 253, 381–387
positively oriented, 59, 65
positive rational function, 196
potential function, 260, 261
potential theory, 260
power of a complex number, 46
power series, 93–102, 123, 132
principal part, 144

- principle**
 branch of $\log z$, 46
 maximum-modulus, 191–193,
 248
 reflection, 291–293, 297–298
product rule, 78
- quadratic formula**, 9, 21
quartic polynomial, 384–385
quotient rule, 78
- radius of convergence**, 95, 106
range, 30
rational function, 37, 78, 147, 181,
 196, 376
ratio test, 42
real axis, 9
real number, 1
real part, 1, 9
reflection, 206
reflection principle, 291–293,
 297–298
removable singularity, 136, 272
residue, 138, 139
Residue Theorem, 154, 169
resonance, 364
response function, 369
reverse orientation, 59
Riemann Mapping Theorem, 225,
 312
Riemann–Lebesgue Lemma, 343
root, 13–14, 178
root test, 42
Rouche's Theorem, 177
Routh–Hurwitz conditions, 188
- scaling**, 22
Schwarz–Christoffel
 transformation, 225, 227–241
Schwarz inequality, 10
Schwarz Lemma, 193
- separation of a point and a convex**
 set, 30
shifting theorem for the
 Z-transform, 369
simple curve, 56
simply-connected domain, 107,
 225, 312
singularity
 at infinity, 152
 essential, 137, 150, 152
 isolated, 135
 removable, 136
sinh z , 54, 126
sink, 202, 265, 269
sin z , 48, 100
smooth curve, 57
source, 202, 264, 269
sourceless flow, 88, 90
spiral vortex, 283
stability, 183–185, 375–378
stable polynomial, 185–188
stagnation point, 91, 275
star-shaped set, 29
straight line, 13, 21, 199
strain energy, 249
stream function, 222, 261
streamlining, 220
stress function, 279
Strum sequence, 386–387
summation of series by residues,
 168
system transfer function, 375
- telegraph equation**, 334
temperature distribution, 254–255
theorem
 Cauchy, 107
 Eneström–Kakaya, 382
 Gauss–Lucas, 381
 Green's, 65–70
 Jordan Curve, 56
- theorem (continued)**
 Kelvin's Minimum-Energy, 281
 Kutta–Joukowski, 274
 Liouville's, 130
 Mean-Value, 193, 248
 Morera's, 129
 Residue, 154
 Riemann mapping, 225
 Riemann–Lebesgue, 343
 Rouche's, 177
 Schwarz Lemma, 193
 shifting for Z-transform, 369
thermal conductivity, 255
transform
 complex Fourier, 336–339
 Fourier, 320–342
 Laplace, 346–352
 Schwarz–Christoffel, 227–241
 Z-, 365–373
translation, 22
traveling wave, 364
triangle inequality, 12
trigonometric functions, 48–52,
 157, 161
- unbounded set**, 28
upper half-plane, 221, 288
- vector field**, 88
Volterra integral equation, 364
vortex, 282, 283
- wave equation**, 333, 365
work, 88
- zeros of an analytic function**, 128,
 171–179
zeros of a polynomial, 154, 179,
 185–188, 381–387
Z-transform, 365–373

(continued from front flap)

- AN INTRODUCTION TO ALGEBRAIC STRUCTURES, Joseph Landin. (65940-2)
FINITE DIFFERENCE EQUATIONS, H. Levy and F. Lessman. (67260-3)
GAMES AND DECISIONS: INTRODUCTION AND CRITICAL SURVEY, R. Duncan Luce and Howard Raiffa. (65943-7)
FIRST ORDER MATHEMATICAL LOGIC, Angelo Margaris. (66269-1)
TOPOLOGY, George McCarty. (65633-0)
INTRODUCTION TO TOPOLOGY, Bert Mendelson. (66352-3)
THE WORLD OF MATHEMATICS, James R. Newman (ed.). (41153-2, 41150-8, 41151-6, 41152-4) Four-volume set
GEOMETRY: A COMPREHENSIVE COURSE, Dan Pedoe. (65812-0)
FUNCTIONAL ANALYSIS, Frigyes Riesz and Béla Sz.-Nagy. (66289-6)
APPLIED PROBABILITY MODELS WITH OPTIMIZATION APPLICATIONS, Sheldon M. Ross. (67314-6)
INTRODUCTION TO THE CALCULUS OF VARIATIONS, Hans Sagan. (67366-9)
MATRICES AND LINEAR ALGEBRA, Hans Schneider and George Phillip Barker. (66014-1)
GROUP THEORY, W. R. Scott. (65377-3)
STATISTICAL METHOD FROM THE VIEWPOINT OF QUALITY CONTROL, Walter A. Shewhart. (65232-7)
LECTURES ON CLASSICAL DIFFERENTIAL GEOMETRY (SECOND EDITION), Dirk J. Struik. (65609-8)
ORDINARY DIFFERENTIAL EQUATIONS, Morris Tenenbaum and Harry Pollard. (64940-7)
MODERN ALGEBRA, Seth Warner. (66341-8)
ADVANCED CALCULUS, David V. Widder. (66103-2)
CHALLENGING MATHEMATICAL PROBLEMS WITH ELEMENTARY SOLUTIONS, A. M. Yaglom and I. M. Yaglom. (65536-9, 65537-7) Two-volume set
INTRODUCTION TO PARTIAL DIFFERENTIAL EQUATIONS WITH APPLICATIONS, E.C. Zachmanoglou and Dale W. Thoe. (65251-3)
USES OF INFINITY, Leo Zippin. (41178-8)

Paperbound unless otherwise indicated. Available at your book dealer, online at www.doverpublications.com, or by writing to Dept. 23, Dover Publications, Inc., 31 East 2nd Street, Mineola, NY 11501. For current price information or for free catalogs (please indicate field of interest), write to Dover Publications or log on to www.doverpublications.com and see every Dover book in print. Each year Dover publishes over 500 books on fine art, music, crafts and needlework, antiques, languages, literature, children's books, chess, cookery, nature, anthropology, science, mathematics, and other areas.

Manufactured in the U.S.A.

COMPLEX VARIABLES

Second Edition

Stephen D. Fisher

The most important topics in the theory and application of complex variables receive a thorough, coherent treatment in this introductory text. Intended for undergraduates or graduate students in science, mathematics, and engineering, this volume features hundreds of solved examples, exercises, and applications designed to foster a complete understanding of complex variables as well as an appreciation of their mathematical beauty and elegance.

Prerequisites are minimal: a three-semester course in calculus will suffice to prepare students for discussions of these topics: the complex plane, basic properties of analytic functions (including a rewritten and reorganized discussion of Cauchy's Theorem), analytic functions as mappings, analytic and harmonic functions in applications, and transform methods. Useful appendices include tables of conformal mappings and Laplace transforms, as well as solutions to odd-numbered exercises.

Students and teachers alike will find this volume, with its well-organized text and clear, concise proofs, an outstanding introduction to the intricacies of complex variables.

Unabridged Dover (1999) republication of the work published by Wadsworth & Brooks, Pacific Grove, CA, 1990. Preface. Appendices. Index. 448pp. 6 $\frac{1}{2}$ x 9 $\frac{1}{4}$. Paperbound.

Free Dover Mathematics and Science Catalog (59065-8) available upon request.

See every Dover book in print at www.doverpublications.com

ISBN 0-486-40679-2