Statistics. Simplified.



ABOUT

COURSE

BASIC STATS

MACHINE LEARNING

SOFTWARE TUTORIALS ✓ TOOLS ✓

Wilcoxon Signed-Rank Test Calculator

BY ZACH MAY 4, 2020

The Wilcoxon Signed-Rank Test is the non-parametric version of the paired t-test. It is used to test whether or not there is a significant difference between two population means.

To perform a Wilcoxon Signed-Rank Test, simply fill in the data values for two samples below and then click the "Calculate" button.

SEARCH

Search ...

Q

ABOUT

Statology is a site that makes learning statistics easy by explaining topics in simple and straightforward ways. Learn more about us.

Sample 1

14, 17, 12, 15, 15, 9, 12, 13, 13, 15, 19, 17, 14, 14, 16

Sample 2

15, 17, 15, 15, 17, 14, 9, 14, 11, 16, 18, 20, 20, 10, 17

CALCULATE

W test statistic = 29.5

Number of non-tied pairs (n) = 13

STATOLOGY STUDY

Statology Study is the ultimate online statistics study guide that helps you study and practice all of the core concepts taught in any elementary statistics course and makes your life so much easier as a student.




INTRODUCTION TO STATISTICS COURSE

Introduction to Statistics is our premier online video course that teaches you all of the topics covered in introductory statistics.

Get started with our course today.

Refer to the critical values table below to decide whether or not the test result is statistically significant based on *n* and your chosen alpha level.

If the test statistic W is less than the value found in the critical values table below, then the test result is statistically significant.



RECENT POSTS

Pandas: How to Apply Function to Multiple Columns

How to Write Pandas Series to CSV (With Example)

How to Use the combine_first()
Function in Pandas

	Alpha value				
n	0.005	0.01	0.025	0.05	0.10
5	_	_	-	_	0
6	-	.=	-	0	2
7	-	-	0	2	3
8	-	0	2	3	5
9	0	1	3	5	8
10	1	3	5	8	10
11	3	5	8	10	13
12	5	7	10	13	17
13	7	9	13	17	21
14	9	12	17	21	25
15	12	15	20	25	30
16	15	19	25	29	35
17	19	23	29	34	41
18	23	27	34	40	47
19	27	32	39	46	53
20	32	37	45	52	60
21	37	42	51	58	67
22	42	48	57	65	75
23	48	54	64	73	83
24	54	61	72	81	91
25	60	68	79	89	100
26	67	75	87	98	110
27	74	83	96	107	119
28	82	91	105	116	130
29	90	100	114	126	140
30	98	109	124	137	151



View all posts by Zach

PREV

Friedman Test: Definition, Formula, and Example

NEXT

How to Perform a One-Way ANOVA by Hand

Leave a Reply

Your email address will not be published. Required fields are marked *

Comment *

Name *	
Email *	
Website	
Website	
POST COMMENT	

© 2023 Statology | Privacy Policy