

Study Notes

An Introduction of Probability Theory and Its Applications

Jianyu Zhou

October 30, 2014

Contents

1	The Sample Space	2
1.1	The Empirical Background	2
1.2	Examples	2
1.3	The Sample Space and Events	3
1.4	Relations among Events	3
1.5	Discrete Sample Spaces	4
1.6	Probabilities in Discrete Sample Spaces: Preparations	4
1.7	The Basic Definitions and Rules	4
1.8	Problems Solution	6
2	Elements of Combinatorial Analysis	7
2.1	Preliminary	7

1 The Sample Space

1.1 The Empirical Background

Events Events are the results of experiments or observations. The events should be distinguished between *compound* (*decomposable*) and *simple* (*indecomposable events*). A compound event is an aggregate of certain simple events.

Sample points Sample points are just the simple events. Every indecomposable result of the (idealized) experiment is represented by one, and only one, sample point.

Sample Space Sample space is the aggregate of all sample points.

1.2 Examples

Notes

1. All intuitive background sampling problems are abstractly equivalent to the scheme of **placing r balls into n cells**, in the sense that the outcomes differ only in their verbal description.
(There are some examples listed on P10.)
2. Here comes a question that how to calculate the size of sampling space (how many sample points in the sample space) when placing r balls into n cells? This question should be answered under different assumption.
 - Placing r **distinguishable** balls into n **distinguishable** cells.

$$N = n^r$$

- Placing r **indistinguishable** balls into n **distinguishable** cells.

?

- Placing r **distinguishable** balls into n **indistinguishable** cells.

?

- Placing r **indistinguishable** balls into n **indistinguishable** cells.

?

When we facing some particular problems, the model of distinguishable or indistinguishable balls is purely a matter of purpose and convenience.

1.3 The Sample Space and Events

Sample space is the universe of the sample points. All sample points are indecomposable and cover all the outcomes of an idealized experiment. The event is an aggregate of some sample points (one or the aggregate of all sample points is also an event).

New events can be defined in terms of two or more given events. Here comes the notation of the formal *algebra of events* (algebra of point sets).

1.4 Relations among Events

We use Ω to denote sample space and capitals to denote events (sets of sample points). The fact that a sample point x is contained in the event A is denoted by $x \in A$. Thus $x \in \Omega$ for every sample point x . We write $A = B$ only if the two events consist of exactly the same points.

Definitions

1. We shall use the notation $A = 0$ to express that the event A contains no sample points. The zero must be interpreted in **a symbolic sense and not as the numeral**.
2. The event consisting of all points not contained in the event A will be called the *complementary event* (or *negation*) and will be denoted by A' . In particular $\Omega' = 0$.
3. With any two events A and B we can associate two new events defined by the conditions “*both A and B occur*” and “*either A or B or both occur*”. These events will be denoted by AB (intersection of A and B) and $A \cup B$ (union of A and B). If A and B exclude each other, then there are no points common to A and B and the event AB is impossible; analytically this situation is described by the equation $AB = 0$ which should be read “ A and B are *mutually exclusive*”.
4. To every collection A, B, C, \dots of events we define some notions as follows.
 - The aggregate of sample points which belong to all the given sets will be denoted by $ABC \dots$ and called the *intersection* (or *simultaneous realization*).
 - The aggregate of sample points which belong to at least one of the given sets will be denoted by $A \cup B \cup C \cup \dots$ and called the union (or realization of at least one).
 - The events A, B, C, \dots are *mutually exclusive* if no two have a point in common (any two of the events are mutually exclusive), that is, if $AB = 0, AC = 0, \dots, BC = 0, \dots$

5. The symbols $A \subset B$ and $B \supset A$ signify that every point of A is contained in B ; they are read, respectively, A *implies* B and B *is implied by* A . **If this is the case**, we shall also write $B - A$ instead of BA' to denote the event that B *but not* A *occurs*. The event $B - A$ contains all those points which are in B but not in A . With this notation we can write $A' = \Omega - A$ and $A - A = 0$.

- If A and B are mutually exclusive, then the occurrence of A implies the non-occurrence of B and vice versa. Thus $AB = 0$ means the same as $A \subset B'$ and as $B \subset A'$.
- The event $A - AB$ means the occurrence of A but not of the A and B . Thus $A - AB = AB'$.

1.5 Discrete Sample Spaces

Discrete Sample Space A sample space is called discrete if it contains only finitely many points or infinitely many points which can be arranged into a simple sequence E_1, E_2, \dots

The probabilities of events in discrete sample spaces are obtained by mere additions, whereas in other spaces integrations are necessary.

1.6 Probabilities in Discrete Sample Spaces: Preparations

The notion of probability is based on an *idealized model*. In a sense, no ideal models exist in reality. But in many applications, it is sufficiently accurate to describe reality and the idealized model can be extremely useful when the experiment is impossible to reconstruct.

Here comes some interesting examples. When placing 3 distinguishable balls into 3 cells, there will be 27 distinguishable possibilities. It appears natural to assume that all sample points are *equally probable*, that is, that each sample point has probability $\frac{1}{27}$. If the 3 balls are indistinguishable, there remain 27 different possibilities, even though only ten different forms are distinguishable. This consideration leads us to attribute the following probabilities to this ten points as following table. Tablename

However, in our case with $r = n = 3$, Bose and Einstein showed that certain particles attribute equal probability $\frac{1}{10}$ to each of the ten sample points.

These examples teach us not to rely too much on a priori arguments and to be prepared to accept new and unforeseen schemes.

1.7 The Basic Definitions and Rules

Fundamental Convention Given a discrete sample space Ω with sample points E_1, E_2, \dots , we shall assume that with each point E_j there is associated

a number, called the *probability* of E_j and denoted by $P\{E_j\}$. It is to be non-negative and such that

$$P\{E_1\} + P\{E_2\} + \dots = 1.$$

Note that the point which probability is zero (which is impossible to occur) is not excluded. In discrete sample spaces, any sample point can be eliminated from the sample space. However, the numerical values of probabilities are not known in advance.

Definition The probability $P\{A\}$ of any event A is the sum of the probabilities of all sample point in it.

For the sample space,

$$P\{\Omega\} = 1.$$

It follows that for any event A ,

$$0 \leq P\{A\} \leq 1.$$

Consider now two arbitrary events A_1 and A_2 . To compute the probability $P\{A_1 \cup A_2\}$ that either A_1 or A_2 or both occur, we have to add the probabilities of all sample points contained either in A_1 or in A_2 , but each point is to be counted only once. Therefore,

$$P\{A_1 \cup A_2\} \leq P\{A_1\} + P\{A_2\}.$$

Theorem For any two events A_1 and A_2 the probability that either A_1 or A_2 or both occur is given by

$$P\{A_1 \cup A_2\} = P\{A_1\} + P\{A_2\} - P\{A_1 A_2\}.$$

If $A_1 A_2 = 0$, that is, if A_1 and A_2 are mutually exclusive, then the formula reduces to

$$P\{A_1 \cup A_2\} = P\{A_1\} + P\{A_2\}.$$

The probability $P\{A_1 \cup A_2 \cup \dots \cup A_n\}$ of the realization of at least one among n events can be computed by a formula analogous to the formula before and will be discussed later. Thus for arbitrary events A_1, A_2, \dots the inequality

$$P\{A_1 \cup A_2 \cup \dots\} \leq P\{A_1\} + P\{A_2\} + \dots$$

holds (Occasionally it's referred to as *Boole's inequality*). In the special case where the events A_1, A_2, \dots are mutually exclusive, we have

$$P\{A_1 \cup A_2 \cup \dots\} = P\{A_1\} + P\{A_2\} + \dots$$

Finally, the probabilities of all sample points in sample spaces should not be same. Although considering them sharing the same probability will simplify the problem, this assumption will be restricted almost entirely the study of games.

1.8 Problems Solution

- 1.

2 Elements of Combinatorial Analysis

2.1 Preliminary

todo...