

Problem 1: Let G be a subgroup of $\mathrm{GL}_n(k)$ and

$$I := \bigoplus_{d>0} P_d^G, \quad I_D := \bigoplus_{d>0} (D_d(\mathbb{R}^n))^G$$

(note that we exclude constants). A polynomial $f \in \mathbb{C}[x_1, \dots, x_n]$ is called G -harmonic if $u(f) = 0$ for all $u \in I_D$.

- (a) Show that f is G -harmonic iff $\langle u_1 u_2, f \rangle = 0$ for all $u_1 \in D(\mathbb{R}^n)$ and $u_2 \in I_D$.
(b) (optional) Let H be the vector space of G -harmonic polynomials. Prove that if G is a subgroup of O_n , then

$$\mathbb{C}[x_1, \dots, x_n] = H \oplus \mathbb{C}[x_1, \dots, x_n]I.$$

- (c) Show that f is SO_n -harmonic iff $\Delta(f) = 0$ and interpret the statement of (ii) in this case.

Proof. (a): If $\langle u_1 u_2, f \rangle = 0$ for all $u_1 \in D(\mathbb{R}^n)$ then in particular for $u_1 = 1$, $\langle u_2, f \rangle = 0$.

Conversely, suppose $u(f) = 0$ for $u \in I_D$. f can be split into $f_0 + f_1 + \dots + f_d$ where $f_k \in P_k$.
to do

(c): □

Problem 2: Let A be a k -algebra and a be algebraic over A with minimal polynomial p_a . Show that $\text{Spec}(a)$ is the set of roots of p_a in k .

Proof. $\text{Spec}(a)$ is defined as the set of $\lambda \in k$ for which $(a - \lambda)$ is invertible. We use the fact that

$$p_a(t) - p_a(\lambda) = (t - \lambda)q(t)$$

for some polynomial $q(t)$ with degree $\deg(p_a) - 1$. In the case $t = a$, this gives

$$p_a(a) - p_a(\lambda) = (a - \lambda)q(a)$$

$$-p_a(\lambda) = (a - \lambda)q(a)$$

$$(a - \lambda)^{-1} = -p_a(\lambda)^{-1}q(a)$$

(note that $q(a) \neq 0$ since p_a is minimal and $\deg(q) < \deg(p_a)$). So as long as $p_a(\lambda) \neq 0$, $\lambda \notin \text{Spec}(a)$. However, if $p_a(\lambda) = 0$, then $(a - \lambda)$ is not invertible because it divides $p_a(\lambda)$ i.e. it is a zero-divisor. \square

Problem 3: Let A be a k -algebra and M a *simple* A -module. Prove that

$$\dim_k(\operatorname{End}_A M) \leq \dim_k M \leq \dim_k A.$$

(here \leq is the order on cardinals).

Proof. Pick some nonzero $m \in M$ and let $f : \operatorname{End}_A M \rightarrow M$ be defined by $f : \varphi \mapsto \varphi(m)$. f is injective as follows: suppose $\varphi, \psi \in \operatorname{End}_A M$ and $\varphi(m) = \psi(m)$. Then define $M' \subseteq M$ by

$$M' := \{m' \in M : \varphi(m') = \psi(m')\}.$$

Because φ and ψ are linear, M' is a submodule of M . But M is simple, so M' is either 0 or M . If it is 0 then no such φ, ψ exist, thus f is injective. If $M' = M$, then $\operatorname{End}_A M$ is trivial. In either case, $\dim_k(\operatorname{End}_A M) \leq \dim_k M$.

For the other inequality, let $g : A \rightarrow M$ be the map $g : a \mapsto a \cdot m$. The image of g is a submodule of M , but M is simple so $\operatorname{im}(g)$ must be 0 or M and it is clearly not 0 because $g(1) = m \neq 0$. Thus g is surjective, establishing that $\dim_k M \leq \dim_k A$. \square

Problem 4: Let A be a ring with nonzero idempotent $e = e^2$. Note that the set eAe forms a ring with unit e . Prove the following:

- (a) For any left A -module M , there is a natural eAe -module structure on eM . Furthermore, for any A -modules M, N , there is a natural morphism of additive groups $f : \text{Hom}_A(M, N) \rightarrow \text{Hom}_{eAe}(eM, eN)$.
- (b) Multiplication on the right gives an algebra isomorphism $(eAe)^{\text{op}} \rightarrow \text{Hom}_A(Ae, Ae)$.
- (c) Assuming that $AeA = A$, the map f in (a) is a bijection.
- (d) Assuming $AeA = A$, an A -module M is simple iff the eAe -module eM is simple (and similarly for direct sums of simple modules).
- (e) (optional) Assuming $AeA = A$, there is an algebra isomorphism $A^{\text{op}} \rightarrow \text{Hom}_{eAe}(eA, eA)$.

Proof. (a): To give a module structure we have to specify how scaling and addition work. They are as follows:

$$eae \cdot em = e(aem) \in eM, \quad em_1 + em_2 = e(m_1 + m_2) \in eM.$$

For the morphism $f : \text{Hom}_A(M, N) \rightarrow \text{Hom}_{eAe}(eM, eN)$, take

$$f : \varphi \mapsto (em \mapsto e\varphi(m) \in eN).$$

To verify that $f(\varphi)$ is indeed eAe -linear,

$$\begin{aligned} f(\varphi)(eae \cdot em) &= e\varphi(aem) = eae\varphi(m) = eae \cdot f(\varphi)(em) \\ f(\varphi)(em_1 + em_2) &= e\varphi(m_1 + m_2) = e\varphi(m_1) + e\varphi(m_2) = f(\varphi)(em_1) + f(\varphi)(em_2). \end{aligned}$$

- (b): For $ebe \in eAe$, we have the associated linear map $Ae \rightarrow Ae$ given by

$$\Phi(ebe) = ae \mapsto ae \cdot ebe = (aeb)e \in Ae.$$

This map $\Phi : (eAe)^{\text{op}} \rightarrow \text{Hom}_A(Ae, Ae)$ is an algebra homomorphism; it is clearly linear, and to show that it is multiplicative,

$$\Phi(ebe) \circ \Phi(ece) = (ae \mapsto ae \cdot ece \cdot ebe = aecebe) = \Phi(ece \cdot ebe)$$

(note that the order is switched, reflecting the oppositeness of $(eAe)^{\text{op}}$). To show that it is an isomorphism, we have the inverse

$$\Phi^{-1} : \varphi \mapsto e\varphi(e)$$

and one can check that Φ, Φ^{-1} are indeed inverses:

$$(\Phi^{-1} \circ \Phi)(ebe) = \Phi^{-1}(ae \mapsto (aeb)e) = eeebe = ebe$$

and

$$(\Phi \circ \Phi^{-1})(\varphi) = \Phi(e\varphi(e)) = (ae \mapsto aee\varphi(e) = \varphi(ae)) = \varphi.$$

- (c): With the assumption that $AeA = A$, there is some linear combination $\sum_{i=1}^n a_i e b_i = 1$ where $a_i, b_i \in A$. Thus, any m can be expressed as a A -linear combination of elements in eM :

$$m = \sum_{i=1}^n a_i \cdot e(b_i m).$$

Using this idea, we can define the inverse of f as

$$f^{-1} : \varphi \mapsto (m \mapsto \sum_{i=1}^n a_i \cdot \varphi(e b_i m)).$$

It is simple to check that these are inverses. □