

## Review

# Systemic inflammatory response syndrome

M. G. DAVIES and P.-O. HAGEN

*Vascular Biology and Atherosclerosis Research Laboratory, Departments of Surgery, Duke University Medical Center, PO Box 3473, Durham, North Carolina 27710, USA*

*Correspondence to: Dr P.-O. Hagen*

**Background** Localized inflammation is a physiological protective response which is generally tightly controlled by the body at the site of injury. Loss of this local control or an overly activated response results in an exaggerated systemic response which is clinically identified as systemic inflammatory response syndrome (SIRS). Compensatory mechanisms are initiated in concert with SIRS and outcome (resolution, multiple organ dysfunction syndrome or death) is dependent on the balance of SIRS and such compensatory mechanisms. No directed therapies have been successful to date in influencing outcome.

**Method** This review examines the current spectrum and pathophysiology of SIRS.

**Results and conclusion** Further clinical and basic scientific research is required to develop the global picture of SIRS, its associated family of syndromes and their natural histories.

One of the most frequent and serious problems confronting clinicians is the management of infection and the systemic responses that it induces. From 100 000 to 300 000 patients develop bacteraemia annually in the USA<sup>1</sup> and shock is a complication of sepsis in almost 50 per cent of these patients, with a mortality rate of 40–60 per cent. The clinical features associated with sepsis were noted by Hippocrates in 400 BC when he remarked that 'in acute diseases, coldness of the extremities is a very bad sign'. Sepsis remains a clinical challenge for the surgeon in and out of the critical care unit. The incidence of sepsis has been increasing for the past 60 years and is the most common cause of death in critical care units in the USA and Europe. Although remarkable progress has been made in defining the pathophysiology of this disease, the terminology associated with research in the field has been confusing. In an attempt to stratify the spectrum of sepsis, a consensus conference of the Society of Critical Care Medicine and the American College of Chest Physicians was held in August 1991 to produce a series of universal definitions for the systemic inflammatory response syndrome (SIRS), sepsis and other clinical conditions related to sepsis<sup>2</sup> (Fig. 1). SIRS was developed to imply a clinical response arising from a non-specific insult and includes two or more defined variables (Table 1). Sepsis is defined as SIRS with a documented infection. The sequela of SIRS/sepsis is multiple organ dysfunction syndrome (MODS) which can be defined as failure to maintain homeostasis without intervention (Figs 2 and 3). Primary MODS is a direct result of a well defined insult, while secondary MODS develops not in direct response to the insult but as a consequence of a host response. MODS is recognized as a sequel of SIRS. While nearly all patients with sepsis develop dysfunction of one organ system, multiple organ dysfunction occurs in about 30 per cent of patients with sepsis. Similarly, MODS can be identified in over 30 per cent of patients with trauma, 24 per cent of those with acute pancreatitis and nearly 40 per cent of patients with burns.

There is a continuum from the development of SIRS to the onset of sepsis and progression to septic shock and multiple organ dysfunction. In a prospective survey of admissions to a tertiary care facility in the USA, 68 per cent of patients met the criteria for SIRS<sup>3</sup>. Of these, 26 per cent developed sepsis, 18 per cent developed severe sepsis and 4 per cent developed septic shock within 28 days. The interval between the identification of SIRS and the development of sepsis was inversely correlated with the number of SIRS criteria initially identified (two, three or four). Mortality rates in that study were 7, 16, 20 and 46 per cent respectively for SIRS, sepsis, severe sepsis and septic shock. Similar trends have been identified in the Italian SEPSIS study<sup>4</sup>. It is important to note that identification of SIRS alone in a patient in the intensive care unit has a poor specificity for predicting the development of sepsis and septic shock<sup>5</sup>. However, there is an increasing incidence of organ system failure as




Fig. 1 The inter-relationship between sepsis, systemic inflammatory response syndrome (SIRS) and infection. Note that bacteraemia may or may not denote sepsis. ▨, Blood-borne infection. (Reproduced from reference 2 with permission from the Society of Critical Care and Williams and Wilkins, Baltimore, Maryland, USA)

patients progress from SIRS to septic shock<sup>3</sup>. Another study has shown that the use of SIRS criteria in patients with trauma to predict the onset of MODS or infection is not reliable<sup>6</sup>.

### Physiology of inflammation


The innate ability of the body to defend itself is based on three elements: external barriers against invasion and tissue injury, non-specific systems against foreign

**Table 1** Criteria for systemic inflammatory response syndrome, sepsis and multiple organ dysfunction syndrome


SIRS	Systemic inflammatory response is a characteristic clinical response manifested by two or more of the following: Temperature above 38°C or below 36°C (rectal) Heart rate above 90 beats per min Respiratory rate above 20 breaths per min or $P_{aCO_2}$ less than 4.3 kPa WBC count above 12 000 cells per mm <sup>3</sup> , below 4000 cells per mm <sup>3</sup> or 10 per cent immature (bands) forms
Sepsis	SIRS with documented infection
Severe sepsis	SIRS with documented infection and haemodynamic compromise
MODS	A state of physiological derangement in which organ function is not capable of maintaining homeostasis

SIRS, systemic inflammatory response syndrome;  $P_{aCO_2}$ , arterial partial pressure of carbon dioxide; WBC, white blood cell; MODS, multiple organ dysfunction syndrome. (Reproduced from reference 2 with permission from the Society of Critical Care and Williams and Wilkins, Baltimore, Maryland, USA.)

pathogens and debris, and antigen-specific responses to foreign pathogens<sup>7</sup>. Inflammation is the body's initial non-specific response to tissue injury produced by mechanical, chemical or microbial stimuli. Inflammation is a rapid highly amplified controlled humoral and cellular response: the complement, kinin, coagulation and fibrinolytic cascades are triggered in tandem with the activation of phagocytes and endothelial cells<sup>8</sup>. This local response may



**Fig. 2** The pendulum and spectrum of systemic inflammatory response syndrome (SIRS), compensatory anti-inflammatory response syndrome (CARS) and mixed antagonist response syndrome (MARS). Tissue insult/injury triggers a triad of systems encompassing the macrophage cytokines and endothelial cells. This results in SIRS/CARS/MARS which results in end-organ dysfunction. This can progress to multiple organ dysfunction syndrome (MODS) particularly when aggravated by a second hit (another tissue insult/injury), or can move towards resolution particularly when second hits are avoided



**Fig. 3** Development of systemic inflammatory response syndrome (SIRS). Stage I: in response to an insult, the local environment produces cytokines which are primarily intended to evoke an inflammatory response, promote wound repair and recruit the cells of the reticuloendothelial system. In stage II, small quantities of cytokines are released into the circulation to enhance the local response. Macrophages and platelets are recruited and growth factor production is stimulated. The acute-phase response is tightly controlled by a simultaneous decrease in proinflammatory mediators and release of endogenous antagonists. This continues until the wound is healed, the infection has resolved and homeostasis is restored. Occasionally homeostasis is not re-established and stage III (SIRS) develops. The sequela of SIRS/sepsis is multiple organ dysfunction syndrome

be considered benign as long as the inflammatory process is regulated appropriately to keep cells and mediators sequestered. There are four major events in the inflammatory process: vasodilatation, increased microvascular permeability, cellular activation/adhesion and coagulation. Vasodilatation and increased microvascular permeability at the site of injury increase locally available oxygen and nutrients, and produce heat, swelling and tissue oedema. The local haemodynamic changes give rise to the four classical symptoms associated with local inflammation: rubor (erythema), tumor (oedema), calor (heat) and dolor (pain). The normal physiological response to stress and injury results in a series of cardiovascular changes (increases in heart rate, contractility and cardiac output) and neuroendocrine changes (increased release of catecholamines, cortisol, antidiuretic hormone, growth hormone, glucagon and insulin). There is an increased fluid requirement due to 'third spacing'. The major metabolic change that occurs in response to inflammation is an initial increase in oxygen consumption. The arteriovenous difference in oxygen content will remain normal provided oxygen delivery is maintained; however, anaerobic metabolism will ensue if the body fails to meet the oxygen deficit<sup>9-11</sup>. Concurrent with the increased metabolic responses, there is a fall in systemic vascular resistance. Unless a second insult occurs, the peak effect of these local and systemic physiological changes occurs within 3-5 days after the initial stimulus and abates by 7-10 days. Clinically, a progressive decrease in 'third space' fluid requirement, and a downward trend in pulse and temperature followed by spontaneous diuresis, herald an uncomplicated and improving clinical course.

Cytokines are the physiological messengers of the inflammatory response and the principal molecules involved are tumour necrosis factor (TNF)  $\alpha$ , interleukins (IL-1 and IL-6), interferons and colony stimulating factors (CSFs). Polymorphonucleocytes (PMNs), monocytes/macrophages and endothelial cells are the cellular effectors of the inflammatory response. Leucocyte activation leads to increased leucocyte aggregation and tissue infiltration within the microcirculation where these leucocytes (PMNs and macrophages) undergo a respiratory burst, and increase their oxygen consumption and production of cytokines and other inflammatory mediators<sup>12</sup>. Endothelial cells exposed to this milieu of humoral and leucocyte-derived factors also become activated, and commence the expression of several adhesion molecules and receptors on their surface along with the synthesis and secretion of additional cytokines and secondary inflammatory mediators, including prostaglandins, leukotrienes, thromboxanes, platelet activating factor (PAF), oxygen free radicals, nitric oxide and proteases (cathepsin, elastase). Many of these secondary inflammatory mediators are also produced by leucocytes. The presence of activated endothelial cells and the enhanced cytokine milieu results in activation of the coagulation cascades which leads to local thrombosis minimizing blood loss and the walling off of injured tissues, thus attempting physiologically to isolate the inflamed areas.


### Pathophysiology of systemic inflammatory response syndrome

Osler said 'patients die not of their disease, they die of the physiological abnormalities of their disease'. Localized

inflammation is a physiological protective response which is generally tightly controlled by the body at the site of injury. Loss of this local control or an overly activated response results in an exaggerated systemic response which is clinically identified as SIRS. SIRS may be initiated by infection (viruses, bacteria, protozoa and fungi) or by non-infectious causes such as trauma, autoimmune reactions, cirrhosis and pancreatitis (*Fig. 1*). In a recent article, Bone<sup>13</sup> proposed that there were three stages in the development of SIRS (*Fig. 3*). In Stage I, in response to an insult, the local environment produces cytokines which are primarily intended to evoke an inflammatory response, promote wound repair and recruit the cells of the reticuloendothelial system. In Stage II small quantities of cytokines are released into the circulation to enhance the local response. Macrophages and platelets are recruited and growth factor production is stimulated. An acute-phase response is initiated and is tightly controlled by a simultaneous decrease in proinflammatory mediators and release of endogenous antagonists. These mediators keep the initial inflammation response in check both by downregulating cytokine production and by counteracting the effects of cytokines already released. This continues until the wound is healed, the infection resolved and homeostasis is restored. Occasionally homeostasis is not re-established and Stage III (SIRS) develops (*Fig. 3*). With the failure of homeostasis, a massive systemic reaction begins. The predominant effects of cytokines become destructive rather than protective. The flood of inflammatory mediators triggers numerous humoral cascades and results in sustained activation of the reticular endothelial system with loss of microcirculatory integrity and insults to various distant end organs (*Figs 3 and 4*).

Although flow and permeability changes at the local area increase nutrient delivery, uncontrolled systemic vasodilatation produces a sustained decrease in systemic vascular resistance and hypotension, while increased systemic vascular permeability results in significant extravascular third spacing. Coupled to these events there is depression of myocardial contractility which may be due to the effect of paracrine nitric oxide production and coronary non-occlusive microvascular damage and myocyte injury<sup>14</sup>. These phenomena make it difficult to volume resuscitate a hypotensive patient adequately. In combination, the loss of peripheral vascular tone and the loss of volume into extravascular spaces negate the normal homeostatic response necessary to maintain oxygen delivery and correct the abnormal arteriovenous difference in oxygen content<sup>10,15</sup>. The inability physiologically to correct these adverse responses results in end-organ hypoperfusion, oedema, initiation of anaerobic metabolism and end-organ dysfunction.

Early in the SIRS process, large numbers of leucocytes become adherent to the activated endothelial cells of the vessel wall and can interrupt the microcirculatory flow<sup>16</sup>. Leucocyte adherence is partially related to an increase in the number of adhesion molecules present on the endothelial cells. TNF- $\alpha$ , IL-1 and many other inflammatory mediators trigger endothelial cells to express new or increased numbers of adhesion molecules. In addition to the mechanical interruption of microcirculatory flow, activated leucocytes can damage adjacent endothelial cells and the surrounding extravascular tissue. TNF- $\alpha$  and IL-1 are considered primary proinflammatory mediators and induce a range of secondary proinflammatory mediators called chemokines.



**Fig. 4** The cellular biology of systemic inflammatory response syndrome. Lipopolysaccharide (LPS) can interact with lipopolysaccharide binding protein (LBP) and bactericidal/permeability increasing protein (BPI). LBP in turn interacts with the LPS receptor CD14 inducing cytokine expression. Tissue insult/injury triggers a triad of effector systems: macrophages, cytokines and endothelial cells. The principal cytokines are tumour necrosis factor (TNF)  $\alpha$ , interleukin (IL) 1 and IL-6. Polymorphonucleocytes (PMNs), macrophages and endothelial cells are the cellular effectors of the inflammatory response. Endothelial cells exposed to this milieu of humoral and leucocyte-derived factors also become 'activated' and commence the expression of several adhesion molecules and receptors on their surface and the synthesis/secretion of additional cytokines and secondary inflammatory mediators. ICE, interleukin converting enzyme; TNF- $\alpha$ R, TNF- $\alpha$  receptor; IL-1R, IL-1 receptor; IL-1RA, IL-1 receptor antagonist; PAF, platelet activating factor; TM, thrombomodulin; TF, tissue factor; tPA, tissue plasminogen activator; PAI-1, plasminogen activator inhibitor 1; VCAM, vascular cell adhesion molecule; ICAM, intercellular adhesion molecule 1; PECAM, platelet-endothelial cell adhesion molecule; COX, cyclooxygenase; XO, xanthine oxidase; iNOS, inducible nitric oxide synthase; SMC, smooth muscle cell

Two major groups of chemokines have been categorized, the CXC subfamily which is chemotactic for neutrophils, and the CC subfamily which is chemotactic for monocytes<sup>17</sup>.

Activated endothelial cells express multiple factors (e.g. tissue factor, platelet-endothelial cell adhesion molecule, thromboxane (TX)  $A_2$ ) which convert their local environment from a neutral coagulant environment to a procoagulant one. In addition, TNF- $\alpha$  triggers the coagulation cascades by activation of the extrinsic pathway<sup>18</sup>. Endotoxin triggers both the coagulation and fibrinolysis cascades<sup>19</sup>. Factor XII has a central role in the pathogenesis of septic shock activated by the peptidoglycan residues and teichoic acid from the cell wall of Gram-positive organisms as efficiently as by lipopolysaccharide (LPS) and lipid A from Gram-negative organisms<sup>20</sup>. Factor XIIa triggers both the intrinsic coagulation pathway through activation of Factor XI, and induces both endothelial cells and macrophages to produce tissue factor, which in turn activates the extrinsic coagulation pathway. Antibodies to tissue factor prevent LPS-induced disseminated intravascular coagulation in rabbits<sup>21</sup>. Endotoxaemia causes an increase in the levels of tissue plasminogen activator (tPA) in plasma which is rapidly counterbalanced by the release of plasminogen activator inhibitor (PAI) 1<sup>22</sup>. In sepsis, plasma levels of protein C and antithrombin III decrease and there are increased plasma levels of PAI-1. Plasma thrombomodulin, which is derived from degradation of endothelial cell membrane thrombomodulin, is also increased in SIRS<sup>23</sup>. The multiple actions of thrombin and the failure

of natural inhibitory mechanisms, such as antithrombin III, protein S, protein C and plasma fibrinolysis inhibitors, also contribute to this process<sup>24</sup>. This procoagulant environment coupled to endothelial cell injury predisposes to the development of excessive microthrombi, further obstructing local blood flow and exacerbating end-organ dysfunction.

These potentially destructive systemic and regional responses in SIRS (increased peripheral vasodilatation, excessive microvascular permeability, accelerated microvascular clotting, leucocyte/endothelial cell activation) contribute to the development of profound pathophysiological changes in the various organs, and are considered the major aetiological factors in the development of septic shock, disseminated intravascular coagulation<sup>25</sup>, adult respiratory distress syndrome (ARDS) and other end-organ dysfunction leading to MODS. Linked to these end-organ failures are the metabolic and nutritional effects of an activated cytokine milieu which produces fever, hypermetabolism, anorexia, protein catabolism, cachexia, and altered fat, glucose and trace mineral metabolism<sup>26</sup>. These processes are accelerated if a second insult such as shock, infection or ischaemia follows the initial injury.

### Mediators of systemic inflammatory response syndrome

The mediator response in SIRS may be divided into four phases based on the cytokine/cellular response: induction,

triggering of cytokine synthesis, evolution of cytokine cascade and elaboration of secondary mediators with ensuing cellular injury. Of the multitude of mediators operating in SIRS/sepsis, the three most influential appear to be TNF- $\alpha$ , IL-1 and IL-6. Raised serum levels of TNF- $\alpha$  and IL-6 are seen in subjects challenged with endotoxin and in patients with sepsis<sup>27,28</sup>. Whereas absolute levels of TNF- $\alpha$  and IL-6 are crudely predictive of death from septic shock, persistence of TNF- $\alpha$  and IL-6 in serum is highly predictive of the development of MODS and death<sup>29,30</sup>. The deleterious host response to infection has been dissected extensively with regard to Gram-negative bacterial sepsis and endotoxaemia leading to cytokine release. Specific attention has been directed toward TNF- $\alpha$ , IL-1 and IL-6 because their release has been most closely associated with morbidity and death<sup>28</sup>. IL-6 also appears to be associated with the host septic response, probably being released in response to secretion of either TNF- $\alpha$ , IL-1 or both cytokines. Abrogation of IL-6 using IL-6 antibodies has been associated with improved survival in animal models of Gram-negative bacterial sepsis or in models in which TNF- $\alpha$  is infused. Infusion of IL-6 itself does not have any effect. Similarly, although direct injection of IL-8 does not produce adverse effects, raised levels of this cytokine have been demonstrated following a challenge of either Gram-negative bacteria or endotoxin in experimental models, but a great deal of information suggests that this cytokine acts to potentiate the effects of other mediators.

The events following endotoxin exposure provide a good model on which to discuss the four phases of SIRS. Endotoxin, shed from the bacteria as they multiply or die, is one of the most powerful triggers of SIRS by stimulating phagocytic cells, particularly macrophages, to synthesize TNF- $\alpha$  and IL-1, to activate the complement/coagulation cascades and to induce endothelial cell activation. Cytokines are not stored and their synthesis requires gene transcription or translation of messenger RNA (mRNA). In the baboon, after exposure to endotoxin, serum levels of TNF- $\alpha$  peak at 1.5 h, IL-1 $\beta$  at 3 h and interferon (IFN)  $\gamma$  at 6 h<sup>31</sup>. In humans, infusions of either endotoxin, TNF- $\alpha$  or IL-1 cause myalgia, chills, headache, nausea and tachycardia following which cardiac output increases and systemic vascular resistance falls<sup>32</sup>. These symptoms begin 90–120 min after endotoxin exposure, concurrent with the time TNF- $\alpha$  levels rise. In contrast, these symptoms occur almost immediately after TNF- $\alpha$  infusion. Although some investigators have identified the presence of raised IL-1 levels after endotoxin exposure, this is not a consistent finding. Similarly, increased IFN- $\gamma$  concentrations have not been demonstrated after an infusion of endotoxin in humans. These studies have been extended to profile the role of cytokines in patients who develop sepsis. Raised levels of TNF- $\alpha$ , IL-1, IL-6, IL-8 and IFN- $\gamma$  have been reported to occur in groups of patients with sepsis, but the pattern of cytokine abnormalities in each individual may not be similar to that which would be expected based on data obtained from numerous animal studies<sup>28</sup>. Furthermore, the correlation of raised specific cytokines with death has not been demonstrated uniformly. In addition to the circulating levels of cytokines there are increased serum concentrations of cytokine antagonists (soluble TNF- $\alpha$  receptor and IL-1 receptor antagonist) and antiendotoxin core antibodies in patients with SIRS; these antagonists were present at concentrations 30–100 000-fold greater than their respective cytokines<sup>33</sup>. There are increased

complement levels in sepsis, and these have been associated with fatal outcome in both Gram-positive and Gram-negative septic shock<sup>34–36</sup>. Survivors of SIRS/sepsis show improvement over the first 72 h in high molecular weight kininogen concentrations and higher than normal Factor V values compared with non-survivors. Persistently low serial Factor XII, high molecular weight kininogen and Factor V are associated with a poor prognosis<sup>37</sup>. Enhanced levels of C-reactive protein are seen in SIRS/sepsis and a decrease in C-reactive protein level precedes clinical resolution of the condition<sup>38</sup>.

### *Endotoxin*

The Gram-negative bacterial wall consists of inner and outer membranes, the latter of which contains many proteins as well as LPS. LPS consists of three regions: the O antigen polysaccharide, the core and the lipid A regions. The lipid A region is considered to be responsible for the majority of the toxicity. LPS can interact with mammalian cell membranes by several different types of receptor, including CD11/18, CD14, the acetyl-low density lipoprotein scavenger receptor, other less well defined membrane proteins and by non-specific cell membrane lipid interactions. There is a family of serum proteins which possesses LPS binding sites, of which lipopolysaccharide binding protein (LBP) and bactericidal/permeability increasing protein (BPI) are the most researched<sup>39,40</sup>. Interaction of LPS with CD14 receptors requires the LPS first to bind to LBP<sup>41</sup>. CD14 response to LPS-LBP complex is enhanced compared with that to LPS alone. The enhancing effect of LBP provides an early warning mechanism for the presence of low concentrations of endotoxin. Blockade of CD14 offers an additional therapeutic option in controlling the cytokine response<sup>42</sup>. BPI binds to LPS and prevents macrophage activation, and is protective in rodent models of lethal endotoxaemia<sup>41</sup>. A soluble CD14 receptor has been identified and probably participates in all phases of SIRS with its ability to induce LPS sensitivity in cells normally insensitive to LPS (e.g. smooth muscle cells)<sup>43</sup>.

### *Tumour necrosis factor $\alpha$*

Various cells of the reticular endothelial cell system, such as monocytes, pulmonary macrophages, hepatic Kupffer cells and peritoneal macrophages produce TNF- $\alpha$ <sup>44</sup>. The expression of TNF- $\alpha$  is tightly controlled both at transcriptional and translational levels. The half-life of circulating TNF- $\alpha$  is short, 14–18 min, and it is degraded in several organs including the liver, skin, gastrointestinal tract and kidney. Specific receptors for TNF- $\alpha$  are found on a wide variety of cells and a maximal biological response is elicited by occupancy of as few as 5 per cent of these receptors<sup>45,46</sup>. The systemic and tissue-specific cellular mechanisms of TNF- $\alpha$  are dependent on its direct effects as well as the release of other soluble mediators from host cells. It elicits the release of neutrophils from the bone marrow and initiates neutrophil margination by inducing expression of adhesion molecules, promoting their transendothelial passage and activation (degranulation, production of superoxides and release of lysozymes). TNF- $\alpha$  promotes differentiation of monocytes and macrophages, and induces the activation of macrophages. It stimulates the synthesis of acute-phase proteins and activates the common pathway of the coagulation and complement systems. TNF- $\alpha$  produces a dose-dependent

increase in endothelial procoagulant activity and may inhibit thrombomodulin expression at the endothelial cell surface. It induces IL-1 release from endothelial cells and macrophages, while IL-1 subsequently stimulates the biosynthesis of other cytokines. The presence of IL-1 and other cytokines appears to enhance the sensitivity of tissues to the effects of TNF- $\alpha$ <sup>47</sup>. Exogenous administration of pharmacological doses of TNF- $\alpha$  to experimental animals evokes the pathophysiological events associated with SIRS<sup>48</sup>. In an overwhelming bacterial sepsis model in baboons, passive immunization with a monoclonal antibody to TNF- $\alpha$  before the bacterial challenge confers complete protection against both shock and death. Levels of IL-1 $\beta$  and IL-6 are also attenuated<sup>31</sup>. Passive immunization of mice with anti-TNF- $\alpha$  prevents lethal endotoxaemia<sup>49</sup>. C3H/HeJ mice, which are genetically deficient in the ability to synthesize TNF- $\alpha$ , tolerate lethal doses of endotoxin with minimal effects<sup>50</sup>.

### Interleukins

IL-1 appears to be released either in parallel or in response to TNF- $\alpha$  secretion<sup>51</sup>. Abrogation of the effects of endotoxin or TNF, by specific antibodies, reduces IL-1 levels and concurrently decreases the mortality rate in experimental animal models. Animal studies in which improvement in survival occurs after blockade of IL-1 activity by use of an IL-1 receptor antagonist (IL-1RA) have provided strong evidence supporting an important role of IL-1 in deleterious effects during Gram-negative bacterial sepsis<sup>52</sup>. Monocytes and tissue macrophages are the primary sources of IL-1. IL-1 consists of two distinct molecules, IL-1 $\alpha$  and IL-1 $\beta$ , that are structurally related polypeptides<sup>53</sup>. Most IL-1 $\alpha$  remains in the cytosol in a precursor form or is associated with the cell membrane in a biologically active form. The presence of a cell-associated form of IL-1 can explain the capability of activated macrophages to induce natural killer cell cytotoxicity, T cell proliferation and other functions by cellular contact in the absence of any releasable IL-1. IL-1 $\beta$  is cleaved by the IL-1 $\beta$  converting enzyme to its mature form within the cell after which it is secreted. IL-1 $\beta$  is also readily degraded from its precursor by trypsin, plasmin and other proteases<sup>51</sup>. There are two classes of high-affinity IL-1 receptors described and tissue distribution varies from 100 to 10 000 receptors per cell. Patients with sepsis have greatly enhanced expression of type II IL-1 receptor mRNA and cell surface receptors, and have raised levels of soluble IL-1 receptors which may represent a mechanism of regulating IL-1 activity in sepsis<sup>54</sup>. Both IL-1 $\alpha$  and IL-1 $\beta$  have a short half-life of 6–10 min. IL-1 $\alpha$  and IL-1 $\beta$  have not been detected in the circulation of human volunteers who received endotoxin intravenously, whereas TNF- $\alpha$  and IL-6 were detected readily<sup>55</sup>. The absence of IL- $\alpha$  during inflammation is consistent with its primary role as a membrane-bound cytokine principally involved in local paracrine and autocrine regulation. IL-1 is a strong inducer of granulocyte/macrophage CSF (GM-CSF), macrophage CSF and hepatic acute-phase protein synthesis. Excessive IL-1 release produces excessive margination of activated neutrophils into the vascular wall, stimulates endothelial cell procoagulant activity and increases leucocyte binding, but decreases heparan sulphate binding<sup>56</sup>. Unlike TNF- $\alpha$ , IL-1 is not directly lethal but IL-1 will reproduce many of the acute haematological and metabolic phenomena

associated with severe sepsis and is equipotent in inducing the synthesis of other cytokines<sup>44</sup>.

IL-6 is a family of at least six differentially modified phosphoglycoproteins which are released rapidly within 60 min in response to injury<sup>28</sup>. They act as a B cell stimulatory factor, a hybridoma/plasmacytoma growth factor, a hepatocyte stimulating factor and a cytotoxic T cell differentiation factor<sup>57</sup>. The prevailing subtype of IL-6 after an endotoxin challenge is a 26-kDa protein. IL-6 interacts synergistically with IL-1 to affect thymocyte proliferation, and in combination with TNF- $\alpha$  augments T cell proliferation and promotes PMN activation and accumulation. The temporal relationship of IL-6 appearance within the cytokine cascade suggests a strong relationship to antecedent TNF- $\alpha$  or IL-1 stimulation<sup>55</sup>. Transcription and production are enhanced in response to TNF- $\alpha$  and IL-1. When TNF- $\alpha$  or IL-1 activity is attenuated, the subsequent IL-6 response is decreased<sup>58</sup>. IL-6 administration does not cause haemodynamic compromise, regardless of the quantity given. IL-6 suppresses LPS-induced TNF- $\alpha$  production and TNF- $\alpha$ -induced IL-1 production. Anti-IL-6 monoclonal antibodies protect mice from lethal *Escherichia coli* infection and lethal exposure to TNF- $\alpha$ <sup>59</sup>.

IL-4 and IL-8 also participate in the response to injury. IL-4 synergistically increases TNF- $\alpha$  or IL-1-induced antigen expression in endothelial cells, but inhibits the increased expression of adhesion molecules by TNF- $\alpha$ , IL-1 or IFN- $\gamma$ . IL-8 is produced by the endothelial cells and is chemotactic for both neutrophils and lymphocytes. Administration of anti-IL-8 antibodies prevents neutrophil-dependent tissue infiltration and damage<sup>60</sup>. IL-4 enhances lymphocyte adhesion to the endothelial cell. It regulates growth and differentiation of T cells. IL-4 induces antigen expression on macrophages and suppresses IL-8 expression from stimulated monocytes but not from stimulated fibroblasts or endothelial cells.

IFN- $\gamma$  promotes the release of TNF- $\alpha$ , IL-1 and IL-6 by augmenting the effects of endotoxin on macrophages, thereby increasing the expression of adhesion molecules and cellular receptors for TNF- $\alpha$ . It may act synergistically with TNF- $\alpha$  to produce cytotoxic and cytostatic activity, synergistically increases IL-2 promotion of TNF- $\alpha$  release and promotes B cell activation to increase antibody production. IFN- $\gamma$  enhances adhesion of lymphocytes to endothelial cells, induces marked morphological changes in endothelial cells, and encourages PMN activation and accumulation. IFN- $\gamma$  enhances the phagocytic activity of PMNs and macrophage microbial function. IFN- $\gamma$  antagonizes the actions of GM-CSF. GM-CSF stimulates PMN phagocytosis, degranulation and cytotoxicity. In addition it promotes maturation of macrophages and enhances their activity.

### Cells

#### Endothelium

Because of its position, the endothelium both mediates and modulates the inflammatory and immunological responses in SIRS. The endothelium regulates the microvasculature, reacting to the metabolic needs of the tissue; it is essential in organ autoregulation and in the responses of these microvasculatures to changes in local blood flow through an extensive array of endogenously produced vasoactive factors<sup>61</sup>. The endothelium regulates intravascular coagulation by its participation in and

separation of procoagulant pathways, inhibition of procoagulant proteins, regulation of fibrinolysis and production of thromboregulatory compounds<sup>62</sup>. Basal secretion of tissue factor, a procoagulant enzyme, is low compared with that of the underlying smooth muscle cells and fibroblasts. However, if stimulated or injured, the endothelial cells can increase tissue factor production by tenfold to 40-fold. The basic barrier function of the endothelium separates intravascular coagulation factors (Factor VIIa) from tissue factor in the subendothelium and also prevents exposure of platelets to the proaggregating constituents of the subendothelium, such as collagen and von Willebrand factor. Furthermore, endothelial cells produce and express on their extracellular surfaces small amounts of the proteoglycan heparan sulphate, which serves to localize and increase the intrinsic activity of antithrombin III and tissue factor pathway inhibitor where it acts as a potent inhibitor of Factor Xa and, through its interactions with Factor Xa, produces feedback inhibition of the Factor VIIa–tissue factor complex. Endothelial cells inhibit procoagulant proteins with the protein C pathway, an autoregulatory pathway that involves protein C, protein S and thrombomodulin. Besides its direct effects on activated coagulant factors, protein C<sub>a</sub> also increases endothelial cell fibrinolytic activity by complexing with and decreasing the activity of PAI-1, thereby increasing fibrinolysis. Endothelial cells synthesize and assemble the plasminogen activators: urokinase (uPA) and tPA. *In vivo*, normal endothelial cells express tPA only. However, if stimulated by a variety of cytokines and circumstances, endothelial cells preferentially synthesize uPA and downregulate tPA synthesis. In addition to these two fibrinolytic enzymes, endothelial cells also secrete two PAIs, PAI-1 and PAI-2. Both are serine protease inhibitors and form equimolar complexes with either active uPA or tPA molecules. PAI-1 requires the presence of fibronectin in the extracellular matrix to maintain its active conformation.

The process of cell adherence, cell activation and cell migration involves an interplay between the expression of adhesion molecules by the endothelial cells, leucocyte activation and local cytokine activity<sup>63</sup>. The adhesion molecules involved in endothelial cell interactions with leucocytes are currently composed of three families: the selectins which govern the interaction of lymphocytes and neutrophils, the immunoglobulins which include antigen-specific receptors for T and B lymphocytes, and the integrins which are important in platelet adhesion and cell migration. The presence of the cytokines IL-1, TNF- $\alpha$  and transforming growth factor (TGF)  $\beta$  or the presence of LPS stimulates endogenous endothelial cell production of IL-1 and IL-6, and induces IL-8 secretion. IL-8 has been shown to regulate transendothelial migration of PMNs through the endothelial barrier. The release of IL-8 is associated with a change in cell adhesion molecule expression from selectins, which are shed into the circulation to integrins which allow firm binding of leucocytes. Moreover, the action of IL-8 is enhanced by the fact that IL-8 is secreted preferentially into the vessel wall and is deposited in the subendothelial matrix by the endothelial cells, giving rise to a transmural chemotactic gradient. When endothelial cells are stimulated by cytokines or thrombin, they express endothelial cell leucocyte adhesion molecule (ELAM) 1 and intercellular adhesion molecule (ICAM) 1. Once activated, endothelial cells also produce an enhancement factor, PAF, which modulates the rapid expression of these adhesion

molecules. The selectin, ELAM-1, is expressed on endothelial cells within hours and binds both PMNs and monocytes. The expression of ICAM-1, a member of the immunoglobulin family, is increased by IFN- $\gamma$ , IL-1 and TNF- $\alpha$ . ICAM-1 facilitates the adhesion of both PMNs and lymphocytes. A second molecule in this series, ICAM-2, which is partly homologous to ICAM-1, mediates the binding of T and B cells to endothelial cells. Vascular cell adhesion molecule (VCAM) 1 is another inducible endothelial surface immunoglobulin which binds both lymphocytes and monocytes. Finally, stimulated endothelial cells can express GMP-140, a surface receptor which preferentially binds platelets. The binding of platelets increases the local availability of PAF and further accelerates the endothelial cell expression of adhesion molecules.

### Leucocytes

Neutrophils play a pivotal role in SIRS/sepsis-associated tissue injury. Numerous stimuli can activate neutrophils via specific receptor systems on their surface (TNF- $\alpha$ , IL-1, IL-8, GM-CSF, IFN- $\gamma$ , leukotriene (LT) B<sub>4</sub>, PAF, ICAM-1 and ELAM-1, C3a and C5a). Neutrophils secrete a wide variety of mediators in response to these stimuli, both proinflammatory (IL-1 $\alpha$  and IL-1 $\beta$ , TNF- $\alpha$ , IL-6, IL-8, IFN- $\gamma$ ) and anti-inflammatory (TGF- $\beta$ ) cytokines and their antagonists (IL-1RA)<sup>64</sup>. PMN activation is manifested by increased  $\beta$ -2 integrin expression and enhanced superoxide radical generation. Upregulation of C11b correlates with serum IL-6 level<sup>65</sup>. Cell surface nicotinamide adenine dinucleotide phosphate oxidase is activated and large quantities of oxygen metabolites are produced. Lipo-oxygenase and phospholipases produce LTB<sub>4</sub> and PAF which result in further accumulation of neutrophils and enhanced endothelial cell cytotoxicity<sup>66,67</sup>. Neutrophils can degrade elastin and collagen I, II, III and VI, fibrinogen, fibronectin and proteoglycans through the release of a neutral serine protease which is a major component of their granular enzymes. The metallo-proteases (MMP-1, MMP-2 and MMP-3) are also secreted and will digest collagen, gelatin and proteoglycans respectively. Neutrophils bear several proteins with antibacterial properties such as BPI, cationic antimicrobial protein and desmins<sup>68</sup>. Neutrophil rolling on the vascular endothelium is regulated in part by the neutrophil cell adhesion molecule L-selectin. In patients with SIRS, neutrophil L-selectin expression is down-regulated in a dose- and time-dependent manner by TNF- $\alpha$ <sup>69</sup>. Neutrophils contribute to a procoagulant environment within the microcirculation by activating platelet aggregation and coagulation cascades, and inhibiting fibrinolysis<sup>70,71</sup>. In a porcine model, a monoclonal antibody against E- and L-selectin (EL246) significantly reduces neutrophil accumulation and tissue injury, but does not attenuate deranged pulmonary and systemic haemodynamics<sup>72</sup>. Similarly, organ-specific injury (acute lung injury) can be attenuated by EL246<sup>73</sup>.

Circulating monocytes move rapidly along the endothelial cell surface (3–5 l/min for the ascending aorta). The first step in monocyte recruitment is the weak attachment of both monocyte and endothelial cell selectin molecules to their opposing corresponding oligosaccharides. The monocyte selectin, L-selectin, is produced constitutively while endothelial selectin, ELAM-1, is induced by IL-1 and TNF- $\alpha$ . After this initial 'weak' attachment process has been initiated the second step,



'firm' attachment, in monocyte recruitment occurs with the adhesion of the monocyte integrin  $\alpha_m\beta_2$  and the endothelial cell receptor ICAM-1. ICAM-1 can be induced by exposure of the endothelial cells to IL-1, TNF- $\alpha$  and IFN- $\gamma$ . Monocyte extravasation requires endothelial cells to express monocyte chemoattractant protein (MCP) 1 at the cell to cell junction to allow diapedesis to occur along a chemoattractant gradient<sup>74</sup>. Once inside the intima, the monocytes amplify the MCP-1 signal by synthesizing and secreting their own MCP-1, a characteristic of tissue macrophages<sup>75-77</sup>. Matrix attachment of the extravasated monocytes occurs through integrin receptors and the tissue macrophages express genes for IL-1, IL-8 and superoxide dismutase<sup>78,79</sup>.

Monocytes/macrophages carry out the fundamental protective functions of ingesting and killing invading micro-organisms. Macrophages play a central role in the immune response by presenting antigens to lymphocytes during the development of specific immunity. Migration of monocytes into different tissues appears to be a random phenomenon during homeostasis, where they undergo transformation into tissue macrophages with morphological and sometimes functional properties that are characteristic for that tissue. The life span of tissue macrophages is believed to be months. The most important functional step in the maturation of bone marrow-derived monocytes to tissue macrophages is the lymphokine-driven conversion of the normal resting macrophage to an activated macrophage<sup>80</sup>. These cells are generally larger, more metabolically active, and able to release soluble substances and oxidative metabolites. Activated macrophages migrate more vigorously in response to chemotactic factors released from invading micro-organisms and enter sites of inflammation more efficiently than unactivated macrophages. Although tissue macrophages are capable of phagocytosis, tissue macrophage-mediated modulation and chemoattraction of non-macrophage inflammatory cells are important activities during sepsis<sup>81</sup>. Once activated, macrophages are the source of numerous cytokines involved in host defence and inflammation. Activation of tissue macrophages results in the generation of reactive oxygen species, and oxidation of arachidonic acid by lipo-oxygenase and cyclo-oxygenase pathways to generate leukotrienes, prostaglandins and thromboxane. Monocytes and tissue macrophages phagocytize particulate material via at least two distinct receptors present in their plasma membrane: the immunoglobulin (Ig) G Fc receptor and the receptor for the alternate complement pathway. Phagocytosis is increased by the presence of C3b. Particles opsonized with C3b react with activated mononuclear phagocyte C3b adherence receptors to promote increased phagocytosis through the Fc and C3b receptors respectively. Phagocytosis is augmented by fibronectin.

## Secondary inflammatory mediators

### *Arachidonic metabolites*

Metabolites of arachidonic acid, particularly those of lipo-oxygenase and cyclo-oxygenase, are significant autocrine and paracrine mediators of the SIRS. Recognized early as potent vasodilators, they play an important role in the low systemic vascular resistance and hypotension that occur in septic shock. LPS, TNF- $\alpha$  and IL-1 all induce the release of prostaglandins from endothelial cells. The major endothelial cell-derived prostaglandin is PGI<sub>2</sub>, a potent

vasodilator. Raised PGI<sub>2</sub> concentrations have been found to correlate with the severity of septic shock. The abluminal release from endothelial cells is small compared with its luminal release. PGI<sub>2</sub> acts on smooth muscle cells via receptor-mediated activation of adenylate cyclase. The ratio of PGI<sub>2</sub>/PGE<sub>2</sub> generation is lower in the microcirculation than in major vessels. With PGI<sub>2</sub> production, endothelial cells also generate a small amount of TXA<sub>2</sub>, a proaggregating vasoconstrictor. A variety of other eicosanoids, such as monohydroxy, dihydroxy and epoxy derivatives of arachidonic acid, which are formed by the cyclo-oxygenase-, lipo-oxygenase- and cytochrome P450-dependent mono-oxygenation pathways, also influence vascular tone. PGI<sub>2</sub> inhibits platelet aggregation and adhesion, reduces thrombus formation and acts synergistically with prostaglandin PGE<sub>2</sub> to increase the effects of serotonin and bradykinin. PGE<sub>2</sub> inhibits both IL-1 production and responsiveness of thymocytes to IL-1. Low concentrations of PGE<sub>2</sub> stimulate TNF- $\alpha$  release, while higher concentrations suppress TNF- $\alpha$  production at a dose-dependent level; PGE<sub>2</sub> inhibits mitogenesis of T and B cells and acts synergistically with prostacyclin to increase the effects of serotonin and bradykinin on vascular permeability. TXA<sub>2</sub> induces platelet aggregation and neutrophil accumulation, increases vascular permeability and enhances permeability of single and double unit membranes. LTB<sub>4</sub> promotes neutrophil chemotaxis and adhesion of neutrophils to endothelium. Neutrophils have specific LTB<sub>4</sub> receptors. LTB<sub>4</sub> is weakly chemotactic for eosinophils and increases vascular permeability, either directly or through interaction with neutrophils and endothelial cells. LTC<sub>4</sub>, LTD<sub>4</sub> and LTE<sub>4</sub> stimulate release of prostacyclin, increase vascular permeability and cause contraction of adjacent endothelial cells and a resulting increase in the diameter of interendothelial cell pores.

### *Nitric oxide*

Synthesis of nitric oxide plays a crucial role in acute and chronic inflammatory processes, in the SIRS and in sepsis<sup>82,83</sup>. Nitric oxide is synthesized from the conversion of L-arginine to citrulline by at least two categories of enzyme: constitutive nitric oxide synthases (cNOS; predominantly membrane bound) and inducible nitric oxide synthases (iNOS; predominantly cytosolic) both of which are calcium- and calmodulin-dependent. Protein phosphorylation by the guanosine 3',5'-cyclic monophosphate (cGMP)-dependent kinases, which are activated by nitric oxide-mediated increases in target cell cGMP, is the basis of many of the effects attributed to nitric oxide. Nitric oxide shares many of the vasoactive properties of prostacyclin in that it can relax smooth muscle and inhibit platelet aggregation. In the appropriate circumstances, nitric oxide can be converted in the endothelial cell to peroxynitrite, a potentially toxic molecule. Peripheral blood monocytes, alveolar macrophages, Kupffer cells and neutrophils, once stimulated, are all capable of synthesizing iNOS and producing nitric oxide<sup>84</sup>. Nitric oxide enhances the vasodilatation, the formation of oedema and the modulation of sensory nerve endings which are hallmarks of the inflammatory response. Inhibition of nitric oxide synthesis will reduce the degree of acute inflammation. Raised concentrations of nitric oxide are likely to come from a combination of sources: activated vascular cells, neutrophils and macrophages<sup>85,86</sup>. These effects can be prevented by treatment with glucocorticoids and NOS inhibitors. In endotoxic shock,



increases in nitric oxide production can be related to the level of hypotension<sup>87</sup>. Bacterial endotoxin induces iNOS and raised nitric oxide synthesis in venous smooth muscle cells, in the cardiomyocytes and in the endocardium, and leads to increased venous pooling and cardiac dysfunction (sepsis-related dilated cardiomyopathy)<sup>86</sup>. Cytokine induction of myocardial iNOS results in increased nitric oxide production and negative inotropy<sup>88</sup>. NOS activity is significantly higher in patients with sepsis<sup>89</sup>.

Studies in animal models on the inhibition of NOS activity have shown a mixed picture of efficacy in the treatment of septic shock<sup>90</sup>. While 30 mg/kg L-nitro-monomethyl arginine will prevent endotoxin shock, 300 mg/kg will accelerate the condition. Early reports suggest that inhibition of nitric oxide is beneficial in septic shock. The benefit of inhibited NOS activity is a sustained increase in systemic blood pressure, while the adverse effects are decreased cardiac output and raised pulmonary vascular resistance<sup>91-94</sup>. Hepatocyte dysfunction appears to be the result of cytokine-induced production of nitric oxide (through iNOS) and cytokine release from macrophages and Kupffer cells<sup>95</sup>. Macrophages synthesize nitric oxide and the high concentrations of nitric oxide are responsible for the cytotoxicity of macrophages to tumour cells and bacteria<sup>94,95</sup>. The cytotoxicity of nitric oxide results from its interaction with iron-containing moieties in enzymes of the respiratory cycle and the DNA synthesis pathways<sup>96</sup>.

#### *Reactive oxygen species*

Endothelial cells contain xanthine dehydrogenase/oxidase and the free radicals produced or transferred from the extracellular space under normal conditions are reduced by endothelial cell superoxide dismutase, catalase and the glutathione redox cycle. Low intracellular levels of oxygen free radicals stimulate cyclo-oxygenation of arachidonic acid. Superoxide radicals can induce vasodilatation in several tissue beds and this is thought to be mediated by the release of prostacyclin from endothelial cells. However, increased intracellular concentrations of oxygen free radicals are able to inactivate nitric oxide and can inhibit the production of prostacyclin in endothelial cells by inhibiting both cyclo-oxygenase (COX-I) and COX-II synthetases. Higher levels will result in the destruction of these enzymes. In contrast, TXA<sub>2</sub> synthetase is resistant to such free radical inhibition and destruction. The net effect of these interactions is vasoconstriction<sup>97</sup>. In addition, oxygen free radicals can lead to the formation of peroxides and, because of the difference in kinetics between the cyclo-oxygenase and peroxidase enzymes, lipid peroxides can accumulate; raised concentrations of these compounds will also destroy both COX-I and COX-II synthetases. This damage can be prevented by antioxidants. In normal physiology, a balance exists between the production of PGI<sub>2</sub> and TXA<sub>2</sub>, and between nitric oxide and oxygen free radicals, which allows for the maintenance of vascular tone<sup>98</sup>.

When tissues are injured by ischaemia or anoxia, their ability to control the metabolism of oxygen is compromised and the species that are generated activate a superoxide-dependent chemoattractant process. This leads to an influx of leucocytes which generates still more reactive oxygen species. Reactive oxygen species are involved in most types of inflammatory tissue injury and are derived predominantly from phagocytic leucocytes<sup>99,100</sup>. Reactive oxygen species produce cellular injury directly by oxidative degradation of essential cellular components

and indirectly by altering the protease/antiprotease balance that exists between cells. Reactive oxygen species can further initiate and amplify the inflammatory process by upregulation of several proinflammatory cytokines (IL-2, IL-6 and TNF- $\alpha$ ) and adhesion molecules (E-selectin, ICAM-1 and VCAM-1)<sup>100,101</sup>. The undisputed contribution of reactive oxygen species to the SIRS has prompted increasing experimental work on means to counteract their local tissue effects<sup>102</sup>. A preliminary report of a randomized trial of N-acetylcysteine in patients with established sepsis-induced ARDS suggests that the antioxidant therapy is useful<sup>103</sup>.

#### *Platelet activating factor*

Endotoxin induces the release of PAF from macrophages, PMNs, platelets and endothelial cells. Systemically, PAF has a negative inotropic effect on the heart and lowers arterial blood pressure. PAF is a potent phospholipid inflammatory mediator that increases cell adhesion, and activates cells by direct effect or through the formation of toxic oxygen species and arachidonic acid metabolites. PAF stimulates the release of TNF- $\alpha$ , leukotrienes, TXA<sub>2</sub>, and promotes leucocyte activation and subsequent free radical formation. There is growing evidence that haematological growth factors and cytokines interact with PAF, leading to amplification of mediator release in septic shock, and that PAF mediates many of the toxicities associated with TNF- $\alpha$  and IL-1<sup>104,105</sup>. Within the microcirculation, PAF encourages platelet aggregation, leading to thrombus formation, and markedly alters microvascular permeability by stimulating calcium efflux in endothelial cells, which results in retraction and loss of reciprocal contact. Specific PAF receptor antagonists provide protection against the fatal complications of endotoxic shock in animal models<sup>106,107</sup>.

#### *Nutrition*

Severe depletion of body protein stores can result from prolonged starvation, or from hormonal or cytokine-mediated effects during critical illness<sup>108-110</sup>. Specialized enteral and parenteral nutrition is now a standard component of care in critically ill patients. This adjunctive therapy corrects and prevents nutrient deficiencies, attenuates the loss of body protein, enhances immune function, and beneficially modifies a body's response to illness, thereby improving clinical outcome in malnourished patients. In patients without severe head injuries, infectious complications are the most common cause of death, and are a frequent cause of morbidity and mortality. Morbidity from sepsis is significantly reduced in critically injured patients when total enteral nutrition (TEN) rather than total parenteral nutrition (TPN) is provided, implying benefits of enteral feeding on host defences<sup>111</sup>. A lack of enteral feeding is thought to lead to a breakdown of the gut mucosal barrier and translocation of bacteria or their products, while the use of TEN or trickle feeding prevents the deterioration in the gut's ability to prevent translocation. Preservation of the barrier function of the gut is essential in controlling sepsis-related morbidity associated with severe injury and stress. Ample experimental evidence suggests that TEN is superior to TPN in preventing many coincidental septic events and modulating the host response to ongoing sepsis that occur in critically ill patients. Immunonutrition is the term being used now to refer to the effects of nutritional

hyperalimentation on the immune system<sup>112</sup>. The components of immunonutrition are glutamine, arginine, omega-3 fatty acids and RNA nucleotides. Glutamine has a significant trophic effect on the gut. It is the preferred fuel source for the intestinal tract in times of stress and is a key component in preventing enterocyte loss, atrophy of the mucosa and loss of barrier function. It is also a known fuel for lymphocytes. Arginine has significant immunostimulant effects, with prevention of thymic atrophy, increased production of natural killer and helper T cells, and increased IL-2 release, which stimulates T cell activation. Long-chain omega-6 fatty acids, which are a common component of TPN and TEN solutions, are more significantly immunosuppressive than omega-3 fatty acids. The presence of omega-3 fatty acids leads to a change in the profile of prostaglandins and leukotrienes produced during stress and sepsis (omega-6 fatty acids increase PGE<sub>2</sub> and LTB<sub>4</sub> synthesis while omega-3 fatty acids increase PGE<sub>3</sub> and LTB<sub>5</sub> synthesis). The absence of RNA nucleotides has been shown to decrease the maturation and phenotypic expression of T lymphocytes, to decrease IL-2 production, to inhibit resistance to infection and to diminish T cell-based immunity. There are mixed opinions that nutritional formulae which contain immunomodulators may convey a small benefit over standard formulae. There are some data to suggest that they may be detrimental in certain subsets of patients<sup>112</sup>.

### Potential therapies

There are three points in the sequence of the pathogenesis of SIRS/sepsis at which therapy can be instituted. First, the nidus of infection can be eradicated with appropriate antimicrobial therapy, surgical drainage, or both. Second, the sepsis-associated cardiovascular metabolic and multiorgan system disturbances can be treated and, third, inhibitors of toxic mediators can be administered<sup>113</sup>.

#### *Anti-lipopolysaccharide*

LPS comprises complex molecules composed of a polysaccharide side chain (O antigen), attached to a glucosamine-based phospholipid (lipid A) by a 'core' polysaccharide. Polyclonal antisera and monoclonal antibodies to lipid A and core regions of mutant *E. coli* have been developed and tested in clinical trials, some of which have had positive results<sup>114</sup> while others have been negative<sup>115,116</sup>. Two prophylactic studies comparing antiserum with preimmune serum in high-risk surgical patients and in neutropenic patients with cancer did not show a reduction in Gram-negative infections by the administration of antiserum<sup>117,118</sup>. The use of the non-toxic derivatives of lipid A, the presumed toxic moiety of the endotoxin molecule, to attenuate the response to LPS, enhances non-specific resistance to infection and induces tolerance to endotoxin<sup>119–122</sup>.

Recently murine and human monoclonal IgM antibodies have been developed using the *E. coli* J5 mutant and have been tested for the treatment of patients with Gram-negative infections in prospective randomized double-blind multicentre trials. In an initial study with E5 murine IgM monoclonal antibody, patients with suspected Gram-negative sepsis were assigned randomly to receive either E5 antibody or placebo<sup>123</sup>. There was no decrease in mortality rates in patients from either group but after

subgroup analysis it did appear there was a decrease in mortality rate in patients without shock at the time of entry into the study. A second study concentrating on patients with Gram-negative sepsis but no shock failed to show any benefit with E5<sup>124</sup>. Treatment with a second antibody, HA-1A (human), did not improve overall population survival at 28 days<sup>125</sup>. A subgroup of patients with Gram-negative bacteraemia showed a significantly improved survival at 28 days and this reduction in mortality rate was more pronounced in patients with shock than in those presenting without shock. Unfortunately there were more patients in the placebo arm than in the HA-1A arm who had considerably more risk factors and they received inadequate antibiotic therapy compared with the experimental patients<sup>126</sup>. Other therapies for blocking the LPS-induced phase are soluble CD14 receptors, anti-LPS-binding protein, anti-CD14 receptor antibodies and bacterial permeability increasing protein.

#### *Anti-tumour necrosis factor $\alpha$*

As TNF- $\alpha$  is a primary mediator of the SIRS response, neutralization is an attractive possibility. Antibodies to TNF- $\alpha$  have decreased the mortality rate in models of lethal bacteraemia and, although there was a decrease in 3-day mortality rate in humans, there was no decrease in the rate between patients treated with placebo and a monoclonal antibody against TNF- $\alpha$  at 28 days<sup>127–129</sup>. There are naturally occurring proteins which represent the extracellular domains of the two TNF- $\alpha$  receptors. They act as natural TNF antagonists and prevent septic shock in *E. coli*-treated baboons and death in mice. A chimaeric molecule in which the soluble TNF- $\alpha$  receptor is linked covalently to the Fc portion of IgG has been designed and produced. With a single administration, the chimaeric molecule can significantly improve septic shock in animal models<sup>130</sup>. In patients with septic shock the chimaeric molecule (TNFR:Fc) did not reduce the mortality rate<sup>131</sup>.

#### *Anti-interleukin 1*

There is a lag time between activation of TNF- $\alpha$  and IL-1 expression. This window of opportunity suggests that targeting IL-1 may be temporally more efficacious than targeting TNF- $\alpha$ . IL-1 receptor antagonists reduced mortality rates in a rabbit model of septic shock even when given after the onset of shock<sup>132</sup>. IL-1RA is a naturally occurring protein which binds to the human IL-1 receptor but has no agonist activity<sup>133,134</sup>. It must be administered in very large molar amounts in order to block IL-1 activity. There is no significant increase in overall survival time in patients treated with IL-1RA<sup>135</sup>. However, there is a significant increase in survival time in patients treated with IL-1RA within the first 2 days if these patients have a 24 per cent or more risk of death within 28 days. There was no benefit if the patient had a risk of death below 24 per cent<sup>136</sup>.

#### *Miscellaneous agents*

Cytokine synthesis can be blocked at a translational level by steroids and at a pretranslational level by pentoxifylline (oxypentifylline) and aminone. Steroids block the translation of TNF- $\alpha$  mRNA in macrophages. They must be administered pre-emptively to have an effect;

administration after the onset of sepsis carries no benefit<sup>30,137–140</sup>. Pentoxifylline and amrinone are phosphodiesterase inhibitors that lead to increased intracellular levels of adenosine 3',5'-cyclic monophosphate which interrupts intracellular signalling<sup>141,142</sup>. Pentoxifylline decreases TNF- $\alpha$  synthesis in a murine model of endotoxic shock, while amrinone has been shown to be a more potent inhibitor of LPS-stimulated TNF- $\alpha$  synthesis<sup>143–145</sup>. In humans, pentoxifylline is able to decrease TNF- $\alpha$  but not IL-6 and IL-8 serum concentrations, and to decrease augmented PMN reactivity during septic shock<sup>146,147</sup>. Biologically-derived glucan infusions improve immune function after trauma and after high-risk abdominal surgery<sup>148,149</sup>. Taurolidine is considered to be a non-specific LPS antagonist. It has no beneficial therapeutic effect on the outcome of patients admitted to the intensive care unit with sepsis syndrome, using progression, resolution of organ failure and 28-day mortality rate as markers of outcome<sup>150</sup>.

Arachidonic acid metabolites help mediate haemodynamic alterations in SIRS/sepsis<sup>36,151–153</sup>. Indomethacin given 1 h before or after a bolus of TNF- $\alpha$  blocks metabolic acidosis, shock and death in rats<sup>154</sup>. Similarly, ibuprofen has also been shown to be effective in animal models<sup>155,156</sup>. However, it has no effect in man<sup>157</sup>. Leukotriene inhibitors have been shown to be beneficial in septic shock<sup>158</sup>. Combined therapies to inhibit several components of the arachidonic acid pathways are more effective than single drug therapy in protecting animals from progressing to MODS<sup>159,160</sup>.

Specific PAF receptor antagonists provide protection against the fatal complications of endotoxic shock in animal models<sup>106,107</sup>. In patients, the use of PAF antagonists has shown no benefit<sup>161</sup>. Activation of the kallikrein-kinin system in sepsis has long been recognized and, experimentally, bradykinin antagonists have been shown to be beneficial in Gram-negative sepsis<sup>162</sup>.

### Newer concepts in systemic inflammatory response syndrome

With failure of the single agent therapies to control SIRS, several authors have suggested that there is a fundamental misconception of the disease process within the human body. SIRS, which is proinflammatory, is only one side of a two-sided response. The other side is an anti-inflammatory response termed the compensatory anti-inflammatory response syndrome (CARS)<sup>163</sup>. The rationale for this theory is that many of the proinflammatory mediators, particularly the interleukins, that participate in SIRS can inhibit immune function by decreasing monocyte/macrophage, B cell and T cell function. In addition, the proinflammatory mediators can inhibit their own synthesis or enhance the synthesis of natural antagonists. Cumulatively, these responses are the body's attempt to re-establish homeostasis, and result in anergy and increased susceptibility to infection. Thus, at any one time there is a battle between the 'Ying' and 'Yang' of the inflammatory system which clinically manifests itself as SIRS, CARS or an intermediate, mixed inflammatory response syndrome (MARS). The spectrum of consequences of these responses has been termed CHAOS (Cardiovascular shock, Homeostasis, Apoptosis, Organ dysfunction and immune Suppression)<sup>163,164</sup>. This new set of concepts suggests that if SIRS is predominant, conventional antimediator or antagonist therapies will be

of value, but if CARS is predominant, then novel immune stimulants or antiantagonist therapies will be required. To enter this algorithm, a fundamental step will be required, that of achieving a diagnosis of the systemic inflammatory state of the patient (SIRS, CARS or MARS).

### Conclusion

The consensus definition of SIRS has allowed translation of several unrelated disease states into a universal process from which it will be possible to categorize, research and treat the body's normal and abnormal inflammatory response to any insult. For surgeons, it forms the basis of the physiology and pathophysiology which their patients experience and which they treat in their practice. In the past decade the cellular and humoral events of inflammation and the systemic inflammatory response that can result have been better defined, yet our overall knowledge in this field is still limited. The newer concepts of MARS, CARS and CHAOS in the natural history of SIRS testify to these limitations and suggest possible new horizons to our understanding (Fig. 5). At present, attempts to intervene in the cascades that participate in SIRS have met with little success. These unifocal attempts are based on our present knowledge and require pre-emptive therapy which cannot at present satisfy the clinical requirement of controlling SIRS after the cascades have been initiated. It is akin 'to closing the barn door after the horse has bolted'<sup>165</sup>. Furthermore, there is no evidence to refute a contention that part of the SIRS




Fig. 5 The inter-relationship of systemic inflammatory response syndrome (SIRS), compensatory anti-inflammatory response syndrome (CARS) and mixed antagonist response syndrome (MARS) and the CHAOS (Cardiovascular shock, Homeostasis, Apoptosis, Organ dysfunction and immune Suppression) theory. Clinical sequelae of the SIRS and the CARS. (Reproduced from reference 163 with permission from the Society of Critical Care and Williams and Wilkins, Baltimore, Maryland, USA)

response is beneficial for the patient in the longer term. Further clinical and basic scientific research will be required to develop the global picture of SIRS, its associated family of syndromes and their natural histories. Modulation of SIRS/CARS remains in its infancy.

## References

- Parker MM, Parrillo JE. Septic shock. Hemodynamics and pathogenesis. *JAMA* 1983; 250: 3324-7.
- Bone RC, Balk RA, Cerra FB *et al.* Definitions for sepsis and organ failure and guidelines for the use of innovative therapies in sepsis. The ACCP/SCCM Consensus Conference Committee. American College of Chest Physicians/Society of Critical Care Medicine. *Chest* 1992; 101: 1644-55.
- Rangel-Frausto MS, Pittet D, Costigan M, Hwang T, Davis CS, Wenzel RP. The natural history of the systemic inflammatory response syndrome (SIRS). A prospective study. *JAMA* 1995; 273: 117-23.
- Salvo I, de Cien W, Musicco M *et al.* The Italian SEPSIS study: preliminary results on the incidence and evolution of SIRS, sepsis, severe sepsis and septic shock. *Intensive Care Med* 1995; 21(Suppl 2): S244-9.
- Pittet D, Rangel-Frausto S, Li N *et al.* Systemic inflammatory response syndrome, sepsis, severe sepsis and septic shock: incidence, morbidities and outcomes in surgical ICU patients. *Intensive Care Med* 1995; 21: 302-9.
- Smail N, Messiah A, Edouard A *et al.* Role of systemic inflammatory response syndrome and infection in the occurrence of early multiple organ dysfunction syndrome following severe trauma. *Intensive Care Med* 1995; 21: 813-16.
- Baker CC, Huynh T. Sepsis in the critically ill patient. *Curr Probl Surg* 1995; 32: 1015-92.
- Collins FM. Cellular antimicrobial immunity. *CRC Critical Review in Microbiology* 1978; 7: 27-91.
- Dantzker D. Oxygen delivery and utilization in sepsis. *Crit Care Clin* 1989; 5: 81-98.
- Dantzker DR, Foresman B, Gutierrez G. Oxygen supply and utilization relationships. A reevaluation. *American Review Respiratory Diseases* 1991; 143: 675-9.
- Tuchschmidt J, Fried J, Astiz M, Rackow E. Elevation of cardiac output and oxygen delivery improves outcomes in septic shock. *Chest* 1992; 102: 216-20.
- Slotman GJ, Burchard KW, Williams JJ, D'Arezzo A, Yellin SA. Interaction of prostaglandins, activated complement, and granulocytes in clinical sepsis and hypotension. *Surgery* 1986; 99: 744-51.
- Bone RC. Toward a theory regarding the pathogenesis of the systemic inflammatory response system: what we do and do not know about cytokine regulation. *Crit Care Med* 1996; 24: 163-72.
- Quezado ZM, Natanson C. Systemic hemodynamic abnormalities and vasopressor therapy in sepsis and septic shock. *Am J Kidney Dis* 1992; 20: 214-22.
- Suffredini AF, Shelhamer JH, Neumann RD, Brenner M, Baltaro RJ, Parrillo JE. Pulmonary and oxygen transport effects of intravenously administered endotoxin in normal humans. *American Review Respiratory Diseases* 1992; 145: 1398-403.
- Hinshaw LB. Sepsis/septic shock: participation of the microcirculation: an abbreviated review. *Crit Care Med* 1996; 24: 1072-8.
- Graves DT, Jiang Y. Chemokines: a family of chemotactic cytokines. *Crit Rev Oral Biol Med* 1995; 6: 109-18.
- Van der Poll T, Buller HR, ten Cate H *et al.* Activation of coagulation after administration of tumor necrosis factor to normal subjects. *N Engl J Med* 1990; 322: 1622-7.
- Jansen PM, Boermeester MA, Fischer E *et al.* Contribution of interleukin-1 to activation of coagulation and fibrinolysis, neutrophil degranulation, and the release of secretory-type phospholipase A<sub>2</sub> in sepsis: studies in nonhuman primates after interleukin-1 $\alpha$  administration and during lethal bacteremia. *Blood* 1995; 86: 1027-34.
- Kalter ES, van Dijk WC, Timmerman A, Verhoef J, Bouma BN. Activation of purified human plasma prekallikrein triggered by cell wall fractions of *Escherichia coli* and *Staphylococcus aureus*. *J Infect Dis* 1983; 148: 682-91.
- Warr TA, Mohan Rao LV, Rapaport SI. Disseminated intravascular coagulation in rabbits induced by administration of endotoxin or tissue factor: effect of anti-tissue factor antibodies and measurement of plasma extrinsic pathway inhibitor activity. *Blood* 1990; 75: 1481-9.
- Suffredini AF, Harpel PC, Parrillo JE. Promotion and subsequent inhibition of plasminogen activation after administration of intravenous endotoxin to normal subjects. *N Engl J Med* 1989; 320: 1165-72.
- Gando S, Kameue T, Nanzaki S, Nakanishi Y. Cytokines, soluble thrombomodulin and disseminated intravascular coagulation in patients with systemic inflammatory response syndrome. *Thromb Res* 1995; 80: 519-26.
- Fourrier F, Chopin C, Goudemand J *et al.* Septic shock, multiple organ failure, and disseminated intravascular coagulation. Compared patterns of antithrombin III, protein C, and protein S deficiencies. *Chest* 1992; 101: 816-23.
- Gando S, Kameue T, Nanzaki S, Nakanishi Y. Disseminated intravascular coagulation is a frequent complication of systemic inflammatory response syndrome. *Thromb Haemost* 1996; 75: 224-8.
- Souba WW. Cytokine control of nutrition and metabolism in critical illness. *Curr Probl Surg* 1994; 31: 577-643.
- Munchie HR, Manogue KR, Spriggs DR *et al.* Detection of circulating tumor necrosis factor after endotoxin administration. *N Engl J Med* 1988; 318: 1481-6.
- Waage A, Brandtzaeg P, Halstensen A, Kierulf P, Espevik T. The complex pattern of cytokines in serum with meningococcal septic shock. Association between interleukin 6, interleukin 1, and fatal outcome. *J Exp Med* 1989; 169: 333-8.
- Pinsky MR, Vincent JL, Deviere J, Alegre M, Kahn RJ, Dupont E. Serum cytokine levels in human septic shock. Relation to multiple-system organ failure and mortality. *Chest* 1993; 103: 565-75.
- Waage A, Halstensen A, Espevik T. Association between tumour necrosis factor in serum and fatal outcome in patients with meningococcal disease. *Lancet* 1987; i: 355-7.
- Tracey KJ, Fong Y, Hesse DG *et al.* Anti-cachectin/TNF monoclonal antibodies prevent septic shock during lethal bacteraemia. *Nature* 1987; 330: 662-4.
- Suffredini AF, Fromm RE, Parker MM *et al.* The cardiovascular response of normal humans to the administration of endotoxin. *N Engl J Med* 1989; 321: 280-7.
- Goldie AS, Fearon KC, Ross JA *et al.* Natural cytokine antagonists and endogenous antiendotoxin core antibodies in sepsis syndrome. The Sepsis Intervention Group. *JAMA* 1995; 274: 172-7.
- Hack CE, Nuijens JH, Felt-Bersma RJF *et al.* Elevated plasma levels of the anaphylatoxins C3a and C4a are associated with a fatal outcome in sepsis. *Am J Med* 1989; 86: 20-6.
- Ognibene FP, Parker MM, Burch-Whitman C *et al.* Neutrophil aggregation activity and septic shock in humans: neutrophil aggregation by a C5a-like material occurs more frequently than complement component depletion and correlates with depression of systemic vascular resistance. *Journal of Critical Care* 1988; 3: 103-11.
- Slotman GJ, Burchard KW, Williams JJ, DiArezzo A, Yellin SA. Interaction of prostaglandins, activated complement, and granulocytes in clinical sepsis and hypotension. *Surgery* 1986; 99: 744-51.
- Pixley RA, Zellis S, Bankes P *et al.* Prognostic value of assessing contact system activation and factor V in systemic inflammatory response syndrome. *Crit Care Med* 1995; 23: 41-51.
- Yentis SM, Soni N, Sheldon J. C-reactive protein as an

- indicator of resolution of sepsis in the intensive care unit. *Intensive Care Med* 1995; 21: 602-5.
- 39 Tobias PS, Mathison JC, Ulevitch RJ. A family of lipopolysaccharide binding proteins involved in responses to Gram-negative sepsis. *J Biol Chem* 1988; 263: 13479-81.
  - 40 Tobias PS, Soldau K, Ulevitch RJ. Isolation of a lipopolysaccharide-binding acute phase reactant from rabbit serum. *J Exp Med* 1986; 164: 777-93.
  - 41 Schumann RR, Leong SR, Flagg GW *et al.* Structure and function of lipopolysaccharide binding protein. *Science* 1990; 249: 1429-31.
  - 42 Opal SM, Fisher CJ, Marra MN, Scott RW, Palardy JE, Brown U. Bactericidal/permeability-increasing protein as a novel therapeutic modality in the treatment of endotoxin shock. *Clin Res* 1991; 39: 351A (Abstract).
  - 43 Parsons PE, Gillespie MK, Moore EE, Moore FA, Worthen GS. Neutrophil response to endotoxin in the adult respiratory distress syndrome: role of CD14. *Am J Respir Cell Mol Biol* 1995; 13: 152-60.
  - 44 Fong Y, Moldawer LL, Shires GT, Lowry SF. The biologic characteristics of cytokines and their implication in surgical injury. *Surg Gynecol Obstet* 1990; 170: 363-78.
  - 45 Tsujimoto M, Vilcek J. Tumor necrosis factor receptors in HeLa cells and their regulation by interferon-gamma. *J Biol Chem* 1986; 261: 5384-8.
  - 46 Aggarwal BB, Eessalu TE, Hass PE. Characterization of receptors for human tumour necrosis factor and their regulation by gamma-interferon. *Nature* 1985; 318: 665-7.
  - 47 Okusawa S, Gelfand JA, Ikejima T, Connolly RJ, Dinarello CA. Interleukin 1 induces a shock-like state in rabbits. Synergism with tumor necrosis factor and the effect of cyclooxygenase inhibition. *J Clin Invest* 1988; 81: 1162-72.
  - 48 Tracey KJ, Beutler B, Lowry SF *et al.* Shock and tissue injury induced by recombinant human cachectin. *Science* 1986; 234: 470-4.
  - 49 Beutler B, Milsark IW, Cerami A. Passive immunization against cachectin/tumor necrosis factor protects mice from lethal effect of endotoxin. *Science* 1985; 229: 869-71.
  - 50 Kawakami M, Cerami A. Studies of endotoxin-induced decrease in lipoprotein lipase activity. *J Exp Med* 1981; 154: 631-9.
  - 51 Dinarello CA. Interleukin-1. *Reviews of Infectious Diseases* 1984; 6: 51-95.
  - 52 Pruitt JH, Copeland EMI, Moldawer LL. Interleukin-1 and interleukin-1 antagonism in sepsis, systemic inflammatory response syndrome, and septic shock. *Shock* 1995; 3: 235-51.
  - 53 March CJ, Mosley B, Larsen A *et al.* Cloning sequence and expression of two distinct human interleukin-1 complementary DNAs. *Nature* 1985; 315: 641-7.
  - 54 Giri JG, Wells J, Dower SK *et al.* Elevated levels of shed type II IL-1 receptor in sepsis. Potential role for type II receptor in regulation of IL-1 responses. *J Immunol* 1994; 153: 5802-9.
  - 55 Fong Y, Moldawer LL, Marano M *et al.* Endotoxemia elicits increased circulating beta 2-IFN/IL-6 in man. *J Immunol* 1989; 142: 2321-4.
  - 56 Nawroth PP, Handley DA, Esmon CT, Stern DM. Interleukin 1 induces endothelial cell procoagulant while suppressing cell-surface anticoagulant activity. *Proc Natl Acad Sci U S A* 1986; 83: 2460-4.
  - 57 May LT, Ghayeb J, Santhanam U *et al.* Synthesis and secretion of multiple forms of 'beta2 interferon/B cell differentiation factor BSF-2/hepatocyte stimulating factor' by human fibroblasts and monocytes. *J Biol Chem* 1989; 263: 7760-6.
  - 58 Fong Y, Tracey KJ, Moldawer LL *et al.* Antibodies to cachectin/tumor necrosis factor reduce interleukin-1 $\beta$  and interleukin-6 appearance during lethal bacteremia. *J Exp Med* 1989; 170: 1627-33.
  - 59 Starnes HF Jr, Pearce MK, Tewari A, Yim JH, Zon JC, Abrams JS. Anti-IL-6 monoclonal antibodies protect against lethal *Escherichia coli* infection and lethal tumor necrosis factor- $\alpha$  challenge in mice. *J Immunol* 1990; 145: 4185-91.
  - 60 Harada A, Sekido N, Akahoshi T, Wada T, Mukaida N, Matsushima K. Essential involvement of interleukin-8 (IL-8) in acute inflammation. *J Leukoc Biol* 1994; 56: 559-64.
  - 61 Davies MG, Hagen P-O. The vascular endothelium. A new horizon. *Ann Surg* 1993; 218: 593-609.
  - 62 Eisenberg PR. Endothelial cell mediators of thrombosis and fibrinolysis: Review in depth. *Coronary Artery Diseases* 1991; 2: 129-66.
  - 63 Ley K. Leukocyte adhesion to vascular endothelium. *J Reconstr Microsurg* 1992; 8: 495-503.
  - 64 Fujishima S, Aikawa N. Neutrophil-mediated tissue injury and its modulation. *Intensive Care Med* 1995; 21: 277-85.
  - 65 Rosenbloom AJ, Pinsky MR, Bryant JL, Shin A, Tran T, Whiteside T. Leukocyte activation in the peripheral blood of patients with cirrhosis of the liver and SIRS. Correlation with serum interleukin-6 levels and organ dysfunction. *JAMA* 1995; 274: 58-65.
  - 66 Cluzel M, Undem BJ, Chilton FH. Release of platelet-activating factor and the metabolism of leukotriene B<sub>4</sub> by the human neutrophil when studied in a cell superfusion model. *J Immunol* 1989; 143: 3659-65.
  - 67 Bussolino F, Aglietta M, Sanavio F, Stacchini A, Lauri D, Camussi G. Alkyl-ether phosphoglycerides influence calcium fluxes into human endothelial cells. *J Immunol* 1985; 135: 2748-55.
  - 68 Pereira HA. CAP37, a neutrophil-derived multifunctional inflammatory mediator. *J Leukoc Biol* 1995; 57: 805-12.
  - 69 Ahmed NA, Yee J, Giannias B, Kapadia B, Christou NV. Expression of human neutrophil L-selectin during the systemic inflammatory response syndrome is partly mediated by tumor necrosis factor  $\alpha$ . *Arch Surg* 1996; 131: 31-5.
  - 70 Gawaz M, Fateh-Moghadam S, Pilz G, Gurland HJ, Werdan K. Platelet activation and interaction with leukocytes in patients with sepsis or multiple organ failure. *Eur J Clin Invest* 1995; 25: 843-51.
  - 71 Haj MA, Neilly IJ, Robbie LA, Adey GD, Bennett B. Influence of white blood cells on the fibrinolytic responses to sepsis: studies of septic patients with and without severe leucopenia. *Br J Haematol* 1995; 90: 541-7.
  - 72 Ridings PC, Windsor AC, Jutila MA *et al.* A dual-binding antibody to E and L-selectin attenuates sepsis-induced lung injury. *Am J Respir Crit Care Med* 1995; 152: 247-53.
  - 73 Ridings PC, Bloomfield GL, Holloway S *et al.* Sepsis-induced acute lung injury is attenuated by selectin blockade following the onset of sepsis. *Arch Surg* 1995; 130: 1199-208.
  - 74 Cybulsky MI, Gimbrone MA Jr. Endothelial expression of a mononuclear leukocyte adhesion molecule during atherogenesis. *Science* 1991; 251: 788-91.
  - 75 Cushing SD, Fogelman AM. Monocytes may amplify their recruitment into inflammatory lesions by inducing monocyte chemotactic protein. *Arteriosclerosis and Thrombosis* 1992; 12: 78-82.
  - 76 Yla-Hetuala S, Lipton BA, Rosenfeld ME *et al.* Expression of monocyte chemoattractant protein 1 in macrophage-rich areas of human and rabbit atherosclerotic lesions. *Proc Natl Acad Sci U S A* 1991; 88: 5252-6.
  - 77 Nelken NA, Coughlin SR, Gordon D, Wilcox JN. Monocyte chemoattractant protein-1 in human atheromatous plaques. *J Clin Invest* 1991; 88: 1121-7.
  - 78 Wang JM, Sica A, Peri G *et al.* Expression of monocyte chemotactic protein and interleukin-8 by cytokine-activated vascular smooth muscle cells. *Arteriosclerosis and Thrombosis* 1991; 11: 1166-74.
  - 79 Sporn SA, Eierman DF, Johnson CE *et al.* Monocyte adherence results in selective induction of novel genes sharing homology with mediators of inflammation and tissue repair. *J Immunol* 1990; 144: 4434-41.
  - 80 Nelson NJ, Belosevic M, Green SJ, Turpin J, Nacy CA. Interleukin-2 and the regulation of activated macrophage cytotoxic activities. *Adv Exp Med Biol* 1992; 319: 77-88.
  - 81 Unanue ER, Allen PM. The basis for the immunoregulatory role of macrophages and other accessory cells. *Science* 1987; 236: 551-7.

- 82 Nussler AK, Billiar TR. Inflammation, immunoregulation, and inducible nitric oxide synthase. *J Leukoc Biol* 1993; 54: 171-8.
- 83 Wolfe TA, Dasta JF. Use of nitric oxide synthase inhibitors as a novel treatment for septic shock. *Ann Pharmacother* 1995; 29: 36-46.
- 84 Nathan CT, Hibbs JB Jr. Role of nitric oxide synthesis in macrophage antimicrobial activity. *Curr Opin Immunol* 1991; 3: 65-70.
- 85 Thiemeermann C. The role of the L-arginine-nitric oxide pathway in circulatory shock. *Adv Pharmacol* 1994; 28: 45-79.
- 86 Morris SM Jr, Billiar TR. New insights into the regulation of inducible nitric oxide synthesis. *Am J Physiol* 1994; 266: E829-39.
- 87 Brady AJ, Poole-Wilson PA. Circulatory failure in septic shock. Nitric oxide: too much of a good thing? *Brit Heart J* 1993; 70: 103-5.
- 88 Finkel MS, Oddis CV, Jacob TD, Watkins SC, Hattler BG, Simmons RL. Negative inotropic effects of cytokines on the heart mediated by nitric oxide. *Science* 1992; 257: 387-9.
- 89 Goode HF, Howdle PD, Walker BE, Webster NR. Nitric oxide synthase activity is increased in patients with sepsis syndrome. *Clin Sci* 1995; 88: 131-3.
- 90 Nava E, Palmer RMJ, Moncada S. Inhibition of nitric oxide synthesis in septic shock: how much is beneficial? *Lancet* 1991; 338: 1555-7.
- 91 Kilbourn RG, Gross SS, Jubran A *et al.* N-methyl L-arginine inhibits tumor necrosis factor-induced hypotension: implications for the involvement of nitric oxide. *Proc Natl Acad Sci U S A* 1990; 87: 3629-32.
- 92 Pastor C, Teisseire B, Vicaute E, Payen D. Effects of L-arginine and L-nitro-arginine treatment on blood pressure and cardiac output in a rabbit endotoxin shock model. *Crit Care Med* 1994; 22: 465-9.
- 93 Petros A, Bennet D, Vallance P. Effect of nitric oxide synthase inhibitors on hypotension in patients with septic shock. *Lancet* 1991; 338: 1557-8.
- 94 Geroulanos S, Schilling J, Cakmakci M, Jung HH, Largiader F. Inhibition of NO synthesis in septic shock. *Lancet* 1992; 339: 435(Letter).
- 95 Hibbs JB Jr, Taintor RR, Vavrin Z, Rachlin EM. Nitric oxide: a cytotoxic activated macrophage effector molecule. *Biochem Biophys Res Commun* 1989; 157: 87-94.
- 96 Moncada S, Palmer RMJ, Higgs EA. Nitric oxide: physiology, pathophysiology, and pharmacology. *Pharmacol Rev* 1991; 43: 109-42.
- 97 Katusic ZS, Vanhoutte PM. Superoxide anion is an endothelium-derived contracting factor. *Am J Physiol* 1989; 257: H33-7.
- 98 Rubanyi GM. Vascular effects of oxygen-derived free radicals. *Free Radic Biol Med* 1988; 4: 107-20.
- 99 Haglund U, Gerdin B. Oxygen-free radicals (OFR) and circulatory shock. *Circ Shock* 1991; 34: 405-11.
- 100 Conner EM, Grisham MB. Inflammation, free radicals, and antioxidants. *Nutrition* 1996; 12: 274-7.
- 101 Lucchesi BR. Complement, neutrophils and free radicals: mediators of reperfusion injury. *Arzneimittelforschung* 1994; 44: 420-32.
- 102 Swank DW, Moore SB. Roles of the neutrophil and mediators in adult respiratory distress syndrome. *Mayo Clin Proc* 1989; 64: 1118-32.
- 103 Bernard GR. N-acetylcysteine in experimental and clinical acute lung injury. *Am J Med* 1991; 91: 54-9S.
- 104 Sun X-M, Hsueh W. Bowel necrosis induced by tumor necrosis factor in rats is mediated by platelet-activating factor. *J Clin Invest* 1988; 81: 1328-31.
- 105 Touvay C, Vilain B, Carre C, Mencia-Huerta JM, Braquet P. Role of platelet-activating factor (PAF) in the bronchopulmonary alterations and beta-adrenoceptor function induced by endotoxin. *Biochem Biophys Res Commun* 1988; 152: 527-33.
- 106 Moore JM, Earnest MA, DiSimone AG, Abumrad NN, Fletcher JR. A PAF antagonist BN 52021, attenuates thromboxane release and improves survival in lethal canine endotoxemia. *Circ Shock* 1991; 35: 53-9.
- 107 Yue TL, Farhat M, Rabinovici R, Perera PY, Vogel SN, Feuerstein G. Protective effect of BN50739, a new platelet-activating factor antagonist, in endotoxin-treated rabbits. *J Pharmacol Exp Ther* 1990; 254: 976-81.
- 108 McClave SA, Lowen CC, Snider HL. Immunonutrition and enteral hyperalimentation of critically ill patients. *Dig Dis Sci* 1992; 37: 1153-61.
- 109 Ziegler TR, Gatzert C, Wilmore DW. Strategies for attenuating protein-catabolic responses in the critically ill. *Annu Rev Med* 1994; 45: 459-80.
- 110 McMahon MM, Farnell MB, Murray MJ. Nutritional support of critically ill patients. *Mayo Clin Proc* 1993; 68: 911-20.
- 111 Bower RH, Cerra FB, Bershadsky B *et al.* Early enteral administration of a formula (impact) supplemented with arginine, nucleotides, and fish oil in intensive care unit patients: results of a multicenter, prospective, randomized, clinical trial. *Crit Care Med* 1995; 23: 436-49.
- 112 Alexander JW. Specific nutrients and the immune response. *Nutrition* 1995; 11(Suppl 2): 229-32.
- 113 Parrillo JE. Management of septic shock: present and future. *Ann Intern Med* 1991; 115: 491-3.
- 114 Ziegler EJ, McCutchan JA, Fierer J *et al.* Treatment of Gram-negative bacteremia and shock with human antiserum to a mutant *Escherichia coli*. *N Engl J Med* 1982; 307: 1225-30.
- 115 J5 Study Group. Treatment of severe infectious purpura in children with human plasma from donors immunized with *Escherichia coli* J5: a prospective double-blind study. *J Infect Dis* 1992; 165: 695-701.
- 116 Calandra T, Glauser MP, Schellekens J, Verhoef J. Treatment of Gram-negative septic shock with human IgG antibody to *Escherichia coli* J5: a prospective, double-blind randomized trial. *J Infect Dis* 1988; 158: 312-19.
- 117 Baumgartner JD, Glauser MP, McCutchan JA *et al.* Prevention of Gram-negative shock and death in surgical patients by an antibody to endotoxin core glycolipid. *Lancet* 1985; ii: 59-63.
- 118 McCutchan JA, Wolf JL, Ziegler EJ. Ineffectiveness of single-dose human antiserum to core glycolipid (*E. coli* J5) for prophylaxis of bacteremic, Gram-negative infections in patients with prolonged neutropenia. *Schweiz Med Wochenschr* 1983; 113(Suppl 14): 40-5.
- 119 Golenbock DT, Leggett JE, Rasmussen P, Craig WA, Raetz CR, Protor RA. Lipid X protects mice against fatal *Escherichia coli* infection. *Infect Immun* 1988; 56: 779-84.
- 120 Chase JJ, Kubey W, Dulek MH *et al.* Effect of monophosphoryl lipid A on host resistance to bacterial infection. *Infect Immun* 1986; 53: 711-12.
- 121 Astiz ME, Rackow EC, Kim YB, Weil MH. Monophosphoryl lipid A induces tolerance to lethal hemodynamic effects of endotoxemia. *Circ Shock* 1991; 33: 92-7.
- 122 Nakamura H, Ishizaka A, Urano T *et al.* Effects of pretreatment with SDZ MRL 953, a novel immunostimulatory lipid A analog, on endotoxin-induced acute lung injury in guinea pigs. *Clinical and Diagnostic Laboratory Immunology* 1995; 2: 672-7.
- 123 Greenman RL, Schein RMH, Martin MA *et al.* A controlled clinical trial of E5 murine monoclonal IgM antibody to endotoxin in the treatment of Gram-negative sepsis. The XOMA Sepsis Study Group. *JAMA* 1991; 266: 1097-102.
- 124 Bone RC, Balk RA, Fein AM *et al.* A second large controlled clinical study of E5, a monoclonal antibody to endotoxin: results of a prospective, multicenter, randomized, controlled trial. The E5 Sepsis Study Group. *Crit Care Med* 1995; 23: 994-1006.
- 125 Ziegler EJ, Fisher CJ Jr, Sprung CL *et al.* Treatment of Gram-negative bacteremia and septic shock with HA-1A



- human monoclonal antibody against endotoxin. A randomized, double-blind placebo-controlled trial. The HA-1A Sepsis Study Group. *N Engl J Med* 1991; 324: 429–36.
- 126 Warren HS, Danner RL, Munford RS. Anti-endotoxin monoclonal antibodies. *N Engl J Med* 1992; 326: 1153–7.
  - 127 Abraham E, Wunderink R, Silverman H *et al.* Efficacy and safety of monoclonal antibody to human tumor necrosis factor alpha in patients with sepsis syndrome. A randomized, controlled, double-blind, multicenter clinical trial. TNF- $\alpha$  MAb Sepsis Study Group. *JAMA* 1995; 273: 934–41.
  - 128 Van Zee KJ, Kohno T, Fischer E, Rock CS, Moldawer LL, Lowry SF. Tumor necrosis factor soluble receptors circulate during experimental and clinical inflammation and can protect against excessive tumor necrosis factor- $\alpha$  *in vitro* and *in vivo*. *Proc Natl Acad Sci U S A* 1992; 89: 4845–9.
  - 129 Reinhart K, Weigand-Lohnert C, Grimminger F *et al.* Assessment of the safety and efficacy of the monoclonal anti-tumor necrosis factor antibody-fragment, MAK 195F, in patients with sepsis and septic shock: a multicenter, randomized, placebo-controlled, dose-ranging study. *Crit Care Med* 1996; 24: 733–42.
  - 130 Mohler KM, Torrance DS, Smith CA *et al.* Soluble tumor necrosis factor (TNF) receptors are effective therapeutic agents in lethal endotoxemia and function simultaneously as both TNF carriers and TNF antagonists. *J Immunol* 1993; 151: 1548–61.
  - 131 Fisher CJ, Agosti JM, Opal SM *et al.* Treatment of septic shock with the tumor necrosis factor receptor:Fc fusion protein. The Soluble TNF Receptor Sepsis Study Group. *N Engl J Med* 1996; 334: 1697–702.
  - 132 Aiura K, Gelfand JA, Wakabayashi G. Interleukin-1 receptor antagonist blocks staphylococcal induced septic shock in rabbits. *Cytokine* 1991; 3: 498.
  - 133 Eisenberg SP, Evans RJ, Arend WP. Primary structure and functional expression from complementary DNA of a human interleukin-1 receptor antagonist. *Nature* 1990; 343: 341–6.
  - 134 Hannum CH, Wilcox CJ, Arend WP. Interleukin-1 receptor antagonist activity of a human interleukin-1 inhibitor. *Nature* 1990; 343: 336–40.
  - 135 Fisher CJ Jr, Dhainaut JF, Opal SM *et al.* Recombinant human interleukin-1 receptor antagonist in the treatment of patients with sepsis syndrome. Results from a randomized, double-blind, placebo-controlled trial. Phase III rhIL-1ra Sepsis Syndrome Study Group. *JAMA* 1994; 271: 1836–43.
  - 136 Knaus WA, Harrell FE Jr, LaBrecque JF *et al.* Use of predicted risk of mortality to evaluate the efficacy of anticytokine therapy in sepsis. The rhIL-1ra Phase III Sepsis Syndrome Study Group. *Crit Care Med* 1996; 24: 46–56.
  - 137 Bone RC, Fisher CJ Jr, Clemmer TP, Slotman GJ, Metz CA, Balk RA. A controlled clinical trial of high-dose methylprednisolone in the treatment of severe sepsis and septic shock. *N Engl J Med* 1987; 317: 653–8.
  - 138 The Veterans Administration Systemic Sepsis Cooperative Study Group. Effect of high-dose glucocorticoid therapy on mortality in patients with clinical signs of systemic sepsis. *N Engl J Med* 1987; 317: 659–65.
  - 139 Sprung CL, Caralis PV, Marcial EH *et al.* The effects of high-dose corticosteroids in patients with septic shock. A prospective, controlled study. *N Engl J Med* 1984; 311: 1137–43.
  - 140 Hinshaw LB, Beller BK, Chang ACK *et al.* Effects of prior administration of steroids upon recovery from lethal sepsis. *Surg Gynecol Obstet* 1986; 163: 335–44.
  - 141 Taffet SM, Singhel KJ, Overholtzer JF, Shurtleff SA. Regulation of tumor necrosis factor expression in a macrophage-like cell line by lipopolysaccharide and cyclic AMP. *Cell Immunol* 1989; 120: 291–300.
  - 142 Strieter RM, Remick DG, Ward PA *et al.* Cellular and molecular regulation of tumor necrosis factor- $\alpha$  production by pentoxifylline. *Biochem Biophys Res Commun* 1988; 155: 1230–6.
  - 143 Schade UF. Pentoxifylline increases survival in murine endotoxin shock and decreases formation of tumor necrosis factor. *Circ Shock* 1990; 31: 171–81.
  - 144 Giroir BP, Beutler B. Effect of amrinone on tumor necrosis factor production in endotoxin shock. *Circ Shock* 1992; 36: 200–7.
  - 145 Graninger W, Wenisch C. Pentoxifylline in severe inflammatory response syndrome. *J Cardiovasc Pharmacol* 1995; 25(Suppl 2): S134–8.
  - 146 Zeni F, Pain P, Vindimian M *et al.* Effects of pentoxifylline on circulating cytokine concentrations and hemodynamics in patients with septic shock: results from a double-blind, randomized, placebo-controlled study. *Crit Care Med* 1996; 24: 207–14.
  - 147 Oismuller C, Mayer N, Micksche M, Steltzer H, Hammerle AF. *In vitro* modulation of human neutrophil function by pentoxifylline in patients with septic syndrome. *Shock* 1995; 4: 161–5.
  - 148 de Felipe J, de Rocha e Silva M Jr, Marcel FMB, de Macedo Soares A, Mendes NF. Infection prevention in patients with severe multiple trauma with the immunomodulator beta 1–3 polyglucose (glucan). *Surg Gynecol Obstet* 1993; 177: 383–8.
  - 149 Babineau TJ, Marcello P, Swails W, Kenler A, Bistrian B, Forse RA. Randomized Phase I/II trial of macrophage-specific immunomodulator (PGG-glucan) in high-risk surgical patients. *Ann Surg* 1994; 220: 601–9.
  - 150 Willatts SM, Radford S, Leitermann M. Effect of the antiendotoxic agent, tauridine, in the treatment of sepsis syndrome: a placebo-controlled double-blind trial. *Crit Care Med* 1995; 23: 1033–9.
  - 151 Lefer AM. Significance of lipid mediators in shock states. *Circ Shock* 1989; 27: 3–12.
  - 152 Harlan JM, Winn RK. The role of phospholipase products in the pathogenesis of vascular injury in sepsis. In: Root RK, Sande MA, eds. *Septic Shock*. Contemporary Issues in Infectious Diseases. Vol. 4. New York: Churchill Livingstone, 1985: 83–104.
  - 153 Reines HD, Halushka PV, Cook JA, Wise WC, Rambo W. Plasma thromboxane concentrations are raised in patients dying with septic shock. *Lancet* 1982; ii: 174–5.
  - 154 Kettelhut IC, Fiers W, Goldberg AL. The toxic effects of tumor necrosis factor *in vivo* and their prevention by cyclooxygenase inhibitors. *Proc Natl Acad Sci U S A* 1987; 84: 4273–7.
  - 155 Fink MP, Morrissey PE, Stein KL, Clement RE, Fiallo V, Gardiner WM. Systemic and regional hemodynamic effects of cyclo-oxygenase and thromboxane synthetase inhibition in normal and hyperdynamic endotoxemic rabbits. *Circ Shock* 1988; 26: 41–57.
  - 156 Jacobs ER, Soulsby ME, Bone RC. Ibuprofen in canine endotoxin shock. *J Clin Invest* 1982; 70: 536–41.
  - 157 Haupt M, Jastremski M, Clemmer T, Metz C, Brown B. Effects of ibuprofen in patients with severe sepsis syndrome. *Chest* 1989; 96: 291S.
  - 158 Smith EF III, Kinter LB, Jugus M, Wasserman MA, Eckardt RO, Newton JF. Beneficial effects of the peptidoleukotriene receptor antagonist, SK and F 104353, on the responses to experimental endotoxemia in the conscious rat. *Circ Shock* 1988; 25: 21–31.
  - 159 Byrne K, Sielaff TD, Michna B *et al.* Increased survival time after delayed histamine and prostaglandin blockade in a porcine model of severe sepsis-induced lung injury. *Crit Care Med* 1990; 18: 303–8.
  - 160 Turner CR, Quinlan MF, Schwartz LW, Wheeldon EB. Therapeutic intervention in a rat model of ARDS. I. Dual inhibition of arachidonic acid metabolism. *Circ Shock* 1990; 32: 231–42.
  - 161 Dhainaut JF, Tenailon A, Le Tulzo Y *et al.* Platelet-activating factor receptor antagonist BS 52021 in the treatment of severe sepsis: a randomized double-blind,



- placebo-controlled multicenter clinical trial. BN 52021 Sepsis Study Group. *Crit Care Med* 1994; 22: 1720–8.
- 162 Ridings PC, Blocher CR, Fisher BJ, Fowler AA III, Sugerman HJ. Beneficial effects of a bradykinin antagonist in a model of Gram-negative sepsis. *J Trauma* 1995; 39: 81–8.
- 163 Bone RC. Sir Isaac Newton, sepsis, SIRS, and CARS. *Crit Care Med* 1996; 24: 1125–8.
- 164 Bone RC. Immunologic dissonance: a continuing evolution in our understanding of the systemic inflammatory response syndrome (SIRS) and the multiple organ dysfunction syndrome (MODS). *Ann Intern Med* 1996; 125: 680–7.
- 165 Bone RC. Why sepsis trials fail. *JAMA* 1996; 276: 565–6.