
detection of bacterium
hydrogen peroxide catabolic process
cellular oxidant detoxification
protein ubiquitination
fruit ripening
purine nucleoside transmembrane transport
plant−type hypersensitive response
pathogen−associated molecular pattern receptor signaling pathway
formation of cytoplasmic translation initiation complex
purine nucleobase transmembrane transport
pectin catabolic process
regulation of hydrogen peroxide metabolic process
response to iron ion starvation
translation
regulation of translation
cytokinin−activated signaling pathway
systemic acquired resistance, salicylic acid mediated signaling pathway
response to oxidative stress
coumarin biosynthetic process
response to ethylene
defense response to fungus
response to hypoxia
cellular response to abscisic acid stimulus
BP
lactoperoxidase activity
heme binding
purine nucleoside transmembrane transporter activity
hyoscyamine (6S)−dioxygenase activity
structural constituent of ribosome
protein phosphatase inhibitor activity
purine nucleobase transmembrane transporter activity
pectate lyase activity
metal ion binding
O−methyltransferase activity
pectinesterase activity
protein heterodimerization activity
oxidoreductase activity, acting on a sulfur group of donors, disulfide as acceptor
iron ion binding
structural constituent of chromatin
MF
extracellular region
plant−type cell wall
SAGA complex
ribosome
SCF ubiquitin ligase complex
ribonucleoprotein complex
membrane
cytosolic large ribosomal subunit
nucleosome
CC
Exp. 2 tol.
Exp. 3 tol.
Exp. 2 tol. + DE
Exp. 3 tol. + DE
+ DE penn only
+ DE tub only
Background (cons. + DE)
−log10(p)
5
4
3
2
HOGs in set
50
100
150
200
250