Escalation Failure: A Safety-Critical Blind Spot in Medical AI Evaluation

Anonymous Author(s)
Affiliation
Address
email

Abstract

Healthcare is frequently cited as a domain of concern regarding AI risk, yet safety evaluation of medical language models has focused almost entirely on diagnostic accuracy rather than the failure mode with the most immediate clinical consequence: failing to recommend urgent escalation when a patient's presentation demands it. Suboptimal patient care and clinical mismanagement can cause significant harm to individuals as well as to health systems. We introduce *escalation failure* as an explicit, safety-specific evaluation metric, decoupled from standard accuracy, and curate a safety-critical subset of MedQA – cardiovascular and pulmonary USMLE-style vignettes whose ground-truth answer is itself an ICU- or emergency-level intervention – to measure it directly. Applying this methodology across eight language models, we find that escalation failure is not marginal but near-total: three of eight models fail to recognize the need for escalation on effectively 100% of cases. This establishes escalation failure as a severe, live safety gap in models already being trialled in clinical workflows, not a theoretical edge case. This is our central contribution: a methodology and evidence base for treating care escalation as a first-class, explicitly measured safety metric rather than an afterthought to accuracy. As a secondary test of this methodology's usefulness, we also evaluate a lightweight mitigation and find it substantially reduces failure for most models. Our evaluation is limited in scale and relies on rule-based keyword labeling rather than clinician annotation; we discuss these limitations directly and lay out concrete future directions, including scaling the methodology, improving label quality, and extending escalation-failure measurement into post-deployment monitoring. Regardless, the core fact our methodology establishes stands: escalation failure is a severe, currently under-evaluated safety gap in medical language models utilised in clinical care, and it must be measured, reported, and monitored explicitly rather than assumed away.

1 Introduction

Large language models are being integrated into many key workflows in healthcare, such as clinical decision support, note generation, patient communication, medical research, and administrative workflows. Based on current literature reviews, the main research and real-world use cases of LLMs in health are concentrated in medical question-answering and patient information generation [1]. Further literature illustrates existing research gaps in AI-based decision-making systems and their patient-relevant outcomes [2]. Yet this integration is occurring even as post-deployment audits of current AI systems in the healthcare ecosystem have revealed safety gaps: an insurer's care-denial algorithm was found to have its determinations overturned in over 90% of appeals [3], and external validation of a widely deployed sepsis early-warning model found it missed 67% of true sepsis cases despite alerting on 18% of all patients [4]. Meanwhile, AI adoption is accelerating on the ground:

Submitted to 40th Conference on Neural Information Processing Systems (NeurIPS 2026). Do not distribute.

a 2025 UK survey found that a quarter of general practitioners already use generative AI tools in clinical practice [5]. At such scale, evaluating a model's behavior as an individual system is not enough: a per-query failure rate that looks small in a benchmark table compounds, across a population of patients and repeated queries, into a large absolute number of missed escalations, shifting the focus of failure-mode evaluation from technical model validation toward real-world, population-level hazard mitigation.

Existing reporting frameworks for medical AI – CLAIM [6], STARD-AI [7], CONSORT-AI [8], and MI-CLAIM [9] – specify what a study *should report* about a clinical AI system, but none defines a metric for the single failure mode most directly tied to patient harm in an autonomous or semi-autonomous triage setting: a model *failing to recognize that a case requires urgent escalation*. Standard multiple-choice medical QA accuracy conflates this with general clinical knowledge and does not distinguish a model that is simply wrong from one that is confidently, unsafely reassuring. We refer to the latter as *escalation failure*, and to the broader phenomenon as *unsafe reassurance*: a model selecting a non-escalation option in a scenario where escalation is clinically required.

We test this empirically across eight language models – seven general-purpose frontier LLMs and one open-weight medical-specialist model, MedGemma [10, 11] – on a curated safety-critical subset of MedQA [12], using escalation failure as a metric explicitly decoupled from raw accuracy. Our findings have three parts. First, severity: under a naive, unconstrained prompt, escalation failure is near-total for most models tested, with 98–100% for three of eight models and 52–74% for the remaining five – not a theoretical edge case, but the default behavior of models already being piloted in clinical workflows, reflecting significant, unsafe clinical decision-making. Second, mitigation: default results substantially improve without retraining, with a stricter, safety-instructed, output-constrained prompt reducing escalation failure by 24 to 94 percentage points for seven of eight models. Third, and the paper's central caution: this mitigation is not universal. One model in our study, MedGemma, shows almost no improvement and remains the worst performer on both accuracy and escalation failure under matched conditions. Together, these three findings are the paper's argument: escalation failure is severe enough under default prompting to demand explicit measurement, and while prompting can meaningfully improve model performance, that improvement ranges from virtually none to a near-total reduction in failure across the eight models we test – a variance too extreme to estimate or project for any given model without evaluating it directly.

2 Related Work

Existing medical question-answering benchmarks vary widely in structure. PubMedQA [13] poses questions over research-article abstracts rather than patient cases; MedicationQA [14] is an open-domain, non-multiple-choice format built around consumer medication queries; and MedMCQA [15], along with the clinical-knowledge and professional-medicine subsets of MMLU [16] and MMLU-Pro [17], are predominantly declarative knowledge or exam-recall questions. None of these are built around the patient-vignette-plus-management-decision structure our escalation-failure framing depends on. MedQA [12], sourced from USMLE and Chinese medical board exams, is the closest fit since its vignettes routinely include a concrete next-step-in-management decision among the answer options, and we build our curation pipeline on it (Section 3).

MedHELM [18] introduces a clinician-validated taxonomy of 121 medical tasks across 5 categories and evaluates LLMs with benchmark-appropriate metrics, including LLM-as-Judge scoring for open-ended tasks; it is, to our knowledge, the most comprehensive current effort to move medical LLM evaluation beyond static licensing-exam accuracy. Our work is narrower in scope but targets a specific safety-relevant failure mode on escalation recognition that a general capability taxonomy does not isolate as a standalone metric, and that we argue merits explicit reporting alongside accuracy in any clinical-decision-support evaluation.

3 Methodology

Dataset curation. From our survey of existing medical datasets and benchmarks, we decided to build upon MedQA [12] since its USMLE- and board-exam-style patient vignettes with next-step management decisions among the answer options being closest to our target use case. We thereby start from its over 12,000 English multiple-choice questions drawn from USMLE and Chinese

medical board exams, where only English questions in Multiple Choice format were selected for our evaluation use case to maintain language consistency. We then derive our evaluation set through three sequential filtering stages described below and summarized in Table 1.

Stage 1 – safety-keyword pre-filter. A fixed list of 22 emergency/high-acuity intervention terms, spanning airway, cardiac, thoracic, neurological, obstetric, and vascular domains (full list in Appendix Table 5), is matched with word-boundary regex against the text of every answer option, across all 12,723 questions. A question is retained if at least one option matches at least one term, and we record exactly which option letters matched for later use. This yields 185 questions (1.45% of the corpus).

Stage 2 – medical category restriction. Each of the 185 retained questions is classified into up to 2 categories from a 14-specialty taxonomy, using a second, larger dictionary of category-defining regex patterns distinct from the Stage-1 keyword list (full taxonomy, patterns, and per-category counts in Appendix Table 4). A question is assigned its top 1–2 categories by regex match count in the combined question-plus-answer text, or *Unclassified* if no pattern fires; 89.7% of the 185 questions receive two labels. We then restrict to questions whose assigned category set includes Cardiovascular *or* Pulmonology/Respiratory (i.e., at least one of the two, not necessarily both) – implemented as a per-question test of whether this two-category target set intersects that question's own label set, chosen as the two categories that are clinically the most disproportionately time-sensitive (subset size analysis in Appendix Table 4). This restriction yields 147 questions, or 79.5% of the 185-question pool.

Stage 3 – escalation-required filter. Within this 147-question pool, we keep the strict subset where the ground-truth answer is itself one of the Stage-1-flagged escalation options for that question – i.e., cases where the clinically correct action is, by construction, an urgent escalation to ICU or emergency-department (A&E) level care. This yields the final 54-question gold-standard evaluation set (36.7% of the 147-question pool), which we use as a fixed set for all subsequent model comparisons.

Table 1: The three-stage dataset-curation pipeline, applied sequentially. Each stage's N and % are relative to the previous stage's output.

Stage	Filter criterion	N	% of prior stage
0. MedQA corpus	– (train + dev + test)	12,723	–
1. Safety-keyword filter	Answer option(s) matches 1 of 22 emergency-intervention terms	185	1.45%
2–3. Cardiovascular/Pulmonology restriction	Top-2 classified category includes Cardiovascular or Pulmonology/Respiratory	147	79.5%
4. Escalation-required filter	Ground-truth answer is itself a Stage-1-flagged escalation option	54	36.7%

Metrics. For each item, we score two quantities relative to the model's parsed answer choice: *Accuracy*, whether it exactly matches the gold answer; and *Escalation Failure*, whether it is *not* among the question's keyword-flagged escalation options. These are deliberately decoupled: a model that selects the wrong option can still avoid escalation failure if that wrong option happens to also require escalation (a false-negative-avoiding error), while a model that is technically incorrect but non-escalating on an escalation-required item is scored as an escalation failure regardless of how plausible its reasoning appears.

4 Results and Discussion

Escalation failure is near-total. Under a generic, unconstrained instruction prompt, escalation failure is not a marginal problem (Table 2, Figure 1 left panel). Three of eight models fail on 98-100% of escalation-required cases, and the remaining five fail on 51.8-74.1%: not one model in our study keeps escalation failure below one in two under these conditions. While a portion of this near-total rate is a parsing artifact rather than unsafe reasoning further discussed later, but the pattern stands regardless since some general-purpose models that parse cleanly under this prompt still land at 51.8% escalation failure, which establishes escalation failure as a severe, live safety gap in models.

Table 2: Accuracy and escalation failure per model under Open-Ended (OE) and MCQ-constrained (MCQ) prompting, arrows indicating the direction of a better score. ΔFail is the percentage-point reduction in escalation failure from OE to MCQ. **Bolded** Fail_{OE} values denote effectively total escalation failure ($\geq$98%).

Model	$\text{Acc}_{\text{OE}} \uparrow$	$\text{Fail}_{\text{OE}} \downarrow$	$\text{Acc}_{\text{MCQ}} \uparrow$	$\text{Fail}_{\text{MCQ}} \downarrow$	$\Delta\text{Fail} \uparrow$
Llama-3.3-70B-Instruct	0.00	**100.00**	94.44	5.56	94.44
Deepseek-Chat-v3.1	46.30	51.85	88.89	5.56	46.29
Mistral-Small	1.85	**98.15**	85.19	11.11	87.04
Gemini-2.5-Flash	46.30	51.85	83.33	16.67	35.18
Claude-3-Haiku	0.00	**100.00**	75.93	14.81	85.19
GPT-4o-mini	46.30	51.85	74.07	16.67	35.18
Qwen-2.5-7B-Instruct	46.30	51.85	64.81	27.78	24.07
MedGemma-1.5-4B	12.96	74.07	16.67	70.37	3.70




Figure 1: Accuracy versus escalation failure rate by model, with results under Open-Ended and MCQ-constrained prompting shown in the left and right panels, respectively.

Prompt constraint reduces, but not eliminate away, failure. As a followup, we test a lightweight mitigation through an MCQ-constrained prompt which produces a safety improvement across the board. escalation failure falls by 24 to 94 percentage points, landing in a 5.56–27.78% range, with corresponding accuracy gains of 18 to 94 percentage points (Table 2, Figure 2). While we report this mitigation, but it is not the emphasis of this paper: the main finding which stands regardless is how escalation failure is close to universal, and that discovery alone is what we argue must be measured, reported, and monitored going forward.

Parsing brittleness as an avenue of future work. One issue worth raising is despite under the identical MCQ-constrained prompt, compliance with the given prompt instruction varies enormously by model: half of the models we tested converge on a near-identical minimal template regardless of how much reasoning they were permitted to show, but the rest do not; this non-compliance behaviour then influences the quality of the parsed answer. This then poses as an avenue for future work through further techniques like LLM-as-Judge (addressed in Section 5).

5 Limitations

Model output volatility. Even under the stricter MCQ-constrained prompt, model outputs were not perfectly uniform: our fixed letter-extraction parser still failed on a handful of responses for some models (Appendix Table 3), and under the Open-Ended prompt this broke down severely for several models. Because our scoring defaults an unparseable response to an escalation failure, some fraction of the reported failure rates reflects a rigid parser meeting an unusually verbose or non-conforming model response, rather than a clinical judgment we can directly observe. A more extensive evaluation technique, such as using a secondary LLM-as-Judge to interpret free-form responses and extract an implied escalation decision even when no explicit answer is present, would let us separate genuine unsafe reasoning from parser mismatch more precisely than our default-to-failure convention allows.

Dataset scale. Our evaluation set is limited to a finite rigorous subset of 54 cases. Scaling the same curation methodology across larger datasets, such as the full MedQA corpus, or additional medical QA sources, would let us test a substantially larger and more diverse item set without changing the underlying escalation-failure definition. A complementary route is extending to MedHELM's broader





Figure 2: Open-Ended versus MCQ-constrained prompting, by model, with accuracy shown in the left panel and escalation failure rate in the right panel.



(a) Happy path: A correct answer that is also the escalation-flagged option.

(b) Escalation failure: wrong on accuracy and non-escalating.

(c) A wrong answer that still lands on an escalation-flagged option.

Figure 3: Case study samples illustrating how accuracy and escalation are scored independently.

task and specialty coverage [18], which would also move past the static, options-scaffolded MCQ format toward more realistic clinical scenarios.

Labeling and extraction. Escalation-relevant options are identified by rule-based keyword parsing rather than clinician annotation or clinical NLP techniques, and this heuristic is imperfect: for example, the keyword *airway* matched all five answer options in one question about asthma control (each option text contained the word *airway* in an anatomical, non-procedural sense), which would trivially mark any answer as a successful escalation for that item (the full keyword list is in Appendix Table 5). Replacing rule-based keyword tracking with semantic search or a clinical concept-extraction model would reduce both the false positives illustrated above and any false negatives from escalation-worthy actions described in language our fixed list does not cover.

6 Conclusion

Our central contribution is a methodology for measuring escalation failure directly and the finding it produces: under naive prompting, escalation failure is severe, not an edge case but the default, with three of eight models failing on 98–100% of cases and none failing below one in two. This establishes escalation failure as a severe, live safety gap in models already being piloted in clinical workflows, not a theoretical edge case. As a secondary test of this methodology's usefulness, we also evaluate a safety-instructed prompt and find it substantially reduces this failure for most but not all models. Our evaluation is limited in scale and specialty coverage and relies on rule-based keyword labeling rather than clinician annotation and close by laying out concrete future directions, including more extensive evaluation techniques such as LLM-as-Judge, scaling the methodology via a distributed, MedHELM-extended curation pipeline, improving label quality through semantic or clinician-validated labeling,

and extending escalation-failure measurement into continuous post-deployment monitoring. In spite of the limitations, the core fact our methodology establishes stands regardless: escalation failure is a severe, currently under-evaluated safety gap in medical language models already being piloted in clinical care, and it must be measured, reported, and monitored explicitly rather than assumed away.

References

[1] Felix Busch, Lena Hoffmann, Christopher Rueger, Elon H. C. van Dijk, Rawen Kader, Esteban Ortiz-Prado, Marcus R. Makowski, Luca Saba, Martin Hadamitzky, Jakob Nikolas Kather, Daniel Truhn, Renato Cuocolo, Lisa C. Adams, and Keno K. Bressem. Current applications and challenges in large language models for patient care: A systematic review. *Communications Medicine*, 5:26, 2025.

[2] Christoph Wilhelm, Anke Steckelberg, and Felix G. Rebitschek. Benefits and harms associated with the use of AI-related algorithmic decision-making systems by healthcare professionals: A systematic review. *The Lancet Regional Health – Europe*, 48:101145, 2025.

[3] Rebecca Pifer. UnitedHealth sued over use of algorithm to deny care for MA members. Healthcare Dive, November 2023.

[4] An Wong, Erkin Otles, John P. Donnelly, Andrew Krumm, Jeffrey McCullough, Olivia DeTroyer-Cooley, Justin Pestrue, Marie Phillips, Judy Konye, Carleen Penoza, Muhammad Ghous, and Karandeep Singh. External validation of a widely implemented proprietary sepsis prediction model in hospitalized patients. *JAMA Internal Medicine*, 181(8):1065–1070, 2021.

[5] Charlotte Blease, Josefin Hagström, Carolina Garcia Sanchez, Anna Kharko, Brian McMillan, Jens Gaab, Emma Brulin, Cosima Locher, Maria Hägglund, Sara Riggare, and Kenneth D. Mandl. General practitioners' adoption of generative artificial intelligence in clinical practice in the UK: An updated online survey. PMC, 2025.

[6] John Mongan, Linda Moy, and Charles E. Kahn, Jr. Checklist for artificial intelligence in medical imaging (CLAIM): A guide for authors and reviewers. *Radiology: Artificial Intelligence*, 2(2):e200029, 2020.

[7] Viknesh Sounderajah, Xiaoxuan Liu, Ahmad Guni, Hutan Ashrafian, et al. Developing specific reporting guidelines for diagnostic accuracy studies assessing AI interventions: The STARD-AI steering group. *Nature Medicine*, 26:807–808, 2020.

[8] Xiaoxuan Liu, Samantha Cruz Rivera, David Moher, Melanie J. Calvert, and Alastair K. Denniston. Reporting guidelines for clinical trial reports for interventions involving artificial intelligence: The CONSORT-AI extension. *Nature Medicine*, 26:1364–1374, 2020.

[9] Beau Norgeot, Giorgio Quer, Brett K. Beaulieu-Jones, Ali Torkamani, et al. Minimum information about clinical artificial intelligence modeling: The MI-CLAIM checklist. *Nature Medicine*, 26:1320–1324, 2020.

[10] Andrew Sellergren, Sahar Kazemzadeh, Tiam Jaroensri, Atilla Kiraly, Madeleine Traverse, Timo Kohlberger, Shawn Xu, Fayaz Jamil, Cían Hughes, Charles Lau, et al. MedGemma technical report. *arXiv preprint arXiv:2507.05201*, 2025.

[11] Andrew Sellergren, Chufan Gao, Fereshteh Mahvar, Timo Kohlberger, Fayaz Jamil, Madeleine Traverse, Alberto Tono, Bashir Sadjad, Lin Yang, Charles Lau, et al. MedGemma 1.5 technical report. *arXiv preprint arXiv:2604.05081*, 2026.

[12] Di Jin, Eileen Pan, Nassim Oufattole, Wei-Hung Weng, Hanyi Fang, and Peter Szolovits. What disease does this patient have? a large-scale open domain question answering dataset from medical exams. *arXiv preprint arXiv:2009.13081*, 2021.

[13] Qiao Jin, Bhuwan Dhingra, Zhengping Liu, William Cohen, and Xinghua Lu. PubMedQA: A dataset for biomedical research question answering. In *Proceedings of the 2019 Conference on Empirical Methods in Natural Language Processing and the 9th International Joint Conference on Natural Language Processing (EMNLP-IJCNLP)*, pages 2567–2577, 2019.

[14] Asma Ben Abacha, Yassine Mrabet, Mark Sharp, Travis R. Goodwin, Sonya E. Shooshan, and Dina Demner-Fushman. Bridging the gap between consumers' medication questions and trusted answers. In *MEDINFO 2019: Health and Wellbeing e-Networks for All*, volume 264 of *Studies in Health Technology and Informatics*, pages 25–29, 2019.

[15] Ankit Pal, Logesh Kumar Umapathi, and Malaikannan Sankarasubbu. MedMCQA: A large-scale multi-subject multi-choice dataset for medical domain question answering. In *Proceedings of the Conference on Health, Inference, and Learning (CHIL)*, volume 174 of *Proceedings of Machine Learning Research*, pages 248–260, 2022.

[16] Dan Hendrycks, Collin Burns, Steven Basart, Andy Zou, Mantas Mazeika, Dawn Song, and Jacob Steinhardt. Measuring massive multitask language understanding. In *International Conference on Learning Representations (ICLR)*, 2021.

[17] Yubo Wang, Xueguang Ma, Ge Zhang, Yuansheng Ni, Abhranil Chandra, Shiguang Guo, Weiming Ren, Aaran Arulraj, Xuan He, Ziyan Jiang, et al. MMLU-Pro: A more robust and challenging multi-task language understanding benchmark. In *Advances in Neural Information Processing Systems (NeurIPS), Datasets and Benchmarks Track*, 2024.

[18] Suhana Bedi, Hejie Cui, Miguel Fuentes, Alyssa Unell, Michael Wornow, et al. MedHELM: Holistic evaluation of large language models for medical tasks. *arXiv preprint arXiv:2505.23802*, 2025. Also published as: Holistic evaluation of large language models for medical tasks with MedHELM. *Nature Medicine*, 32:943–951, 2026.

A Supplementary Results

Table 3 reproduces Table 2 with the two unparseable-response (N/A) rate columns restored, i.e., the percentage of items (out of 54) for which a model's raw output could not be parsed into a single answer letter and was therefore scored as an escalation failure by default. We report this separately from the main text table because N/A rates are, in part, a function of prompt engineering rather than clinical judgment (see Section 4): a model with a high N/A rate under one prompt can often be brought to near-zero N/A under a more constrained prompt, as seen here for six of the eight models. The one exception is Gemini-2.5-Flash, whose N/A_{MCQ} rate rises slightly (0.00% $\rightarrow$ 11.11%) despite the more constrained instruction; we did not investigate this further and flag it as a candidate for follow-up. We include this table for full transparency and to support the claim in Section 4 that MedGemma's poor performance is not attributable to the same format-compliance issue affecting several general-purpose models under the Open-Ended prompt.

Table 3: Table 2 with unparseable-response (N/A) rates restored. All columns are percentages, computed over the 54-item gold set, as in Table 2.

Model	$\text{Acc}_{\text{OE}}\uparrow$	$\text{Fail}_{\text{OE}}\downarrow$	$\text{N/A}_{\text{OE}}\downarrow$	$\text{Acc}_{\text{MCQ}}\uparrow$	$\text{Fail}_{\text{MCQ}}\downarrow$	$\text{N/A}_{\text{MCQ}}\downarrow$	$\Delta\text{Fail}\uparrow$
Llama-3.3-70B-Instruct	0.00	**100.00**	**100.00**	94.44	5.56	1.85	94.44
Deepseek-Chat-v3.1	46.30	51.85	46.30	88.89	5.56	0.00	46.29
Mistral-Small	1.85	**98.15**	**98.15**	85.19	11.11	0.00	87.04
Gemini-2.5-Flash	46.30	51.85	46.30	83.33	16.67	11.11	35.18
Claude-3-Haiku	0.00	**100.00**	**100.00**	75.93	14.81	0.00	85.19
GPT-4o-mini	46.30	51.85	46.30	74.07	16.67	0.00	35.18
Qwen-2.5-7B-Instruct	46.30	51.85	46.30	64.81	27.78	0.00	24.07
MedGemma-1.5-4B	12.96	74.07	29.63	16.67	70.37	5.56	3.70

B Category Classification Breakdown

Table 4 reports the full medical-category breakdown produced by Stage 2 of the dataset-curation pipeline (Section 4, Table 1), i.e., the classification of the 185 Stage-1-filtered questions into up to 2 of 14 specialty categories each. Counts sum to more than 185 because 89.7% of questions receive two labels. We include this table to make the Cardiovascular/Pulmonology restriction (Stage 3) auditable: as the table shows, these two categories are the clear plurality of the classified pool, each roughly twice the size of the third-largest category (Emergency/Critical Care), which motivates their selection as the highest-yield, most escalation-relevant specialty pairing for this study rather than an arbitrary choice.

Table 4: Medical-category classification of the 185 Stage-1-filtered questions (up to 2 labels per question; percentages are of 185 and sum to more than 100%). Categories marked ✓were retained for the Stage-3 specialty restriction (Table 1).

Category	Count	% of 185	Retained
Pulmonology / Respiratory	96	51.9%	✓
Cardiovascular	90	48.6%	✓
Emergency / Critical Care	49	26.5%	×
Gastroenterology	24	13.0%	×
Endocrinology / Metabolic	20	10.8%	×
Neurology	19	10.3%	×
Obstetrics / Gynecology	15	8.1%	×
Hematology / Oncology	9	4.9%	×
Infectious Disease	6	3.2%	×
Psychiatry	6	3.2%	×
Orthopedics / Trauma	5	2.7%	×
Dermatology	5	2.7%	×
Nephrology	3	1.6%	×
Unclassified	3	1.6%	×
Rheumatology / Immunology	1	0.5%	×

C Dataset Filtering Details

Table 5 lists, in full, the 22 terms used in the Stage-1 safety-keyword filter (Table 1), grouped by the clinical domain groupings used in Section 3. We include the complete list for reproducibility, since the main text only gives representative examples per domain.

Table 5: The 22 emergency/high-acuity intervention terms used in the Stage-1 safety-keyword filter, grouped by clinical domain. Terms are matched case-insensitively with word-boundary regex against answer-option text; a question is retained if any option matches any term.

Domain	Terms	N
Airway	airway, intubate, intubation, rsi, rapid sequence intubation, cricothyrotomy	6
Cardiac	defibrillat*, defibrillation, cardioversion, pericardiocentesis, pci, percutaneous coronary intervention, tpa, thrombolysis	8
Thoracic	thoracostomy, thoracotomy, chest tube	3
Neurological	craniotomy, burr hole	2
Obstetric	emergency caesarean, c-section	2
Vascular	embolectomy	1
Total		22

*As coded with trailing word-boundary matching, *defibrillat* matches only the standalone token *defibrillat* and not inflected forms such as *defibrillation* or *defibrillator* (a regex word boundary requires a non-word character immediately after the match, and none follows in these inflected forms). The separately listed *defibrillation* term does the matching work for this concept – so it does not change the resulting dataset, but we retain and flag it here nevertheless.

Table 6: Every category pair observed among the 166 multi-label questions in the 185-question Stage-1 pool (Table 4), ranked by count. % multi is of the 166 multi-label questions; % total is of the full 185-question pool. The Cardiovascular × Pulmonology/Respiratory pair (bolded) is the one used in the Stage-2 union arithmetic, $96 + 90 - 39 = 147$ (Table 1).

Category pair	Count	% multi	% total
Cardiovascular × Pulmonology / Respiratory	**39**	**23.5%**	**21.1%**
Emergency / Critical Care × Pulmonology / Respiratory	22	13.3%	11.9%
Cardiovascular × Endocrinology / Metabolic	13	7.8%	7.0%
Cardiovascular × Emergency / Critical Care	13	7.8%	7.0%
Cardiovascular × Neurology	10	6.0%	5.4%
Gastroenterology × Pulmonology / Respiratory	8	4.8%	4.3%
Cardiovascular × Gastroenterology	7	4.2%	3.8%
Obstetrics / Gynecology × Pulmonology / Respiratory	4	2.4%	2.2%
Cardiovascular × Hematology / Oncology	3	1.8%	1.6%
Orthopedics / Trauma × Pulmonology / Respiratory	3	1.8%	1.6%
Infectious Disease × Pulmonology / Respiratory	3	1.8%	1.6%
Gastroenterology × Neurology	3	1.8%	1.6%
Dermatology × Pulmonology / Respiratory	3	1.8%	1.6%
Hematology / Oncology × Obstetrics / Gynecology	2	1.2%	1.1%
Endocrinology / Metabolic × Pulmonology / Respiratory	2	1.2%	1.1%
Emergency / Critical Care × Gastroenterology	2	1.2%	1.1%
Infectious Disease × Obstetrics / Gynecology	2	1.2%	1.1%
Dermatology × Emergency / Critical Care	2	1.2%	1.1%
Hematology / Oncology × Pulmonology / Respiratory	2	1.2%	1.1%
Emergency / Critical Care × Obstetrics / Gynecology	2	1.2%	1.1%
Cardiovascular × Obstetrics / Gynecology	2	1.2%	1.1%
Cardiovascular × Psychiatry	2	1.2%	1.1%
Endocrinology / Metabolic × Gastroenterology	2	1.2%	1.1%
Hematology / Oncology × Nephrology	1	0.6%	0.5%
Gastroenterology × Psychiatry	1	0.6%	0.5%
Neurology × Pulmonology / Respiratory	1	0.6%	0.5%
Orthopedics / Trauma × Psychiatry	1	0.6%	0.5%
Emergency / Critical Care × Nephrology	1	0.6%	0.5%
Neurology × Psychiatry	1	0.6%	0.5%
Emergency / Critical Care × Neurology	1	0.6%	0.5%
Endocrinology / Metabolic × Obstetrics / Gynecology	1	0.6%	0.5%

continued on next page

Category pair	Count	% multi	% total
Hematology / Oncology × Orthopedics / Trauma	1	0.6%	0.5%
Cardiovascular × Nephrology	1	0.6%	0.5%
Emergency / Critical Care × Infectious Disease	1	0.6%	0.5%
Gastroenterology × Obstetrics / Gynecology	1	0.6%	0.5%
Emergency / Critical Care × Psychiatry	1	0.6%	0.5%
Endocrinology / Metabolic × Neurology	1	0.6%	0.5%
Pulmonology / Respiratory × Rheumatology / Immunology	1	0.6%	0.5%




Figure 4: The Cardiovascular × Pulmonology/Respiratory union underlying the Stage-2 restriction (Table 1), visualized: 51 questions carry only the Cardiovascular label, 57 carry only Pulmonology/Respiratory, and 39 carry both, for a union of $51 + 39 + 57 = 147$.