

# ON THE DIRECTIONS OCCURRING IN LATTICE-LINE COVERINGS OF THE INTEGER PLANE

JAN SNELLMAN 

Matematiska Institutionen, Linköpings Universitet, 581 83 Linköping, Sweden  
`jan.snellman@liu.se`

**ABSTRACT.** We consider families of lines that cover every point of the integer lattice  $\mathbb{Z}^2$ , subject to the constraint that no two lines of different direction in the family meet at a lattice point. Restricting to *lattice lines* (lines containing at least two, hence infinitely many, lattice points, equivalently of rational direction), we show that the set of directions occurring in such a covering can be made dense in the space of line directions. The construction is a recursive splitting of  $\mathbb{Z}^2$  into nested rank-2 sublattice cosets, each handed off to a freshly chosen direction; the key technical point is a steering lemma showing that at every stage of the recursion a new direction arbitrarily close to any prescribed target can still be realized, via an elementary sieve bound.

*Note.* An earlier version of this work appeared as the technical report [10]. The present manuscript extends it; see `changes-wrt-report.md` in the source repository [9] for a running account of what has been added or altered since.

## 1. INTRODUCTION

Consider the integer lattice  $\mathbb{Z}^2 \subset \mathbb{R}^2$ . We wish to cover every point of  $\mathbb{Z}^2$  by a family  $\mathcal{F}$  of lines, subject to: any two lines of  $\mathcal{F}$  may cross, but never at a point of  $\mathbb{Z}^2$ . Which sets of line-directions can occur across such a family?

*Remark 1* (The lattice-line hypothesis). As posed, with no further restriction on the lines allowed, the question is trivial: for each  $z \in \mathbb{Z}^2$  take the line  $L_z$  through  $z$  of irrational slope  $\alpha_z$ . Then

$$L_z \cap \mathbb{Z}^2 = \{z\} \tag{1}$$

(a second lattice point on  $L_z$  would force  $\alpha_z \in \mathbb{Q}$ ), so distinct such lines never share a lattice point, and choosing the  $\alpha_z$  to range over a dense (indeed, all) set of irrational directions already realizes an uncountable dense set of directions. Everything of interest in this manuscript therefore concerns the restriction to *lattice lines*: lines containing at least two (hence, by the same argument, infinitely many) points of  $\mathbb{Z}^2$ , equivalently lines of rational direction. This restriction is in force throughout.

We now formalize the objects of study as explicit definitions.

**Definition 2** (Primitive direction). A *primitive direction* is a pair  $d = (p, q) \in \mathbb{Z}^2$  with  $\gcd(p, q) = 1$ . Two primitive directions  $d, d'$  are *equal as directions* if  $d' = \pm d$ . Write

$$\varphi_d(x, y) = qx - py, \tag{2}$$

a surjective homomorphism  $\mathbb{Z}^2 \rightarrow \mathbb{Z}$  depending only on  $d$  up to sign of the whole map (in particular  $\ker \varphi_d = \ker \varphi_{-d}$ ). The space of directions is  $\mathbb{RP}^1 = S^1/\{\pm 1\}$ , parametrized by  $\theta \in [0, \pi)$ ; we metrize it by

$$d(\theta, \theta') = \min(|\theta - \theta'|, \pi - |\theta - \theta'|), \quad (3)$$

so “within  $\varepsilon$  of  $\theta$ ” always means within  $\varepsilon$  in this metric. Since  $[0, \pi) \rightarrow \mathbb{RP}^1$  is a continuous surjection, a set dense in  $[0, \pi)$  is automatically dense in  $\mathbb{RP}^1$ .

**Definition 3** (Lattice lines). For a primitive direction  $d = (p, q)$  and  $c \in \mathbb{Z}$ , the *lattice line*  $\ell_{d,c}$  is the level set

$$\ell_{d,c} = \varphi_d^{-1}(c) = \{(x, y) \in \mathbb{Z}^2 : qx - py = c\}. \quad (4)$$

Every line in  $\mathbb{R}^2$  containing at least two points of  $\mathbb{Z}^2$  equals  $\ell_{d,c}$  for a unique primitive direction  $d$  (up to sign) and a unique  $c \in \mathbb{Z}$ ; we call such a line a lattice line of direction  $d$ .

**Definition 4** (Covering family, valid family). A *covering family* is a set  $\mathcal{F}$  of lattice lines with  $\bigcup \mathcal{F} \supseteq \mathbb{Z}^2$ . A covering family  $\mathcal{F}$  is *valid* if no two lines of  $\mathcal{F}$  of different direction share a point of  $\mathbb{Z}^2$ ; equivalently, for every pair of distinct primitive directions  $d, d'$  occurring in  $\mathcal{F}$ , and every  $\ell_{d,c}, \ell_{d',c'} \in \mathcal{F}$ , we have  $\ell_{d,c} \cap \ell_{d',c'} \cap \mathbb{Z}^2 = \emptyset$ .




FIGURE 1. Two lattice lines of different directions  $d, d'$  may cross *off* the lattice (diamond marker) without violating validity (Definition 4): only crossings *at* points of  $\mathbb{Z}^2$  (dots) are forbidden between distinct directions.

**Lemma 5** (The everywhere-disjoint variant is trivial). *If the lines of  $\mathcal{F}$  are required to be pairwise disjoint everywhere (not merely off the lattice), then every line in  $\mathcal{F}$  shares a single direction, and any one direction already suffices (Example 6 below).*

*Proof.* Two distinct lines in  $\mathbb{R}^2$  are either parallel or meet at exactly one point, so pairwise disjointness forces every line in the family to share a single direction.  $\square$

All the content is in the “crossings allowed off the lattice” variant of Definition 4, which is our subject here.

**Example 6.** For a primitive direction  $d = (p, q)$ , the lines  $\ell_{d,c}$ ,  $c \in \mathbb{Z}$ , partition  $\mathbb{Z}^2$  (they are the fibers of the surjective homomorphism  $\varphi_d$ ):

$$\mathbb{Z}^2 = \bigsqcup_{c \in \mathbb{Z}} \ell_{d,c}. \quad (5)$$

So a single direction always suffices to cover  $\mathbb{Z}^2$  by pairwise-disjoint (in particular lattice-point-disjoint) lattice lines.

Our main result:

**Theorem 7.** *There is a valid covering family  $\mathcal{F}$  (Definition 4) whose set of realized directions is dense in  $\mathbb{RP}^1$ .*




FIGURE 2. A preview of Theorem 7: the first 100 directions realized by the construction of Section 3 below, each drawn as a diameter of the unit circle. See Figure 7 for the same data with the construction's progress (order and consecutive jumps) made visible.

## 2. TWO LEMMAS ON LATTICE LINES

**Lemma 8** (Rigidity). *Let  $d_1 = (p_1, q_1)$ ,  $d_2 = (p_2, q_2)$  be two distinct primitive directions, and set*

$$\Delta = p_1 q_2 - p_2 q_1 \neq 0. \quad (6)$$

*The lines  $\ell_{d_1, c_1}$  and  $\ell_{d_2, c_2}$  meet at a lattice point if and only if*

$$\Delta \mid (p_1 c_2 - p_2 c_1) \quad \text{and} \quad \Delta \mid (q_1 c_2 - q_2 c_1). \quad (\text{R})$$

*In particular, if  $|\Delta| = 1$ , condition (R) holds vacuously for every choice of  $c_1, c_2$ , so  $d_1, d_2$  can never both occur in a valid family (Definition 4).*

*Proof.* A lattice point  $(x, y)$  lies on both lines exactly when the linear system

$$\begin{aligned} q_1 x - p_1 y &= c_1, \\ q_2 x - p_2 y &= c_2 \end{aligned} \quad (7)$$

holds. Eliminating  $y$  gives

$$\Delta x = p_1 c_2 - p_2 c_1, \quad (8)$$

and eliminating  $x$  symmetrically gives

$$\Delta y = q_1 c_2 - q_2 c_1. \quad (9)$$

Since  $\Delta \neq 0$ , equations (8)–(9) determine  $x, y \in \mathbb{Q}$  uniquely (this is exactly Cramer’s rule applied to (7)), and  $(x, y) \in \mathbb{Z}^2$  if and only if  $\Delta$  divides both right-hand sides, which is precisely (R). Conversely, if (R) holds, the quotients in (8)–(9) are integers and one checks directly, by substituting back into (7), that they solve the system. If  $|\Delta| = 1$ , both divisibility conditions in (R) hold automatically for every  $c_1, c_2 \in \mathbb{Z}$ .  $\square$

**Lemma 9** (Coset). *Let  $p, q$  be coprime integers,  $x_0, y_0 \in \mathbb{Z}$ , and set*

$$c_0 = qx_0 - py_0. \quad (10)$$

*For  $k \in \mathbb{Z}$ , write  $r_k = c_0 + pqk$ . Then*

$$\bigcup_{k \in \mathbb{Z}} \ell_{(p,q), r_k} = \{(x, y) \in \mathbb{Z}^2 : x \equiv x_0 \pmod{|p|}, y \equiv y_0 \pmod{|q|}\}. \quad (11)$$

*Proof.* Write  $C$  for the right-hand side of (11). Since  $\gcd(p, q) = 1$  implies in particular  $p, q \neq 0$ , congruence modulo  $p$  (resp.  $q$ ) is well defined, and  $C$  admits the parametrization

$$C = \{(x_0 + pu, y_0 + qv) : u, v \in \mathbb{Z}\}. \quad (12)$$

For a point of the form (12),

$$\varphi_{(p,q)}(x_0 + pu, y_0 + qv) = q(x_0 + pu) - p(y_0 + qv) = c_0 + pq(u - v), \quad (13)$$

using (10) and cancelling the  $pqu$  terms. By (4), this point lies on  $\ell_{(p,q), r_k}$  if and only if  $u - v = k$ . As  $u$  ranges over  $\mathbb{Z}$  (with  $v = u - k$  then determined), every point of  $C$  with that particular value of  $u - v$  is visited exactly once, and (13) shows every point of  $\ell_{(p,q), r_k} \cap C$  has  $u - v = k$ : hence

$$\ell_{(p,q), r_k} \cap C = \{(x_0 + pu, y_0 + q(u - k)) : u \in \mathbb{Z}\}, \quad (14)$$

and moreover every point of  $\ell_{(p,q), r_k}$  manifestly lies in  $C$  (take  $u, v$  with  $u - v = k$  so that (13) matches  $r_k$ ; such  $(x, y)$  automatically satisfies  $x \equiv x_0 \pmod{p}$ ,  $y \equiv y_0 \pmod{q}$ ). So  $\ell_{(p,q), r_k} \subseteq C$  for every  $k$ , with equality of the union to all of  $C$  once  $k$  ranges over  $\mathbb{Z}$ , since every pair  $(u, v) \in \mathbb{Z}^2$  has *some* value of  $u - v$ .  $\square$

Recall that by *Farey neighbors* is meant two primitive directions

$$d_1 = (p_1, q_1), \quad d_2 = (p_2, q_2)$$

(equivalently, fractions  $p_1/q_1, p_2/q_2$  in lowest terms) with

$$|p_1 q_2 - p_2 q_1| = 1.$$

This is exactly the  $|\Delta| = 1$  case of Lemma 8; see [4, Ch. III] for the classical theory. The rigidity lemma shows that a family realizing many directions cannot simply throw them together: directions related by a determinant of  $\pm 1$  (“unimodular pairs”, e.g. any pair of Farey neighbors read as slopes) are permanently incompatible. The coset lemma is the tool that lets us build a family avoiding all bad pairs at once: applied recursively below, it splits the




FIGURE 3. Lemma 9 for  $d = (p, q) = (3, 2)$  and  $(x_0, y_0) = (1, 1)$ , so that  $c_0 = qx_0 - py_0 = -1$  and  $r_k = -1 + 6k$ . Small grey dots are  $\mathbb{Z}^2$ ; large dots are the coset  $C = \{(x, y) : x \equiv 1 \pmod{3}, y \equiv 1 \pmod{2}\}$ , a translate of the sublattice  $3\mathbb{Z} \times 2\mathbb{Z}$ . Each line  $\ell_{(3,2),r_k}$  meets  $\mathbb{Z}^2$  *only* in  $C$ , and consecutive points along a line differ by  $(p, q) = (3, 2)$ ; as  $k$  ranges over  $\mathbb{Z}$  the lines partition  $C$ . The levels are equally spaced with gap  $pq = 6$ : a smaller gap would give lines missing  $\mathbb{Z}^2$  altogether, a larger one would leave points of  $C$  uncovered.

current sub-coset into finer sub-cosets, one of which is handed off to each freshly chosen direction.

### 3. THE CONSTRUCTION

**Definition 10** (Rank-2 coset). For coprime positive integers  $n_1, n_2$  and  $x_0, y_0 \in \mathbb{Z}$ , define

$$R(n_1, n_2, x_0, y_0) = \{(x, y) \in \mathbb{Z}^2 : x \equiv x_0 \pmod{n_1}, y \equiv y_0 \pmod{n_2}\}. \quad (15)$$

Since  $n_1, n_2 > 0$ , every  $(x, y) \in R(n_1, n_2, x_0, y_0)$  is of the form  $x = x_0 + n_1u$ ,  $y = y_0 + n_2w$  for a *unique* pair  $(u, w) \in \mathbb{Z}^2$ ; call  $(u, w)$  the *internal coordinates* of the point.

We state the two halves of the original Splitting lemma as two separate lemmas: the first identifies each finer sub-coset with a plain coset of a new direction, and the second records that these sub-cosets genuinely partition  $R$ .

**Lemma 11** (Splitting: coset identity). *Let  $R = R(n_1, n_2, x_0, y_0)$ , let  $s, t$  be nonzero integers with*

$$\gcd(s, n_2) = \gcd(t, n_1) = \gcd(s, t) = 1, \quad (\text{H})$$

*and put*

$$P = n_1s, \quad Q = n_2t. \quad (16)$$

For  $u_0, w_0 \in \mathbb{Z}$ , let

$$R_{u_0, w_0} = \{(x, y) \in R : x = x_0 + n_1 u, y = y_0 + n_2 w \text{ for some } u \equiv u_0 \pmod{|s|}, w \equiv w_0 \pmod{|t|}\} \quad (17)$$

(a sub-coset of  $R$ , picked out via the internal coordinates of Definition 10). Then

$$\bigcup_{k \in \mathbb{Z}} \ell_{(P, Q), c_0 + PQk} = R_{u_0, w_0}, \quad c_0 = Q(x_0 + n_1 u_0) - P(y_0 + n_2 w_0). \quad (S)$$

In particular  $(P, Q)$  is itself a primitive direction:  $\gcd(P, Q) = 1$ .

*Proof. Primitivity of  $(P, Q)$ .* A prime dividing both  $P = n_1 s$  and  $Q = n_2 t$  would have to divide one of  $n_1, n_2$  (excluded, as  $\gcd(n_1, n_2) = 1$ ), or  $n_1$  and  $t$  (excluded by  $\gcd(t, n_1) = 1$  in (H)), or  $s$  and  $n_2$  (excluded by  $\gcd(s, n_2) = 1$ ), or  $s$  and  $t$  (excluded by  $\gcd(s, t) = 1$ ). These four cases are exhaustive, so no such prime exists.

The identity (S). For  $(x, y) = (x_0 + n_1 u, y_0 + n_2 w)$ ,

$$Qx - Py = Q(x_0 + n_1 u) - P(y_0 + n_2 w) = (Qx_0 - Py_0) + n_1 n_2 (tu - sw), \quad (18)$$

using  $Q = n_2 t, P = n_1 s$ . Put  $c_0^{\text{loc}} = tu_0 - sw_0$ . Then, by Lemma 9 applied to the coprime pair  $(s, t)$  in the *internal* coordinates  $(u, w)$ ,

$$\bigcup_{k \in \mathbb{Z}} \{(u, w) : tu - sw = c_0^{\text{loc}} + stk\} = \{u \equiv u_0 \pmod{|s|}, w \equiv w_0 \pmod{|t|}\}. \quad (19)$$

Substituting the local level  $c_0^{\text{loc}} + stk$  into (18) via  $x_0, y_0$  translates it into the global level

$$\begin{aligned} (Qx_0 - Py_0) + n_1 n_2 (c_0^{\text{loc}} + stk) &= c_0 + n_1 n_2 st k \\ &= c_0 + PQk, \end{aligned} \quad (20)$$

where the first equality uses  $c_0 = (Qx_0 - Py_0) + n_1 n_2 c_0^{\text{loc}}$  (a direct computation confirming this matches the formula for  $c_0$  stated in (S)), and the second uses  $PQ = n_1 n_2 st$ . Combining (19) and (20) with (18) gives exactly (S).  $\square$

**Example 12.** Take  $n_1 = 2, n_2 = 3, x_0 = y_0 = 1$ , so

$$R = R(2, 3, 1, 1) = \{(x, y) : x \text{ odd}, y \equiv 1 \pmod{3}\}.$$

Take  $s = 2, t = 3$ ; one checks directly that (H) holds:

$$\gcd(s, n_2) = \gcd(2, 3) = 1, \quad (21)$$

$$\gcd(t, n_1) = \gcd(3, 2) = 1, \quad (22)$$

$$\gcd(s, t) = \gcd(2, 3) = 1. \quad (23)$$

Then  $P = n_1 s = 4, Q = n_2 t = 9$  (and indeed  $\gcd(4, 9) = 1$ ). Take the residue pair  $(u_0, w_0) = (1, 2)$ ; then

$$c_0 = Q(x_0 + n_1 u_0) - P(y_0 + n_2 w_0) = 9 \cdot (1 + 2) - 4 \cdot (1 + 6) = 27 - 28 = -1. \quad (24)$$

Three points of  $R_{1,2}$ , at three different values of  $k$  in (S):

$$(u, w) = (1, 2) \mapsto (x, y) = (3, 7) : \quad \varphi_{(4,9)}(3, 7) = 27 - 28 = -1 = c_0 + 36 \cdot 0, \quad (25)$$

$$(u, w) = (3, 2) \mapsto (x, y) = (7, 7) : \quad \varphi_{(4,9)}(7, 7) = 63 - 28 = 35 = c_0 + 36 \cdot 1, \quad (26)$$

$$(u, w) = (1, 5) \mapsto (x, y) = (3, 16) : \quad \varphi_{(4,9)}(3, 16) = 27 - 64 = -37 = c_0 + 36 \cdot (-1), \quad (27)$$

using  $PQ = 36$  throughout.




FIGURE 4. Example 12. Small grey dots are  $\mathbb{Z}^2$ ; medium grey dots the coarse coset  $R = R(2, 3, 1, 1) = \{x \text{ odd}, y \equiv 1 \pmod{3}\}$ ; large dots the fine sub-coset  $R_{1,2} = \{x \equiv 3 \pmod{4}, y \equiv 7 \pmod{9}\}$  singled out by the residue pair  $(u_0, w_0) = (1, 2)$ . By Lemma 11 that sub-coset is again a plain coset, now for the direction  $(P, Q) = (4, 9)$ , and Lemma 9 applies to it verbatim: the lines shown are  $\ell_{(4,9),r_k}$  with  $r_k = -1 + 36k$ . Consecutive points along a line differ by  $(4, 9)$ ; the arrow marks  $(3, 7) \mapsto (7, 16)$ , two of the three points computed in the example. Note that the lines pass through *no* point of  $R$  outside  $R_{1,2}$  — that confinement is the content of the lemma, and what makes the recursion safe.

**Lemma 13** (Splitting: partition). *With  $R, s, t$  as in Lemma 11,*

$$R = \bigsqcup_{\substack{0 \leq u_0 < |s| \\ 0 \leq w_0 < |t|}} R_{u_0, w_0} \quad (28)$$

(a disjoint union over the  $|s||t|$  residue pairs).

*Proof.* Fix  $(x, y) \in R$ , with internal coordinates  $(u, w)$  as in Definition 10. By the division algorithm,

$$u = s \left\lfloor \frac{u}{s} \right\rfloor + u_0, \quad 0 \leq u_0 < |s|,$$

for a unique such  $u_0$  ( $u_0$  is the unique representative of  $u$  modulo  $s$  lying in  $[0, |s|)$ , and similarly  $w_0 \in [0, |t|)$  for  $w$  modulo  $t$ ). By construction,

$$u \equiv u_0 \pmod{|s|} \quad \text{and} \quad w \equiv w_0 \pmod{|t|},$$

so  $(x, y) \in R_{u_0, w_0}$  for this particular pair  $(u_0, w_0)$ , establishing

$$R \subseteq \bigcup_{u_0, w_0} R_{u_0, w_0};$$

the reverse inclusion is immediate from (17). For disjointness (equivalently, uniqueness of  $(u_0, w_0)$  for each point): suppose

$$(x, y) \in R_{u_0, w_0} \cap R_{u'_0, w'_0}$$

via internal coordinates  $(u, w)$  and  $(u'', w'')$  respectively. Since internal coordinates are unique (Definition 10),  $u'' = u$  and  $w'' = w$ , so

$$u_0 \equiv u \equiv u'_0 \pmod{|s|};$$

as both  $u_0, u'_0 \in [0, |s|)$ , an interval of length  $|s|$ , and their difference is a multiple of  $s$  of absolute value  $< |s|$ , we get  $u_0 = u'_0$ , and symmetrically  $w_0 = w'_0$ .  $\square$

A naive approach to this construction might attempt to realize an *arbitrary* target direction by using a multiple of it known to lie in the current sublattice, without checking that the multiple's own geometric line (which follows its *primitive* direction, not the multiple used to construct it) stays confined to the intended coset; this is exactly the pitfall Lemma 11 avoids, via the primitivity argument built into its own proof.

**Lemma 14** (Steering). *Let  $n_1, n_2$  be coprime positive integers. For every  $\theta \in [0, \pi)$  and every  $\varepsilon > 0$  there exist nonzero integers  $s, t$ ,  $|s|, |t| \geq 2$ , satisfying (H), such that the direction of  $(n_1 s, n_2 t)$  lies within  $\varepsilon$  of  $\theta$ .*

*Proof.* By symmetry (swap the roles of  $s$  and  $t$  to treat  $\theta = \pi/2$ , which targets reciprocal slope 0 and is handled by the same argument below with the roles of  $n_1, n_2$  exchanged, subject to the same window shift introduced at (33)) it suffices to treat  $\theta \neq \pi/2$ , i.e. a finite target slope  $\mu = \tan \theta$ ; the case  $\mu < 0$  is identical with  $t \mapsto -t$ , so assume  $\mu \geq 0$ .

Set

$$\delta = \frac{1}{2} \prod_{r|n_1} \left(1 - \frac{1}{r}\right) > 0, \quad E = 2^{\omega(n_1)+1}, \quad (29)$$






FIGURE 5. Lemma 13 for the  $R$ ,  $s = 2$ ,  $t = 3$  of Example 12. Every point of the coarse coset  $R$  carries exactly one of the  $|s||t| = 6$  residue pairs  $(u_0, w_0)$ : shape records  $u_0 \in \{0, 1\}$ , colour records  $w_0 \in \{0, 1, 2\}$ . The six classes are the subcosets  $R_{u_0, w_0}$ , and the picture is the lemma: no point carries two labels and none carries none, so the classes are disjoint and exhaust  $R$ . Circled is the class  $R_{1,2}$  of Figure 4 — one of the six, handed to a single new direction, while the other five remain available to later steps of the recursion.

both depending only on  $n_1$ , where  $\omega(n_1)$  is the number of distinct prime divisors of  $n_1$ . For a prime  $p \nmid n_1 n_2$  and any interval  $I \subset \mathbb{Z}$  of length  $L$ , inclusion–exclusion over the divisors of  $n_1 p$  gives

$$\#\{t \in I : \gcd(t, n_1 p) = 1\} \geq L \prod_{r|n_1} \left(1 - \frac{1}{r}\right) \left(1 - \frac{1}{p}\right) - 2^{\omega(n_1)+1} \geq \delta L - E, \quad (30)$$

using  $p \geq 2$  for the last factor. Fix

$$L_0 = \left\lceil \frac{E+1}{\delta} \right\rceil, \quad (31)$$

independent of  $p$  — and, crucially, independent of  $\mu$  as well. Choose a prime  $p \nmid n_1 n_2$  large enough that

$$L_0 + 2 < \frac{\varepsilon n_1 p}{n_2} \quad (32)$$

(possible since the right side  $\rightarrow \infty$  as  $p \rightarrow \infty$ , for fixed  $\varepsilon, n_1, n_2$ ; note (32) does not involve  $\mu$  at all, unlike  $t_0$  below), set  $s = p$  and

$$t_0 = \left\lfloor \frac{\mu n_1 p}{n_2} \right\rfloor + 2. \quad (33)$$

The  $+2$  shift in (33) is what makes the forthcoming  $|t| \geq 2$  requirement hold *unconditionally* rather than only “for  $p$  large”: since  $\mu \geq 0$ , already  $t_0 \geq 2$  by (33), so every integer in the window  $I$  below inherits  $|t| \geq 2$  directly, with no degenerate behaviour at  $\mu = 0$  (where, without the shift,  $t_0 = 0$  for *every*  $p$ , the window never moves away from the origin as  $p \rightarrow \infty$ , and  $t = 1$ , always coprime to  $n_1 p$ , is a witness that nothing rules out being the *only* one, violating  $|t| \geq 2$ ). The interval  $I = [t_0, t_0 + L_0]$  then contains, by (30) with  $L = L_0$  and (31), at least  $\delta L_0 - E \geq 1$  integer  $t$  with  $\gcd(t, n_1 p) = 1$  — equivalently  $\gcd(t, n_1) = \gcd(t, s) = 1$ , as  $s = p$  is prime — and, as just noted,  $|t| \geq 2$  automatically. For any such  $t$ ,

$$\left| \frac{n_2 t}{n_1 s} - \mu \right| = \frac{n_2}{n_1 p} \left| t - \frac{\mu n_1 p}{n_2} \right| < \frac{n_2(L_0 + 2)}{n_1 p} < \varepsilon, \quad (34)$$

using (32) for the last step and  $|t - \mu n_1 p / n_2| < L_0 + 2$ , which follows from  $t \in [t_0, t_0 + L_0]$  and  $t_0 - \mu n_1 p / n_2 \in (1, 2]$  by (33), so  $(P, Q) = (n_1 s, n_2 t)$  has slope within  $\varepsilon$  of  $\mu$ , hence (as  $\arctan$  is 1-Lipschitz, and  $\arctan \mu = \theta - \pi$  when  $\theta > \pi/2$  identifies the same point of  $\mathbb{RP}^1$ ) direction within  $\varepsilon$  of  $\theta$ .  $\square$

The point of Lemma 14 is that it is *not* enough to know that *some* admissible  $(s, t)$  exists at every stage: repeatedly reusing, say,  $(s, t) = (2, 3)$  is admissible forever (all three coprimality conditions of (H) hold trivially against the resulting  $n_1 = 2^k, n_2 = 3^k$ ), yet gives directions converging to a single point of  $\mathbb{RP}^1$ , not a dense set. What is needed, and what Lemma 14 provides, is the ability to steer toward an *arbitrary prescribed* target at every stage.

*Remark 15* (Tails of a dense sequence remain dense). If  $X$  is a  $T_1$  topological space with no isolated points,  $D \subseteq X$  is dense, and  $F \subset X$  is finite, then  $D \setminus F$  is still dense in  $X$ . This follows since any nonempty open  $U$  is itself infinite, as no point of  $X$  is isolated; so  $U \setminus F$  is nonempty and open, hence meets  $D$ , giving a point of  $D \setminus F$  in  $U$ . Applied with  $X = \mathbb{RP}^1$ , which is compact, metric, hence  $T_1$ , and has no isolated points (it is a circle), and  $D = \{\theta_k\}_{k \geq 1}$ : every tail  $\{\theta_k\}_{k \geq N}$  is dense, since it contains  $D \setminus \{\theta_1, \dots, \theta_{N-1}\}$ . This is the fact the density argument below needs beyond mere density of  $\{\theta_k\}$  itself — without “no isolated points” it can fail (e.g. a sequence dense in  $\{0\} \cup [1, 2]$  that visits 0 only once).

We now have all pieces needed to prove Theorem 7.

*Proof of Theorem 7.* Fix an enumeration of  $\mathbb{Z}^2$ ,  $(z_k)_{k \geq 1}$ , and a sequence  $(\theta_k)_{k \geq 1}$  dense in  $[0, \pi)$  (e.g.  $\theta_k = k\alpha \bmod \pi$  for  $\alpha/\pi$  irrational, by equidistribution of the irrational rotation). Set  $R_0 = \mathbb{Z}^2$ , i.e.  $(n_1, n_2, x_0, y_0) = (1, 1, 0, 0)$ .

Given  $R_{k-1} = R(n_1, n_2, x_0, y_0)$ , apply Lemma 14 to find  $s, t$  (possibly negative — Lemma 14 allows this, needed to reach both signs of slope) realizing a direction within  $1/k$  of  $\theta_k$ , and apply Lemmas 11–13 to split  $R_{k-1}$

into its  $|s||t| \geq 4$  sub-cosets, each fully coverable by direction  $(P_k, Q_k) = (n_1s, n_2t)$  restricted to one residue class of levels. Put into  $\mathcal{F}$  all such lines for every sub-coset except one, reserved as follows: if  $z_k \in R_{k-1}$ , reserve the lexicographically-least  $(u_0, w_0)$  whose sub-coset does not contain  $z_k$  (possible since there are  $\geq 4$  sub-cosets and only one need be avoided); otherwise reserve the lexicographically-least  $(u_0, w_0)$  outright. Let  $R_k = R_{u_0, w_0}$  be that reserved sub-coset; Figure 6 shows the first two stages of the resulting chain  $\mathbb{Z}^2 = R_0 \supset R_1 \supset R_2$ . The recursion rests on two selection rules, of rather different characters; see Remark 16.

The two quantities that matter going forward are genuinely different: the *achieved direction* recorded for the density argument below is the (possibly signed) pair  $(P_k, Q_k) = (n_1s, n_2t)$ , while  $R_k$  itself, to serve as the input  $R(n'_1, n'_2, x'_0, y'_0)$  to the *next* stage's application of Lemma 14 (whose hypotheses require positive  $n'_1, n'_2$ , as does Definition 10), must use the *positive* pair

$$n'_1 = n_1|s|, \quad n'_2 = n_2|t|, \quad x'_0 = x_0 + n_1u_0, \quad y'_0 = y_0 + n_2w_0 :$$

a direct check from the definition of  $R_{u_0, w_0}$  (17) (writing  $u = u_0 + |s|u'$ ,  $w = w_0 + |t|w'$  for the internal coordinates, since “ $u \equiv u_0 \pmod{|s|}$ ” means exactly this) shows  $R_{u_0, w_0} = R(n_1|s|, n_2|t|, x'_0, y'_0)$  on the nose.

Write  $R_k = R(n_1^{(k)}, n_2^{(k)}, x_0^{(k)}, y_0^{(k)})$  for this representation, i.e.  $n_1^{(k)} = n_1|s|$ ,  $n_2^{(k)} = n_2|t|$  (with  $n_1^{(0)} = n_2^{(0)} = 1$  from the base case), the notation used below. This keeps

$$\gcd(n_1^{(k)}, n_2^{(k)}) = \gcd(n_1|s|, n_2|t|) = \gcd(P_k, Q_k) = 1$$

(Lemma 11), so the invariant needed for the next iteration is preserved.

*Coverage invariant.* Stage  $i$  claims exactly  $R_{i-1} \setminus R_i$ , and  $R_i \subseteq R_{i-1}$  for every  $i \geq 1$ . Telescoping from  $R_0 = \mathbb{Z}^2$  gives, for every  $k$ ,

$$\mathbb{Z}^2 \setminus R_{k-1} = \bigcup_{i < k} (R_{i-1} \setminus R_i) = \bigcup_{i < k} (\text{stage-}i \text{ claims}), \quad (35)$$

i.e. a point lies outside the current region  $R_{k-1}$  exactly when some earlier stage already claimed it. This is what justifies the reservation rule above only ever needing to avoid  $z_k$  when  $z_k \in R_{k-1}$ : whenever  $z_k \notin R_{k-1}$ , (35) shows  $z_k$  was already claimed by an earlier stage, so no action is needed for it at stage  $k$ .

*Coverage.* By (35), every  $z_k$  is eventually claimed: if  $z_k \in R_{k-1}$ , the reservation rule ensures  $z_k$  lies in one of the claimed sub-cosets at stage  $k$  (not the reserved  $R_k$ ); if  $z_k \notin R_{k-1}$ , it was already claimed at some earlier stage. So

$$\bigcup_k (\text{stage-}k \text{ claims}) = \mathbb{Z}^2. \quad (36)$$

Since the stage- $k$  claims are exactly  $R_{k-1} \setminus R_k$ , and  $R_0 = \mathbb{Z}^2$ , this says precisely that

$$\bigcap_{k \geq 0} R_k = \emptyset,$$




FIGURE 6. The first two stages of the nested chain  $\mathbb{Z}^2 = R_0 \supset R_1 \supset R_2$  built in the proof of Theorem 7. Small grey dots are  $R_0 = \mathbb{Z}^2$ ; medium dots  $R_1 = R(2, 3, 0, 1) = \{x \equiv 0 \pmod{2}, y \equiv 1 \pmod{3}\}$ ; large dots  $R_2 = R(4, 9, 0, 1) = \{x \equiv 0 \pmod{4}, y \equiv 1 \pmod{9}\}$ . Each stage claims everything it does *not* reserve, so the region shrinks while the claimed set grows to fill  $\mathbb{Z}^2$ . The run shown takes  $(s, t) = (2, 3)$  at both stages and the enumeration point  $z_1 = (0, 0)$  (crossed). Since  $z_1 \in R_0$ , stage 1 may not reserve the sub-coset containing it: the lexicographically least *eligible* residue pair is  $(0, 1)$  rather than  $(0, 0)$ , which is why  $R_1$  consists of  $y \equiv 1$  rather than  $y \equiv 0$  modulo 3. That reservation rule is exactly the one the Lean formalization computes; the pairs  $(s, t)$  are illustrative only — see Remark 16.

that is: every point leaves the chain at some *finite* stage. This is what makes the recursion self-contained at order type  $\omega$ , with no step beyond those indexed by  $k \in \mathbb{N}$ .

Nesting alone would not give it. The rule “always reserve the sub-coset containing the origin” produces a chain that is strictly decreasing, with the same densities  $1/(n_1^{(k)} n_2^{(k)}) \rightarrow 0$ , and yet retains  $(0, 0)$  in every  $R_k$  — a point that no line of  $\mathcal{F}$  would ever cover. What excludes this is the reservation rule’s dependence on the enumeration  $(z_k)$ , and that works only because every point of  $\mathbb{Z}^2$  carries a finite index.

*Disjointness.* Every individual line placed into  $\mathcal{F}$  at stage  $k$  has all of its lattice points inside the sub-coset  $R_{u_0, w_0}$  it came from (a subset of  $R_{k-1} \setminus R_k$ ,

since  $R_{u_0, w_0} \neq R_k$ ): this is exactly (S) of Lemma 11 (each line on the left is a subset of the union), the step the earlier, flawed version of this construction got wrong (see the remark following Lemma 11). So stage  $k$ 's claimed points lie in

$$R_{k-1} \setminus R_k \subset R_{k-1}.$$

For  $j > k$ ,  $R_{j-1} \subseteq R_k$  (as  $R_i \subseteq R_{i-1}$  for every  $i$ ), so stage  $j$ 's points, lying in  $R_{j-1}$ , are disjoint from stage  $k$ 's. Hence distinct lines from different stages never share a lattice point, and distinct lines of the same direction from the same stage are disjoint level sets of the same  $\varphi_{(P_k, Q_k)}$ , so never share one either. (Different stages in fact always realize different directions:  $|P_k| = n_1^{(k)}$  is strictly increasing in  $k$ , since

$$n_1^{(k)} = n_1^{(k-1)} |s_k| \geq 2n_1^{(k-1)};$$

though the argument above does not need this.)

*Density.* Fix  $\theta \in \mathbb{RP}^1$  and  $\varepsilon > 0$ . By Remark 15, every tail  $\{\theta_k\}_{k \geq N}$  is dense in  $\mathbb{RP}^1$ ; so for any  $N$  there is some  $k \geq N$  with  $\theta_k$  within  $\varepsilon/2$  of  $\theta$ . Taking  $N > 2/\varepsilon$ , the corresponding achieved direction  $(P_k, Q_k)$  — within  $1/k < \varepsilon/2$  of  $\theta_k$  — then lies within  $\varepsilon$  of  $\theta$ . (Each  $(P_k, Q_k)$  is genuinely realized in  $\mathcal{F}$ : at least  $|s||t| - 1 \geq 3$  of the  $\geq 4$  sub-cosets at stage  $k$  are claimed, all nonempty, each contributing lines of that direction.) So every  $\theta \in \mathbb{RP}^1$  is a limit of directions realized in  $\mathcal{F}$ : the realized direction set is dense. □

*Remark 16* (The two selection rules). The recursion in the proof above is not a definition until two things are pinned down: which admissible pair  $(s, t)$  to take among those Lemma 14 merely asserts to exist, and which sub-coset to reserve. Neither affects the conclusion — the argument uses only the *properties* of the chosen objects, never their identity — but a recursion is a well-defined function only once both are fixed. Informally one writes “choose  $s, t$ ” and moves on; a formalization cannot, since the recursion has to be an actual term.

The two are settled in genuinely different ways, and the contrast is worth noticing. The reservation rule is *computable*: the reserved pair is the lexicographically least element of the finite set of residue pairs  $(u_0, w_0) \in [0, |s|) \times [0, |t|)$  whose sub-coset avoids  $z_k$  — a condition that is vacuous when  $z_k \notin R_{k-1}$ , which is exactly the “otherwise” clause of the rule stated in the proof. That set is nonempty because there are at least  $|s||t| \geq 4$  sub-cosets and, by Lemma 13,  $z_k$  lies in exactly one of them, so at least three remain. The steering pair, by contrast, is obtained from the existence statement by classical choice: Lemma 14 singles out no canonical  $(s, t)$ , and the proof needs none.

Why only one of the two is computable is a question of order type rather than of principle. The reservation is a minimum over the *finite* set  $[0, |s|) \times [0, |t|)$ , so “lexicographically least” commits us to nothing: every total order on a finite set has finite order type, and a graded order would serve equally well — it would generally reserve a different sub-coset, but the proof never uses which one. The steering pair, by contrast, is drawn from an *infinite* subset of  $\mathbb{Z}^2$ , and there the choice of order does matter. Ordered

lexicographically,  $\mathbb{Z}^2$  has order type  $\omega^2$ ; a nonempty subset still has a least element, so the selection is well defined — but locating it may require deciding whether an entire infinite column meets the set, which no terminating search can do. Under a *graded* order — by  $|s| + |t|$ , say, with lexicographic tie-breaking — the order type is  $\omega$ , every pair has only finitely many predecessors, and the least admissible pair is found by running through candidates in that order until one passes the coprimality tests and the strict inequality of Lemma 14. That search halts precisely because the order type is  $\omega$ .

So classical choice here is a convenience, not a necessity: taking the graded-least admissible pair makes the steering rule computable too, and with it the recursion as a whole. We keep the existential form because that is what the argument uses, and no canonical pair is needed for the theorem. (For term orders on  $\mathbb{N}^n$ , which are required in addition to respect addition — a constraint of no use here — the possible order types are exactly  $\omega^1, \dots, \omega^n$ . This is a consequence of the classification of such orders: by Robbiano’s theorem [8] every one of them is given by a real matrix acting on exponent vectors and compared lexicographically, and Robbiano settles as well *when two matrices determine the same order*, so the representation is genuinely understood rather than merely available. Eisenbud puts the classification in the equivalent form that every monomial order is a lexicographic product of at most  $n$  weight orders [7, Exercise 15.13]; the number of factors is then the exponent of the order type. The moral survives in the present looser setting: what a terminating search needs is not compatibility with the group structure, but order type  $\omega$ .)

So the construction is a definite object once an enumeration  $(z_k)$ , a dense sequence  $(\theta_k)$  and a selection of steering pairs are fixed; only the last of these is genuinely arbitrary. Figure 6 shows one such run.

#### 4. PICTURES OF THE CONSTRUCTION

Figure 6 above shows the nested regions  $R_k$  that drive the recursion; this section instead looks at what the construction *achieves*. Figure 7 plots the first 10, 20, and 100 directions realized by an actual run of the construction (code in `verification-code/`, targeting the equidistributed sequence  $\theta_k = k\pi(\sqrt{5} - 1)/2 \bmod \pi$ , an irrational rotation by the golden ratio conjugate), each direction shown as an antipodal pair of points on the unit circle (a line’s direction is unsigned) and colored by its position in the sequence.

*Remark 17* (Sequence order against angular order). The points in Figure 7 are indexed by the stage  $k$  that produced them — that index is what the color ramp shows — and not by where they sit on  $\mathbb{RP}^1$ . The two orders are necessarily different, and not merely incidentally so: an enumeration of a dense subset of  $[0, \pi)$  cannot be monotone. If  $\theta_1 < \theta_2 < \dots$ , then no term at all lies in the nonempty open interval  $(\theta_1, \theta_2)$ , since every later term is at least  $\theta_2$ ; the set of terms therefore misses an open set and is not dense. A decreasing enumeration fails for the same reason. Density thus forces the enumeration to jump about, and an equidistributed target sequence makes it jump about as evenly as it can. The scatter in the figure is the construction’s order type  $\omega$  lying transverse to the geometry of  $\mathbb{RP}^1$ , not an artifact of the particular run.




FIGURE 7. Realized directions after 10, 20, and 100 steps of the construction, colored by order (dark = early, light = late), with a thin path tracing consecutive jumps. The steering argument scatters the directions rather than sweeps them around the circle, since each target angle is chosen from an equidistributed sequence rather than visited in angular order.

As a purely exploratory aside, Figure 8 redraws 1000 realized directions (extending the run behind Figure 7) as point and line masses on the closed unit disc: each direction contributes a dot at its two antipodal boundary points plus its diameter chord, weighted by  $w_k = (k + 5)^{-0.6} e^{-k/250}$ . Unlike a weight such as  $1/k$  or  $1/(k + 5)$ , this one is *summable*, so the resulting measure genuinely converges as  $k \rightarrow \infty$  to a fixed, permanently non-uniform limit — no renormalization needed, and none of the visible structure is a finite- $k$  artifact fading toward the uniform measure on  $\mathbb{RP}^1$ . The exponent and cutoff were chosen so that no small handful of directions dominates the picture: the first 5 directions carry only 9.3% of the total mass, the first 20 carry 26%, the first 100 carry 62%; at  $k = 1000$ , 99.6% of the full  $k \rightarrow \infty$  total is already accounted for, so the figure is an honest rendering of the actual limit rather than an arbitrary truncation of it. Nothing in this manuscript rests on it — see the Acknowledgements for how it came about.

Two further exploratory pictures, again purely illustrative, follow the proof's own mechanism rather than the directions it realizes. The *coverage invariant* established above ((35)–(36)) says that a point  $(u, v) \in \mathbb{Z}^2$  is claimed by exactly one stage  $k$ , namely the unique  $k$  with  $(u, v) \in R_{k-1} \setminus R_k$ ; write  $\text{stage}(u, v) = k$  for that index. The enumeration  $(z_k)$  used throughout the proof to decide reservations can be any enumeration of  $\mathbb{Z}^2$  in which every point has finite index; the run behind Figure 9 takes the concrete choice of ordering points by  $L^\infty$ -distance from the origin, breaking ties lexicographically.

Stage  $k$  claims  $R_{k-1} \setminus R_k$  all at once, via every line  $\ell_{(P_k, Q_k), c}$  of that stage's direction  $(P_k, Q_k)$  needed to cover it (Lemma 13) — so stage conflates many distinct lines into one value. A finer invariant tracks the individual line rather than the stage: enumerate the set  $S = \{\ell_{(P_k, Q_k), c}\}$  of lines actually used by the construction as  $S = \bigcup_j \{s_j\}$ , indexing each line  $s_j = (P, Q, c)$  the first time the same  $L^\infty$  enumeration  $(z_k)$  lands on one of its points, and

Tempered sqrt-harmonic weight  $w_k = (k+5)^{-0.6}e^{-k/250}$  (summable -- a genuine, permanent limiting measure, not a step toward uniform)  
 first 5 points: 9.3% of total mass -- first 20: 26% -- first 100: 62% -- N=1000 already captures 99.6% of the true limit  
 density reading stained-glass reading




FIGURE 8. 1000 realized directions redrawn as point and line masses on the closed unit disc, weighted by the summable  $w_k = (k+5)^{-0.6}e^{-k/250}$  so the picture shows a genuine limiting measure rather than a step toward the uniform measure on  $\mathbb{RP}^1$  (see main text). Left: a smoothed density reading. Right: an order-colored overlay of the same weighted points and chords (dark purple = early, yellow-green = late). Purely exploratory — not used anywhere else in this manuscript.

set

$$\text{hit}(u, v) = \min\{j : (u, v) \in \ell_{s_j}\}.$$

Unlike stage, this has no direct role in the proof — the construction only ever needs the coarser, stage-level disjointness — but it exposes structure the stage-level picture cannot: at  $N = 25$ , 887 distinct lines are used, of which 423 meet the window in exactly 3 points and 270 in exactly 4, so  $\text{hit}$  is a genuinely fine-grained ordinal, plotted in Figure 10 on a continuous scale rather than the discrete palette of Figure 9.

*Remark 18.* Both the splitting mechanism (Lemmas 11–13) and the sign-generality of Lemma 14 were checked by direct brute-force simulation (exhaustive crossing and coverage checks on finite windows up to  $81 \times 81$ , including adversarial diagonal and sign-alternating schedules) before being trusted; a splitting step without the primitivity safeguard built into Lemma 11 was found this way to be genuinely flawed rather than merely under-justified. The full argument was also read independently by a second reader (another Claude agent) working from the statement of the lemmas alone, who found a real gap: restricting  $s, t$  to be positive only reaches half of  $\mathbb{RP}^1$ , and existence of *some* admissible move at each stage does not imply the ability to steer toward an *arbitrary* target — both addressed in the versions of Lemmas 11–13 and 14 given here). A machine-checked (Lean/Mathlib) formalization of this argument has also been completed, as a separate, complementary effort, available at [9]; brute-force simulation and independent reading catch different classes of error than a machine-checked proof does, and vice versa.






FIGURE 9.  $\text{stage}(u, v)$  over a  $51 \times 51$  window ( $N = 25$ ) of the run behind Figure 7. Left: the full window — stage 1 alone (gold) already claims 97.0% of it, since the first direction excludes only a single residue class. Right: stage 1 hidden, revealing that of the three residual columns visible on the left, only the one nearest the origin keeps recursing (through stages 3–7, telescoping inward as  $|y|$  grows), while the other two are fully resolved by stage 2 alone and never revisited — a direct consequence of  $z_k$  being enumerated by distance from the origin, so whichever residual region sits closest is exactly the one reopened next.

## 5. RELATED WORK

The reformulation via  $\varphi_d$  and the resulting rigidity phenomenon (Lemma 8) form the natural two-dimensional analogue of the classical theory of *covering systems of congruences*, introduced by Erdős [3] to show that a positive proportion of the integers are not of the form  $2^k + p$  for  $p$  prime. A very close relative of our covering problem — covering  $\mathbb{Z}^2$  by finitely many finite-index *sublattices* (necessarily containing the origin, unlike the general affine cosets considered here) — is studied by Cremona and Koymans [1], who give it the same “projective/homogeneous covering congruences” framing; their refinement machinery (splitting one lattice into several of prime-power relative index) is structurally reminiscent of the splitting used here, one rank up.

A different classical result addresses a related but distinct question about how many directions a covering needs. Corzatt [2] conjectured that if a *finite, convex* set of lattice points is covered by  $n$  lines (with no constraint on where the lines may cross), the lines can always be chosen to use at most four distinct slopes. Our setting is the opposite regime: an infinite, unbounded point set, with a crossing-avoidance constraint standing in for a bound on the number of lines, and correspondingly the answer is qualitatively different: densely many directions are both necessary to consider and available.

Lattice-line-covers: which individual line reaches each point of  $\mathbb{Z}^2$  first?

Line hitting time ( $N = 25$ ): index of the first individual line  $ax - by = c$  covering each point  
(887 distinct lines used in this window)




FIGURE 10.  $\text{hit}(u, v)$  over the same  $51 \times 51$  window. The concentric *square* (not circular) rings are the  $L^\infty$  enumeration made visible. The three low-hit blobs aligned at  $x = -15, 0, 15$  are a single mechanism repeated: stage 1's shallow direction  $(15, -2)$  places three window-points of the *same* line 15 apart in  $x$  (for instance  $(0, 2)$ ,  $(-15, 4)$ ,  $(15, 0)$  all lie on  $\ell_{(15, -2), -30}$ ); since  $\text{hit}$  is assigned at a line's first discovery and one representative always sits near the origin, its far echoes inherit the same tiny index for free, landing at nearly the same  $|y|$  because the direction is so shallow. The thin seam exactly at  $x = 0$  is a different mechanism: those points are almost always the sole window-representative of a deep-recursion line (coefficients up to  $P \approx 2.6 \times 10^6$  at  $N = 25$ ), so  $\text{hit}$  there tracks  $|y|$  alone rather than  $\max(|x|, |y|)$ , a slower growth law that reads as a seam cutting through the square rings.

The unimodular pairs forbidden by Lemma 8 are exactly the Farey-neighbor pairs of slopes; see e.g. Hardy and Wright [4, Ch. III] for the classical theory of Farey sequences and the closely related Stern–Brocot tree. This is precisely the toolkit used in the digital-geometry literature on digitized straight lines, where continued-fraction and Farey-tree techniques describe digital approximations to a line of given (rational or irrational) slope; see Kiselman [5], and, for the continued-fraction approach specifically, Uscka-Wehlou’s dissertation [6].

*Open problem 19.* Is it possible to generalize Theorem 7 to  $\mathbb{Z}^n$  for  $n \geq 3$ ? In this formulation, one would study configurations of “lattice hyperplanes” or “lattice lines”, such that no two different directions meet at a lattice point, and determine which sets of directions can occur.

#### ACKNOWLEDGEMENTS

This problem occurred to the author during an idle conversation with colleagues F.B. and T.E. (in the department coffee room) about covering the lattice  $\mathbb{Z}^2$  with lines. It was considered as a useful test for LLM-assisted proof discovery and proof formalization, since the author conjectured that the proof of the result, if it was indeed true, should only use elementary methods, which are already in lean (Mathlib).

Figure 8 was inspired by a remark — certainly misunderstood by the author — about a relation to Besicovitch–Orponen theory, made by his colleague B.E.; the keyword “measure” stuck.

The results in this manuscript: the reformulation, the rigidity and coset lemmas, the recursive construction, and both rounds of error-correction (one caught by numerical simulation, one by an independent review), were found by Claude (Anthropic), working with the author in an interactive session. The additional references collected in the repository’s extended-reference list [9] were located and verified by Claude via web search. A list of “certificates” showing how each reference is used in this manuscript (this “certificate of non-hallucination”), together with a more detailed explanation of the whole process of producing this manuscript, from inception to exploration to proof discovery to proof formalization and verification, are described in the gitlab repository [9]; the formalization and verification code is also archived, with a citable DOI, on Zenodo [11].

This manuscript was likewise produced by the author and Claude jointly, with the author mainly directing Claude toward clearer exposition for human readers: restructuring the proofs into short, explicitly labeled lemmas, and breaking up long equations for readability. The author then checked the final manuscript to the best of his ability, made final edits, and assumes full responsibility for any errors, mistakes, or gaps that remain.

#### FUNDING

This research received no external funding. It was carried out as part of the author’s regular duties at Linköping University, within the fraction of that employment allocated to research, the remainder being teaching and other duties.

## CONFLICTS OF INTEREST

The author declares no financial or non-financial conflicts of interest. The nature and extent of LLM assistance in producing this manuscript is disclosed in full in the Acknowledgements above.

## REFERENCES

- [1] J. E. Cremona and P. Koymans, *Lattice Coverings and Homogeneous Covering Congruences*, arXiv:2601.03212v2 (2026).
- [2] C. E. Corzatt, *Covering convex sets of lattice points with straight lines*, *Congressus Numerantium* **50** (1985), 129–135.
- [3] P. Erdős, *On integers of the form  $2^k + p$  and some related problems*, *Summa Brasil. Math.* **2** (1950), 113–123.
- [4] G. H. Hardy and E. M. Wright, *An Introduction to the Theory of Numbers*, 6th ed., revised by D. R. Heath-Brown and J. H. Silverman, Oxford University Press, Oxford, 2008.
- [5] C. O. Kiselman, *Elements of Digital Geometry, Mathematical Morphology, and Discrete Optimization*, World Scientific, Singapore, 2022.
- [6] H. Uscka-Wehlou, *Digital Lines, Sturmian Words, and Continued Fractions*, Ph.D. thesis, Uppsala Dissertations in Mathematics **65**, Uppsala University, 2009.
- [7] D. Eisenbud, *Commutative Algebra with a View Toward Algebraic Geometry*, Graduate Texts in Mathematics **150**, Springer-Verlag, New York, 1995.
- [8] L. Robbiano, *Term orderings on the polynomial ring*, in: *Proceedings of EUROCAL '85*, Vol. 2 (Linz, 1985), *Lecture Notes in Computer Science* **204**, Springer, Berlin, 1985, pp. 513–517.
- [9] J. Snellman and Claude (Anthropic), *lattice-line-covers* (source repository, including the Lean/Mathlib formalization and reference certificates), 2026, <https://gitlab.liu.se/jansn19/lattice-line-covers>.
- [10] J. Snellman, *On the directions occurring in lattice-line coverings of the integer plane*, LiTH-MAT-R, ISSN 0348-2960, No. 2026:02, Linköping University Electronic Press, 2026. doi:10.3384/LiTH-MAT-R-2026-02. Open access via DiVA, urn:nbn:se:liu:diva-226766.
- [11] J. Snellman, *lattice-line-covers: Lean/Mathlib formalization and verification code*, Zenodo, Linköping University community, 2026. Concept DOI (all versions) doi:10.5281/zenodo.21916621; this snapshot doi:10.5281/zenodo.21916622.