



# Change of basis

---


## Linear Algebra

Department of Computer Engineering

Sharif University of Technology

Hamid R. Rabiee [rabiee@sharif.edu](mailto:rabiee@sharif.edu)

Maryam Ramezani [maryam.ramezani@sharif.edu](mailto:maryam.ramezani@sharif.edu)



# Basis Review

---



## Example

- Find the coordinate vector of  $2 + 7x + x^2 \in \mathcal{P}^2$  with respect to the basis  $B = \{x + x^2, 1 + x^2, 1 + x\}$ .
- If  $C = \{1, x, x^2\}$  is the standard basis of  $\mathcal{P}^2$  then we have  $[2 + 7x + x^2]_C = (2, 7, 1)$ .



We want to find scalars  $c_1, c_2, c_3 \in \mathbb{R}$  such that

$$2 + 7x + x^2 = c_1(x + x^2) + c_2(1 + x^2) + c_3(1 + x).$$

By matching coefficients of powers of  $x$  on the left-hand and right-hand sides above, we arrive at following system of linear equations:

$$c_2 + c_3 = 2$$

$$c_1 + c_3 = 7$$

$$c_1 + c_2 = 1$$

This linear system has  $c_1 = 3, c_2 = -2, c_3 = 4$  as its unique solution, so our desired coordinate vector is

$$[2 + 7x + x^2] = (c_1, c_2, c_3) = (3, -2, 4)$$

# Change of Basis

---



□  $B = \{v_1, \dots, v_n\}$  are basis of  $\mathbb{R}^n$ .

□  $P = [v_1 \ v_2 \ \dots \ v_n]$

□  $P[a]_B = a$



## Theorem


Let  $B = \{b_1, b_2, \dots, b_n\}$  and  $C = \{c_1, c_2, \dots, c_n\}$  be bases of a vector space  $V$ . Then there is a unique  $n \times n$  matrix  $P_{C \leftarrow B}$  such that

$$[x]_C = P_{C \leftarrow B} [x]_B$$

The columns of  $P_{C \leftarrow B}$  are the  $C$ -coordinate vectors of the vectors in basis  $B$ .

That is ,

$$P_{C \leftarrow B} = [[b_1]_C \quad [b_2]_C \quad \dots \quad [b_n]_C]$$



$$(P_{C \leftarrow B})^{-1} = P_{B \leftarrow C}$$

$$P_B[x]_B = x, \quad P_C[x]_C = x, \quad \text{and} \quad [x]_C = P_C^{-1}x$$

$$[x]_C = P_C^{-1}x = P_C^{-1}P_B[x]_B$$





## Example

Find the change-of-basis matrices  $P_{C \leftarrow B}$  and  $P_{B \leftarrow C}$  for the bases  $B = \{x + x^2, 1 + x^2, 1 + x\}$  and  $C = \{1, x, x^2\}$  of  $\mathcal{P}^2$ . Then find the coordinate vector of  $2 + 7x + x^2$  with respect to B.



## Example

Let  $b_1 = \begin{bmatrix} 1 \\ -3 \end{bmatrix}$ ,  $b_2 = \begin{bmatrix} -2 \\ 4 \end{bmatrix}$ ,  $c_1 = \begin{bmatrix} -7 \\ 9 \end{bmatrix}$ ,  $c_2 = \begin{bmatrix} -5 \\ 7 \end{bmatrix}$ , the bases for  $\mathbb{R}^2$  given by  $B = \{b_1, b_2\}$ ,  $C = \{c_1, c_2\}$ .

- Find the change-of-coordinates matrix from  $C$  to  $B$ .
- Find the change-of-coordinates matrix from  $B$  to  $C$ .



## Example

Find the change-of-basis matrix  $P_{C \leftarrow B}$ , where

$$B = \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \right\}, C = \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \right\}.$$

# Linear Transformation and Change of Basis

---



## Example

We have  $B = \{x^3, x^2, x, 1\}$  and  $B' = \{x^2, x, 1\}$  are bases for  $\mathcal{P}_3(x)$  and  $\mathcal{P}_2(x)$ , respectively. Find the matrix of transformation  $T: \mathcal{P}_3(x) \rightarrow \mathcal{P}_2(x)$ .



Since  $\begin{bmatrix} a_3 \\ a_2 \\ a_1 \\ a_0 \end{bmatrix}$  the vector representation of  $a_3x^3 + a_2x^2 + a_1x + a_0 \in \mathcal{P}^3(x)$ , we have

$$\begin{aligned} \left[ \frac{d}{dt} \right]_{\{B, B'\}} &= \begin{bmatrix} \frac{d}{dt}(x^3) & \frac{d}{dt}(x^2) & \frac{d}{dt}(x) & \frac{d}{dt}(1) \end{bmatrix} \\ &= \begin{bmatrix} 3 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix} \end{aligned}$$

# Matrix representation of linear function




## Important

Let  $T: \mathbb{R}_n \rightarrow \mathbb{R}_m$  be a linear function and  $u = \begin{bmatrix} u_1 \\ \vdots \\ u_n \end{bmatrix} \in \mathbb{R}_n$ .

The matrix  $[A(e_1) \ \dots \ A(e_n)]$  is called the matrix representation of linear function (transformation)  $T$  which is denoted by  $[A]_E$ .




What is the relation between  $[A]_B$  and  $[A]_E$ ?



- $B = \{v_1, v_2, \dots, v_n\}$  are basis of  $\mathbb{R}^n$ .
- $P = [v_1 \ v_2 \ \dots \ v_n]$
- $[T(x)]_B = P^{-1}AP[x]_B$





$$[A]_B = P^{-1}[A]_E P$$



## Definition

Suppose  $V$  and  $W$  are vector spaces over the same field. We say that  $V$  and  $W$  are **isomorphic**, denoted by  $V \cong W$ , if there exists an invertible linear transformation  $T: V \rightarrow W$  (called an **isomorphism** from  $V$  to  $W$ ).

- If  $T: V \rightarrow W$  is an isomorphism then so is  $T^{-1}: W \rightarrow V$ .
- If  $T: V \rightarrow W$  and  $S: W \rightarrow X$  are isomorphism then so is  $S \circ T: V \rightarrow X$ .  
in particular, if  $V \cong W$  and  $W \cong X$  then  $V \cong X$ .

## Example

Show that the vector space  $V = \text{span}(e^x, xe^x, x^2e^x)$  and  $\mathbb{R}^3$  are isomorphic.



The standard way to show that two space are isomorphic is to construct an isomorphism between them. To this end, consider the linear transformation  $T: \mathbb{R}^3 \rightarrow V$  defined by

$$T(a, b, c) = ae^x + bxe^x + cx^2e^x.$$

It is straightforward to show that this function is linear transformation, so we just need to convince ourselves that it is invertible. We can construct the standard matrix  $[T]_{B \leftarrow E}$ , where  $E = \{e_1, e_2, e_3\}$  is the standard basis of  $\mathbb{R}^3$ :

$$\begin{aligned} [T]_{B \leftarrow E} &= [[T(1, 0, 0)]_B, [T(0, 1, 0)]_B, [T(0, 0, 1)]_B] \\ &= [[e^x]_B, [xe^x]_B, [x^2e^x]_B] = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \end{aligned}$$

Since  $[T]_{B \leftarrow E}$  is clearly invertible (the identity matrix is its own inverse),  $T$  is invertible too and is thus an isomorphism.



- ❑ Chapter 1: Advanced Linear and Matrix Algebra, Nathaniel Johnston
- ❑ Chapter 6: Linear Algebra David Cherney
- ❑ Linear Algebra and Optimization for Machine Learning
- ❑ Introduction to Applied Linear Algebra Vectors, Matrices, and Least Squares