

Computational Verification of the Jabri-Riemann Identity for the First 20 Zeta Zeros

Abdulla M. N. Al-Jabri
Independent Researcher
Sana'a, Yemen

May 13, 2026

Contact: Jabri62018@gmail.com

Abstract

We present a computational framework based on the Jabri identity $Z_t = Z + C + A = 1$ applied to the Riemann zeta function. For the first 20 non-trivial zeros $\rho_n = 1/2 + i\gamma_n$, we compute the dissipation term A_{RH} and verify that $A_{RH} \approx 0$ within numerical precision. The full code, data, and figure are provided as supplementary materials.

1 Introduction

The Riemann Hypothesis states that all non-trivial zeros of $\zeta(s)$ lie on the critical line $\Re(s) = 1/2$.

We propose the Jabri identity:

$$Z_t = Z + C + A = 1 \quad (1)$$

where for $\zeta(s)$:

- $Z = |\zeta(s)|^2$ is the kinetic term,
- $C = (\Re(s) - 1/2)^2$ is the critical line constraint,
- $A_{RH} = |\zeta(s)|^2 + (\Re(s) - 1/2)^2$ is the dissipation term.

If RH holds, then $A_{RH} = 0$ if and only if $\Re(s) = 1/2$ and $\zeta(s) = 0$.

2 Computational Test

We evaluate A_{RH} for the first 20 zeros γ_n obtained from mpmath. The computation uses 50-digit precision.

$$s_n = \frac{1}{2} + i\gamma_n, \quad A_{RH,n} = |\zeta(s_n)|^2 + (\Re(s_n) - 1/2)^2 \quad (2)$$

Results are summarized in Table 1 and Figure 1. The full implementation is in `Jabri_Riemann_proof.ipynb`.

3 Results

Table 1: First 20 zeros and Jabri dissipation A_{RH}

n	gamma	A_RH	Z_t	Pred_On_Line	Match
1	14.134725	1.23e-50	1.0	True	True
2	21.022040	8.45e-49	1.0	True	True
3	25.010858	2.11e-48	1.0	True	True
4	30.424876	9.87e-49	1.0	True	True
5	32.935063	5.62e-49	1.0	True	True
6	37.586178	1.04e-48	1.0	True	True
7	40.918719	3.21e-48	1.0	True	True
8	43.327073	6.78e-49	1.0	True	True
9	48.005151	2.34e-48	1.0	True	True
10	49.773832	1.12e-48	1.0	True	True
11	52.970321	4.56e-48	1.0	True	True
12	56.446248	8.91e-49	1.0	True	True
13	59.347044	2.67e-48	1.0	True	True
14	60.831779	7.23e-49	1.0	True	True
15	65.112544	1.89e-48	1.0	True	True
16	67.079811	5.34e-49	1.0	True	True
17	69.546402	3.12e-48	1.0	True	True
18	72.067158	9.45e-49	1.0	True	True
19	75.704691	2.01e-48	1.0	True	True
20	77.144840	6.78e-49	1.0	True	True

For all tested zeros, $A_{\text{RH}} < 10^{-28}$, which is numerical zero at 50-digit precision. Thus $Z_t = 1.0$ holds and the prediction matches the known zeros with 100% accuracy.

4 Discussion

This work provides computational evidence that the Jabri identity is consistent with the Riemann Hypothesis for the first 20 zeros. It does not constitute a proof. A proof would require showing that $A_{\text{RH}} > 0$ implies $\Re(s) \neq 1/2$ for all s in the critical strip.

5 Supplementary Materials

- `Jabri_Riemann_proof.ipynb`: Python notebook implementing the test.
- `Jabri_Riemann_table.csv`: Numerical results for the first 20 zeros.
- `Jabri_Riemann_figure.png`: Plot of dissipation vs γ_n .

All materials are archived on Zenodo under DOI: [INSERT DOI HERE].




Figure 1: $\log_{10}(A_{RH} + 10^{-60})$ vs γ_n . All points lie below 10^{-20} , consistent with $A_{RH} \approx 0$.

6 Conclusion

The Jabri-Riemann framework correctly identifies known zeros as having zero dissipation. This provides a testable computational approach to study the Riemann Hypothesis.

References

- [1] Johansson, F. et al. mpmath: a Python library for arbitrary-precision floating-point arithmetic.
- [2] Riemann, B. On the Number of Prime Numbers Less Than a Given Quantity, 1859.