

Stiff Differential Equations Supplementary Material

Tomas Mathews 40328403, Jacob Simpson 40331806

February 9, 2024

Recall equation (1) from the report.

$$\frac{dx}{dt} = -30x \quad (1)$$

with initial condition $x(0) = 1$

To solve such an equation, we can use the method of **variable separable**.

Multiplying both sides of 1 by dt and dividing by x yields.

$$\frac{dx}{x} = -30dt$$

Integrating over both sides then gives us that:

$$\ln(x) = -30t + c \implies x(t) = e^{-30t+c} = x(t) = Ce^{-30t}$$

Using our initial condition, we get that.

$$x(0) = 1 = Ce^{-30(0)} = C$$

Hence, we get equation (3) from the report ie:

$$x(t) = e^{-30t} \quad (2)$$

Now, recall the more complicated equation (2) from the report.

$$\frac{dx}{dt} = -20x + 20 \sin(\omega t) + \omega \cos(\omega t) \quad (3)$$

To solve this equation we will use the method of the **Integrating Factor** Firstly, Rearranging equation (2) gives us:

$$\frac{dx}{dt} + 20x = 20 \sin(\omega t) + \omega \cos(\omega t)$$

Let $\mu(t) = e^{\int 20dt}$ and multiply both sides by $\mu(t)$ gives:

$$e^{20t} \frac{dx}{dt} + 20e^{20t}x = -e^{20t}(-20 \sin(\omega t) - \omega \cos(\omega t))$$

Since $20e^{20t} = \frac{d}{dt}e^{20t}$, we get that:

$$e^{20t} \frac{dx}{dt} + \frac{d}{dt}(e^{20t})x = -e^{20t}(-20 \sin(\omega t) - \omega \cos(\omega t))$$

Applying the reverse product rule, we get:

$$\frac{d}{dt}(xe^{20t}) = -e^{20t}(-20 \sin(\omega t) - \omega \cos(\omega t))$$

Integrating both sides with respect to t , we get that:

$$x(t)e^{20t} = e^{20t} \sin(\omega t) + C$$

Finally, dividing both sides by $\mu(t)$ we get:

$$x(t) = \sin(\omega t) + Ce^{-20t}$$

Lastly, using are initial conditions $x(0) = 1$, we get that $C = 1$ and hence we obtain equation (4):

$$x(t) = \sin(\omega t) + e^{-20t} \quad (4)$$

Below, we can see the two plots for equation 3 and 4.




Figure 1: The plot above shows the solution to equation 1.




Figure 2: The plot above shows the solution to equation 2