

Notes for ECE 369 - Discrete Mathematics for Computer Engineering

Ezekiel Ulrich

September 7, 2023

These are lecture notes for fall 2023 ECE 36900 at Purdue. Modify, use, and distribute as you please.

Contents

Course Introduction	1
Equations	1
Propositional Logic	3
Rules and proofs	4
Predicate logic	7
Proofs	9

Course Introduction

This course introduces discrete mathematical structures and finite-state machines. Students will learn how to use logical and mathematical formalisms to formulate and solve problems in computer engineering. Topics include formal logic, proof techniques, recurrence relations, sets, combinatorics, relations, functions, algebraic structures, and finite-state machines. For more information, see the syllabus.

Equations

1. De Morgan's Theorem:

$$\neg(P \wedge Q) \equiv \neg P \vee \neg Q$$

$$\neg(P \vee Q) \equiv \neg P \wedge \neg Q$$

2. Modus ponens (mp)

$$p$$

$$p \rightarrow q$$

$$\therefore q$$

3. Modus tonens (mt)

$$p \rightarrow q$$

$$\neg q$$

$$\therefore \neg p$$

Propositional Logic

We often wish that others would be more logical, tell the truth, or shower. While studying formal logic cannot help with the latter (in fact, studies have shown a negative correlation between hygiene and studying formal logic) it is a useful way to define what the first mean two. In a formal logic model, we have two constructs:

- **Statements/proposition:** A statement or a proposition is a sentence that is either true or false. Propositions are often represented with letters of the alphabet. For example: " q : the more time you spend coding, the less time you have to buy deodorant."
- **Logical connectives:** Used to connect statements. For example, "and" is a logical connective in English. It can be used to connect two statements, e.g. "the person next to me smells like dog *and* looks like a dog" to obtain a new statement with its own truth value.

Here are common logical connectives in Boolean logic:

Logical Connective	Symbol
Negation (NOT)	\neg or $'$
Conjunction (AND)	\wedge
Disjunction (OR)	\vee
Exclusive OR (XOR)	\oplus
Implication	\rightarrow
Biconditional	\leftrightarrow

Table 1: Logical Connectives in Boolean Logic

Truth table: Defines how each of the connectives operate on truth values. Every connective has one. For example, consider \wedge AND:

p	q	$p \wedge q$
T	T	T
T	F	F
F	T	F
F	F	F

Table 2: Truth table for \wedge

We see that p AND q is only true when p is true and q is true. Similarly, p or q is only true when p is true or q is true (or both). An important connective for discovering new truths is the implication \rightarrow , which basically says "if the first letter is true, then so is the second". Let p : "I live in Wiley" and q : "I have no AC". In English, the statement $p \rightarrow q$ would be stated as "If I live in Wiley, then I have no AC".

Table 3 shows the truth table for \rightarrow . It may not seem immediately clear why, for instance, if p and q are false, then $p \rightarrow q$ is true. If we

p	q	$p \rightarrow q$
T	T	T
T	F	F
F	T	T
F	F	T

Table 3: Truth table for \rightarrow

consider what this means in English, then all we know is that I don't live in Wiley. Perhaps I live in Tarkington and still don't have AC, or perhaps I live in Honors and I do. In any case the first letter isn't true, so "if the first letter is true then so is the second" stands as true. If we have the statement $p \rightarrow q$, then we call q a *necessary condition* for p .

Conversely, p is a *sufficient condition* for q .

Say we have a statement such as $\neg A \vee B \rightarrow C$. This is ambiguous, since we can interpret it as either $(A \vee B) \rightarrow C$ or $A \vee (B \rightarrow C)$. The truth tables will differ in each case, so it becomes necessary to specify in what order we should apply logical connectives.

1. Parentheses "()"
2. Negation " \neg "
3. AND and OR " \wedge, \vee "
4. Implication " \rightarrow "
5. Biconditional " \leftrightarrow "

Rules and proofs

With each additional variable in your truth table, the number of choices grows exponentially. Specifically, if you have n statement letters, you would have 2^n choices for your truth table.

Tautology: A formula that is true in every model. Example: the Bible is infallible because the Bible itself says it is infallible.

Contradiction: A formula that is false in every model Examples: "it is raining and it is not raining", "I am sleeping and I am awake", "IU is a good school".

Confusion often arises when negating a sentence such as "the book is thick and boring". A natural inclination is to negate it thus: "the book is not thick and not boring". However, consider the truth table for this: p : "the book is thick", q : "the book is boring". We can see the last two rows are not identical, therefore the negation of "the book is not thick and not boring" is not "the book is not thick and not boring". For p to be false, either the book must not be thick *or* the book must not be boring. This is summarized by **De Morgan's Theorem**:

$$\neg(P \wedge Q) \equiv \neg P \vee \neg Q$$

Interestingly, it is possible to prove any statement in a system where a contradiction exists. This is known as the *principle of explosion*. To see how it works, consider the following example:

1. p : Donuts are good for you.
2. q : Unicorns exist.

I'll now assume the contradictory statement "donuts are good for you and donuts are not good for you".

$$\neg p \wedge p \quad (\text{Contradiction})$$

$$p \vee q \quad (\text{Addition})$$

$$\neg p$$

$$\therefore q$$

Ergo, unicorns exist :)

p	q	$p \wedge q$	$\neg(p \wedge q)$	$\neg p \wedge \neg q$
T	T	T	F	F
T	F	F	T	F
F	T	F	T	F
F	F	F	T	T

$$\neg(P \vee Q) \equiv \neg P \wedge \neg Q$$

We now have a sufficient understanding of truth tables and logical connectives to come up with some useful rules. First of these are

Modus ponens (mp):

$$\begin{array}{l} p \\ p \rightarrow q \\ \hline \therefore q \end{array}$$

and

Modus tollens (mt):

$$\begin{array}{l} p \rightarrow q \\ \neg q \\ \hline \therefore \neg p \end{array}$$

Below are two tables for commonly used rules.

Expression	Equivalent to	Name - abbreviation
$p \vee q$ $p \wedge q$	$q \vee p$ $q \wedge p$	Commutative - comm
$(p \vee q) \vee r$ $(p \wedge q) \wedge r$	$p \vee (q \vee r)$ $p \wedge (q \wedge r)$	Associative - ass
$\neg(p \wedge q)$ $\neg(p \vee q)$	$\neg p \vee \neg q$ $\neg p \wedge \neg q$	De Morgan's Laws - De Morgan
$p \rightarrow q$	$\neg p \vee q$	Implication - imp
p	$\neg(\neg p)$	Double negation - dn
$p \leftrightarrow q$	$(p \rightarrow q) \wedge (q \rightarrow p)$	Def'n of equivalence - equ

Table 4: Equivalence rules

At this point, let us formally define an

Argument: An argument can be symbolized as

$$P_1 \vee P_2 \vee P_3 \vee \dots \vee P_n \rightarrow Q$$

where P_i is called a hypothesis and Q is the conclusion. If this statement is a tautology, then the argument is *valid*. There are multiple ways we could prove a given argument is a tautology. For instance, we

From	Can derive	Name - abbreviation
$p, p \rightarrow q$	q	Modus ponens - mp
$p \rightarrow q, \neg q$	$\neg p$	Modus tollens - mt
p, q	$p \vee q$	Conjunction - con
$p \vee q, \neg p$	q	Disjunction - dis
$p \wedge q$	p, q	Simplification - sim
p	$p \vee q$	Addition - add

Table 5: Inference rules

could create a truth table and brute force an answer. However, with even four hypotheses this process is tedious, and with each additional hypothesis it becomes exponentially harder. Therefore we instead often turn to the

Proof sequence: a sequence of well-formed formulas in which each formula is either a hypothesis or the result of applying a derivation rule to earlier well-formed formulas. In practice this looks like

$$\begin{array}{l}
 P_1 \text{ (hypothesis)} \\
 P_2 \text{ (hypothesis)} \\
 P_3 \text{ (hypothesis)} \\
 \dots \\
 P_n \text{ (hypothesis)} \\
 \text{(formula) 1 (obtained from derivation rule)} \\
 \text{(formula) 2 (obtained from derivation rule)} \\
 \dots \\
 \text{(formula) } n \text{ (obtained from derivation rule)} \\
 \therefore Q
 \end{array}$$

Let's use all this new information in a simple proof.

$$\begin{array}{l}
 A \text{ (hypothesis)} \\
 A \rightarrow B \text{ (hypothesis)} \\
 B \rightarrow C \text{ (hypothesis)} \\
 B \text{ (1, 2, mp)} \\
 C \text{ (4, 3, mp)} \\
 \hline
 \therefore C
 \end{array}$$

If we wish to apply our knowledge of logic to the real world, some practice in translating natural language to formal logic is necessary. Let's test it with this statement: "If chicken is on the menu, then don't order fish, but you should have either fish or salad. So if chicken is on the menu, have salad." Let C: "Chicken is on the menu", F: "You order fish", and S: "You have salad". We know that if chicken is on the

menu you don't order fish, that you should have either fish or salad, and we'd like to show that if chicken is on the menu you should have salad.

C	(hypothesis)
$C \rightarrow \neg F$	(hypothesis)
$F \vee S$	(hypothesis)
$\neg F$	(1, 2, mp)
S	(3, 4, dis)
<hr/>	
$\therefore S$	

Predicate logic

Predicate logic: Capable of making statements about entire groups instead of individual letters. In predicate logic, propositions are expressed in terms of predicates, variables and quantifiers, the latter of which propositional logic lacks.

Quantifier: How many objects have a certain property: "for every" or "for some".

Predicate: Property that a variable may have.

Domain of interpretation: Collection of objects from which the variable is taken.

Universal quantifier: "For all": \forall . States that a certain property holds for all objects in a domain.

Existential quantifier: "There exists": \exists . States that a certain property holds for at least one object in a domain.

As an example of a predicate well-formed formula: $(\forall x)[(\exists y)x > y]$. We would read this statement as "for all x there exists a y such that $x > y$." At first glance it may seem obvious that this statement is true, but consider the domain. What if the domain is all natural numbers? Then we could let $x = 1$ (or zero depending on your definition of natural numbers) and there would be no corresponding lesser y . We can see from this example that the truth value of a predicate logic formula depends on the domain as well as quantifiers and predicates.

Just as with propositional logic, we often need to translate English statements into predicate logic. Take the statement "every movie made by George Lucas is great". We can rephrase this as "for any movie, if the movie is made by George Lucas, it is great". We would write this formula as

$$(\forall x)(GL(x) \rightarrow Great(x))$$

(Author's note: no value judgement is associated with this English statement).

Let's examine some rules in predicate logic. First, negation:

$$\neg[\forall x A(x)] \leftrightarrow (\exists x) \neg A(x)$$

Some rules from propositional logic still apply in predicate logic. Take modus ponens as an example:

$$\begin{array}{l} (\forall x)(\forall y)L(x,y) \rightarrow [(\exists x)H(x)] \quad \text{(hypothesis)} \\ \neg[(\exists x)H(x)] \quad \text{(hypothesis)} \\ \neg[(\forall x)(\forall y)L(x,y)] \quad (1, 2, \text{mt}) \\ (\exists x)(\exists y)\neg L(x,y) \quad (3, \text{DM}) \\ \hline \therefore (\exists x)(\exists y)\neg L(x,y) \end{array}$$

Table 6 holds predicate inference rules. These rules hold given certain

Name	Abrv.	Given	Can conclude
Existential generalization	eg	$P(a)$	$(\exists x)P(x)$
Existential instantiation	ei	$(\exists x)P(x)$	$P(a)$
Universal generalization	ug	$P(x)$	$(\forall x)P(x)$
Universal instantiation	ui	$(\forall x)P(x)$	$P(a)$

Table 6: Predicate inference rules

conditions. Namely:

(eg) x not in $P(a)$

(ei) Must be the first rule that introduces a

(ug) $P(x)$ not derived from a hypothesis with x as a free variable, and $P(x)$ is not derived by ei from wff with x as a free variable.

(ui) a is a constant.

Let's see some of these rules in action by with a predicate logic proof. Say we have the statement "every ECE student works harder than somebody, and everyone who works harder than any other person gets less sleep than that person. Maria is an ECE student. Ergo, Maria gets less sleep than someone. Let $E(x)$: " x is an ECE student", $W(x,y)$: " x works harder than y ", $S(x,y)$: " x gets less sleep than y ", and m :

A *free variable* is a variable not bound by a quantifier. For example, in the formula

$$(\forall x)(\forall y)P(x,y)$$

both x and y are bound by quantifiers. Constrast this with the formula

$$(\exists x)(\forall y)q(x,y,z)$$

In this example, z is a free variable, since it is not associated with any quantifiers.

Maria. We want to prove $\exists a(S(m, a))$.

$\forall x, E(x) \rightarrow (\exists y)(W(x, y))$ (hypothesis)

$\forall x, \forall y(W(x, y) \rightarrow S(x, y))$ (hypothesis)

$E(m)$ (hypothesis)

$\exists y(E(m) \rightarrow S(m, y))$ (1, ui)

$E(m) \rightarrow W(m, a)$ (4, ei)

$\forall y(W(m, y) \rightarrow S(m, y))$ (3, ui)

$W(m, a) \rightarrow S(m, a)$ (6, ui)

$W(m, a)$ (3,5 mp)

$S(m, a)$ (7,8, mp)

$\exists a(S(m, a))$ (9, eg)

$\therefore \exists a(S(m, a))$

Proofs

List of common proof techniques:

1. Exhaustive proof: In this kind of proof, the statement to be proved is split into a finite number of cases or sets of equivalent cases, and where each type of case is checked to see if the proposition in question holds
2. Refuting by counter-example: If we have a universal statement such as $\forall x(P(x) \rightarrow Q(x))$, we may show it to be false by finding a single a such that $\neg P(a)$.
3. Proof by deduction: Trying to prove $p \rightarrow q$, start by assuming p and then show q .
4. Proof by contraposition: The contrapositive of $p \rightarrow q$ is $\neg q \rightarrow \neg p$. A statement and its contrapositive are logically equivalent, so if proving $p \rightarrow q$ is too difficult we may try to prove $\neg q \rightarrow \neg p$ instead.
5. Proof by contradiction: Suppose I have to prove $p \rightarrow q$. I can begin by saying p is true and $\neg q$ is true. If by a series of steps I arrive a contradiction, then I may say p implies q .
6. Proof by induction: employs a neat trick which allows you to prove a statement about an arbitrary number n by first proving it is true when $n = 1$ (or some other base case), assuming it is true for $n = k$, and then showing it is true for $n = k + 1$.

Below are some example proofs using each of these techniques.

1. Proof by deduction: say we wish to prove the product of two even integers is even. We first need to translate this English sentence to a mathematical statement, which we can do in this case like so:

$$x = 2a, a \in \mathbb{Z}, y = 2b, b \in \mathbb{Z} \rightarrow x \times y = 2c, c \in \mathbb{Z}$$

Our proof is below.

$$\begin{aligned} x &= 2a, a \in \mathbb{Z} \\ y &= 2b, b \in \mathbb{Z} \\ z &= x \times y \\ &= 2a \times 2b \\ &= 2(2ab) \\ &= 2c, c \in \mathbb{Z} \quad \square \end{aligned}$$

Since c is an integer, $2c$ is even and the proof is complete. Try proving the product of two odds is odd in a similar fashion.

2. Proof by contradiction: say we wish to prove $\sqrt{2}$ is irrational. In a theme that will become common as we see more proofs by contradiction, assume the opposite. That is, assume $\sqrt{2}$ is rational. By the definition of rational, we can then write

$$\sqrt{2} = \frac{a}{b}, a, b \in \mathbb{Z}$$

Where a and b share no common factors. We can then perform the following series of steps.

$$\begin{aligned} \sqrt{2} &= \frac{a}{b} \\ b\sqrt{2} &= a \\ 2b^2 &= a^2 \end{aligned}$$

This means that a^2 is even. It can be easily shown that if a^2 is even then a is even. That means that a^2 will actually be divisible by 4. We can rearrange to get

$$\begin{aligned} 2b^2 &= 4c, c \in \mathbb{Z} \\ b^2 &= 2c \end{aligned}$$

So b is likewise even. But if both a and b are even, then they share a common factor and our original supposition is false. Ergo $\sqrt{2}$ is irrational. \square .