

Notes for ECE 30200 - Probabilistic Methods in Electrical and Computer Engineering

Shubham Saluja Kumar Agarwal

January 16, 2025

These are lecture notes for Fall 2025 ECE 30200 by professor Mary Comer at Purdue. Modify, use, and distribute as you please.

Contents

Modelling Random Experiments 2

Set Theory 2

Special Sets 3

Set Operations 3

Set Algebra 4

Modelling Random Experiments

Examples:

- Flip a coin
- Rolling dice
- Generate a bit from a random binary source
- Generate a sequence of n bits from a random binary source
- Count packets arriving at a router
- Measure the voltage at a point in a circuit

All of these examples could be at a fixed time, or over an interval of time.

There is a very precise framework that can be used to model any random experiment.

Overview:

- The outcome that occurs each time a random experiment is run is not known in advance, but **the set of all possible outcomes is assumed to be known**.
- Subsets of the set of all possible outcomes are called events.
- Probabilities are assigned to events, not outcomes, using a probability measure.

We need to use set theory to work with these sets.

Set Theory

A set is an **unordered** collection of elements denoted by $\{\}$.

Example:

$$\{1, 2, 3\} = \{3, 2, 1\} = \{2, 1, 3, 1\}$$

Notation:

- $w \in A$ means w is in set A
- $w \notin A$ means w is not in set A

There are two ways to specify a set:

1. Comma-separated list:

- $A = \{1, 3, A, C\}$

- $A = \{x_1, x_2, \dots, x_n\}$ (that is, n elements for some finite $n \geq 1$ and the i th element is x_i)
- $A = \{x_1, x_2, \dots\}$ (an infinite number of elements)

2. Rule for membership or set builder notation:

- $A = \{w \in \mathbb{Z} : 1 \leq w \leq 6\}$
- Special notation for intervals $\in \mathbb{R}$:
 - (a, b) represents $\{x \in \mathbb{R} : a < x < b\}$
 - $[a, b]$ represents $\{x \in \mathbb{R} : a \leq x \leq b\}$

Note: $(a, b) \subset \mathbb{R}$ is a set, but $(a, b) \in \mathbb{R}^2 = \mathbb{R} \times \mathbb{R}$ is also a point on the Cartesian plane or an ordered pair.

Definition: sets A and B are equal if they contain exactly the same elements, and is denoted by $A = B$.

Definition: $(w \in A \implies w \in B) \implies (A \subset B)$ for sets A and B where $A \subset B$ means A is a subset of B .

Note: we will not distinguish between proper subsets $((A \subset B) \& (A \neq B))$ and subsets.

$$A = B \iff (A \subset B) \& (B \subset A)$$

Special Sets

The set with no elements is called the empty set or the null set. Denoted by \emptyset or $\{\}$ which is not the same as $\{\emptyset\}$.

The set containing all possible elements of interest is called the universal set S . It is known as the sample space within probability, and will contain all possible outcomes of a random experiment.

Set Operations

- Intersect (\cap): $A \cap B = \{w \in S : (w \in A) \& (w \in B)\}$
- Union (\cup): $A \cup B = \{w \in S : (w \in A) \vee (w \in B)\}$
- Complement ($\square^c, \square', \bar{\square}$): $A^c = \{w \in S : w \notin A\}$
- Disjoint: $A \cap B = \emptyset$
- Difference: $A - B = A \cap B^c$

Set Algebra

Set algebra provides us with a second method to prove that two sets A, B are equal.

It uses the fact that both union and intersection are commutative and associative to prove the statement.

The following are a few properties that come from commutativity and associativity:

- $A \cup B = B \cup A$
- $A \cap B = B \cap A$
- $(A \cap B) \cap C = A \cap (B \cap C)$
- $(A \cup B) \cup C = A \cup (B \cup C)$

\cup is distributive over \cap and vice versa:

- $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$
- $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$