



## Tansley review

# The molecular–physiological functions of mineral macronutrients and their consequences for deficiency symptoms in plants

Author for correspondence:  
*Jan K. Schjoerring*

Received: 27 August 2020

Accepted: 15 September 2020

**Thomas Christian de Bang<sup>1\*</sup> , Søren Husted<sup>1\*</sup> , Kristian Holst Laursen<sup>1\*</sup> , Daniel Pergament Persson<sup>1\*</sup> and Jan Kofod Schjoerring<sup>1\*</sup>**

<sup>1</sup>Plant and Soil Science Section, Department of Plant and Environmental Sciences, Faculty of Science, University of Copenhagen, Thorvaldsensvej 40, Frederiksberg C DK-1871, Denmark

## Contents

Summary	2446	VI. Calcium	2459
I. Introduction	2446	VII. Magnesium	2462
II. Nitrogen	2447	VIII. Conclusion and perspectives	2465
III. Phosphorus	2451	Acknowledgements	2465
IV. Sulphur	2454	References	2466
V. Potassium	2457		

## Summary

*New Phytologist* (2021) **229**: 2446–2469  
doi: 10.1111/nph.17074

**Key words:** calcium, magnesium, mineral element, nitrogen, nutritional disorder, phosphorus, potassium, sulphur.

The visual deficiency symptoms developing on plants constitute the ultimate manifestation of suboptimal nutrient supply. In classical plant nutrition, these symptoms have been extensively used as a tool to characterise the nutritional status of plants and to optimise fertilisation. Here we expand this concept by bridging the typical deficiency symptoms for each of the six essential macronutrients to their molecular and physiological functionalities in higher plants. We focus on the most recent insights obtained during the last decade, which now allow us to better understand the links between symptom and function for each element. A deep understanding of the mechanisms underlying the visual deficiency symptoms enables us to thoroughly understand how plants react to nutrient limitations and how these disturbances may affect the productivity and biodiversity of terrestrial ecosystems. A proper interpretation of visual deficiency symptoms will support the potential for sustainable crop intensification through the development of new technologies that facilitate automated management practices based on imaging technologies, remote sensing and in-field sensors, thereby providing the basis for timely application of nutrients via smart and more efficient fertilisation.

## I. Introduction

The 14 inorganic elements required by plants to complete a full life cycle are coined the *essential plant nutrients*, and grouped into macronutrients and micronutrients on the basis of their

concentration in plant dry matter. The macronutrients are comprised of nitrogen (N), phosphorus (P), sulphur (S), potassium (K), calcium (Ca) and magnesium (Mg). In situations in which a nutrient is not present in sufficient amounts to support its functional roles, it will lead to a state of deficiency, with specific responses characteristic for each nutrient (van Maarschalkerweerd & Husted, 2015). The severity of a nutrient deficiency can range

\*These authors contributed equally to this work.

from mild and transient to severe and chronic, and plants may experience multiple deficiencies during their lifespan, some even occurring simultaneously. In addition, widespread interactions with a range of different abiotic and biotic growth factors such as water, light, pests and pathogens, may lead to atypical symptoms of nutrient-related disorders.

On a global scale, various nutrient deficiencies have major negative consequences for crop production, resulting in reduced yields and a poor quality of food and feed. At the same time, inappropriate use of fertiliser may lead to pollution of terrestrial and aquatic environments, while fertiliser production itself is highly energy demanding, thus contributing to climate change (Sharpley *et al.*, 2018; Thompson *et al.*, 2019). In natural terrestrial ecosystems, nutrient availability is, next to that of water, the environmental factor that most strongly affects the responses of plants to climate change and their ability to cope with environmental stress, thus having a major effect on species biodiversity (Isbell *et al.*, 2013; Lambers & Oliveira, 2019; Terrer *et al.*, 2019).

A large body of knowledge concerning the functional properties of macronutrients in plants already exists (Hawkesford *et al.*, 2012). However, in the recent years significant new knowledge on the molecular processes underlying the direct responses of plants to limitation of a given nutrient has appeared. In addition, starvation networks, describing how individual nutrients interact and cross-talk along the pathway from uptake to integration in metabolism, have been unravelled (Kopriva *et al.*, 2015; Dong *et al.*, 2017; Maeda *et al.*, 2018; Courbet *et al.*, 2019; Hu *et al.*, 2019; Medici *et al.*, 2019).

In this review, we highlight recent insights into the physiological and molecular functions of plant macronutrients. Special emphasis is given to linking their physiological functions with the typical deficiency symptoms that may appear when nutrient availability is suboptimal for proper growth. Understanding the coupling between visual deficiency symptoms and their interactions with the functional properties of the triggering nutrient(s) is relevant for breeding of new nutrient-efficient genotypes that are able to withstand unfavourable climatic events and adverse soil conditions. Moreover, a proper understanding and interpretation of visual deficiency symptoms will support the potential for sustainable intensification of crop production systems by enabling development of new technologies that support automated and data-driven management practices based on bioimaging, remote sensing, in-field spectrometers and/or sensors, thereby providing a basis for timely application of nutrients to crops.

## II. Nitrogen

Nitrogen is an integral constituent of purines and pyrimidines that make up nucleic acids (DNA and RNA). The amine group of amino acids in all peptides, proteins and enzymes also have N as an essential component. The majority of shoot N is present in the chloroplasts as RuBisCo, photosystem proteins and chlorophyll molecules that contain four N atoms in the tetrapyrrole (chlorin) ring. The N containing ions/molecules nitrate ( $\text{NO}_3^-$ ), ammonium ( $\text{NH}_4^+$ ) and nitric oxide (NO) are important signalling molecules (Table 1).

Plants predominantly take up N in the form of  $\text{NO}_3^-$  and  $\text{NH}_4^+$ , but also organic N in the form of peptides and amino acids may be absorbed (Ganeteg *et al.*, 2017). The uptake of the different N forms is mediated via transport proteins with different specificity and affinity. In addition, a substantial amount of N may be provided by N-fixing microbial associations of which the best studied is that between legumes and rhizobia, the latter fixing  $\text{N}_2$  into  $\text{NH}_4^+$  in special root organs called nodules.

Once taken up, N is either reduced for assimilation directly in the roots or translocated to the shoot via the xylem as amino acids, amides,  $\text{NO}_3^-$ ,  $\text{NH}_4^+$  or as ureides in tropical legumes. In the shoot, N is stored, metabolised or further translocated via the phloem to N sinks, such as developing leaves, roots, fruits and seeds. N is stored as protein, primarily RuBisCo, or in vacuoles as  $\text{NO}_3^-$ , while translocation to sinks primarily occurs as amino acids (Tegeder & Masclaux-Daubresse, 2018). Recent studies have shown that N translocation in the phloem of *A. thaliana* may also occur as  $\text{NO}_3^-$  (Chen *et al.*, 2020) or ureides (Takagi *et al.*, 2018).

Primary N metabolism begins with assimilation of  $\text{NO}_3^-$ , which is first reduced to  $\text{NO}_2^-$  by nitrate reductase (NR) in the cytosol, then further reduced and converted to  $\text{NH}_4^+$  by nitrite reductase (NiR) in plastids or chloroplasts. The formed  $\text{NH}_4^+$  is, together with  $\text{NH}_4^+$  generated by other processes such as photorespiration, protein catabolism and lignin biosynthesis, assimilated into glutamine by the enzyme glutamine synthetase (GS). Glutamine may subsequently react with 2-oxoglutarate to yield two glutamate molecules, a reaction catalysed by glutamate synthase (GOGAT). Alternatively, glutamine may react with aspartate to yield asparagine and glutamate as catalysed by asparagine synthetase (AS).

### 1. Molecular and physiological responses to N deficiency

Plants inadequately supplied with N induce a myriad of changes to their transcriptome for example affecting root architecture, shoot development and flowering. A central component in this response is  $\text{NO}_3^-$  and its function as signalling molecule.

### 2. The nitrate transceptor NRT1.1 is essential for adaptation to fluctuating N conditions

Plants respond within minutes to changes in external  $\text{NO}_3^-$  levels by changing the expression of hundreds to thousands of genes as part of the primary nitrate response (PNR), including Nir, NR,  $\text{NO}_3^-$  transporters and genes involved in organic acid metabolism. Nitrate is sensed by the  $\text{NO}_3^-$  transceptor NRT1.1 (NPF6.3). The CBL-interacting protein kinase (CIPK) CIPK8 is involved in dephosphorylation of NRT1.1 at high  $\text{NO}_3^-$ , whereas CIPK23 phosphorylates NRT1.1 under low  $\text{NO}_3^-$  conditions (Y. Y. Wang *et al.*, 2018). In the nonphosphorylated state, NRT1.1 dimerises to a homo-dimer with low  $\text{NO}_3^-$  affinity, while phosphorylation leads to a monomeric and high-affinity state (Fig. 1.1) (Sun *et al.*, 2014). The effects of CIPK23 on the  $\text{NO}_3^-$  transport properties of NRT1.1 are counteracted by ABI2, a phosphatase that is inhibited by abscisic acid (ABA) (Leran *et al.*, 2015). CIPK23 is also involved

**Table 1** Functional roles and foliar deficiency symptoms of the macronutrients.

Nutrient	Dry matter conc. (%)	Functional roles	Phloem mobile	Deficiency symptoms	Stratification
Nitrogen	1–5	Major constituent of: Macromolecules (e.g. proteins, nucleic acids, coenzymes, membrane components, chlorophyll). Organic metabolites (e.g. amino acids, amines, phytohormones, secondary metabolites). Signalling ions/molecules (e.g. $\text{NO}_3^-$ , $\text{NH}_4^+$ , NO).	Yes	General chlorosis on oldest leaves. Stunted growth, small leaves, reduced shoot branching and early flowering. Often anthocyanosis on leaf and stem.	Acropetal
Phosphorus	0.3–0.5	Structural element of nucleic acids and phospholipids. Energy metabolism (ATP and NADPH). Signalling and enzyme activation via phosphorylation and dephosphorylation.	Yes	Anthocyanosis. Dark-green and/or purple leaves.	Acropetal
Sulphur	0.1–0.5	Constituent of organic metabolites and cellular components (e.g. cysteine, methionine, glutathione, ferredoxin and secondary metabolites like glucosinolates and alliins). Electron transport.	Conditional	Chlorosis of young leaves. Stunted growth. Anthocyanosis.	Basipetal
Potassium	1–6	Osmotic regulation and provision of turgor for cell growth. Regulation of stomatal opening and plant movement. Cation-anion balance, electro-neutrality, biochemical pH stat. Stabilisation of binding between biomolecules.	Yes	Chlorosis on tip of oldest leaves that develop into marginal necrosis. Bronzing. Slack appearance due to poor turgor and stomatal control.	Acropetal
Calcium	0.2–2	Structural element. Stabilisation of membranes, cell walls and cytoskeleton. Signalling ( $\text{Ca}^{2+}$ ).	No	Disintegration of root tissue. Necrotic lesions on leaf edges and tips. Meristem death. Necrotic spots on fruits and vegetables. Leaf deformity.	Basipetal
Magnesium	0.1–1	Central element in chlorophyll. Electrostatic interactions and complex formation with enzymes and substrates, for example ATP activation through Mg-ATP complexation essential for energy-requiring transport processes.	Yes	Intervenous chlorosis on oldest leaves that eventually develop into necrosis. Accumulation of sucrose and starch in chloroplast.	Acropetal


Stratification: Basipetal, symptoms first appear on youngest leaves; acropetal, symptoms first appear on oldest leaves.

in  $\text{NH}_4^+$  uptake as it phosphorylates and inhibits the ammonium transporter AMT1 at high  $\text{NH}_4^+$  concentrations (Straub *et al.*, 2017). Nitrate sensing by NRT1.1 is concentration dependent and triggers  $\text{Ca}^{2+}$  signalling by the three calcium-sensor protein kinases (CPKs) CPK10, CPK30 and CPK32, that translocate to the nucleus in response to  $\text{NO}_3^-$  (K.-h. Liu *et al.*, 2017). The CPKs then phosphorylate Nin-like protein 7 (NLP7), which appears to be a master regulator of the primary N response (Marchive *et al.*, 2013).

### 3. Low N induces morphological changes to root systems architecture

Mild N deficiency leads to elongation of lateral roots and the primary root, while severe or prolonged N deficiency inhibits primary root growth and total root length (Fig. 1.2) (Gruber *et al.*, 2013). Stimulation of lateral root growth during mild N deficiency has been shown to be auxin-dependent (Ma *et al.*, 2014). During prolonged N deficiency, inhibition of lateral root growth is controlled by NRT1.1 which removes auxin from lateral root primordia inhibiting their growth. By contrast, high  $\text{NO}_3^-$  inhibits auxin transport away from lateral root primordia, thereby stimulating root growth (Kruck *et al.*, 2010; Bouguen *et al.*, 2015). The local inhibition of

lateral root growth during N starvation is, in addition to auxin, also regulated by CLAVATA3 (CLV3)/EMBRYO SURROUNDING REGION-related (CLE) peptides, which are induced to specifically inhibit emergence of lateral roots through a CLAVATA1-dependent signalling pathway (Fig. 1.2) (Araya *et al.*, 2014). Two systemic signalling pathways known as the N-demand and N-supply pathways promote root growth and N uptake in root parts exposed to high N (Poitout *et al.*, 2018). In the N-demand pathway, C-TERMINALLY ENCODED PEPTIDES (CEP) peptides are produced in low N root parts that, via shoot-acting receptors and phloem-mobile Class III glutaredoxin polypeptides (CEPD1/CEPD2), promote root growth and N uptake in high N root parts (Tabata *et al.*, 2014; Ohkubo *et al.*, 2017). The N-supply pathway is initiated by synthesis and translocation of cytokinin from the root to the shoot, which both induces N-dependent leaf expansion and increased expression of  $\text{NO}_3^-$  transporters in high N roots (Poitout *et al.*, 2018).  $\text{NH}_4^+$  toxicity inhibits primary root growth, while local  $\text{NH}_4^+$  may stimulate lateral root branching (Liu & von Wirén, 2017). N-deficient plants exposed to a local supply of  $\text{NH}_4^+$  respond by developing more second and third order lateral roots through a mechanism dependent on the  $\text{NH}_4^+$  transporter AMT1;3 (Lima *et al.*, 2010).



#### 4. Transition from vegetative to reproductive stage

In general, low N leads to earlier flowering compared with high N conditions, however prolonged N starvation may result in a

flowering time comparable with that of high N in *A. thaliana* (Lin & Tsay, 2017). Several mechanisms are responsible for linking N status and flowering, of which a central one revolves round the balance between the microRNAs miR156 and miR172 (Fig. 1.3).

**Fig. 1** Schematic model showing how nitrogen (N) affects plant growth and physiology. (1) NRT1.1 (NPF6.3) is a dual-affinity nitrate transporter that senses external nitrate concentrations. At high  $\text{NO}_3^-$ , the calcineurin B-like interacting protein kinase 8 (CIPK8) regulates dephosphorylation of NRT1.1, causing dimerisation in a low-affinity state. Cytosolic  $\text{Ca}^{2+}$  concentrations increase and the  $\text{Ca}^{2+}$  signal is further transmitted by three calcium-sensor protein kinases (CPK): CPK10, CPK30 and CPK32 that phosphorylate Nin-like protein 7 (NLP7) in the nucleus. NLP7 induces expression of hundreds of genes as part of the primary nitrate response (PNR). At low  $\text{NO}_3^-$ , CIPK23 phosphorylates NRT1.1 resulting in a high-affinity monomeric state. (2) Spatial distribution and concentration of N affects root development. Mild N deficiency results in a root foraging response, that is existing lateral roots grow longer and deeper, a response caused by high auxin in root tips. At severe deficiency, root growth is inhibited through auxin removal from root tips. In addition, CLAVATA3/Embryo Surrounding Region-Related (CLE) peptides inhibit lateral root growth locally. Under conditions with heterogeneous N distribution, systemic signalling via C-TERMINALLY ENCODED PEPTIDES (CEP-CEPD) from low N roots induces lateral root growth and N uptake in high N roots. Cytokinin promotes root proliferation in high N roots via root-shoot-root signalling. (3) N levels affect flowering time. (a) Mild N deficiency promotes flowering compared with severe starvation and high N levels. (b) Nitrogen availability and flowering are correlated via different pathways related to ageing, photoperiod, gibberellic acid (GA) and vernalisation, as well as the autonomous pathway. Central to this regulation is the flowering gene FLOWERING LOCUST (FT). In the photoperiod pathway, low and high  $\text{NO}_3^-$  oppositely affect expression of Ferredoxin-NADP<sup>+</sup>-oxidoreductase 1 (FNR1) that positively affects FT via Constans (CO) and Cryptochrome 1 (CRY1). In the ageing pathway, the balance between miR156 and miR172 oppositely affects FT via expression of APATELA2 (AP2) and SPL. In the GA pathway,  $\text{NO}_3^-$  modulates GA1 levels and DELLA protein activity that in turn affects flowering via AP2 and SPL in the ageing pathway. The vernalisation pathway and the autonomous pathway both inhibit FLOWERING LOCUS C (FLC) that in turn inhibits FT. (4) (left) Glutamate is the precursor for chlorophyll synthesis and is stimulated by nitrogen; (right) Nitrogen deficiency results in proteolysis and released amino acids are translocated through the phloem from old source leaves to young sink leaves. (5) Leaf growth and shoot branching are regulated by cytokinin in a  $\text{NO}_3^-$ -dependent manner. Biosynthesis of the root cytokinin trans-zeatin (tZ) and the cytokinin precursor trans-zeatin-riboside (tZR) is correlated with nitrate levels. Root-derived cytokinin positively regulates shoot apical meristem (SAM) size and activity, and breaks auxiliary bud dormancy. Collectively this leads to more shoot branches and more leaves. In addition, cytokinin delays senescence and activates photosynthesis resulting in improved carbohydrate assimilation. (6) In some species, N deficiency induces anthocyanin biosynthesis in stems and leaves. Low N induces the expression of PRODUCTION OF ANTHOCYANIN PIGMENT (PAP) transcription factors via miR165-mediated downregulation of SQUAMOSA PROMOTER BINDING PROTEIN-LIKE 9 (SPL9). High N represses the expression of PAP genes via NLP7-mediated expression of LATERAL ORGAN BOUNDARY DOMAIN (LBD) genes. Furthermore, DELLA proteins positively affect anthocyanin accumulation during N deficiency through repression of gibberellic acid (GA) on anthocyanin biosynthesis.

During plant ageing, miR156 levels decrease relative to miR172, promoting transition from the vegetative stage to the reproductive stage. miR156 represses expression of several SQUAMOSA PROMOTER BINDING PROTEIN-LIKE (SPL) transcription factors, which in turn affects expression of both miR172 and several flowering genes (Spanudakis & Jackson, 2014). Low N conditions induce expression of miR156 and repress miR172, hence affecting flowering time and stage transition depending on developmental stage (Liang *et al.*, 2012; Lin & Tsay, 2017). Another recently discovered mechanism linking N status and flowering is based on the intensification of circadian rhythms controlled by the blue light photoreceptor cryptochrome 1 (CRY1). Here, low N induces ferredoxin-NADP<sup>+</sup>-oxidoreductase (FNR1), which in turn controls activity of CRY1, leading to earlier flowering (Yuan *et al.*, 2016). Three other pathways are known to link N status and flowering, namely the autonomous, gibberellic acid (GA) and the vernalisation pathways (Fig. 1.3).


## 5. Coupling of visual leaf symptoms with functional roles

Low N leads to developmental responses such as stunted growth, early flowering, reduced tillering in cereals, shoot branching and smaller leaves due to decreased cell division and expansion (Table 1). The most prominent visual symptom of N deficiency is chlorotic leaves, where the chlorosis spreads uniformly (general chlorosis) across the entire leaf due to either reduced chlorophyll synthesis or breakdown of existing chlorophyll binding proteins in the photosystems (Table 1; Fig. 2). Nitrogen assimilation in the chloroplast is tightly coupled with chlorophyll biosynthesis via the GS/GOGAT pathway as glutamate is the precursor for all chlorophylls produced in the tetrapyrrole biosynthetic pathway (Fig. 1.4). Hence, there is a strong correlation between the N status

of plants and the chlorophyll content, which can be used to optimise N fertiliser application (Hudson *et al.*, 2011). Although the mutual regulation of chlorophyll biosynthesis and breakdown largely remains unknown, it is well established that N deficiency induces chlorophyll breakdown via proteolysis leading to the release of amino acids, amides and  $\text{NH}_4^+$  (Fig. 1.4) (Havé *et al.*, 2016). All released N compounds are highly mobile in the phloem. Older leaves consequently act as source tissue during low N situations supplying young and developing tissue like leaves, flowers and seeds with N. As a consequence, visual symptoms are first visible on the oldest leaves (acropetal stratification).

Reduced shoot branching, stunted growth and inhibition of leaf expansion constitute central physiological responses to N deficiency. These responses are closely linked to cytokinin produced in roots in response to  $\text{NO}_3^-$  (Rahayu *et al.*, 2005) (Fig. 1.5). Subsequent root-to-shoot translocation of trans-zeatin (a cytokinin) and trans-zeatin-riboside (a cytokinin precursor) affects leaf growth through stimulation of leaf expansion and meristem activity (Osugi *et al.*, 2017). In addition, root-derived trans-zeatin-riboside directly stimulates shoot apical meristem (SAM) growth and thereby overall shoot biomass (Landrein *et al.*, 2018). Shoot branching is controlled by formation and activation of axillary buds. Axillary buds are activated by  $\text{NO}_3^-$  due to a stimulatory effect on cytokinin production, whereas low  $\text{NO}_3^-$  maintains bud dormancy and thereby reduce branching (Fig. 1.5) (Müller *et al.*, 2015). This effect further relies on interaction with auxin and strigolactones (de Jong *et al.*, 2014).

Another prominent visual symptom of N deficiency is reddish coloration due to anthocyanin production (Fig. 1.6). This is only seen in some species and may also be influenced by other abiotic factors such as high light intensity, low temperatures and P deficiency. In *A. thaliana*, low N induces expression of key



**Fig. 2** Nitrogen (N) deficiency always appears on the oldest leaves first and is characterised by a general chlorosis. (a) Leaf of healthy maize plant to the left and N-deficient leaf with general chlorosis to the right. (b) Anthocyanin accumulation in a N-deficient oilseed rape leaf with general chlorosis. (c) General chlorosis in tomato, with anthocyanosis on veins and on the abaxial side of leaves.

transcription factors positively regulating anthocyanin synthesis, including PRODUCTION OF ANTHOCYANIN PIGMENT1 (PAP1), PAP2 and GL3 (Xu *et al.*, 2015). Repression of GA signalling by DELLA proteins is required for the induction of several biosynthesis genes, including genes related to anthocyanin accumulation during N deficiency (Y. Zhang *et al.*, 2017). By contrast, three members of the LATERAL ORGAN BOUNDARY DOMAIN (LBD) protein family LBD37/38/39, which are  $\text{NO}_3^-$  induced, act as negative regulators of anthocyanin synthesis by repressing PAP1 and PAP2 expression (Rubin *et al.*, 2009). In parallel, the SQUAMOSA PROMOTER BINDING PROTEIN-LIKE (SPL) transcription factor, SPL9, is a negative regulator of anthocyanin synthesis, as it represses PAP1, PAP2 and other transcription factors that positively regulate anthocyanin production (Fig. 1.6). SPL9 is repressed by miR156, which itself is induced by N deficiency, thus linking low N conditions to anthocyanin production by increasing expression of PAP1 and the biosynthetic gene DFR (Cui *et al.*, 2014).

### III. Phosphorus

Phosphorus is a structural element of essential biomolecules involved in energy metabolism (ATP, NADPH), of nucleic acids (DNA, RNA) and of phospholipids in cell membranes (Table 1). The hydrolysis of the P–O–P bond in adenosine triphosphate (ATP) to yield ADP and inorganic phosphate (Pi) is the principal source of biochemical energy within the cell (Kamerlin *et al.*, 2013). Plants predominantly take up P as the inorganic orthophosphate ions  $\text{H}_2\text{PO}_4^-$  and  $\text{HPO}_4^{2-}$ , depending on soil pH.

A wide range of enzymes has evolved to catalyse reactions involving phosphates. These enzymes catalyse phosphorylation and dephosphorylation processes affecting the activity of proteins, stabilising them or labelling them for breakdown, facilitating or inhibiting their movement between subcellular compartments or regulating protein–protein interactions. The importance of protein phosphorylation/dephosphorylation in cellular signalling is emphasised by the fact that one third of all proteins in the cell are phosphorylated at any given time (Manning *et al.*, 2002).

#### 1. Functional properties of P in photosynthesis


Phosphorus is an essential element in the light reaction of photosynthesis, where absorbed light drives the electron transport chain in the thylakoid membrane to generate ATP and NADPH. Phosphorus also plays an essential role in the dark reactions (Calvin–Benson cycle), in which ATP and NADPH are used to convert  $\text{CO}_2$  to carbohydrates in the chloroplast stroma (Fig. 3.1). When plants are exposed to P deficiency, lowered chloroplast Pi levels lead to a reduction in ATP production, because Pi together with  $\text{CO}_2$  and  $\text{H}_2\text{O}$ , are the primary substrates for photosynthesis. Moreover, during P deficiency a larger fraction of  $\text{NADP}^+$  remains in the reduced form (NADPH), because it cannot be utilised in the Calvin–Benson cycle due to ATP limitation (Fig. 3.2) (Carstensen *et al.*, 2018). A prominent feature of P deficiency is therefore a marked decrease in  $\text{CO}_2$  assimilation and reduced biomass production.

#### 2. Molecular and physiological responses to P deficiency

In response to P limitation, plants trigger a series of biochemical adjustments, including activation of organic acid (OA) exudation to mobilise inorganic P trapped in precipitated Al, Fe and Ca salts in the rhizosphere. OA are mainly produced in the mitochondria through the tricarboxylic acid (TCA) cycle (Krebs cycle) and to a lesser extent in the glyoxysomes as part of the glyoxylate cycle. OA have a high affinity for Al, Fe and Ca and can displace Pi from insoluble complexes through ligand exchange, increasing the P solubility and make soil Pi available for plant uptake.

The activity of key enzymes such as citrate synthase (CS) and PEP carboxylase involved in OA biosynthesis increases in response to P starvation and OA are subsequently released to the rhizosphere as a response. The proteins responsible for OA secretion belong predominately to the ALMT (aluminium-activated malate transporters) and the MATE/AACT (multidrug and toxic compound extrusion/aluminium-activated citrate transporter) families (Delhaize *et al.*, 2012).

There is a strong correlation between the P status of plants and cellular respiration (Wang *et al.*, 2015). Plant mitochondria possess



**Fig. 3** Schematic model showing how phosphorus (P) affects plant growth and physiology. (1) P the chloroplast stroma affects CO<sub>2</sub> assimilation through the carboxylation rate of RuBisCo and the regeneration of ribulose-1,5-bisphosphate (RuBP). 3-PGA, 3-phosphoglycerate; G3P, glyceraldehyde-3-phosphate. (2) Lowered chloroplast PO<sub>4</sub><sup>3-</sup> (Pi) levels lead to a reduction in ATP production. The reduced ATP synthesis impacts the flow of protons from the thylakoid lumen to the chloroplast stroma, causing lumen acidification, which triggers energy dissipation via nonphotochemical quenching. (3) When the Pi concentration of the cell decreases, less substrate is available for oxidative phosphorylation (ATP synthesis) via the conventional cytochrome oxidase pathway (COX). This triggers the alternative oxidase (AOX) pathway (no ATP synthesis), ensuring that electron transport in the respiration process continues under P starvation. Moreover, P deficiency leads to NO accumulation, which impacts the TCA cycle and leads to aconitase inhibition. As a consequence, citrate accumulates, leading to stimulation of AOX relative to COX and promoting citrate exudation from the root to the rhizosphere. (4) Root system architecture is affected during P deficiency. In cereals, the response is complex and much more variable than observed in Arabidopsis, in which the architecture is shaped to allow a more thorough exploration of the top soil where most P is present. P deficiency may enhance/repress primary root and lateral root growth, whereas the root density and length generally are enhanced. (5) During P deficiency, strigolactone biosynthesis increases in the root leading to increased strigolactone transport to the shoot and higher levels of strigolactone in the bud, which initiate degradation of the auxin efflux protein PIN1, inhibiting auxin transport away from the bud. When the auxin level in the bud is high, the cytokinin level is low and bud outgrowth is inhibited. (6) Anthocyanosis: photoabatement by anthocyanin shields photosynthetic systems from light stress.

two distinct pathways for the transfer of electrons to molecular oxygen during respiration: the cytochrome *c* oxidase (COX) and the alternative oxidase (AOX) pathways (Fig. 3.3). Generally, the

AOX expression is very low until a stress condition emerges. When the Pi concentration of the cytosol goes down, less substrate is available for oxidative phosphorylation (ATP synthesis) via the

conventional COX pathway, triggering the AOX pathway, which ensures that electron transport in the respiration process continues under P starvation. Simultaneously, the root exudation of citrate increases. The interactions between Pi, AOX and citrate exudation are not yet fully resolved at the mechanistic level (Del-Saz *et al.*, 2018). However, P deficiency leads to accumulation of the messenger molecule NO in roots (Wang *et al.*, 2010), which inhibits aconitase and thereby prevents the conversion of citrate to iso-citrate (Gupta *et al.*, 2012). The resulting accumulation of citrate subsequently triggers the AOX pathway and allows continuation of respiration when the COX pathway is hampered. By linking these processes, P sensing is related to the TCA cycle intermediate citrate, modulation of mitochondrial respiration, OA exudation and eventually mobilisation of P in the rhizosphere, allowing for an increased Pi uptake.

Plants respond to P deficiency by increasing the root-to-shoot ratio and by changing the root architecture to facilitate a more thorough exploration of soil P resources, particularly in the top soil where most P tends to accumulate. The root response to P limitation differs across the plant kingdom. Some plants, including *A. thaliana*, respond to P starvation by a process called root apical meristem (RAM) exhaustion, where cells within the root tip loose meristematic activity and cells in the elongation zone are arrested, eventually leading to a reduction in growth of the primary root (Fig. 3.4) (Gutiérrez-Alanís *et al.*, 2018). The RAM process is triggered by direct exposure of the root tip to low P levels in the soil. The local response appears to be independent of the general root P status and is not caused by metabolic disturbances triggered by the P deficiency. It has been shown that accumulation of Fe<sup>3+</sup> within the root tip is essential for the inhibition of primary root growth under low P conditions (Ward *et al.*, 2008). In P-starved *A. thaliana*, Fe accumulates in specific zones of the primary root, which triggers the STOP1-ALMT1 module, resulting in exudation of malate within the apoplast of the root tip. As malate forms strong Fe complexes, this leads to Fe accumulation and Fe toxicity in RAM tissue, causing generation of reactive oxygen species (ROS) though Fenton reactions. The accumulation of ROS appears to stimulate callose formation and a stiffening of the cell walls, preventing cell elongation in RAM. Mutations in STOP1-ALMT1 restore the growth of the primary root under P limitation (Balzergue *et al.*, 2017).

Compared with *A. thaliana*, the root system of, for example, cereals have a more complex response to P deficiency and marked species and genotype differences have been observed (Péret *et al.*, 2014). Maize plants tend to allocate relatively more carbon resources to their roots during P deficiency and shows no reduction or even a slight enhancement in root elongation. Also *japonica* rice roots respond with elongation of the primary root under P deficiency (Dai *et al.*, 2012).

Apart from changes to primary root growth under P deficiency, plants respond to P limitation by increasing the root hair density and/or the frequency and length of lateral roots. The length and density of root hairs and fine roots are important hot spots for P acquisition as they provide the largest surface area for P uptake with the lowest carbon investment in root biomass. Transcriptomic

analyses of root hair and tips have shown that accumulation of Pi transporters belonging to the PHT1 family are markedly more frequent at the interface between the newly developed root tissue and the soil to be explored. In *A. thaliana* seedlings, only the cap seems to play an important role in P acquisition, whereas the elongation zone above the cap seems to play a minor role (Kanno *et al.*, 2016).

Auxin is a key regulator of root system architecture, and plays an important role in shaping the architecture of roots during P starvation. Especially RAM appears to be important in that respect as auxin accumulates here when P becomes limiting, which inhibits further growth of the root tip, irrespective of the shoot P status. Strikingly, this inhibitory response of root tip growth to auxin is opposite to the elongation of lateral roots, which are promoted by auxin accumulation. During P deficiency, the expression of the auxin receptor TIR1 is upregulated, thereby increasing the sensitivity of the pericycle cells towards auxin. This leads to activation of the auxin response factor (ARF), a transcription factor that promote lateral root emergence. The auxin accumulation in RAM tissue during P deficiency is also involved in triggering the formation of trichoblast cells to form root hairs when the auxin levels are upregulated at the root tip. At low P levels, auxin is mobilised and enters the root hair differentiation zone via the auxin influx carrier (AUX1). Elevated auxin levels in trichoblasts subsequently trigger a gene expression cascade, mediated by a series of transcription factors, to promote root hair formation and elongation (Bhosale *et al.*, 2018).

Also, shoot branching is modified during P limitation (Fig. 3.5). Even moderate- and short-term P deficiency in barley is known to affect grain yields significantly via a reduction in the number of fertile tillers (Carstensen *et al.*, 2019). This effect is mediated via strong interactions between tissue P levels and the hormones strigolactones, auxin and cytokinin. During P deficiency, strigolactone biosynthesis is increased in the roots, promoting strigolactone transport towards the shoot (Kohlen *et al.*, 2011). When the level of strigolactone increases in the bud, degradation of the auxin efflux carrier (PIN1) is initiated, which inhibits export of auxin from the bud to the stem (Kebrom *et al.*, 2013). When the auxin levels in the bud are high, the cytokinin levels are low and bud outgrowth is prevented. During P sufficiency, strigolactone levels are not increased in the bud and auxin is transported away from the bud by the PIN1 proteins. At low auxin levels, the cytokinin levels increase, which in turn triggers bud outgrowth.

### 3. Coupling of functional roles and visual deficiency symptoms for P

A reddish-purple cast to the leaves due to anthocyanin accumulation (anthocyanosis) is a marked symptom of P deficiency across a wide range of plant species (Table 1; Fig. 4). Anthocyanins are flavonoid pigments synthesised on the cytoplasmic face of the endoplasmic reticulum and stored in vacuoles where they are protected from oxidation by the low pH. The anthocyanins typically accumulate on a dark-green coloured background (Fig. 3.6), due to a high chlorophyll concentration caused by a



**Fig. 4** Phosphorus (P) deficiency always appears on the oldest leaves first and is characterised by anthocyanosis on a nonchlorotic background. (a) Leaf of P-deficient maize plant showing anthocyanosis on a dark nonchlorotic background. (b) Anthocyanosis on the abaxial side of P-deficient tomato leaf.

reduced cell division and extension in P-deficient plants (Hughes and Lev-Yadum, 2015).

However, not all plant species develop anthocyanosis when exposed to P deficiency, important exceptions are for example potato, sugar beet and rice (Hughes and Lev-Yadum, 2015). It is also striking that symptoms develop very differently across species (Fig. 4). Tomato develops symptoms on the abaxial side, whereas maize seems to express symptoms on the adaxial side and other species tend to accumulate anthocyanins on the stems or along the leaf margins. The majority of species, however, seem to accumulate anthocyanins across the whole leaf or in major parts of the leaf. As the chlorophyll concentration on a leaf area basis is maintained or even increased during P deficiency this typically leads to the development of a red-purple-bluish hue across the leaf surface. The severity of P symptoms appears to be heavily influenced by abiotic factors, such as leaf temperature and light intensity.

Phosphorus deficiency triggers an increased expression of genes in the anthocyanin biosynthetic pathway. Some of these genes appear to be GA dependent. When plants are exposed to P deficiency, transcript levels of genes encoding enzymes activated by GA are reduced, whereas those that deactivate GAs are increased. This leads to a decrease in bioactive GA levels, which in turn triggers the synthesis of specific growth-regulating proteins (DELLAs). Increased levels of DELLA in P-deficient plants upregulate transcripts encoding for enzymes involved in anthocyanin biosynthesis, which include *F3H* and *LDOX*, but not *UF3GT* and *PAP1* (Jiang *et al.*, 2007).

Currently, it is believed that the primary role of anthocyanins is photoprotection of the light-harvesting complexes (LHCs) of PSII, which theoretically should provide a functional advantage to red leaves when the capacity for thermal energy dissipation is exceeded by excessive absorption of photons (Gould *et al.*, 2018). As a consequence, anthocyanosis can be seen as a feedback response to P deficiency in order to prevent over-excitation and damage to PSII. However, using closely related genotypes of maize and coleus (*Solenostemon scutellarioides*) with different

regulation of anthocyanin production, Henry *et al.* (2012) showed over a range of light intensities that leaves with anthocyanosis did not better tolerate the impacts of P deficiency than green leaves. Thus, the putative adaptive significance of anthocyanin accumulation under P deficiency stress still remains enigmatic.

#### IV. Sulphur

Atmospheric S (e.g. sulphur dioxide, SO<sub>2</sub>) can be taken up and utilised by aerial plant parts, but the main source of S is inorganic sulphate (SO<sub>4</sub><sup>2-</sup>), which is taken up by roots and assimilated into a variety of key metabolites, including the amino acids cysteine and methionine and various prosthetic groups.


Plant species differ considerably in their demand for S, which generally decreases in the order: Cruciferae > Leguminosae > Graminae. The occurrence of S deficiency has increased since the 1990s, mainly because of decreased deposition of atmospheric SO<sub>2</sub>. Consequently, research on S sensing, uptake, assimilation and functional properties have increased substantially during the past decades (Weissert & Kehr, 2017).

##### 1. Sulphur sensing, uptake and assimilation

Sulphur-deficient plants sense the S status of the soil–root environment via the tissue level of cysteine precursors (Dong *et al.*, 2017). Intermediates in SO<sub>4</sub><sup>2-</sup> and NO<sub>3</sub><sup>-</sup> reduction (SO<sub>3</sub><sup>2-</sup> and NO<sub>2</sub><sup>-</sup>) are toxic and S fluxes must be balanced relative to the fluxes of N and C. Sulphate assimilation is therefore downregulated when C or N is limiting (Long *et al.*, 2015; Kopriva *et al.*, 2019).

When plants are exposed to S deficiency, high-affinity SO<sub>4</sub><sup>2-</sup> transporters in the SULTR family, which function by co-transporting SO<sub>4</sub><sup>2-</sup> along with protons, are upregulated to facilitate uptake from the rhizosphere (Fig. 5.1). An ethylene insensitive-like (EIL) transcription factor, SLIM1, has been isolated and shown to be involved in this response (Koprivova & Kopriva, 2016). In *A. thaliana*, SULTR1;1 and SULTR1;2 are responsible for the initial SO<sub>4</sub><sup>2-</sup> uptake from the soil. SULTR1;1 is mainly expressed in the epidermis, including root hairs, while SULTR1;2 is mainly expressed in the cortex (Kimura *et al.*, 2019).

Plants incorporate SO<sub>4</sub><sup>2-</sup> into organic molecules after reduction to sulphide (S<sup>2-</sup>). Among the macronutrients, S is remarkable due to its ability to change between the oxidation states +6 and -2. The change in oxidation state from +6 to -2 is highly energy dependent and consumes 732 kJ mol<sup>-1</sup>. In comparison, NO<sub>3</sub><sup>-</sup> and C assimilation requires only 347 and 478 kJ mol<sup>-1</sup>, respectively. SO<sub>4</sub><sup>2-</sup> is first activated to adenosine 5'-phosphosulphate (APS) by catalytic reaction with ATP sulphurylase. APS can then either be channelled into cysteine synthesis or used for formation of secondary metabolites. Cysteine may be further converted to methionine, glutathione and all other metabolites containing reduced S (Long *et al.*, 2015; Maruyama-Nakashita, 2017).



**Fig. 5** Schematic model showing how sulphur (S) affects plant growth and physiology. (1) The main plant available S source is inorganic sulphate ( $\text{SO}_4^{2-}$ ), which is taken up by membrane localised transporters (SULTRs). Plants incorporate  $\text{SO}_4^{2-}$  into biorganic molecules after reduction to sulphide, much like  $\text{NO}_3^-$  reduction and assimilation. (2) S is a key component of essential metabolites and cellular constituents in plants, including the amino acid cysteine and the low molecular soluble compound glutathione. (3) Plants contain numerous S-containing secondary metabolites, for example glucosinolates (GSLs), of which many serve as defences against microbes and pests. During S deficiency, GSH and GSLs are degraded in order to up-regulate primary S metabolism (APS, adenosine 5'-phosphosulphate; PAPS, 3'-phosphoadenosine-5'-phosphosulfate). (4) S deficiency reduces the protein content and the content of S-containing amino acids. This has consequences for the quality and nutritional value of food and feed. (5) S deficiency leads to reduced protein synthesis, functionality and quality, anthocyanin production, stunted growth, chlorotic young leaves and premature/reduced flowering. (6)  $\text{SO}_4^{2-}$  uptake is coordinated alongside the uptake and assimilation of N. S deficiency leads to reduced nitrate uptake and NR activity. N deficiency leads to reduced sulphate uptake and APS reductase activity. (7) S deficiency symptoms appear on youngest leaves first (basipetal stratification). S remobilisation is increased by low N availability. Adapted from Long *et al.* (2015) and Maruyama-Nakashita (2017).

## 2. Functional roles of S

Sulphur deficiency interacts massively with the transcriptome and ionome of plants (Courbet *et al.*, 2019) resulting in a decrease in photosynthetic carbon assimilation and a reprogramming of

metabolism to ensure seed production (Jobe *et al.*, 2019). In reduced form, S is a key component of various prosthetic groups in Fe–S proteins, such as ferredoxin, which mediate electron transfer in the photosynthetic electron transport chain (Forieri *et al.*, 2013; Zheng & Leustek, 2017). Thus, a key feature of S deficiency is a

marked suppression of the photosynthetic efficiency and rapid development of chlorotic leaves. In S-deficient rice, the chlorophyll content was 50% lower due to a general decrease in the chlorophyll binding proteins of the light-harvesting complex of both PSII and PSI (Lunde *et al.*, 2008).

Glutathione (GSH) is a low molecular soluble S compound with complex and multifaceted functions associated with its reactive thiol group (Noctor *et al.*, 2012) (Fig. 5.2). This thiol group allows glutathione to donate electrons for enzymatic catalysis, to control redox homeostasis and to function as an antioxidant and a signalling molecule in stress responses and development. Glutathione is also involved in storage and long-distance transport of reduced S from shoots to roots (Zheng & Leustek, 2017) and acts as substrate for phytochelatin synthesis, thereby playing an important role in metal hyperaccumulation (Ahmed *et al.*, 2019).

The primary S compounds are complemented by numerous secondary S-containing metabolites of which many are involved in the defence against pathogens (Fig. 5.3). Examples of important S-containing metabolites are glucosinolates, alliins, camalexin and S-rich peptides of the thionin and defensin groups. Glucosinolates are a diverse group of compounds consisting of a thioglucose core with two S atoms ligated to different amino acids resulting in  $> 100$  different compounds. They constitute a large pool of S, up to 30% of the total S, in *Brassica* tissues and the glucosinolate content of plants directly reflects their S status, being increased at high-S nutrition and decreased under S-limited growth (Aghajanzadeh *et al.*, 2014; Long *et al.*, 2015).

### 3. Sulphur and plant quality

Sulphur deficiency is known to have significant effects on the quality of plant-based food and feed (Fig. 5.4). Methionine is an essential amino acid in human nutrition and often represents a limiting nutritional factor in seed-rich diets (Dung Tien *et al.*, 2016). Sulphur has a pronounced effect on the protein composition of cereal grain as opposed to N, which more generally affects the protein content (Yu ZT *et al.*, 2018). Reduced S availability favours the synthesis and accumulation of S-poor or low-S storage proteins

such as  $\omega$ -gliadin and high-molecular-weight subunits of glutenin. This affects the baking quality of wheat, because disulphide bridging during dough preparation is essential for polymerisation of the glutelin fraction to create optimum loaf volume (Liu *et al.*, 2011; Järvan *et al.*, 2017).

S-containing secondary metabolites such as glucosinolates and alliins are not only important defence compounds, but are also the basis of smell and taste of cruciferous vegetables, onion and garlic. The positive influence of S nutritional status on the content of secondary S-containing metabolites has been documented for a range of crop species, in some cases showing a positive interaction with N status, in others not (Bloem *et al.*, 2005; Poisson *et al.*, 2019) (Fig. 5.3).

### 4. Sulphur deficiency symptoms linked to functional properties

Plants growing at suboptimal S supply display stunted growth, chlorosis, anthocyanosis, as well as premature and reduced flowering, and reduced seed setting (Table 1; Figs 5.5, 6). These symptoms are most pronounced in S-demanding Cruciferae species (Schnug & Haneklaus, 2005). Sulphur-deficient plants become chlorotic and stay chlorotic for an extended period until developing necrosis. Symptoms are typically very similar to the general chlorosis found in N-deficient plants, but in some species intervenous chlorosis is prevalent, similar to symptoms occurring in Fe-deficient plants. The differences in leaf appearance are not fully understood, but are probably related to extensive cross-talk with N and Fe during S deficiency (Fig. 5.6) (Forieri *et al.*, 2017). During S limitation, plants are unable to fully induce their Fe-uptake machinery, which includes relevant transport proteins and the release of phytosiderophores in strategy II plants such as barley, maize, and wheat (Courbet *et al.*, 2019). A characteristic feature across different plant species is the marked accumulation of  $\text{NO}_3^-$  in root tissue combined with a reduction in NR activity, showing that S-deficient plants are unable to utilise N in their metabolism and consequently becomes N deficient (Kaur *et al.*, 2011) and display N deficiency symptoms.



**Fig. 6** Sulphur (S) deficiency always appears on the youngest leaves first and spreads rapidly in basipetal direction. (a) Left, leaf of S-deficient tomato plant showing interveinal chlorosis with anthocyanin accumulation on veins and petiole. Right, a healthy control plant without any symptoms. (b) Left, S-deficient leaf of oilseed rape plant. Right, healthy control.

The chlorosis will in most cases be interveinal due to a relatively higher S supply around leaf veins (Fig. 5.5). However, high anthocyanin content may, in some cases, mask this symptom. Chlorosis induced by S deficiency rarely develops into necrosis, as is for example the case for Mg, K and N, but the stratification of symptoms is dependent on N status. Mature (source) leaves becomes chlorotic under N deficiency due to protein degradation and amide export to young sink leaves (acropetal stratification) (Dubouset *et al.*, 2009). By contrast, S starvation first and strongest affects the young leaves (basipetal stratification). The reason for this difference is a much higher mobility of  $\text{NO}_3^-$  between vacuole and cytosol upon demand than is the case for  $\text{SO}_4^{2-}$  (Miller *et al.*, 2009). In addition, organically bound S only becomes exported to the phloem under N deficiency, because the latter is required to trigger protease activity leading to remobilisation of S-containing amino acids (conditional mobility) (Fig. 5.7). In oilseed rape, S deficiency induces ‘white blooming’, where white flower colour develops from an overload of petal cells with carbohydrates caused by disorders in the protein metabolism. This finally ends up in the formation of leuco-anthocyanins that often occurs during periods of high photosynthetic activity (Schnug & Haneklaus, 2005).

## V. Potassium

Potassium forms no covalent bonds and the  $\text{K}^+$  ion is pivotal for maintaining healthy cellular conditions by affecting for example electro-neutrality, osmotic regulation, anion–cation balancing and the biochemical pH stat (Table 1). The majority of  $\text{K}^+$  is found in vacuoles, where it acts as a turgor provider.  $\text{K}^+$  also facilitates protein and enzyme stability through optimal hydration, hence a large number of enzymes ( $> 70$ ) are either stabilised, activated and/or inhibited by  $\text{K}^+$  as a co-factor, for example pyruvate kinase, phosphofructokinase and starch synthase (Anschutz *et al.*, 2014).

All plant movements are conveyed by  $\text{K}^+$  fluxes. Guard cells increase their osmotic potential through an influx of  $\text{K}^+$ , which results in an accompanying water uptake from adjacent cells, leading to stomatal opening. This process is light induced and starts with  $\text{H}^+$ -ATPase driven acidification of the stomatal apoplast, stimulating  $\text{K}^+$  influx via polarisation of the plasmamembrane potential. Electro-neutrality during  $\text{K}^+$  influx is maintained by  $\text{Cl}^-$ ,  $\text{NO}_3^-$  and malate ions. Conversely, during darkness,  $\text{K}^+$  efflux is correlated with water efflux and stomatal closure. Abscisic acid (ABA) induces stomatal closure through  $\text{H}_2\text{O}_2$  and NO along with subsequent ion channel-mediated loss of  $\text{K}^+$  and anions (Chen *et al.*, 2016). In barley, SLAC1 (Slowly Activating Anion Channel) transports  $\text{NO}_3^-$  and mediates stomatal closure via interaction with ABA (Schäfer *et al.*, 2018), hence linking N and K dependent metabolic processes.

$\text{K}^+$  is highly mobile in the phloem and vital for phloem loading of photoassimilates, which at the site of source cells is ATP driven. The phloem is loaded with sucrose via  $\text{H}^+$ -coupled sucrose transporters (SUT), requiring channel-mediated  $\text{K}^+$  transport from the mesophyll apoplast into the phloem in order to maintain the activity of  $\text{H}^+$ -ATPases transporting  $\text{H}^+$  in the opposite direction. As much as 50–80% of K in the xylem may be recycled to the roots and sink


tissues via the phloem (Noa *et al.*, 2014). This recycling is increasingly being understood as a sophisticated system for whole-plant communication of nutritional demands, while at the same time providing a distribution system for sink tissues with variable demands for sucrose and nutrients along the phloem pathway (Gajdanowicz *et al.*, 2011; Dreyer *et al.*, 2017).

In exchange for  $\text{H}^+$ ,  $\text{K}^+$  is vital for establishing the transmembrane pH gradient required for ATP production during photosynthesis.  $\text{K}^+$  is also necessary for photosynthetic  $\text{CO}_2$  fixation through proper stomatal functioning and maintenance of chloroplast conditions with respect to turgor, pH and enzyme activities. In chloroplasts of *A. thaliana*, the  $\text{K}^+$  antiporter KEA3 is important for the ability to maintain high photosynthetic efficiency under fluctuating light conditions (Kunz *et al.*, 2014; Armbruster *et al.*, 2017).

### 1. Molecular–physiological responses to K deficiency

Root epidermal cells, particularly root hairs, represent the main sites for  $\text{K}^+$  uptake (Fig. 7.1). High-affinity  $\text{K}^+$  uptake at low external  $\text{K}^+$  concentrations is mediated by  $\text{K}^+/\text{H}^+$  symporters in the HAK/KUP/KT family, whereas voltage-gated  $\text{K}^+$  channels, for example AtAKT1 in *A. thaliana*, mediate low-affinity  $\text{K}^+$  uptake at higher external concentrations. The flexibility of  $\text{K}^+$  transport is also reflected in a diversity of functionally different channels, for example outward rectifiers ( $\text{K}^+$  release channels) and weak-rectifiers allowing both  $\text{K}^+$  efflux and uptake (Hedrich *et al.*, 2011). In xylem parenchyma cells in the stele,  $\text{K}^+$  loading is controlled by SKOR-like (Stellar  $\text{K}^+$  Outward Rectifier) channels, which release  $\text{K}^+$  into the xylem stream (Gaymard *et al.*, 1998). Stomatal closure is conveyed by  $\text{K}^+$  efflux via GORK (Guard Cell Outwards Rectifying  $\text{K}^+$ ) channels (Ivashikina *et al.*, 2005).

Plants have evolved sophisticated sensing and signalling systems that respond to K deficiency. These systems involve specific signal molecules, ROS,  $\text{Ca}^{2+}$ , phytohormones and microRNAs. Hyperpolarisation of the plasma membrane is the first detected response, occurring within a few minutes after a significant decrease in external  $\text{K}^+$  concentration. After prolonged deficiency (6–30 h), ROS generation and ethylene signalling are typical K deficiency responses (Behera *et al.*, 2017). K deficiency also triggers two successive and distinct  $\text{Ca}^{2+}$  signals involving a calcineurin B-like (CBL1)-CIPK23 Ca sensor–kinase complex activating AKT1  $\text{K}^+$  channels in postmeristematic tissues in the stele. This activation happens rapidly (minutes) after onset of low external  $\text{K}^+$  concentration. After several hours of deficiency, activation of AKT1 in the elongation and root hair differentiation zones takes place mediated via another  $\text{Ca}^{2+}$  signal (Behera *et al.*, 2017). MicroRNAs are likely to play a role in the  $\text{K}^+$  starvation responses linked to  $\text{Ca}^{2+}$ . By high-throughput sequencing and degradome analysis, ata-miR1432-5p was identified as a regulator participating in  $\text{Ca}^{2+}$  signalling associated with the low K response in barley (Zeng *et al.*, 2019). Transcriptional regulation of HAK1 and HAK5 in rice and *A. thaliana* involves the transcription factor ARF2, which after phosphorylation initiates HAK5 transcription at low K<sup>+</sup> but suppresses it during K sufficiency (Wang & Wu, 2017). A number of additional hormonal interactions are also activated during K



**Fig. 7** Schematic model showing key functions of potassium (K) in plant growth and physiology. (1) Root growth is dependent on sucrose loading into the phloem, which is a process that requires K. Hence, during K deficiency, lateral root growth is reduced. In addition, the endodermis of K-deficient roots become more suberised than that of nondeficient roots. This increased suberisation improves translocation of K to the shoots, as it improves water conductance and reduces leakage of K out of the stele. (2) Influx of K is necessary for guard cells to become fully turgid so that the stomatal pore opening can maintain CO<sub>2</sub> conductance. (3) K sustains photosynthesis by maintaining mesophyll cell and chloroplast integrity. (4) K is necessary for sucrose loading into the phloem, hence K deficiency leads to sucrose accumulation in source leaves. (5) Due to poor sucrose loading into the phloem, K deficiency leads to reduced sucrose supply to sink tissues like seeds, grains and fruits. (6) By maintaining proper CO<sub>2</sub> conductance, K suppresses the number of electrons available for side reactions with O<sub>2</sub>, which would otherwise create reactive oxygen species (ROS). (7) K facilitates proper xylem vessel maturation and functioning. K-deficient plants hence have reduced water conductance due to poor interaction between K and the pectin matrix of intercellular pectin. These effects tend to give K-deficient plants a slack appearance, as well as a lowered capability to handle water-related stress, that is drought and salinity.

stress, including ethylene, auxin, cytokinin and jasmonic acid (Schachtman, 2015; Wang & Wu, 2017; G. Li *et al.*, 2017). In *A. thaliana*, K starvation induces ethylene production, which also increases ROS levels, ultimately leading to upregulation of low-affinity K transporters. Some proteins and enzymes related to the jasmonic pathway are upregulated in wheat plants during K stress (G. Li *et al.*, 2017).

## 2. Coupling of visual symptoms and K functionalities

Inadequate K supply has a range of negative effects on plant growth and vigour as well as on plant resistance towards environmental stresses like drought and salinity. During K stress, organic acids are exuded (Zöorb *et al.*, 2014) and lateral root growth is inhibited. The root endodermis tends to become more suberised under K deficiency (Fig. 7.1), a process triggered by ABA (Barberon *et al.*, 2016) and affecting the translocation of K<sup>+</sup> (Chen *et al.*, 2019).

In the shoot, the earliest visual symptoms of K deficiency are chloroses at the tip of the oldest leaves, which rapidly turn into marginal necrosis spreading from the tip along the leaf margins (Table 1; Fig. 8). Leaves with these symptoms contain lower K levels in the leaf tips, compared with the basal parts of the leaf, and nonnecrotic symptoms can be reverted by re-supply of K. Some jasmonate-related genes are expressed in K-deficient leaf tips and may promote leaf senescence (Ueno *et al.*, 2018). These symptoms often appear together with bronzing, which is due to accumulation of polyamines (PAs), for example putrescine (Gupta *et al.*, 2013). The polyvalency of PAs, along with the stimulation of their synthesis during low pH, suggests that PAs may aid the cellular charge and pH buffering during K deficiency. The PAs, however, seem to play multiple additional roles in plants, including scavenging of ROS, activation of the antioxidant machinery, alleviation of stress-induced membrane injury and electrolyte leakage (Pottosin *et al.*, 2014).


Tip and marginal chlorosis/necrosis is closely linked to ROS production, or more precisely the handling of ROS in the chloroplasts. Light-exposed chloroplasts are the most potent

ROS generating organelles, producing up to 20-fold more ROS than for example mitochondria (Pottosin & Shabala, 2016). During K deficiency, photosynthesis is strongly decreased as a result of reduced stomatal conductance (Fig. 7.2), increased mesophyll resistance (Fig. 7.3) and lowered ribulose bisphosphate carboxylase activity. Furthermore, efficient photosynthesis is dependent on the export and utilisation of photoassimilates, and a typical feature of K deficiency is sucrose accumulation in the leaves (Fig. 7.4), typically accompanied by sucrose deficiency in the roots, which offer an explanation for the decrease in lateral root growth (Cakmak, 2005). K deficiency also leads to reduced sucrose supply to sink tissues like seeds, grains and fruits (Fig. 7.5). The severe reduction in CO<sub>2</sub> fixation caused by K deficiency results in an excess of electrons, ultimately leading to an increase in ROS via an intensified electron transfer to O<sub>2</sub> and ROS production (Fig. 7.6). As a result, K-deficient plants are sensitive to light intensity, which drives the development of K deficiency symptoms (Cakmak, 2005).


K-deficient plants are sensitive to water limitation, and the leaves of affected plants typically appear slack. Low K status negatively effects stomatal closure, as complete closure of stomata requires backpressure from fully turgid epidermal cells in order to close properly (Fig. 7.2). Another reason for the poor control of stomatal closure is increased ethylene production, which interferes with ABA signalling during K deficiency. Potassium deficiency leads to swelling of the pectin matrix in xylem cell walls and a reduction in the porosity of xylem pit membranes, restricting transpiration via a decreased hydraulic conductance (Fig. 7.7) (Anschatz *et al.*, 2014).

## VI. Calcium

Calcium is essential for higher plants because it is required as a structural element and plays an essential roles in cellular signalling. The role of Ca in signalling is achieved by transducing, integrating and multiplying incoming signals, thus linking environmental stimuli with physiological responses (Thor, 2019). By far the major part of Ca is present in the cell walls, on the outside of cell membranes and in cell organelles. The concentration of free Ca<sup>2+</sup>



**Fig. 8** Potassium (K) deficiency always appears on the oldest leaves first and spreads as a necrosis from the tip along the margins (marginal necrosis) and subsequently appears as interventional necrotic spots on the entire leaf. (a) Marginal necrosis in white clover. (b) Marginal necrosis in maize. (c) Intervenous necrosis in tomato.



ions in the cytosol has an extremely low ( $< 0.1 \mu\text{M}$ ) resting value, but with transient spikes. Ca is virtually immobile in the phloem, implying that there is very limited translocation of Ca from old plant parts (source organs) to young leaves, shoots and fruits (sink organs). These organs therefore depend on the xylem for delivery of Ca.

### 1. The role of Ca as a structural element

The functions of Ca as structural element are primarily related to the fact that Ca confers rigidity to the cell wall system and stabilises it (Fig. 9.1). During the biosynthesis of cell walls, acidic pectin residues (e.g. galacturonic acid) are secreted from the cells as methyl

**Fig. 9** Schematic model showing how calcium (Ca) affects plant growth and physiology. (1) Structural element in shoots and roots. Ca is bound to pectic carboxyl groups in the cell wall and to negative charges on cell membranes, thereby stabilising them. Ca also plays an essential role in the regulation of the microtubule (MT) cytoskeleton mediated via the plant-specific IQ67 DOMAIN (IQD) family in interaction with calmodulin (CaM)-dependent Ca signalling. (2) Signal transduction in shoots and roots. Stimuli are perceived via different membrane receptors leading to transient perturbations of the cytosolic free  $\text{Ca}^{2+}$  concentration via influx mediated by cyclic nucleotide-gated channels (CNGCs), ionotropic glutamate receptor homologues (GLRs), mid1-complementing activity proteins (MCAs), and calcium permeable stress-gated cation channels (CSCs)/reduced hyperosmolarity-induced  $[\text{Ca}^{2+}]$  increase channels (OSCas). P-type  $\text{Ca}^{2+}$ -ATPases mediate  $\text{Ca}^{2+}$  removal from the cytosol via efflux or via sequestration in different organelles (endoplasmic reticulum, Golgi apparatus and plastids).  $\text{Ca}^{2+}$  can also be sequestered in the vacuole by  $\text{Ca}^{2+}$ -ATPases or  $\text{Ca}^{2+}/\text{H}^+$  exchangers (CAXs), the latter driven by ATP or pyrophosphate (PPi).  $\text{Ca}^{2+}$  is released from the vacuole via  $\text{Ca}^{2+}$  permeable channels. Changes in cytosolic  $\text{Ca}^{2+}$  trigger a cascade of downstream processes mediated via sensor relays, viz. calmodulins (CaM) and CaM-like proteins, and responders, namely  $\text{Ca}^{2+}$ -dependent protein kinases (CPKs/CDPKs) and calcineurin B-like (CBL) proteins, which control the activity of CBL-interacting protein kinases (CIPKs). (3) Nutrient sensing. Changes in external nutrient availability affect the cytosolic  $\text{Ca}^{2+}$  concentration and activate or repress nutrient transporters via the  $\text{Ca}^{2+}$ -dependent CBL–CIPK pathway with CIPK23 activating transporters/channels for  $\text{K}^+$  (HAK5, AKT1/AtKC1),  $\text{NO}_3^-$  (NRT1.1/NPF6.3) and  $\text{Fe}^{2+}$  (IRT) upon starvation, while inhibiting  $\text{NH}_4^+$  transport via AMT1;2 in response to high  $\text{NH}_4^+$ . Plant phosphorus starvation may negatively affect the oscillations in cytosolic  $\text{Ca}^{2+}$  concentration caused by mechanical, salt, osmotic, and oxidative stress (adapted from Kudla *et al.*, 2018). (4) Translocation. The Ca concentration in the phloem is very low. Root-to-shoot Ca translocation is therefore determined by the xylem Ca concentration and xylem sap flow related to transpiration and root pressure. With respect to fruits, Ca import is controlled by xylem sap inflow as determined by fruit water uptake and loss driven by cell expansion and transpiration. Xylem sap flow into fruits is affected by xylem vessel development, vessel diameter and connectivity. Ca is bound in the cell walls of xylem vessels and also in intervessel pits, where de-esterification of pectin enables formation of Ca cross-linked gels affecting pit porosity, xylem flow and Ca distribution. There may likely also be some phloem Ca transport in fruits where xylem dysfunction occurs early during fruit development. Adapted from Hocking *et al.* (2016). (5) Examples of Ca deficiency symptoms: blossom end rot in tomato and pepper, tipburn in lettuce and bitter bit in apple.

esters, which are subsequently de-esterified by pectin methylesterase, thereby providing carboxyl groups which bind Ca via covalent and ionic bonds (Hocking *et al.*, 2016). When associated with Ca, the cell wall system is protected from degradation by polygalacturonases (pectin depolymerase). Dicot plant species have a much higher Ca requirement than monocots due to their higher content of pectate with a higher density of binding sites for  $\text{Ca}^{2+}$  (White *et al.*, 2018). In addition to the cell wall system,  $\text{Ca}^{2+}$  stabilises cell membranes via the formation of bridges with phosphate and carboxylate groups in lipids and proteins (Fig. 9.1). Calcium deficiency therefore leads to increased plasma membrane permeability, resulting in cell death in the apical meristem and growth cessation.

## 2. Functions of Ca in signalling and nutrient sensing

The function of  $\text{Ca}^{2+}$  as secondary messenger in signalling processes is mediated via transient perturbations of the cytosolic free  $\text{Ca}^{2+}$  concentration. Various external factors such as light, temperature, gravity, touch and pathogen infection in combination with internal hormonal signals, are perceived via different membrane receptors, leading to specific spatio-temporal patterns of increasing cytosolic  $\text{Ca}^{2+}$  concentrations that trigger a cascade of downstream processes (Fig. 9.2). The generation of these  $\text{Ca}^{2+}$  signals relies on influx of  $\text{Ca}^{2+}$  via coordinated actions of distinct  $\text{Ca}^{2+}$  channels (Tang & Luan, 2017; Demidchik *et al.*, 2018) (Fig. 9.2). In parallel with the actions of  $\text{Ca}^{2+}$  influx channels,  $\text{Ca}^{2+}$  efflux across the plasma membrane and  $\text{Ca}^{2+}$  sequestration into intracellular organelles are important to maintain the transient nature of the  $\text{Ca}^{2+}$  signals. P-type  $\text{Ca}^{2+}$ -ATPases either pump  $\text{Ca}^{2+}$  out of the cell across the plasma membrane or sequester it into organelles so that  $\text{Ca}^{2+}$  is removed from the cytosol.

The specific  $\text{Ca}^{2+}$  signatures formed in response to various environmental stimuli lead in turn to some downstream processes, which amplify and integrate the  $\text{Ca}^{2+}$  signal via interaction with a number of  $\text{Ca}^{2+}$ -binding proteins in the cytosol acting as  $\text{Ca}^{2+}$

sensors (Fig. 9.2) (Bender *et al.*, 2018). These sensors can be classified into sensor relays and sensor responders (Kudla *et al.*, 2018). Sensor relays include calmodulins (CaMs) and CaM-like proteins (CMLs), which upon  $\text{Ca}^{2+}$  binding undergo a conformational change that is subsequently relayed to an interacting protein (La Verde *et al.*, 2018). Two classes of  $\text{Ca}^{2+}$  sensor responders have been characterised, namely the  $\text{Ca}^{2+}$ -dependent protein kinases (CPKs/CDPKs) and the calcineurin B-like (CBL) proteins, which control the activity of CIPKs. Members of the CDPK and CIPK protein families may target receptors responsive to abscisic acid, thereby linking to signal transduction mediated by hormones (Waadt *et al.*, 2017; Kudla *et al.*, 2018; Vighi *et al.*, 2019).

$\text{Ca}^{2+}$  plays a central role in nutrient sensing and in adaptation to changes in nutrient status. The link between external nutrient availability and transport activities is constituted by the  $\text{Ca}^{2+}$ -dependent CBL–CIPK pathway with CIPK23 functioning as a key master regulator of  $\text{K}^+$  and  $\text{NO}_3^-$ , as well as Fe, Mg and  $\text{NH}_4^+$  homeostasis (Fig. 9.3) (Riveras *et al.*, 2015; K.-h. Liu *et al.*, 2017; Kudla *et al.*, 2018; X. Wang *et al.*, 2018). As also mentioned in Section V, CIPK23 phosphorylates and activates the low-affinity  $\text{K}^+$  transporter AKT1 as well as the high-affinity HAK5  $\text{K}^+$  transporter. With respect to  $\text{NO}_3^-$  uptake, CIPK-mediated changes in phosphorylation status affect both high- and low-affinity uptake via the membrane-located  $\text{NO}_3^-$  transceptor NRT1.1/NPF6.3. Downstream of NRT1.1/NPF6.3, CPK Ca-dependent kinases may induce  $\text{NO}_3^-$  assimilatory and regulatory genes, thus constituting the molecular link between major players in the  $\text{NO}_3^-$  signalling pathway. CIPK23 also activates Fe transport in *A. thaliana* roots, while inhibiting AMT1;2 (Ammonium transporter 1;2)-mediated  $\text{NH}_4^+$  uptake (Tian *et al.*, 2016; Straub *et al.*, 2017).

Calcium signalling is not only involved in intracellular regulation, but may also contribute to long-range signal propagation and regulation of physiological responses (Kudla *et al.*, 2018). Besides mobile small molecules (e.g. hormones) and macromolecules (e.g. proteins, small RNAs and mRNAs), electrical signals linked to

waves of  $\text{Ca}^{2+}$  and ROS may traverse the plant and trigger systemic responses (Choi *et al.*, 2016). Over the past decade there has been limited progress in revealing the molecular mechanisms involved in regulating the transport and loading/unloading of  $\text{Ca}^{2+}$  in the phloem (Fig. 9.4). It needs to be further revealed why  $\text{Ca}^{2+}$  concentrations generally are low in the phloem and what the physiological implications of high  $\text{Ca}^{2+}$  would be.  $\text{Ca}^{2+}$  loading of sieve tubes might occur through plasmodesmata. However, increasing cytoplasmic  $\text{Ca}^{2+}$  tends to close plasmodesmata and the transport of chelated  $\text{Ca}^{2+}$  from the symplast of the phloem companion cells through plasmodesmata also seems restricted (Sager & Lee, 2014).

### 3. Functional properties of Ca in relation to plant growth and development

Proper responses of plants to changes in environmental conditions will, in most cases, require reprogramming of the growth rate as determined by cell division and subsequent cell expansion. The adjustment in growth rate involves dynamic reorganisation of the cytoskeleton, that is the complex network of interlinked protein filaments that extends from the cell nucleus to the cell membrane of all cells. Part of the cytoskeleton consists of microtubules (polymers of tubulin) that provide structure and shape to the cytoplasm, thus coordinating and controlling plant cell shape and cell growth. Ca plays an essential role in the regulation of the microtubule cytoskeleton and  $\text{Ca}^{2+}$ -mediated regulation of microtubule-associated proteins provides hubs for cross-talk with other signalling pathways (Fig. 9.1) (Kölling *et al.*, 2019). The regulation of cell function, shape and growth has been suggested to occur via the plant-specific IQ67 DOMAIN (IQD) family, which may act as platform proteins for integration of CaM-dependent  $\text{Ca}^{2+}$ -signalling (Bürstenbinder *et al.*, 2017; Kölling *et al.*, 2019).

Via its role in signalling and stabilisation of cell walls and membranes, Ca plays a crucial role for root growth and adaptation. Starvation of P, but not N, strongly dampens oscillations in cytosolic  $\text{Ca}^{2+}$  concentration caused by mechanical, salt, osmotic and oxidative stress (Matthus *et al.*, 2019). Gradients and oscillations in  $\text{Ca}^{2+}$  concentration also play a key role in the growth of root hairs (S. S. Zhang *et al.*, 2017; Kwon *et al.*, 2018) as well as for establishment of symbiotic associations (Miller *et al.*, 2013).

### 4. Calcium deficiency symptoms linked to functional properties

As Ca in the phloem is maintained at a very low concentrations, different parts of the shoot depend on transpiration or root pressure to provide Ca via the xylem (Fig. 9.4). For organs with low transpiration, such as fruits and internal heart leaves in vegetables, root pressure may be insufficient leading to a range of unique Ca deficiency symptoms (Song *et al.*, 2018). These symptoms reflect the essential role of Ca for proper formation of cell walls and for protecting them from degradation by cell wall-degrading enzymes. Typical foliar symptoms of Ca deficiency are necrotic lesions on leaf

margins and tips, brownish leaf veins and leaf deformities (Table 1). In dicots, this leads to cup-shaped leaves while, in grasses, leaf spirals are a typical deformity. Upon continued Ca deficiency, apical meristems die.

Special problems with Ca deficiency occur in fruits. Fruit Ca uptake is believed to be determined by the  $\text{Ca}^{2+}$  concentration in the xylem sap, as well as xylem sap influx, which is related to fruit stomatal density, transpiration, root pressure and growth (Fig. 9.4) (Hocking *et al.*, 2016). However, in many fruits, permanent loss of xylem functionality occurs. Under these circumstances, the phloem may constitute a transport pathway for Ca mobilised from the pedicel (Song *et al.*, 2018).

The deficiency symptoms for Ca are so characteristic that they carry specific names in specific crops (Figs 9.5, 10). Examples are blossom end rot in tomato, pepper and zucchini, in which sunken, necrotic pits appear at the blossom end of the fruits (Djangsou *et al.*, 2019). The leaf edges of cabbage, lettuce and brussels sprouts experiencing Ca deficiency are likely to develop necrotic edges, called tipburn. Lee *et al.* (2016) demonstrated that the relationship between intracellular Ca concentration and tipburn in *Brassica oleracea* differed between genotypes and were correlated with differential expression of a number of transporter and stress response genes. In apple, Ca deficiency is called bitter bit, reflecting that the fruit skin develops pits and that brown spots appear on the skin and/or in flesh, resulting in a bitter taste (Buti *et al.*, 2018; Yu XM *et al.*, 2018). Calcium deficiency in carrot leads to cavity spot, characterised by oval necrotic spots developing into craters. The proportion of Ca pectate in the cell wall has great importance for resistance against fungal and bacterial pathogens and thereby for the storage and postharvest quality of fruits (Fig. 9.5) (Winkler & Knoche, 2019).

## VII. Magnesium

Depending on the nutritional status, about 5–25% of Mg in plants is present in the chloroplasts, where Mg constitutes the central element of the tetrapyrrole ring in chlorophyll (Fig. 11.1), fine tuning its electric properties. Mg thus participates in photon capturing and transfer of excitation energy from the LHCs to the PSII reaction core.

Besides the mainly covalent binding in chlorophyll, Mg forms ion bonds with negative charges in the cell walls. Differences in cell wall chemistry and cation exchange capacity of the cell walls of different species results in a distinct stoichiometric relationship between shoot concentrations of Mg and Ca in major groups of angiosperm species (White *et al.*, 2018).

The key functionalities of Mg are mainly related to its strong electropositivity, which provides capacity to interact electrostatically with nucleophilic ligands. The  $\text{Mg}^{2+}$  ion forms ternary complexes with enzymes, thereby establishing a specific geometry between the enzyme and the substrate and increasing the efficiency of the catalytic reaction (Fig. 11.2). Magnesium ions exhibit fast ligand exchange kinetics, which means they can easily be replaced by other divalent metal ions, including  $\text{Mn}^{2+}$ ,  $\text{Ca}^{2+}$ ,  $\text{Fe}^{2+}$ ,  $\text{Co}^{2+}$ ,  $\text{Cu}^{2+}$ , and  $\text{Zn}^{2+}$ . Substitution of  $\text{Mg}^{2+}$  with other ions typically changes the catalytic rate of the enzyme and in many cases it also



**Fig. 10** During the vegetative stage, calcium(Ca) deficiency always affects the roots before the shoot. Roots turn brown and the tissue disintegrates. Fruits are also a main target for Ca deficiency. (a) Brown and disintegrating roots in barley. (b) Blossom end rot necrotic plaques in tomato.

changes the functional role of the enzyme. RuBisCo (Ribulose-1,5-bisphosphate carboxylase/oxygenase) represents an illustrative example of this, as differential binding of  $Mg^{2+}$  and  $Mn^{2+}$  changes both the catalytic rate and the substrate preference of the protein (Schmidt & Husted, 2019). When RuBisCo binds to  $Mg^{2+}$ , carboxylation is favoured and proceeds 4–11 times faster than oxygenation, but when RuBisCo binds to  $Mn^{2+}$ , carboxylation and oxygenation occur at similar rates. The ability of  $Mg^{2+}$  to perform ligand exchange is essential for the coupling of ATP with substrates and enzymes in phosphorylation processes, energy-demanding transport processes and ATP synthesis. An important example is  $Mg\text{-ATP}$ , which is the substrate for the  $H^+$ -ATPase that drives the export of sucrose from the leaf cells into the phloem. This makes  $Mg^{2+}$  critical for the phloem loading of sucrose and thus for long-distance transport of photosynthates and carbohydrate partitioning between source and sink tissues (Fig. 11.3) (Träkner *et al.*, 2018).


### 1. Molecular and physiological responses to Mg deficiency

Reflecting its importance in many central reactions, cytosolic  $Mg^{2+}$  levels are tightly regulated. The concentration of free  $Mg^{2+}$  in the cytosol has been estimated to be around 0.4 mM compared with around 4 mM total Mg, the latter including  $Mg^{2+}$  complexed with ATP and other ligands. The major family of  $Mg^{2+}$  transport proteins in higher plants belongs to homologues of the bacterial CorA  $Mg^{2+}$  transporters, the so-called MGTs (Fig. 11.4) (Tang & Luan, 2017). In *A. thaliana*, rice and maize, 10–12 members of the MGT family have been identified (Li *et al.*, 2016; Sun *et al.*, 2017). A higher number of MGT genes, viz. 36, were identified in *B. napus*, clustering in five groups and showing clear homology with MGT genes in *A. thaliana* and rice (Zhang *et al.*, 2019). AtMGT6, and its homologue ZmMGT10 in maize, appear to mediate high-affinity  $Mg^{2+}$  uptake, hence is upregulated under Mg-limited conditions. Knockout of AtMGT6 impaired *A. thaliana* growth under Mg-limited conditions (Yan *et al.*, 2018). AtMGT7 is


preferentially expressed in the roots and also important for plant adaptation to low Mg conditions. The double mutant *mgt6 mgt7* showed more severe phenotypes compared with single mutants under both low- and high-Mg conditions, indicating that these two MGT-type transporters play additive roles in controlling plant  $Mg^{2+}$  homeostasis under a wide range of external Mg concentrations (Yan *et al.*, 2018). Transgenic *A. thaliana* plants overexpressing *ZmMGT10* exhibited longer root length, higher shoot weight, chlorophyll content and  $Mg^{2+}$  uptake under low Mg conditions, thus suggesting that *ZmMGT10* is essential for plant growth and development (H. Li *et al.*, 2017).

After uptake from the soil,  $Mg^{2+}$  is loaded into the xylem for long-distance transport to the shoots.  $Mg^{2+}$  is also mobile in the phloem and therefore easily translocated to fruits, seeds and tubers, as well as remobilised from old to young tissues, the latter promoting reutilisation under Mg-limited conditions. The phloem mobility also implies continuous recycling of  $Mg^{2+}$  between shoots and roots. Relevant transporters for these processes have not been identified. AtMGT2 and AtMGT3 are assumed to be involved in  $Mg^{2+}$  transport into vacuoles and may be assisted by AtMHX, which is an  $Mg^{2+}/H^+$  exchanger (Tang & Luan, 2017).

In order for Mg to fulfil its role as a central atom in chlorophyll, it needs to be transported into the chloroplasts. In *A. thaliana*, MGT10 is localised in the envelope of chloroplasts and expressed mainly in vascular bundles, where it is required for the transport of  $Mg^{2+}$  into the stroma, particularly under high light conditions (Sun *et al.*, 2017). In the chloroplasts, the first step of chlorophyll biosynthesis involves insertion of  $Mg^{2+}$  into the porphyrin ring, a process catalysed by Mg chelatase. The Mg-dechelatases NYE1/SGR1 and NYE2/SGR2 (NON-YELLOWING/STAY-GREEN) are responsible for the breakdown of chlorophyll and release of  $Mg^{2+}$  (Shimoda *et al.*, 2016). Loss of function of both *NYE1* and *NYE2* resulted in a nearly complete retention of chlorophyll during leaf senescence while at the same time enhancing photooxidative damage to maturing seeds in *A. thaliana* and soybean (Z. Li *et al.*,



**Fig. 11** Schematic model showing how magnesium (Mg) affects plant growth and physiology. (1) Mg is a constituent of chlorophyll and also plays an important role in activation of enzymes, especially RuBisCo, catalysing CO<sub>2</sub> fixation in the Calvin cycle. (2) Mg is involved in a range of phosphorylation processes via creating a ternary complex between ATP, substrate and enzyme. Mg also bridges ADP and ATP synthase, leading to synthesis of ATP. (3) Phloem loading of sucrose is mediated by proton-sucrose transport where Mg couples ATP and the H<sup>+</sup>-ATPase. Physiological consequences of Mg deficiency include reduced CO<sub>2</sub> fixation and increased accumulation of sucrose and starch leading to increased production of reactive oxygen species and photo-oxidation of the photosynthetic apparatus. (4) Mg<sup>2+</sup> transport proteins include homologues of the bacterial CorA Mg<sup>2+</sup> transporters, the so-called MGTs. In maize, ZmMGT10, a homologue of Arabidopsis AtMGT6, appears to mediate high-affinity Mg<sup>2+</sup> uptake and is upregulated under Mg limited conditions. Cations such as K<sup>+</sup>, NH<sub>4</sub><sup>+</sup>, Ca<sup>2+</sup>, Mn<sup>2+</sup>, Al<sup>3+</sup>, H<sup>+</sup> and Na<sup>+</sup> may severely inhibit root uptake of Mg<sup>2+</sup>, thus leading to deficiency. Adapted from Tang & Luan (2017). (5) Mg deficiency leads to strong inhibition of root growth due to the reduced translocation of photosynthates. (6) Mg is highly phloem mobile and is remobilised from old to young tissues, thus promoting reutilisation under Mg-limited conditions. Visible symptoms of Mg deficiency include chlorosis gradually developing from the tip of fully expanded older leaves. Patches of interveinal chlorosis may appear in different patterns, for example light, irregular transverse bands on the leaves of cereal species, eventually accompanied by necrosis between the leaf veins.



**Fig. 12** Magnesium (Mg) deficiency always appears on oldest leaves first and initially as interveinal chlorosis which slowly develops into necrotic plaques. (a) Barley. (b) Maize. (c) Oilseed rape. (d) Tomato.

2017). The Mg chelatase gene *STAY-GREEN* (*OsSGR*) is involved in chlorophyll degradation and internal remobilisation of Mg in rice. Under Mg-limited conditions, the expression of *OsSGR* is feedback regulated by ROS, fine tuning chlorophyll degradation in mid-aged leaves with relatively high Mg content and photosynthetic capacity, thereby contributing to photooxidative protection (Peng *et al.*, 2019).

In Mg-deficient plants, photoassimilates accumulate in source leaves before photosynthesis is affected by Mg deficiency. This results in an excessive accumulation of carbohydrates and enhanced production of ROS in the chloroplasts. The impaired allocation of photoassimilates to roots implies a significant reduction of root growth under Mg deficiency (Fig. 11.5). Besides the depletion of carbohydrates, Mg deficiency may increase the production of NO and ethylene in roots, leading to increased root hair development (M. Liu *et al.*, 2017). A recent metabolomics study showed distinct C and N metabolic responses to Mg deficiency in soybean leaves and roots (Yang *et al.*, 2017). Most amino acids showed pronounced accumulation in the leaves, while most organic acids were significantly decreased in the roots.

## 2. Coupling of visual symptoms and nutrient functionalities

The first visual leaf symptom of Mg deficiency is interveinal chlorosis of fully expanded, older leaves (Table 1; Fig. 12). The chlorotic patches may appear in different patterns, for example forming light, irregular transverse bands on the leaves of cereal species ('tiger striped' leaves). The chlorosis typically develops gradually and interveinously from the tip of fully expanded old leaves and is eventually accompanied by purpling and brown-withering (necrosis) between the leaf nerves (Fig. 11.6). Visual symptoms of Mg deficiency typically first appear on the older leaves (acropetal stratification), reflecting that Mg<sup>2+</sup> is highly phloem mobile.

At the physiological and biochemical levels, the responses to Mg deficiency include accumulation of sucrose and starch and reduced CO<sub>2</sub> fixation. These changes reflect the essential functions of Mg in chlorophyll biosynthesis (Fig. 11.1), as well as in the export of photosynthates from the chloroplasts and leaf cells (Fig. 11.3). The accumulation of sucrose and starch in the chloroplasts in turn leads to increased production of ROS, photo-oxidation of the

photosynthetic apparatus and upregulation of photoprotective mechanisms (Tränkner *et al.*, 2018). An accompanying consequence of the reduced export of photoassimilates is the strong inhibition of root growth (Fig. 11.5). Mg deficiency also causes accumulation of low molecular N compounds, which is due to the essential function of Mg in protein synthesis, where Mg constitutes a bridging element for the aggregation of ribosome subunits.

## VIII. Conclusion and perspectives

The macronutrients each have their characteristic deficiency symptoms that can be linked to specific physiological functions. The challenge ahead will be to utilise this mechanistic knowledge to develop sensitive and robust techniques for early diagnosis and correction of nutrient deficiencies. Advancement in imaging technology has greatly accelerated the application of image-based plant nutrition analysis (Li *et al.*, 2020) based on chlorophyll content, foliage reflectance and transmittance (Tan *et al.*, 2018; Zheng *et al.*, 2018). However, interactions between several nutrient deficiencies (Hou *et al.*, 2020) and other environmental biotic and abiotic factors may bias imaging techniques. In such cases, diagnostic methods based on a functional role of the nutrient under consideration may potentially be more specific and robust. An example using mechanistic knowledge for diagnostic purposes is the use of time-resolved chlorophyll a fluorescence as a unique tool to monitor bioactive P in plants and to detect latent P deficiency (Frydenvang *et al.*, 2015; Carstensen *et al.*, 2018, 2019). Early detection of nitrogen status in plants using Raman spectroscopy was recently reported by Huang *et al.* (2020). Further developments within these areas may pave the road for a future where crops can be fertilised according to their physiological demand rather than based on the classical practice, where bulk fertilisation often leads to application of either too little or too much, with negative impacts on both environment, climate and the economy of farmers.

## Acknowledgements

This work was supported by grants from the Independent Research Fund Denmark, Technology and Production Sciences (grant numbers DFF-7017-00082 and DFF-9041-00022).

## ORCID

- Thomas Christian deBang  <https://orcid.org/0000-0002-5019-6422>
- Kristian Holst Laursen  <https://orcid.org/0000-0001-7900-3324>
- Søren Husted  <https://orcid.org/0000-0003-2020-1902>
- Daniel Pergament Persson  <https://orcid.org/0000-0003-3976-190X>
- Jan Kofod Schjoerring  <https://orcid.org/0000-0002-2852-3298>

## References

- Aghajanzadeh T, Hawkesford MJ, De Kok LJ. 2014. The significance of glucosinolates for sulfur storage in *Brassicaceae* seedlings. *Frontiers in Plant Science* 5: 704.
- Ahmed I, Sebastian A, Prasad MNV, Kirti PB. 2019. Emerging trends in transgenic technology for phytoremediation of toxic metals and metalloids. In: Prasad MNV, ed. *Transgenic plant technology for remediation of toxic metals and metalloids*. London, UK: Academic Press Ltd - Elsevier Science Ltd, 43–62.
- Anschutz U, Becker D, Shabala S. 2014. Going beyond nutrition: regulation of potassium homoeostasis as a common denominator of plant adaptive responses to environment. *Journal of Plant Physiology* 171: 670–687.
- Araya T, Miyamoto M, Wibowo J, Suzuki A, Kojima S, Tsuchiya YN, Sawa S, Fukuda H, von Wieren N, Takahashi H. 2014. CLE-CLAVATA1 peptide-receptor signaling module regulates the expansion of plant root systems in a nitrogen-dependent manner. *Proceedings of the National Academy of Sciences, USA* 111: 2029–2034.
- Armbuster U, Galvis VC, Kunz HH, Strand DD. 2017. The regulation of the chloroplast proton motive force plays a key role for photosynthesis in fluctuating light. *Current Opinion in Plant Biology* 37: 56–62.
- Balzergue C, Darteville T, Godon C, Laugier E, Meirimler C, Teulon J-M, Creff A, Bissler M, Brouchoud C, Hagège A et al. 2017. Low phosphate activates STOP1-ALMT1 to rapidly inhibit root cell elongation. *Nature Communications* 8: 15300.
- Barberon M, Vermeer JEM, De Bellis D, Wang P, Naseer S, Andersen TG, Humbel BM, Nawrath C, Takano J, Salt DE et al. 2016. Adaptation of root function by nutrient-induced plasticity of endodermal differentiation. *Cell* 164: 447–459.
- Behera S, Long Y, Schmitz-Thom I, Wang XP, Zhang CX, Li H, Steinhorst L, Manishankar P, Ren XL, Offenborn JN et al. 2017. Two spatially and temporally distinct  $\text{Ca}^{2+}$  signals convey *Arabidopsis thaliana* responses to  $\text{K}^+$  deficiency. *New Phytologist* 213: 739–750.
- Bender KW, Zielinski RE, Huber SC. 2018. Revisiting paradigms of  $\text{Ca}^{2+}$  signaling protein kinase regulation in plants. *Biochemical Journal* 475: 207–223.
- Bhosale R, Giri J, Pandey BK, Giehl RFH, Hartmann A, Traini R, Truskina J, Leftley N, Hanlon M, Swarup K et al. 2018. A mechanistic framework for auxin dependent *Arabidopsis* root hair elongation to low external phosphate. *Nature Communications* 9: 1409.
- Bloem E, Haneklaus S, Schnug E. 2005. Influence of nitrogen and sulfur fertilization on the alliin content of onions and garlic. *Journal of Plant Nutrition* 27: 1827–1839.
- Bougueny E, Brun F, Meynard D, Kubeš M, Pervent M, Leren S, Lacombe B, Krouk G, Guiderdoni E, Zážimalová E et al. 2015. Multiple mechanisms of nitrate sensing by *Arabidopsis* nitrate transceptor NRT1.1. *Nature Plants* 1: 15015.
- Bürstenbinder K, Möller B, Plötner R, Stamm G, Hause G, Mitra D, Abel S. 2017. The IQD family of calmodulin-binding proteins links calcium signaling to microtubules, membrane subdomains, and the nucleus. *Plant Physiology* 173: 1692–1708.
- Buti M, Sargent DJ, Bianco L, Magnago P, Velasco R, Colgan RJ. 2018. A study of gene expression changes at the Bp-2 locus associated with bitter pit symptom expression in apple (*Malus pumila*). *Molecular Breeding* 38: 85.
- Cakmak I. 2005. The role of potassium in alleviating detrimental effects of abiotic stresses in plants. *Journal of Plant Nutrition and Soil Science* 168: 521–530.
- Carstensen A, Herdean A, Schmidt SB, Sharma A, Spetea C, Pribil M, Husted S. 2018. Mapping the impacts of phosphorus deficiency on the photosynthetic electron transport chain in barley. *Plant Physiology* 177: 1–14.
- Carstensen A, Szameitat AE, Frydenvang J, Husted S. 2019. Chlorophyll a fluorescence analysis can detect phosphorus deficiency under field conditions and is an effective tool to prevent grain yield reductions in spring barley (*Hordeum vulgare*, L.). *Plant and Soil* 434: 79–91.
- Chen A, Husted S, Salt DE, Schjoerring JK, Persson DP. 2019. The intensity of manganese deficiency strongly affects root endodermal suberization and ion homeostasis. *Plant Physiology* 181: 729–742.
- Chen KE, Chen HY, Tseng CS, Tsay YF. 2020. Improving nitrogen use efficiency by manipulating nitrate remobilization in plants. *Nature Plants* 6: 1126–1135.
- Chen ZH, Wang YZ, Wang JW, Babla M, Zhao CC, Garcia-Mata C, Sani E, Differ C, Mak M, Hills A et al. 2016. Nitrate reductase mutation alters potassium nutrition as well as nitric oxide-mediated control of guard cell ion channels in *Arabidopsis*. *New Phytologist* 209: 1456–1469.
- Choi WG, Hilleary R, Swanson SJ, Kim SH, Gilroy S. 2016. Rapid long-distance electrical and calcium signaling in plants. *Annual Review in Plant Biology* 67: 287–307.
- Courbet G, Gallardo K, Vigani G, Brunel-Muguet S, Trouverie J, Salon C, Ourry A. 2019. Disentangling the complexity and diversity of crosstalk between sulfur and other mineral nutrients in cultivated plants. *Journal of Experimental Botany* 70: 4183–4196.
- Cui LG, Shan JX, Shi M, Gao JP, Lin HX. 2014. The miR156-SPL9-DFR pathway coordinates the relationship between development and abiotic stress tolerance in plants. *The Plant Journal* 80: 1108–1117.
- Dai X, Wang Y, Yang A, Zhang WH. 2012. OsMYB2P-1, an R2R3 MYB transcription factor, is involved in the regulation of phosphate-starvation responses and root architecture in rice. *Plant Physiology* 159: 169–183.
- Delhaize E, Ma JF, Ryan PR. 2012. Transcriptional regulation of aluminum tolerance genes. *Trends in Plant Science* 17: 341–348.
- Del-Saz NF, Romero-Munar A, Cawthray GR, Palma F, Aroca R, Baraza E, Florez-Sarasua I, Lambers H, Ribas-Carbó M. 2018. Phosphorus concentration coordinates a respiratory bypass, synthesis and exudation of citrate, and the expression of high-affinity phosphorus transporters in *Solanum lycopersicum*. *Plant, Cell & Environment* 41: 865–875.
- Demidchik V, Shabala S, Isayenkov S, Cuin TA, Pottosin I. 2018. Calcium transport across plant membranes: mechanisms and functions. *New Phytologist* 220: 49–69.
- Djangsou H, Enrico F, Domenico R, Matteo B. 2019. Blossom end-rot in tomato (*Solanum lycopersicum* L.): a multi-disciplinary overview of inducing factors and control strategies. *Scientia Horticulturae* 249: 49–58.
- Dong YH, Silbermann M, Speiser A, Forieri I, Linster E, Poschet G, Samami AA, Wanatabe M, Sticht C, Teleman AA et al. 2017. Sulfur availability regulates plant growth via glucose-TOR signalling. *Nature Communications* 8: 1174.
- Dreyer I, Gomez-Porras JL, Riedelsberger J. 2017. The potassium battery: a mobile energy source for transport processes in plant vascular tissues. *New Phytologist* 216: 1049–1053.
- Dubouset L, Abdallah M, Desfeux AS, Etienne P, Meuriot F, Hawkesford MJ, Gombert J, Segura R, Bataille MP, Reze S et al. 2009. Remobilization of leaf S compounds and senescence in response to restricted sulphate supply during the vegetative stage of oilseed rape are affected by mineral N availability. *Journal of Experimental Botany* 60: 3239–3253.
- Dung Tien L, Ha Duc C, Ngoc Quynh L. 2016. Improving nutritional quality of plant proteins through genetic engineering. *Current Genomics* 17: 220–229.
- Forieri I, Sticht C, Reichelt M, Grerz N, Hawkesford MJ, Malagoli M, Wirtz M, Hell R. 2017. System analysis of metabolism and the transcriptome in *Arabidopsis thaliana* roots reveals differential co-regulation upon iron, sulfur and potassium deficiency. *Plant, Cell & Environment* 40: 95–107.
- Forieri I, Wirtz M, Hell R. 2013. Toward new perspectives on the interaction of iron and sulfur metabolism in plants. *Frontiers in Plant Science* 4: 357.
- Frydenvang J, van Maarschalkerweerd M, Carstensen A, Mundus S, Schmidt SB, Pedas PR, Laursen KH, Schjoerring JK, Husted S. 2015. Sensitive detection of phosphorus deficiency in plants using chlorophyll a fluorescence. *Plant Physiology* 169: 353–361.

- Gajdanowicz P, Michard E, Sandmann M, Rocha M, Correa LGG, Ramirez-Aguilar SJ, Gomez-Porras JL, Gonzalez W, Thibaud J-B, van Dongen JT et al. 2011. Potassium ( $K^+$ ) gradients serve as a mobile energy source in plants. *Proceedings of the National Academy of Sciences, USA* 108: 864–869.
- Ganeteg U, Ahmad I, Jamtgard S, Aguetoni-Cambui C, Inselsbacher E, Svennerstam H, Schmidt S, Nasholm T. 2017. Amino acid transporter mutants of *Arabidopsis* provides evidence that a non-mycorrhizal plant acquires organic nitrogen from agricultural soil. *Plant, Cell & Environment* 40: 413–423.
- Gaymard F, Pilot G, Lacombe B, Bouchez D, Bruneau D, Boucherez J, Michaux-Ferriere N, Thibaud JB, Sentenac H. 1998. Identification and disruption of a plant shaker-like outward channel involved in  $K^+$ -release into the xylem sap. *Cell* 94: 647–655.
- Gould KS, Jay-Allemand C, Logan BA, Baissac Y, Bidel LPR. 2018. When are foliar anthocyanins useful to plants? Re-evaluation of the photoprotection hypothesis using *Arabidopsis thaliana* mutants that differ in anthocyanin accumulation. *Environmental and Experimental Botany* 154: 11–22.
- Gruber BD, Giehl RFH, Friedel S, von Wieren N. 2013. Plasticity of the *Arabidopsis* root system under nutrient deficiencies. *Plant Physiology* 163: 161–179.
- Gupta K, Dey A, Gupta B. 2013. Plant polyamines in abiotic stress responses. *Acta Physiologiae Plantarum* 35: 2015–2036.
- Gupta KJ, Shah JK, Brotman Y, Jahnke K, Willmitzer L, Kaiser WM, Bauwe H, Igamberdiev AU. 2012. Inhibition of aconitase by nitric oxide leads to induction of the alternative oxidase and to a shift of metabolism towards biosynthesis of amino acids. *Journal of Experimental Botany* 63: 1773–1784.
- Gutiérrez-Alanís D, Ojeda-Rivera JO, Yong-Villalobos L, Cárdenas-Torres L, Herrera-Estrella L. 2018. Adaptation to phosphate scarcity: tips from *Arabidopsis* roots. *Trends in Plant Science* 23: 721–730.
- Havé M, Marmagne A, Chardon F, Masclaux-Daubresse C. 2016. Nitrogen remobilization during leaf senescence: lessons from *Arabidopsis* to crops. *Journal of Experimental Botany* 68: 2513–2529.
- Hawkesford M, Horst W, Kichey T, Lambers H, Schjoerring JK, Moller IS, White P. 2012. Functions of macronutrients. In: Marschner P, ed. *Marschner's mineral nutrition of higher plants, 3<sup>rd</sup> edn.* San Diego, CA, USA: Academic Press, 135–189.
- Hedrich R, Anschutz U, Becker D. 2011. Biology of plant potassium channels. In: Murphy AS, Peer W, Schulz B, eds. *Plant plasma membrane, vol. 16.* Berlin, Germany: Springer Verlag, 253–274.
- Henry A, Chopra S, Clark DG, Lynch JP. 2012. Responses to low phosphorus in high and low foliar anthocyanin coleus (*Solenostemon scutellarioides*) and maize (*Zea mays*). *Functional Plant Biology* 39: 255–265.
- Hocking B, Tyerman SD, Burton RA, Gillham M. 2016. Fruit calcium: transport and physiology. *Frontiers in Plant Science* 29: 569.
- Hou W, Tränkner M, Lu J, Yan J, Huang S, Ren T, Cong R, Li X. 2020. Diagnosis of nitrogen nutrition in rice leaves influenced by potassium levels. *Frontiers in Plant Science* 11: 165.
- Hu B, Jiang Z, Wang W, Qiu Y, Zhang Z, Liu Y, Li A, Gao X, Liu L, Qian Y et al. 2019. Nitrate-NRT1.1B-SPX4 cascade integrates nitrogen and phosphorus signalling networks in plants. *Nature Plants* 5: 401–413.
- Huang CH, Singh GP, Park SH, Chua N-H, Ram RJ, Park BS. 2020. Early diagnosis and management of nitrogen deficiency in plants utilizing Raman spectroscopy. *Frontiers in Plant Science* 11: 663.
- Hudson D, Guevara D, Yaish MW, Hannam C, Long N, Clarke JD, Bi YM, Rothstein SJ. 2011. GNC and CGA1 modulate chlorophyll biosynthesis and glutamate synthase (GLU1/Fd-GOGAT) expression in *Arabidopsis*. *PLoS ONE* 6: e26765.
- Hughes NM, Lev-Yadun S. 2015. Red/purple leaf margin coloration: potential ecological and physiological functions. *Environmental and Experimental Botany* 119: 27–39.
- Isbell F, Reich PB, Tilman D, Hobbie SE, Polasky S, Binder S. 2013. Nutrient enrichment, biodiversity loss, and consequent declines in ecosystem productivity. *Proceedings of the National Academy of Sciences, USA* 110: 11911–11916.
- Ivashikina N, Deeken R, Fischer S, Ache P, Hedrich R. 2005. AKT 2/3 subunits render guard cell  $K^+$  channels  $Ca^{2+}$  sensitive. *Journal of Genetic Physiology* 125: 483–492.
- Järvan M, Lukme L, Adamson A, Akk A. 2017. Responses of wheat yield, quality and bread-making properties on the sulphur fertilization. *Acta Agriculturae Scandinavica: Section B, Soil & Plant Science* 67: 444–452.
- Jiang C, Gao X, Liao L, Harberd NP, Fu X. 2007. Phosphate starvation root architecture and anthocyanin accumulation responses are modulated by the gibberellin-DELLA signaling pathway in *Arabidopsis*. *Plant Physiology* 145: 1460–1470.
- Jobe TO, Zenzen I, Karvansara PR, Kopriva S. 2019. Integration of sulfate assimilation with carbon and nitrogen metabolism in transition from  $C_3$  to  $C_4$  photosynthesis. *Journal of Experimental Botany* 70: 4211–4221.
- de Jong M, George G, Ongaro V, Williamson L, Willetts B, Ljung K, McCulloch H, Leyser O. 2014. Auxin and strigolactone signaling are required for modulation of *Arabidopsis* shoot branching by nitrogen supply. *Plant Physiology* 166: 384–395.
- Kamerlin SCL, Sharma PK, Prasad RB, Warshel A. 2013. Why nature really choose phosphates. *Quarterly Reviews of Biophysics* 46: 1–132.
- Kanno S, Arrighi JF, Chiarenza S, Bayle V, Berthome R, Peret B, Javot H, Delannoy E, Marin E, Nakanishi TM et al. 2016. A novel role for the root cap in phosphate uptake and homeostasis. *eLife* 5:e14577.
- Kaur G, Chandna R, Pandey R, Abrol YP, Iqbal M, Ahmad A. 2011. Sulfur starvation and restoration affect nitrate uptake and assimilation in rapeseed. *Protoplasma* 248: 299–311.
- Kebrom TH, Spielmeyer W, Finnegan EJ. 2013. Grasses provide new insights into regulation of shoot branching. *Trends in Plant Science* 18: 42–48.
- Kimura Y, Ushiwatari T, Suyama A, Tominaga-Wada R, Wada T, Maruyama-Nakashita A. 2019. Contribution of root hair development to sulphate uptake in *Arabidopsis*. *Plants* 8: 106.
- Kohlen W, Charnikhova T, Liu Q, Bours R, Domagalska MA, Beguerie S, Verstappen F, Leyser O, Bouwmeester H, Ruyter-Spira C. 2011. Strigolactones are transported through the xylem and play a key role in shoot architectural response to phosphate deficiency in nonarbuscular mycorrhizal host *Arabidopsis*. *Plant Physiology* 155: 974–987.
- Kölling M, Kumari P, Bürstenbinder K. 2019. Calcium- and calmodulin-regulated microtubule-associated proteins as signal-integration hubs at the plasma membrane–cytoskeleton nexus. *Journal of Experimental Botany* 70: 387–396.
- Kopriva S, Calderwood A, Weckopp SC, Koprivova A. 2015. Plant sulfur and big data. *Plant Science* 241: 1–10.
- Kopriva S, Malagoli M, Takahashi H. 2019. Sulfur nutrition: impacts on plant development, metabolism, and stress responses. *Journal of Experimental Botany* 70: 4069–4073.
- Koprivova A, Kopriva S. 2016. Hormonal control of sulfate uptake and assimilation. *Plant Molecular Biology* 91: 617–627.
- Krouk G, Lacombe B, Bielach A, Perrine-Walker F, Malinska K, Mounier E, Hoyera K, Tillard P, Leon S, Ljung K et al. 2010. Nitrate-regulated auxin transport by NRT1.1 defines a mechanism for nutrient sensing in plants. *Developmental Cell* 18: 927–937.
- Kudla J, Becker D, Grill E, Hedrich R, Hippler M, Kummer U, Parniske M, Romeis T, Schumacher K. 2018. Advances and current challenges in calcium signaling. *New Phytologist* 218: 414–431.
- Kunz HH, Gierth M, Herdean A, Satoh-Cruz M, Kramer DM, Spetea C, Schroeder JI. 2014. Plastidial transporters KEA1, -2, and -3 are essential for chloroplast osmoregulation, integrity, and pH regulation in *Arabidopsis*. *Proceedings of the National Academy of Sciences, USA* 111: 7480–7485.
- Kwon T, Sparks JA, Liao FQ, Blancaflor EB. 2018. ERULUS is a plasma membrane-localized receptor-like kinase that specifies root hair growth by maintaining tip-focused cytoplasmic calcium oscillations. *Plant Cell* 30: 1173–1177.
- La Verde V, Dominici P, Astegno A. 2018. Towards understanding plant calcium signaling through calmodulin-like proteins: a biochemical and structural perspective. *International Journal of Molecular Sciences* 19: 1331.
- Lambers H, Oliveira RS. 2019. *Plant physiological ecology, 3<sup>rd</sup> edn.* Cham, Switzerland: Springer Nature.
- Landrein B, Formosa-Jordan P, Malivert A, Schuster C, Melnyk CW, Yang W, Turnbull C, Meyerowitz EM, Locke JCW, Jönsson H. 2018. Nitrate modulates stem cell dynamics in *Arabidopsis* shoot meristems through cytokinins. *Proceedings of the National Academy of Sciences, USA* 115: 1382–1387.
- Lee J, Kim J, Choi J-P, Lee M, Kim MK, Lee YH, Hur Y, Nou I-S, Park SU, Min SR et al. 2016. Intracellular  $Ca^{2+}$  and  $K^+$  concentration in *Brassica oleracea* leaf

- induces differential expression of transporter and stress-related genes. *BMC Genomics* 17: 211.
- Leran S, Edel KH, Pervent M, Hashimoto K, Corrige-Faillie C, Offenborn JN, Tillard P, Gojon A, Kudla J, Lacombe B. 2015. Nitrate sensing and uptake in Arabidopsis are enhanced by ABI2, a phosphatase inactivated by the stress hormone abscisic acid. *Science Signaling* 8: ra43.
- Li D, Li C, Yao Y, Li M, Liu L. 2020. Modern imaging techniques in plant nutrition analysis: A review. *Computers and Electronics in Agriculture* 174: 105459.
- Li G, Wu Y, Liu G, Xiao X, Wang P, Gao T, Xu M, Han Q, Wang Y, Guo T et al. 2017. Large-scale proteomics combined with transgenic experiments demonstrates an important role of jasmonic acid in potassium deficiency response in wheat and rice. *Molecular & Cellular Proteomics* 16: 1889–1905.
- Li H, Du H, Huang K, Chen X, Liu T, Gao S, Liu H, Tang Q, Rong T, Zhang S. 2016. Identification, and functional and expression analyses of the CorA/MRS2/MGT-type magnesium transporter family in maize. *Plant and Cell Physiology* 57: 1153–1168.
- Li H, Wang N, Ding J, Liu C, Du H, Huang K, Cao M, Lu Y, Gao S, Zhang S. 2017. The maize CorA/MRS2/MGT-type Mg transporter, ZmMGT10, responses to magnesium deficiency and confers low magnesium tolerance in transgenic Arabidopsis. *Plant Molecular Biology* 95: 269–278.
- Li Z, Wu S, Chen J, Wang X, Gao J, Ren G, Kuai B. 2017. NYEs/SGRs-mediated chlorophyll degradation is critical for detoxification during seed maturation in Arabidopsis. *The Plant Journal* 92: 650–661.
- Liang G, He H, Yu DQ. 2012. Identification of nitrogen starvation-responsive microRNAs in *Arabidopsis thaliana*. *PLoS ONE* 7: e48951.
- Lima JE, Kojima S, Takahashi H, von Wirén N. 2010. Ammonium triggers lateral root branching in Arabidopsis in an AMMONIUM TRANSPORTER1;3-dependent manner. *Plant Cell* 22: 3621–3633.
- LinYL, Tsay YF. 2017. Influence of differing nitrate and nitrogen availability on flowering control in Arabidopsis. *Journal of Experimental Botany* 68: 2603–2609.
- Liu K-h, Niu Y, Konishi M, Wu Y, Du H, Sun Chung H, Li L, Boudsocq M, McCormack M, Maekawa S et al. 2017. Discovery of nitrate-CPK-NLP signalling in central nutrient-growth networks. *Nature* 545: 311–316.
- Liu M, Liu XX, He XL, Liu LJ, Wu H, Tang CX, Zhang YS, Jin CW. 2017. Ethylene and nitric oxide interact to regulate the magnesium deficiency-induced root hair development in Arabidopsis. *New Phytologist* 213: 1242–1256.
- Liu Y, Ohm J-B, Harelund G, Wiersma J, Kaiser D. 2011. Sulfur, protein size distribution, and free amino acids in flour mill streams and their relationship to dough rheology and breadmaking traits. *Cereal Chemistry* 88: 109–116.
- Liu Y, von Wirén N. 2017. Ammonium as a signal for physiological and morphological responses in plants. *Journal of Experimental Botany* 68: 2581–2592.
- Long SR, Kahn M, Seefeldt L, Tsay YF, Kopriva S. 2015. Nitrogen and sulphur. In: Buchanan BB, Grussem W, Jones RL, Vickers K, eds. *Biochemistry and molecular biology of plants*. Chichester, West Sussex; Hoboken, NJ: Wiley Blackwell, 711–767.
- Lunde C, Zygalda A, Simonsen HT, Nielsen PL, Blennow A, Haldrup A. 2008. Sulfur starvation in rice: the effect on photosynthesis, carbohydrate metabolism, and oxidative stress protective pathways. *Physiologia Plantarum* 134: 508–521.
- Ma WY, Li JJ, Qu BY, He X, Zhao XQ, Li B, Fu XD, Tong YP. 2014. Auxin biosynthetic gene TAR2 is involved in low nitrogen-mediated reprogramming of root architecture in Arabidopsis. *The Plant Journal* 78: 70–79.
- Maeda Y, Konishi M, Kiba T, Sakuraba Y, Sawaki N, Kurai T, Ueda Y, Sakakibara H, Yanagisawa S. 2018. A NIGT1-centred transcriptional cascade regulates nitrate signalling and incorporates phosphorus starvation signals in Arabidopsis. *Nature Communications* 9: 1376.
- Manning G, Whyte DB, Martinez R, Hunter T, Sudarsanan S. 2002. The protein kinase complement of the human genome. *Science* 298: 1912–1934.
- Marchive C, Roudier F, Castaings L, Bréhaut V, Blondet E, Colot V, Meyer C, Krapp A. 2013. Nuclear retention of the transcription factor NLP7 orchestrates the early response to nitrate in plants. *Nature Communications* 4: 1713.
- Maruyama-Nakashita A. 2017. Metabolic changes sustain the plant life in low-sulfur environments. *Current Opinion in Plant Biology* 39: 144–151.
- Matthus E, Wilkins KA, Swarbreck SM, Doddrell NH, Docula FG, Costa A, Davies JM. 2019. Phosphate starvation alters abiotic-stress-induced cytosolic free calcium increases in roots. *Plant Physiology* 179: 1754–1767.
- Medici A, Szponarski W, Dangeville P, Safi A, Dissanayake IM, Saenchai C, Emanuel A, Rubio V, Lacombe B, Ruffel S et al. 2019. Identification of molecular integrators shows that nitrogen actively controls the phosphate starvation response in plants. *Plant Cell* 31: 1171–1184.
- Miller AJ, Shen Q, Xu G. 2009. Freeways in the plant: transporters for N, P and S and their regulation. *Current Opinion in Plant Biology* 12: 284–290.
- Miller JB, Pratap A, Miyahara A, Zhou L, Bornemann S, Morris RJ, Oldroyd GED. 2013. Calcium/calmodulin-dependent protein kinase is negatively and positively regulated by calcium, providing a mechanism for decoding calcium responses during symbiosis signaling. *Plant Cell* 25: 5053–5066.
- Müller D, Waldie T, Miyawaki K, To JPC, Melnyk CW, Kieber JJ, Kakimoto T, Leyser O. 2015. Cytokinin is required for escape but not release from auxin mediated apical dominance. *The Plant Journal* 82: 874–886.
- Noa W, Menachem M, Moran N. 2014. Is the leaf bundle sheath cells a “smart flux valve” for K<sup>+</sup> nutrition. *Plant Physiology* 171: 715–722.
- Noctor G, Mhamdi A, Chaouch S, Han Y, Neukermans J, Marquez-Garcia B, Queval G, Foyer CH. 2012. Glutathione in plants: an integrated overview. *Plant, Cell & Environment* 35: 454–484.
- Ohkubo Y, Tanaka M, Tabata R, Ogawa-Ohnishi M, Matsubayashi Y. 2017. Shoot-to-root mobile polypeptides involved in systemic regulation of nitrogen acquisition. *Nature Plants* 3. doi: 10.1038/nplants.2017.29.
- Osugi A, Kojima M, Takebayashi Y, Ueda N, Kiba T, Sakakibara H. 2017. Systemic transport of trans-zeatin and its precursor have differing roles in Arabidopsis shoots. *Nature Plants* 3: 17112.
- Peng YY, Liao LL, Liu S, Nie MM, Li J, Zhang LD, Ma JF, Chen ZC. 2019. Magnesium deficiency triggers SGR-mediated chlorophyll degradation for magnesium remobilization. *Plant Physiology* 181: 262–275.
- Péret B, Desnos T, Jost R, Kanno S, Berkowitz O, Nussaume L. 2014. Root architecture responses: in search of phosphate. *Plant Physiology* 166: 1713–1723.
- Poisson E, Trouverie J, Brunel-Muguet S, Akmouche Y, Pontet C, Pinochet X, Avice JC. 2019. Seed yield components and seed quality of oilseed rape are impacted by sulfur fertilization and its interactions with nitrogen fertilization. *Frontiers in Plant Science* 10: 458.
- Poitout A, Crabos A, Petrik I, Novák O, Krouk G, Lacombe B, Ruffel S. 2018. Responses to systemic nitrogen signaling in Arabidopsis roots involve trans-zeatin in shoots. *Plant Cell* 30: 1243–1257.
- Pottosin I, Shabala S. 2016. Transport across chloroplast membranes: optimizing photosynthesis for adverse environmental conditions. *Molecular Plant* 9: 356–370.
- Pottosin I, Velarde-Buendía AM, Bose J, Fuglsang AT, Shabala S. 2014. Polyamines depolarize the membrane and initiate a cross-talk between plasma membrane Ca<sup>2+</sup> and H<sup>+</sup> pumps. *Biophysical Journal* 106: 586a.
- Rahayu YS, Walch-Liu P, Neumann G, Römhild V, von Wirén N, Bangerth F. 2005. Root-derived cytokinins as long-distance signals for NO<sub>3</sub><sup>-</sup>-induced stimulation of leaf growth. *Journal of Experimental Botany* 56: 1143–1152.
- Riveras E, Alvarez JM, Vidal EA, Osés C, Vega A, Gutiérrez RA. 2015. The calcium ion is a second messenger in the nitrate signaling pathway of Arabidopsis. *Plant Physiology* 169: 1397–1404.
- Rubin G, Tohge T, Matsuda F, Saito K, Scheible WR. 2009. Members of the LBD Family of transcription factors repress anthocyanin synthesis and affect additional nitrogen responses in Arabidopsis. *Plant Cell* 21: 3567–3584.
- Sager R, Lee JY. 2014. Plasmodesmata in integrated cell signalling: insights from development and environmental signals and stresses. *Journal of Experimental Botany* 65: 6337–6358.
- Schachtman DP. 2015. The role of ethylene in plant responses to K<sup>+</sup> deficiency. *Frontiers in Plant Science* 6: 1153.
- Schäfer N, Maierhofer T, Herrmann J, Jørgensen ME, Lind C, von Meyer K, Lautner S, Fromm J, Felder M, Hetherington AM et al. 2018. A tandem amino acid residue motif in guard cell SLAC1 anion channel of grasses allows for the control of stomatal aperture by nitrate. *Current Biology* 28: 1370–1379.
- Schmidt SB, Husted S. 2019. The biochemical properties of manganese in plants. *Plants-Basel* 8: 381.
- Schnug E, Haneklaus S. 2005. Sulphur deficiency symptoms in oilseed rape (*Brassica napus* L.) - The aesthetics of starvation. *Phyton Annales Rei Botanicae* 45: 79–95.
- Sharpley A, Jarvie H, Flaten D, Kleinman P. 2018. Celebrating the 350th Anniversary of phosphorus discovery: a conundrum of deficiency and excess. *Journal of Environmental Quality* 47: 774–777.

- Shimoda Y, Ito H, Tanaka A. 2016. Arabidopsis *STAY-GREEN*, Mendel's green cotyledon gene, encodes magnesium-dechelatase. *Plant Cell* 28: 2147–2160.
- Song WP, Yi JW, Kurniadina OF, Wang HC, Huang XM. 2018. Linking fruit Ca uptake capacity to fruit growth and pedicel anatomy, a cross-species study. *Frontiers in Plant Science* 9: 575.
- Spanudakis E, Jackson S. 2014. The role of microRNAs in the control of flowering time. *Journal of Experimental Botany* 65: 365–380.
- Straub T, Ludewig U, Neuhauser B. 2017. The kinase CIPK23 inhibits ammonium transport in *Arabidopsis thaliana*. *Plant Cell* 29: 409–422.
- Sun J, Bankston JR, Payandeh J, Hinds TR, Zagotta WN, Zheng N. 2014. Crystal structure of the plant dual-affinity nitrate transporter NRT1.1. *Nature* 507: 73.
- Sun Y, Yang R, Li L, Huang J. 2017. The magnesium transporter MGT10 is essential for chloroplast development and photosynthesis in *Arabidopsis thaliana*. *Molecular Plant* 10: 1584–1587.
- Tabata R, Sumida K, Yoshii T, Ohyama K, Shinohara H, Matsubayashi Y. 2014. Perception of root-derived peptides by shoot LRR-RKs mediates systemic N-demand signaling. *Science* 346: 343–346.
- Takagi H, Watanabe S, Tanaka S, Matsuura T, Mori IC, Hirayama T, Shimada H, Sakamoto A. 2018. Disruption of ureide degradation affects plant growth and development during and after transition from vegetative to reproductive stages. *BMC Plant Biology* 18: 287.
- Tan C, Du Y, Zhou J, Wang D, Luo M, Zhang Y, Guo W. 2018. Analysis of different hyperspectral variables for diagnosing leaf nitrogen accumulation in wheat. *Frontiers in Plant Science* 9: 674.
- Tang RJ, Luan S. 2017. Regulation of calcium and magnesium homeostasis in plants: from transporters to signaling network. *Current Opinion in Plant Biology* 39: 97–105.
- Tegeder M, Masclaux-Daubresse C. 2018. Source and sink mechanisms of nitrogen transport and use. *New Phytologist* 217: 35–53.
- Terrer C, Jackson RB, Prentice IC, Keenan TF, Kaiser C, Vicca S, Fisher JB, Reich PB, Stocker BD, Hungate BA et al. 2019. Nitrogen and phosphorus constrain the CO<sub>2</sub> fertilization of global plant biomass. *Nature Climate Change* 9: 684–689.
- Thompson RL, Lassaletta L, Patra PK, Wilson C, Wells KC, Gressent A, Koffi EN, Chipperfield MP, Winiarwita W, Davidson EA et al. 2019. Acceleration of global N<sub>2</sub>O emissions seen from two decades of atmospheric inversion. *Nature Climate Change* 9: 993–998.
- Thor K. 2019. Calcium – nutrient and messenger. *Frontiers in Plant Science* 10: 440.
- Tian Q, Zhang X, Yang A, Wang T, Zhang WH. 2016. CIPK23 is involved in iron acquisition of *Arabidopsis* by affecting ferric chelate reductase activity. *Plant Science* 246: 70–79.
- Tränkner M, Tavakol E, Jáklík B. 2018. Functioning of potassium and magnesium in photosynthesis, photosynthate translocation and photoprotection. *Physiologia Plantarum* 163: 414–431.
- Ueno H, Maeda T, Katsuyama N, Katou Y, Matsuo S, Yano K, Ando A, Nagasuga K, Yamada M, Imanishi S. 2018. Cation measurements and gene expression analysis suggest tomato leaf marginal necrosis is caused by a jasmonate signal induced by K<sup>+</sup> starvation in the tip region of leaflets. *Horticulture Journal* 87: 206–213.
- van Maarschalkerweerd M, Husted S. 2015. Recent developments in fast spectroscopy for plant mineral analysis. *Frontiers in Plant Science* 6: 169.
- Vighi IL, Crizel RL, Perin EC, Rombaldi CV, Galli V. 2019. Crosstalk during fruit ripening and stress response among abscisic acid, calcium-dependent protein kinase and phenylpropanoid. *Critical Reviews in Plant Sciences* 38: 99–116.
- Waadt R, Krebs M, Kudla J, Schumacher K. 2017. Multiparameter imaging of calcium and abscisic acid and high-resolution quantitative calcium measurements using R-GECO1-mTurquoise in *Arabidopsis*. *New Phytologist* 216: 303–320.
- Wang BL, Tang XY, Cheng LY, Zhang AZ, Zhang WH, Zhang FS, Liu JQ, Cao Y, Allan DL, Vance CP et al. 2010. Nitric oxide is involved in phosphorus deficiency-induced cluster-root development and citrate exudation in white lupin. *New Phytologist* 187: 1112–1123.
- Wang X, Hao L, Zhu B, Jiang Z. 2018. Plant calcium signaling in response to potassium deficiency. *International Journal of Molecular Sciences* 19: 3456.
- Wang YY, Cheng YH, Chen KE, Tsay YF. 2018. Nitrate transport, signaling, and use efficiency. *Annual Review of Plant Biology* 69: 85–122.
- Wang Y, Wu WH. 2017. Regulation of potassium transport and signaling in plants. *Current Opinion in Plant Biology* 39: 123–128.
- Wang ZQ, Huang H, Deng JM, Liu JQ. 2015. Scaling the respiratory metabolism to phosphorus relationship in plant seedlings. *Scientific Reports* 5: 6377.
- Ward JT, Lahner B, Yakubova E, Salt DE, Raghothama KG. 2008. The effect of iron on the primary root elongation of *Arabidopsis* during phosphate deficiency. *Plant Physiology* 147: 1181–1191.
- Weissert C, Kehr J. 2017. Macronutrient sensing and signaling in plants. In: Hossain MA, Kamiya T, Burritt DJ, Tran LSP, Fujiwara T, eds. *Plant macronutrient use efficiency*. London, UK: Academic Press Ltd – Elsevier Science Ltd, 45–64.
- White PJ, Broadley MR, El-Serehy HA, George TS, Neugebauer K. 2018. Linear relationships between shoot magnesium and calcium concentrations among angiosperm species are associated with cell wall chemistry. *Annals of Botany* 122: 221–226.
- Winkler A, Knoche M. 2019. Calcium and the physiology of sweet cherries: a review. *Scientia Horticulturae* 245: 107–115.
- Xu W, Dubos C, Lepiniec L. 2015. Transcriptional control of flavonoid biosynthesis by MYB–bHLH–WDR complexes. *Trends in Plant Science* 20: 176–185.
- Yan YW, Mao DD, Yang L, Qi JL, Zhang XX, Tang QL, Li YP, Tang RJ, Luan S. 2018. Magnesium transporter MGT6 plays an essential role in maintaining magnesium homeostasis and regulating high magnesium tolerance in *Arabidopsis*. *Frontiers in Plant Science* 9: 274.
- Yang N, Jiang J, Xie H, Bai M, Xu Q, Wang X, Yu X, Chen Z, Guan Y. 2017. Metabolomics reveals distinct carbon and nitrogen metabolic responses to magnesium deficiency in leaves and roots of soybean [*Glycine max* (Linn.) Merr.]. *Frontiers in Plant Science* 8: 2091.
- Yu XM, Wang JZ, Nie PX, Xue XM, Wang GP, An M. 2018. Control efficacy of Ca-containing foliar fertilizers on bitter pit in bagged 'Fuji' apple and effects on the Ca and N contents of apple fruits and leaves. *Journal of the Science of Food and Agriculture* 98: 5435–5443.
- Yu ZT, Juhasz A, Islam S, Diepeveen D, Zhang JJ, Wang PH, Ma WJ. 2018. Impact of mid-season sulphur deficiency on wheat nitrogen metabolism and biosynthesis of grain protein. *Scientific Reports* 8: 13.
- Yuan S, Zhang ZW, Zheng C, Zhao ZY, Wang Y, Feng LY, Niu GQ, Wang CQ, Wang JH, Feng H et al. 2016. *Arabidopsis* cryptochrome 1 functions in nitrogen regulation of flowering. *Proceedings of the National Academy of Sciences, USA* 113: 7661–7666.
- Zeng JB, Ye ZL, He XY, Zhang GP. 2019. Identification of microRNAs and their targets responding to low-potassium stress in two barley genotypes differing in low-K tolerance. *Journal of Plant Physiology* 234: 44–53.
- Zhang L, Wen A, Wu X, Pan X, Wu N, Chen X, Chen Y, Mao D, Chen L, Luan S. 2019. Molecular identification of the magnesium transport gene family in *Brassica napus*. *Plant Physiology and Biochemistry* 136: 204–214.
- Zhang SS, Pan YJ, Tian W, Dong MQ, Zhu HF, Luan S, Li LG. 2017. *Arabidopsis* CNGC14 mediates calcium influx required for tip growth in root hairs. *Molecular Plant* 10: 1004–1006.
- Zhang Y, Liu Z, Liu J, Lin S, Wang J, Lin W, Xu W. 2017. GA-DELLA pathway is involved in regulation of nitrogen deficiency-induced anthocyanin accumulation. *Plant Cell Reports* 36: 557–569.
- Zheng H, Cheng T, Li D, Yao X, Tian Y, Cao W, Zhu Y. 2018. Combining Unmanned Aerial Vehicle (UAV)-based multispectral imagery and ground-based hyperspectral data for plant nitrogen concentration estimation in rice. *Frontiers in Plant Science* 9: 936.
- Zheng ZL, Leustek T. 2017. Advances in understanding sulfur utilization efficiency in plants. In: Hossain MA, Kamiya T, Burritt DJ, Tran LSP, Fujiwara T, eds. *Plant macronutrient use efficiency*. London, UK: Academic Press, 215–232.
- Zörb C, Senbayram M, Peiter E. 2014. Potassium in agriculture – status and perspectives. *Journal of Plant Physiology* 171: 656–669.