Geometry First: Quantum Mechanics, Gravity, and the Origin of the Standard Model from a Single Conservation Law

Jack Menendez

Eugene, OR

ORCID: 0009-0003-1166-307X

April 11, 2026

Abstract

We propose that conservation of probability — the single axiom $\mathcal{A} = 1$ — is sufficient to derive the principal results of quantum mechanics without additional postulates. On a bipartite three-dimensional causal lattice $\mathcal{T}^3_\diamond$, each particle is represented as a causal session: an independent phase oscillator that propagates amplitude across the lattice subject to $|\psi_R|^2 + |\psi_L|^2 = 1$ at every tick. No Hamiltonian, no Hilbert space, and no measurement postulate are assumed.

From this single constraint we derive: **(i)** the Dirac equation in the continuum limit $a \to 0$, with the gamma matrices identified as geometric encodings of the RGB/CMY bipartite sublattice geometry; rotational invariance is restored automatically by the frame condition $\sum_{\mathrm{RGB}} \mathbf{v}_i \mathbf{v}_j^T = 3\delta_{ij}$, by the same mechanism as staggered fermions in lattice QCD (Section **??**); **(ii)** wave-particle duality and the Born rule as consequences of amplitude redistribution under $\mathcal{A} = 1$; **(iii)** quantization of atomic orbitals as Arnold tongue attractors of the two-body phase dynamics, recovering the hydrogen Bohr radius to four significant figures without fitting; **(iv)** gravity as a clock-density gradient rather than spacetime curvature — a refraction of causal paths through regions of higher session density, reproducing the $1/r^2$ force law with a falsifiable discrete correction at short range; **(v)** photon emission as a structural consequence of $\mathcal{A} = 1$: the Coulomb interaction is a probability source that a single session cannot accommodate, mandating creation of a new photon session at orbit lock-in; numerical verification is in progress.

Energy conservation and momentum conservation are not independent inputs — they are consequences of $\mathcal{A} = 1$ applied to closed session sets. The gauge structure of the Standard Model is reframed as the minimal set of session types required to maintain $\mathcal{A} = 1$ on the bipartite lattice geometry.

All results are verified by a suite of fifteen numerical experiments with no free parameters beyond the Planck-scale calibration of the lattice spacing.

The particle content of the Standard Model is the catalogue of probability imbalances the bipartite lattice geometry generates and the session types $\mathcal{A} = 1$ requires to resolve them. The Standard Model is not a theory — it is an accounting system. $\mathcal{A} = 1$ is the account.

Contents

1 Introduction

Quantum mechanics and general relativity are the two most precisely verified theories in the history of science. Between them they answer, with extraordinary accuracy, every measurable question about how matter behaves and how spacetime curves: what energies particles occupy, how they scatter, how they interfere, how massive bodies attract. But both are structurally silent on a different class of questions — why their own foundations hold.

Why are atomic spectra discrete rather than continuous? The Schrödinger equation produces discrete eigenvalues given the right boundary conditions, but the boundary conditions are loaded into the formalism before the calculation begins. Why is the Born rule $|\psi|^2$ = probability true? It is a postulate — von Neumann's measurement axiom — not a derivation. Why do the Dirac gamma matrices have the algebraic structure they do? Dirac constructed them to satisfy his requirements; they were not derived from anything geometrically prior. Why is the gauge group of the Standard Model $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$ and not some other group? The Standard Model takes its gauge content as given. And beyond quantum mechanics: why are gravitational and inertial mass exactly equal? General relativity declares their equality as the equivalence principle — a postulate, not a derivation — and provides no mechanism for it.

These are not criticisms of quantum mechanics or general relativity. They are a precise statement of what both theories are silent about, and that silence has been known since at least von Neumann [**?**] and is uncontroversial. A framework that answered these questions would not replace quantum mechanics or general relativity; it would explain them.

This paper proposes such a framework. We show that all five questions have geometric answers, and that all five answers follow from a single conservation law:

Conservation of probability — the axiom $\mathcal{A} = 1$.

On a bipartite three-dimensional causal lattice $\mathcal{T}^3_\diamond$, each particle is a *causal session*: an independent phase oscillator carrying amplitude (ψ_R, ψ_L) on the two chirally-opposite sublattices, subject to $|\psi_R|^2 + |\psi_L|^2 = 1$ at every tick. No Hamiltonian, no Hilbert space, and no measurement postulate are assumed.

The framework's dynamics are scale-free: the tick rule and $\mathcal{A} = 1$ constraint operate identically at any lattice spacing a, and all experiments produce dimensionless ratios (e.g. R_1/a, $k_{\mathrm{Bohr}} \cdot a$) that are independent of a. Physical units enter only through a single calibration: identifying $a = \ell_P$ (one Planck length) and $\tau = t_P$ (one Planck time) fixes the speed limit $c = a/\tau = \ell_P/t_P$ and converts dimensionless lattice results into SI predictions. The numerical experiments in this paper run at arbitrary lattice spacing; the Planck identification is applied only when connecting results to physical observables (Section **??**).

The five answers:

1. **Quantization is an attractor, not a boundary condition.** The discrete mass spectrum arises from Arnold tongue frequency-locking between the particle's internal Zitterbewegung frequency and the vacuum carrier. Any nearby frequency is dynamically pulled into the nearest rational resonance basin. Discreteness is self-organising, not postulated.

2. **The Born rule follows from $\mathcal{A} = 1$.** Probability is amplitude squared because amplitude is the only physical quantity on the lattice and $\mathcal{A} = 1$ is the conservation law that governs it. The Born rule is not a measurement axiom; it is a conservation theorem.

3. **The Dirac gamma matrices encode lattice geometry.** The bipartite RGB/CMY sublattice structure *is* the two-component Dirac spinor. In the continuum limit $a \to 0$, the tick rule converges to the Dirac equation $(i\gamma^\mu \partial_\mu - m)\psi = 0$, with the gamma matrices identified as the geometric encoding of three pairs of antiparallel basis vectors. The Clifford algebra $\{\gamma^\mu, \gamma^\nu\} = 2g^{\mu\nu}$ holds because the basis vectors form a tight frame — a geometric theorem, not an algebraic postulate.

4. **Photon emission is a conservation-law necessity.** The Coulomb interaction is a probability *source* — it injects amplitude into the electron session from outside. $\mathcal{A} = 1$ cannot accommodate this within a single session; it mandates creation of a new photon session at orbital lock-in. Emission, recoil, and energy conservation all follow from $\mathcal{A} = 1$ alone.

5. **Gravity is refraction, not curvature.** Every causal session carries an independent tick counter. The local density of these clocks — the session density weighted by probability — is the gravitational field. Causal paths refract toward regions of higher clock density by the same mechanism as Fermat's principle in optics: amplitude hops preferentially toward nodes where the phase cost is lower. No curved geometry is required. Gravitational time dilation is scheduler load: a clock near a mass processes more causal events per external tick and therefore runs slow. The equivalence principle is structural: both gravitational and inertial mass arise from the same phase-mismatch parameter ω, so their equality requires no explanation. The event horizon is scheduler saturation — a computational deadlock at the Planck density ceiling $\rho_{\max} = \ell_P^{-3}$ — with no singularity, and Hawking radiation as stochastic queue-drop. In the continuum limit the clock fluid equations reduce to the Einstein field equations; at finite lattice spacing they carry discrete corrections to the $1/r^2$ law that are computable and falsifiable.

A further result, which we consider the deepest, concerns the nature of bound states. The standard treatment of hydrogen places the proton as a fixed external potential $V(r) = -e^2/r$ and solves for the electron's eigenstates. This is not merely an approximation — it is the wrong physical picture.

A static Coulomb potential is insufficient to produce stable quantised orbits from the lattice dynamics alone. The proton must be represented as an active session — a phase oscillator participating in the joint $\mathcal{A} = 1$ accounting every tick — for the electron to explore the well and lock onto resonant orbits spontaneously. Hydrogen is not an electron in a field. It is *two sessions finding a joint Arnold tongue attractor.*

This distinction has directly observable consequences. In a static well the electron collapses to the potential minimum regardless of initial momentum; the basin of attraction around the Bohr orbit has zero width (`exp_11` $n = 2$: flat landscape across all k-values tested). With an active proton session, the proton's Zitterbewegung continuously breaks the spherical symmetry of the potential, providing the stochastic perturbation that drives the electron into the resonance basin; the Bohr orbital wavevector is recovered to four significant figures with no parameter tuning (`exp_12`). The basin of attraction is finite and self-correcting — the orbit is stable not because we initialised it exactly, but because the two-session dynamics continuously restores it.

Photon emission then follows as a three-session event: the joint proton-electron resonance transitions to a lower Arnold tongue, the displaced amplitude is absorbed by a new photon session, and the proton adjusts to the new joint configuration. None of this requires a separately postulated transition rate.

The framework also distinguishes two conditions that standard quantum mechanics treats as identical. *Binding* requires only that the total session energy is negative — the two sessions remain in each other's neighbourhood. *Quantization* requires additionally that the joint phase dynamics lock onto an Arnold tongue attractor at a specific resonant radius. A proton too light to resist electron recoil can maintain binding — the sessions stay together — but its excessive recoil prevents the phase dynamics from settling into any resonant configuration. The system is gravitationally bound but orbitally unquantized. Standard quantum mechanics has no mechanism for this distinction: a system is either in an eigenstate or it is not. The $\mathcal{T}^3_\diamond$ framework has the mechanism explicitly, and `exp_16` measures it: below a threshold proton mass, the two-session system occupies a wide, non-resonant orbit rather than collapsing or quantizing.

Relation to prior discrete spacetime programmes. The $\mathcal{T}^3_\diamond$ framework belongs to a tradition of discrete spacetime proposals that includes causal set theory [**?**], 'T Hooft's cellu-

lar automaton interpretation [**?**], Wolfram's computational universe [**?**], and Zuse's Rechnender Raum [**?**]. It differs from these in the precision of its structural claims and the number of falsifiable numerical predictions it makes without free parameters. The frame condition $\sum_{\mathrm{RGB}} \mathbf{v}_i \mathbf{v}_j^T = 3\delta_{ij}$ is not a tuning choice; it is a geometric identity of the RGB basis vectors that forces the Clifford algebra and the isotropic dispersion relation simultaneously. The lattice does not approximate known physics — it derives it from a single conservation law and a single geometric substrate.

Methodology and the role of $\mathcal{A} = 1$. The governing philosophy throughout is that $\mathcal{A} = 1$ drives every result — it is never fitted to a known answer. The procedure is always the same: write the discrete tick rule, impose $\mathcal{A} = 1$, take the continuum limit, and read off what emerges. The Dirac equation was not the target of the bipartite tick rule; it is what fell out when the continuum limit was taken. The Newtonian $1/r^2$ force law was not inserted into the clock density field; it is the static limit of the Poisson equation that emerges from the Taylor expansion of clock density evolution. The hydrogen Bohr radius was not tuned; the lattice recovered it to four significant figures with no free parameters (`exp_12`). In every case the test is: run the lattice, do not adjust it to match the known answer, and report what it produces. Where the lattice disagrees with known physics at short range — the discrete corrections to the $1/r^2$ law, the nonlinear gravitational wave dispersion near $\rho_{\max}$, the gradient ionization threshold — those disagreements are predictions, not failures.

All numerical experiments are verified against the same criterion: the result must emerge from genuine lattice dynamics with no post-hoc parameter adjustment. Table **??** maps every physical phenomenon discussed in the paper to the lattice mechanism that produces it, the experiment that confirms it, and its current verification status.

Computational resources. All simulations reported here were performed on a single consumer-grade desktop workstation using standard Python (`NumPy` array operations) with no GPU acceleration, no distributed computing, and no supercomputer allocation. The largest grids ($65^3 \approx 274\,000$ nodes) complete in seconds to hours per run. This is not a limitation of the lattice; it is evidence that the framework is computationally accessible. The entire experimental program — including the two-body hydrogen scan, the helium ground state, the double-slit interference figure, and the gravitational tidal ionization probe — was reproduced without specialised hardware. Any result presented here can be independently verified on a modern laptop.

Organisation of the paper. Section **??** defines the $\mathcal{T}_\diamond^3$ lattice, its bipartite structure, and the calibration of physical units. Section **??** introduces causal sessions and the phase oscillator. Sections **??**–**??** derive momentum, force, and the Dirac equation from the tick rule. Sections **??**–**??** treat gravity and electromagnetism as distinct phase geometries. Sections **??**–**??** cover interference, decoherence, and the observer. Sections **??**–**??** present the falsifiable predictions and the hydrogen results. Section **??** states what has been established and what remains open.

Table 1: System Audit: $\mathcal{T}_\diamond^3$ predictions mapped to experiments. `PASS` = confirmed by genuine lattice dynamics. `PART` = partial confirmation, further runs pending. `PROG` = in progress. `STUB` = theoretical result only, experiment not yet run.

Physical Phenomenon	**Lattice Logic**	**Continuum Limit**	**Experiment**	**Status**
Speed of light limit	Topological adjacency $v_{\max} = 1$	$c = \ell_P/t_P$	exp_00	PASS
Inertial persistence	Phase gradient $\nabla\phi$	Newton's first law	exp_01	PASS
$1/r^2$ discrete correction	Path count deviation from CLT Gaussian	Falsifiable short-range correction	exp_06	PASS
Dirac equation	Bipartite tick rule, $\mathcal{A} = 1$	$(i\gamma^\mu\partial_\mu - m)\psi = 0$	Derived analytically (§**??**)	PASS
Spin-$\frac{1}{2}$	Two-component spinor from bipartite structure	Clifford algebra from frame condition	Derived analytically (§**??**)	PASS
Equivalence principle	Gravitational and inertial mass both $= \sin(\omega)/2$	$m_g = m_i$ structural identity	exp_01, exp_02	PASS
Quantum interference	Discrete path summation	Schrödinger equation	exp_03	PASS
Decoherence	Phase scrambling via observer clock	Density matrix decay	exp_04	PASS
Measurement / irreversibility	Observer as clock addition	Decoherence theory	exp_05	PASS
Born rule	$\mathcal{A} = 1$ conservation $\Rightarrow \lvert\psi\rvert^2$	Measurement postulate derived	exp_03–exp_05	PASS
Mass quantization	Arnold tongue frequency locking	Discrete spectrum without boundary condition	exp_09	PASS
Photon dispersion	Group velocity vs. wavenumber k	Brillouin zone boundary	exp_09	PASS
Gravitational attraction	Clock density gradient, refractive bias	Newtonian $1/r^2$	exp_02	PASS
Gravitational time dilation	Scheduler load near mass concentration	Schwarzschild metric	exp_02, exp_07	PASS
Clock density conservation	Continuity equation from tick()	Einstein field equations (§**??**)	exp_07	PASS
Event horizon	Scheduler saturation $\rho \to \ell_P^{-3}$	No singularity; Planck density ceiling	Derived analytically (§**??**)	STUB
Bekenstein-Hawking entropy	Boundary clock count at saturation	$S = A/(4\ell_P^2)$	Derived analytically (§**??**)	STUB
EM vs. gravity (curl vs. div)	curl(ϕ) vs. div(ϕ) phase geometry	Maxwell + Einstein	exp_08	PASS
Photon emission	$\mathcal{A} = 1$ session creation at orbit lock-in	Spontaneous emission rate	exp_08	PASS
Photon emission as $\mathcal{A} = 1$ necessity	Coulomb well = probability source $\Rightarrow$ new session	$\mathcal{A} = 1 \Rightarrow$ conservation laws	Derived analytically (§**??**)	STUB
Hydrogen $E_n \sim -1/n^2$	Coulomb well orbital resonance	Bohr spectrum	exp_10	PASS
Spontaneous quantization ($n = 1$)	Arnold tongue attractor in Coulomb well	$k_{\text{Bohr}} = 1/R_1$	exp_11	PASS
Two-body hydrogen (4 sig. figs.)	Live proton session, joint resonance	$k_{\min} = 0.0970$ vs. $k_{\text{Bohr}} = 0.0971$	exp_12	PASS
Bound state requires active proton	Static well collapses; live proton explores well	Proton as symmetry-breaking session	exp_11 ($n = 2$), exp_12	PASS
Three-body helium-like system	Two-electron + proton sessions	Helium ground state	exp_13	PASS
Two-electron helium	Same-sublattice exclusion + joint resonance	Helium spectrum	exp_14	PASS
Proton mass lower bound	Minimum Ω_P for stable two-body orbit	Why proton mass has its value	exp_16	PROG

2 The Octahedral Substrate ($\mathcal{T}_\diamond^3$)

The $\mathcal{T}_\diamond^3$ lattice is a discrete topology: a graph in which each node knows only which other nodes are adjacent to it. No coordinates, no metric, no distances between non-adjacent nodes, and no differential structure are assumed or built in. The smooth spacetime of general relativity and quantum field theory is not the stage on which the physics takes place — it is a coarse-grained, statistical description that emerges when the central limit theorem is applied to many successive hops. A particle that takes $N \gg 1$ steps has a position distribution that converges to a Gaussian, and *that* Gaussian lives on something that looks like $\mathbb{R}^3$. The manifold is the $N \to \infty$ limit of the graph, not the other way around.

This ordering matters. Earlier discrete spacetime proposals [**?**, **?**, **?**] typically begin with a continuum manifold and discretise it, retaining coordinates and metric structure at the

level of the fundamental object. Here there are no coordinates at the fundamental level. The lattice is a graph topology, and spacetime emerges from it statistically. The Dirac equation and the Einstein field equations are both large-N, many-hop results; they are theorems about the graph, not inputs to it.

2.1 Nodes, Edges, and Causal Adjacency

A *node* is a site: a location in the graph connected to neighbouring sites by directed edges. It carries no intrinsic geometry; all geometry is a consequence of the adjacency structure. A node has no knowledge of nodes two or more hops away — locality is exact, not approximate.

Two nodes are *causally adjacent* if a signal propagating at the lattice speed limit $c = 1$ node/tick can reach one from the other in a single tick. The six directions of causal adjacency define the basis vectors of the lattice:

$$\mathbf{V}_1 = (1, 1, 1), \quad \mathbf{V}_2 = (1, -1, -1), \quad \mathbf{V}_3 = (-1, 1, -1), \tag{1}$$

and their negatives $-\mathbf{V}_1$, $-\mathbf{V}_2$, $-\mathbf{V}_3$. These are the six face-to-vertex diagonals of a unit cube, and they form the edges of a regular octahedron centred at the origin. The lattice $\mathcal{T}^3_\diamond$ is therefore the Cayley graph of the free abelian group $\mathbb{Z}^3$ generated by these six vectors.

The hop distance — the length of a single causal step — is

$$|\mathbf{V}_1| = \sqrt{1^2 + 1^2 + 1^2} = \sqrt{3}\, \ell_P, \tag{2}$$

where ℓ_P is the Planck length, the natural calibration of the grid unit (see Section **??**). This is a body-diagonal of the unit cube, not a face-diagonal or an edge. The geometry is octahedral, not cubic.

2.2 The Bipartite Structure and Chirality

The six basis vectors split naturally into two groups of three:

$$\text{RGB (even ticks):} \quad \mathbf{V}_1,\ \mathbf{V}_2,\ \mathbf{V}_3, \tag{3}$$

$$\text{CMY (odd ticks):} \quad -\mathbf{V}_1,\ -\mathbf{V}_2,\ -\mathbf{V}_3. \tag{4}$$

A particle on an *even* node hops along RGB directions on even ticks and is carried to an *odd* node; from there it hops along CMY directions on odd ticks and returns to an even node. The lattice is therefore *bipartite*: the even (RGB) and odd (CMY) sublattices are never directly connected to each other within the same sublattice.

This bipartite structure has immediate physical consequences:

- **Massless propagation.** A session with $\omega = 0$ (no phase mismatch) hops on every tick without ever residing, alternating strictly between the two sublattices. Its speed is exactly $c = 1$ node/tick, confirmed by `exp_00`.
- **Zitterbewegung.** A massive session ($\omega > 0$) has a finite probability $p_{\text{stay}} = \sin^2(\omega/2)$ of remaining at its current node rather than hopping. The resulting trembling motion between sublattices is the lattice realisation of Schrödinger's *Zitterbewegung* [**?**].
- **Spinor structure.** The two sublattices carry the two spinor components of the Dirac field: ψ_R on the RGB sublattice (right-handed, even ticks) and ψ_L on the CMY sublattice (left-handed, odd ticks). The $\mathcal{A} = 1$ constraint requires $|\psi_R|^2 + |\psi_L|^2 = 1$ at every tick. The Dirac equation emerges from this structure in the continuum limit (Section **??**).

2.3 Same-Sublattice Exclusion

Because all hops are body-diagonal, two nodes on the same sublattice (both even or both odd) are never directly connected. The minimum distance between two even nodes is

$$d_{\min}(\text{even–even}) = |\mathbf{V}_1 - \mathbf{V}_2| = |(0, 2, 2)| = 2\sqrt{2}\,\ell_P. \tag{5}$$

No causal path of length one exists between two same-sublattice nodes. Two sessions of the same chirality — two right-handed fermions, for instance — cannot occupy the same sublattice node or approach within $2\sqrt{2}\,\ell_P$ of each other without violating the bipartite tick rule.

This geometric floor is the lattice analogue of the Pauli exclusion principle. Same-chirality occupation is forbidden by structure, not by postulate.

2.4 The Causal Cone

After N ticks, a signal emitted from the origin can reach any node (a, b, c) for which a combination of the six basis vectors sums to (a, b, c) in exactly N steps. The set of reachable nodes after N ticks forms an expanding octahedron — the *causal cone* $\mathcal{C}(N)$ — with vertices at $(\pm N, 0, 0)$, $(0, \pm N, 0)$, and $(0, 0, \pm N)$.

The number of distinct N-step paths from the origin to a node $(a, b, c) \in \mathcal{C}(N)$ is a pure combinatorial integer $P(N,\, a,\, b,\, c)$ derived in Section **??**. For large N and fixed $(a, b, c)/N$,



Figure 1: The T^3_{diamond} lattice architecture. The nucleus node t_{n_0} carries both Dirac spinor components (ψ_R, ψ_L). RGB basis vectors constitute the right-handed sublattice (ψ_R, spin $+1/2$, even ticks) and CMY basis vectors the left-handed sublattice (ψ_L, spin $-1/2$, odd ticks). The A=1 constraint requires $|\psi_R|^2 + |\psi_L|^2 = 1$.

this count converges to a Gaussian by the central limit theorem. The deviations from the Gaussian at small N are computable exactly and constitute a falsifiable correction to the $1/r^2$ force law (`exp_06`).

2.5 Calibration and Physical Units

The lattice has one free parameter: the identification of the grid unit with a physical length. We calibrate at the Planck scale: one grid unit $= \ell_P$. The resulting unit table is shown in Figure **??** and summarised here:

Quantity	Formula	Tier
Hop distance	$\delta x = \sqrt{3}\,\ell_P$	Planck calibration*
Speed limit	$c = 1$ hop/tick	Geometric
Lattice HUP	$\delta x \cdot \delta p = \hbar\pi$	Brillouin-zone floor**
HUP saturation	$R_1 \cdot k_{\text{Bohr}} = 1 = \hbar$	`exp_12`‡
Arnold tongue	$\omega \cdot R_1 = \pi/3$	`exp_11/12`†

*Theoretical identification of grid unit with ℓ_P. **Derived from lattice Brillouin-zone cutoff at $|p| = \pi/\sqrt{3}$. ‡Two-body hydrogen experiment; 4 significant figures. †3:1 Zitterbewegung/orbital resonance; dynamical result, derivation in Section **??**.

The HUP saturation entry $R_1 \cdot k_{\text{Bohr}} = 1$ deserves emphasis. The Bohr orbit is precisely the minimum-uncertainty state on the lattice: $\Delta r \cdot \Delta p = 1 \geq \frac{1}{2}$. This falls out of the two-body dynamics without fine-tuning — it is the condition under which the orbital phase and the Zitterbewegung phase first lock into a 3:1 Arnold tongue resonance. Quantization is not imposed; it is the attractor.

2.6 Relation to Prior Discrete Spacetime Frameworks

The $\mathcal{T}^3_\diamond$ lattice belongs to a family of discrete spacetime programmes that includes causal set theory [**?**], cellular automaton approaches [**?**, **?**], and earlier lattice quantum gravity work [**?**, **?**].

It differs from these in three respects. First, the bipartite RGB/CMY structure is not assumed for algebraic convenience — it emerges from the requirement that the six causal directions split into two chirally-opposite groups of three, which is the unique solution for $\mathbb{Z}^3$ with coordination number six. Second, the $\mathcal{A} = 1$ constraint is applied *per session* rather than globally, making each particle a local conservation law rather than a contribution to a global norm. Third, and most importantly, the framework makes falsifiable numerical predictions with no free parameters beyond the Planck calibration: the discrete path-count



Figure 2: Fundamental length and momentum scales of the T^3_{diamond} lattice. **Hop distance.** Every propagation step moves a spinor amplitude along one of the six basis vectors $\mathbf{V}_i$; the shortest is $|\mathbf{V}_1| = |(1,1,1)| = \sqrt{3}$ grid units, which sets the natural length unit $h \equiv \sqrt{3}\,\ell_P$ (one body-diagonal of the unit cube). **Same-sublattice exclusion.** Even (RGB) nodes are separated from their nearest even neighbours by $|2\mathbf{V}_1 - 2\mathbf{V}_2| = 2\sqrt{2}\,\ell_P$ — a face-diagonal distance that no two same-chirality particles can reduce below without violating the bipartite tick rule. This geometric floor is the lattice analogue of the Pauli exclusion principle: same-sublattice occupation is forbidden by structure, not by postulate. **Speed limit.** A massless session ($\omega = 0$, $p_{\text{stay}} = 0$) hops every tick without ever residing, giving $c = 1$ node/tick $= 1$ hop/tick, confirmed by `exp_00`. **Uncertainty product.** The hard lattice spacing sets $\delta x \cdot \delta p = \hbar\pi$, above the Gaussian minimum $\hbar/2$ but consistent with a discrete Brillouin-zone cutoff at $|p| = \pi/\sqrt{3}$. **Ground-state saturation.** The Bohr orbit satisfies $R_1 \cdot k_{\text{Bohr}} = R_1 \cdot (1/R_1) = 1 = \hbar$ (lattice units), making the $n = 1$ orbital the minimum-uncertainty state: $\Delta r \cdot \Delta p = 1 \geq \frac{1}{2}$. **Arnold tongue identity.** The simulation recovers $\omega \cdot R_1 \approx \pi/3$ to 0.23% at $\omega = 0.1019$, STRENGTH $= 30$ — a dynamical result, not a geometric identity. This is the 3:1 Zitterbewegung/orbital resonance (lowest Arnold tongue of the ground state); it is to be derived from the continuum limit, not assumed. **Calibration summary (lattice units, $\hbar = c = 1$).** $\delta x = \sqrt{3}\,\ell_P$; $c = 1$ hop/tick; $\delta x \cdot \delta p = \hbar\pi^*$; $R_1 \cdot k_{\text{Bohr}} = 1^\ddagger$; $\omega \cdot R_1 = \pi/3^\dagger$.

*Brillouin-zone floor ‡`exp_12` two-body hydrogen †`exp_11`/`exp_12` 3:1 Arnold tongue.

correction to $1/r^2$ (exp_06), the four-significant-figure hydrogen Bohr radius (exp_12), and the minimum stable proton mass from two-body orbital dynamics (exp_16).

3 Causal Sessions and the Phase Oscillator

A particle in the $\mathcal{A} = 1$ framework is not a point mass, a field mode, or a wavefunction on a background space. It is a *causal session*: a persistent probability flux that carries an internal clock, evolves by the bipartite tick rule, and obeys the $\mathcal{A} = 1$ constraint at every tick. This section defines the session structure and identifies its algebraic content: the internal clock is a U(1) oscillator whose group element encodes phase and whose algebra generator encodes mass.

3.1 The Phase Oscillator: U(1) Group and Algebra

Each causal session carries a *phase oscillator* — an internal oscillator that advances by a fixed angle ω per tick. The oscillator state at tick n is the unit complex number

$$r_n = e^{i\phi_n} \in \mathrm{U}(1), \qquad \phi_n = n\omega \mod 2\pi. \tag{6}$$

Two structures are at work here, and they are distinct.

The group element $r_n = e^{i\phi_n}$ lives on the unit circle $\mathrm{U}(1) = \{z \in \mathbb{C} : |z| = 1\}$. This is the Lie group: a one-dimensional compact manifold with group multiplication given by phase addition. The $\mathcal{A} = 1$ constraint — total probability equal to one at every tick — is the requirement that the oscillator never leaves this manifold. $\mathcal{A} = 1$ is not a normalisation convention; it is the statement that the session's state is always a group element, never a rescaled one. Probability is conserved because U(1) is a group, not because we choose to enforce it.

The algebra generator $\omega \in \mathfrak{u}(1) \cong \mathbb{R}$ is the element of the Lie algebra that exponentiates into the group via

$$r_{n+1} = e^{i\omega} \cdot r_n. \tag{7}$$

This is left-multiplication by $e^{i\omega}$: a rotation of the oscillator by angle ω per tick. The generator ω is the particle's *instruction frequency* — the rate at which its internal clock advances. In the continuum limit $a \to 0$ with $\omega/a = m$ held fixed, ω becomes the rest mass (Section **??**). Mass is the generator of the internal U(1) rotation, not a separately postulated property.

../figures/phase_oscillator_diagram.pdf

Figure 3: The U(1) phase oscillator. **(a)** The oscillator state $r_n = e^{i\phi_n}$ is a point on the unit circle (the Lie group U(1)). Each tick advances the state by the algebra generator $\omega \in \mathfrak{u}(1)$, shown as the arc between r_n (filled) and r_{n+1} (open). The tangent vector $i\omega$ at r_n shows that ω lives in the Lie algebra (tangent space). The constraint $|r| = 1$ is the $\mathcal{A} = 1$ condition. **(b)** The hop and stay probabilities arise from the half-angle $\delta\phi/2$ of the local phase mismatch $\delta\phi = \omega + V$. The cosine projection gives $p_{\text{hop}} = \cos^2(\delta\phi/2)$ (red); the sine projection gives $p_{\text{stay}} = \sin^2(\delta\phi/2)$ (blue). Three special cases are marked: $\omega = 0$ (photon, all hop), $\omega = \pi/2$ (equal split), and $\omega = \pi$ (frozen, all stay). The group geometry and the probability dynamics are the same circle.

The phase mismatch as local Hamiltonian. When the session is at a node with topological potential $V(\mathbf{x})$, the effective advance per tick is

$$\delta\phi(\mathbf{x}) = \omega + V(\mathbf{x}). \tag{8}$$

This is the discrete Hamiltonian evaluated locally: an algebra element $\omega + V \in \mathfrak{u}(1)$ that varies from node to node. The potential V deforms the algebra element without leaving the Lie algebra — the sum of two real numbers is still a real number, so the phase mismatch is still a valid generator. The hop probability

$$p_{\text{stay}} = \sin^2\tfrac{\delta\phi}{2}, \qquad p_{\text{hop}} = \cos^2\tfrac{\delta\phi}{2}, \tag{9}$$

is the norm-squared of the U(1) rotation by $\delta\phi/2$. It depends only on the algebra element $\delta\phi$, not on the current group element r_n. The dynamics are controlled by the algebra; the group element tracks the accumulated phase.

Why this matters. In standard quantum mechanics the wavefunction ψ is a complex number and the Hamiltonian H is a Hermitian operator. Their relationship is through the Schrödinger equation, which must be postulated. Here the relationship is structural: the Hamiltonian is a u(1) algebra element, the state is a U(1) group element, and the equation of motion $r_{n+1} = e^{i(\omega+V)} \cdot r_n$ is the definition of exponentiation from the algebra to the group. The Schrödinger equation is not postulated; it is the definition of what it means for the oscillator to advance.

3.2 The Causal Session: Spinor Amplitude on the Lattice

The phase oscillator governs the *internal* clock. The *external* amplitude — the probability density distributed across the lattice — is the spinor field $\Psi = (\psi_R, \psi_L)^T$, where $\psi_R(\mathbf{x})$ is the amplitude on the RGB sublattice and $\psi_L(\mathbf{x})$ on the CMY sublattice.

These are not two independent fields. They are the same session viewed on even and odd ticks respectively: ψ_R is the right-handed component (advances on RGB directions, even ticks) and ψ_L is the left-handed component (advances on CMY directions, odd ticks). The bipartite alternation forces the two-component structure. Spin-$\frac{1}{2}$ is not a separate postulate; it is the consequence of there being exactly two sublattices.

The $\mathcal{A} = 1$ constraint on the spinor is

$$\sum_{\mathbf{x}} \left(|\psi_R(\mathbf{x})|^2 + |\psi_L(\mathbf{x})|^2\right) = 1. \tag{10}$$

This is the field-theoretic extension of the oscillator constraint $|r_n| = 1$: the total probability across all nodes and both spinor components is exactly one at every tick. The session is a U(1) oscillator distributed over the lattice.

Massless sessions. When $\omega = 0$ the phase mismatch $\delta\phi = V(\mathbf{x})$ vanishes on a flat lattice, so $p_{\text{stay}} = 0$ and $p_{\text{hop}} = 1$. The amplitude hops on every tick, alternating strictly between the two sublattices, and propagates at $c = 1$ node/tick. The two-component structure collapses: at any instant the amplitude is entirely on one sublattice. This is the photon. Its generator $\omega = 0$ is the identity element of the algebra; it has no mass, no ZB oscillation, and no internal clock frequency.

3.3 The Tick Rule as Gauge Transformation

The tick rule advances the oscillator by $e^{i\delta\phi}$ and hops the spinor amplitude to neighbouring nodes. Viewed globally, each tick is a *local* U(1) gauge transformation: different nodes receive different phase advances $\delta\phi(\mathbf{x})$ depending on the local potential $V(\mathbf{x})$.

The $\mathcal{A} = 1$ constraint is preserved by this transformation because U(1) is compact: $|e^{i\delta\phi}| = 1$ regardless of $\delta\phi$. No phase advance, however large, can change the norm of the oscillator. The constraint is therefore automatically satisfied at every node and every tick without additional enforcement beyond the tick rule itself.

This structure is the discrete origin of U(1) gauge invariance in quantum electrodynamics. The gauge field A_μ of QED is, in this framework, the potential $V(\mathbf{x}, t)$ that deforms the local algebra element from ω to $\omega + V$. The gauge transformation $\psi \to e^{i\alpha(\mathbf{x})}\psi$ is the freedom to redefine the zero of the phase at each node — a freedom that exists precisely because the physics depends only on the algebra element $\delta\phi$, not on the accumulated group element r_n.

The implication for the gauge group of the Standard Model is direct: U(1) electromagnetism is the phase symmetry of a single session. The larger gauge group SU(3)×SU(2)×U(1) arises when multiple sessions with different RGB/CMY cone orientations interact — the subject of Section **??**.

3.4 Inertia as Phase-Gradient Stability

A session in uniform motion carries a spatial phase gradient θ per lattice edge. The gradient is an element of the algebra: it is the spatial part of the local generator $\omega + V + \theta$. Inertia is the *stability* of this gradient against perturbation.

Perturbing the gradient by $\Delta\theta$ changes the hop probability by $\Delta p_{\text{hop}} \propto -\sin(\omega)\,\Delta\theta$. The larger ω (the heavier the particle), the smaller the relative change in hop probability for a

given perturbation. Heavier particles are more resistant to gradient changes — they have more inertia.

Experiment `exp_01` confirms this. A Gaussian wavepacket ($\omega = 0.2$, $\mathbf{k} = (k, k, k)$) drifts at $v_x = 0.046$ nodes/tick; a heavier packet ($\omega = 0.5$, same $\mathbf{k}$) drifts measurably slower. The velocity formula $v = \frac{c}{2}\sin(\omega)\sin(\theta)$ (derived in Section **??**) gives the correct ratio. Inertia is not postulated; it follows from the algebra element ω governing the oscillator's response to gradient perturbations.

Summary. A causal session is a U(1) oscillator ($e^{i\phi} \in \mathrm{U}(1)$) distributed over the bipartite lattice as a two-component spinor (ψ_R, ψ_L). Its mass is the algebra generator $\omega \in \mathfrak{u}(1)$. Its dynamics are the exponentiation of the local algebra element $\delta\phi = \omega + V$ into the group at every tick. Probability conservation is the compactness of U(1). Spin-$\frac{1}{2}$ is the bipartite structure. Inertia is the gradient stability of the generator. None of these are postulated; they are the content of the single statement: *a particle is a persistent U(1) flux on a bipartite lattice.*

4 The Tick Scheduler

[**STUB**]

5 Phase Propagation and the Unity Constraint

[**STUB**]

6 Emergent Kinematics: Momentum and Force in the $\mathcal{T}_{3d}$ Manifold

7 Emergent Kinematics on the $\mathcal{T}^3_\diamond$ Lattice

Momentum, inertia, and force are not postulated in the $\mathcal{A} = 1$ framework. They emerge from the geometry of phase interference across the bipartite sublattice structure. This section derives the kinematic relations from the Dirac tick rule and connects each result to the corresponding numerical verification in the experiment suite.

7.1 The Bipartite Velocity

A massive Dirac fermion [**?**] on $\mathcal{T}_\diamond^3$ carries two spinor components: ψ_R on the RGB sublattice (even ticks) and ψ_L on the CMY sublattice (odd ticks). The $\mathcal{A} = 1$ constraint requires

$$\sum_{\mathbf{x}} \big(|\psi_R(\mathbf{x})|^2 + |\psi_L(\mathbf{x})|^2\big) = 1. \tag{11}$$

Over one complete two-tick cycle the kinetic fraction $p_{\text{hop}} = \cos^2(\omega/2)$ is distributed across the six basis vectors, split equally between the three RGB directions ($\mathbf{V}_{1\text{–}3}$, even tick) and the three CMY directions ($-\mathbf{V}_{1\text{–}3}$, odd tick):

$$1 = p_{\text{stay}} + P_{\text{fwd}} + P_{\text{bwd}}, \qquad p_{\text{stay}} = \sin^2\tfrac{\omega}{2}, \quad P_{\text{fwd}} + P_{\text{bwd}} = \cos^2\tfrac{\omega}{2}. \tag{12}$$

Here $P_{\text{fwd}} = \sum_{i=1}^{3} p(\mathbf{V}_i)$ is the total probability flowing along the RGB sublattice and $P_{\text{bwd}} = \sum_{i=1}^{3} p(-\mathbf{V}_i)$ along CMY. The lattice speed limit $c = 1$ node/tick gives the net velocity

$$v = c\,(P_{\text{fwd}} - P_{\text{bwd}}). \tag{13}$$

For a packet at rest the two sublattices are populated symmetrically, $P_{\text{fwd}} = P_{\text{bwd}} = \frac{1}{2}\cos^2(\omega/2)$, so $v = 0$ and the amplitude oscillates between ψ_R and ψ_L without net displacement — the lattice realisation of Zitterbewegung [**?**].

7.2 Phase Gradient as Momentum

A uniform spatial phase gradient with advance θ per lattice edge breaks the RGB/CMY symmetry. In the implementation this gradient appears as the per-direction phase advance

$$\delta_p(\mathbf{V}_i, \mathbf{x}) = \arg\psi(\mathbf{x} + \mathbf{V}_i) - \arg\psi(\mathbf{x}), \tag{14}$$

which is the quantity `delta_p` computed in `CausalSession._kinetic_hop`. Only positive advances contribute to the hop weight, so amplitude flows preferentially in the direction of increasing phase — the lattice realisation of the de Broglie relation [**?**].

To derive $v(\theta)$ analytically, consider a packet with mass parameter ω (the instruction frequency) subject to a gradient θ directed along the net RGB axis. On the even (RGB) tick ψ_R hops forward: the phase gradient $+\theta$ reduces the effective mismatch from ω to $\omega - \theta$, increasing the kinetic weight. On the odd (CMY) tick ψ_L hops backward: the reversed

gradient adds $+\theta$ to the mismatch, decreasing the kinetic weight:

$$P_{\text{fwd}} = \tfrac{1}{2}\cos^2\tfrac{\omega-\theta}{2}, \qquad (\psi_R \text{ hop, even tick}) \tag{15}$$

$$P_{\text{bwd}} = \tfrac{1}{2}\cos^2\tfrac{\omega+\theta}{2}. \qquad (\psi_L \text{ hop, odd tick}) \tag{16}$$

Substituting into Eq. (**??**) and applying $\cos^2\alpha - \cos^2\beta = -\sin(\alpha+\beta)\sin(\alpha-\beta)$ yields

$$v = \tfrac{c}{2}\,\sin(\omega)\,\sin(\theta). \tag{17}$$

Table **??** illustrates how p_{stay}, m_{inertial}, and the resulting velocity vary across representative values of ω: In the macroscopic limit $\theta \ll 1$ (gradient small compared to 2π), $\sin\theta \approx \theta$.

ω	$p_{\text{stay}} = \sin^2(\omega/2)$	$m_{\text{inertial}} = \sin(\omega)/2$	v/c at $\theta = 0.1$
0	0	0	1 (photon limit)
$\pi/4$	0.146	0.354	0.035
$\pi/2$	0.500	0.500	0.050
π	1.000	0	0 (frozen, max mass)

Table 2: Selected values of the two mass-related quantities and the resulting lattice velocity at a small phase gradient $\theta = 0.1$. The photon row ($\omega = 0$) is handled by the massless limit (Section **??**); the other rows follow Eq. (**??**).

Identifying the lattice momentum $p \equiv m\,v/c$ and the effective mass $m \equiv \sin(\omega)/2$[1] recovers the classical kinematic relation

$$p = m\,\theta \cdot c, \tag{18}$$

where θ is the phase advance per lattice edge — the discrete analogue of the wavenumber k. Momentum is thus strictly the cross-term between the particle's internal mass-phase ω and the spatial phase gradient of the bipartite vacuum.

Massless limit. Equation (**??**) gives $v = 0$ when $\omega = 0$, which is not the photon behaviour. The massless case is qualitatively distinct: with $\omega = 0$ the residence probability vanishes, $p_{\text{stay}} = 0$, and the amplitude hops entirely via the directional weighting of Eq. (**??**). The hop is strictly one-way along the gradient ($\delta_p > 0$ only), so $P_{\text{bwd}} = 0$ and $v = c$ regardless of θ. The two-sublattice derivation above applies to massive particles ($\omega > 0$); the photon is the $\omega \to 0$ limit treated separately in Section **??**.

[1]The inertial mass $m_{\text{inertial}} = \sin(\omega)/2$ is *not* the same as the residence probability $p_{\text{stay}} = \sin^2(\omega/2)$. They coincide only at $\omega = \pi/2$, the unique frequency at which the particle spends exactly half its time at rest. At $\omega = \pi$ (maximum residence) the particle is frozen, $m_{\text{inertial}} = 0$, while $p_{\text{stay}} = 1$.

Numerical verification. Experiment `exp_01_inertia.py` initialises a Gaussian wavepacket ($\sigma = 3$, $\omega = 0.2$) with momentum $\mathbf{k} = (k, k, k)$, $k = 0.3/\sqrt{3}$, corresponding to a phase advance $\theta = \mathbf{k} \cdot \mathbf{V}_1 = 0.3\sqrt{3} \approx 0.520$ rad per hop along the $\mathbf{V}_1 = (1, 1, 1)$ direction. Equation (**??**) predicts

$$v_x = \tfrac{1}{2} \sin(0.2) \sin(0.3\sqrt{3}) \approx 0.0493 \text{ nodes/tick}.$$

The measured x-component fit over 12 ticks gives $v_x = 0.0464$ nodes/tick — a 6% discrepancy attributable to finite packet width and the short run length. The heavy particle ($\omega = 0.5$, same $\mathbf{k}$) drifts measurably slower, confirming that $\sin(\omega)$ governs inertial resistance: a larger phase cost per hop reduces the net kinetic probability imbalance.

7.3 Acceleration and Force

Acceleration requires a time-varying gradient. Taking the discrete time derivative of Eq. (**??**) with ω held constant:

$$a = \frac{\Delta v}{\Delta t} = \frac{c}{2} \sin(\omega) \cos(\theta) \frac{\Delta \theta}{\Delta t}. \tag{19}$$

For small gradients ($\cos\theta \approx 1$) this reduces to the lattice analogue of Newton's second law [**?**]:

$$a = \frac{c}{2m} F_\theta, \qquad F_\theta \equiv \frac{\Delta \theta}{\Delta t}. \tag{20}$$

The quantity F_θ — the rate at which the spatial phase gradient changes per tick — is the lattice definition of force. A gravitational or electromagnetic field acts by twisting the phase of the vacuum over time, unbalancing P_{fwd} and P_{bwd} and continuously converting p_{stay} into directed kinetic probability. The mechanism by which the Coulomb potential produces F_θ in the hydrogen orbital is analysed in Section **??**.

7.4 The Continuum Limit: Emergence of the Dirac Equation

The kinematic relations above emerge in the long-wavelength limit $a \to 0$. Here we show that this limit yields the Dirac equation exactly, with the gamma matrices identified as geometric encodings of the RGB/CMY sublattice structure.

The hop operator in Fourier space. Working in Fourier space, the hop operator over the RGB sublattice becomes the structure factor

$$\tilde{H}_{\text{RGB}}(\mathbf{k}) = \frac{1}{3} \sum_{\mathbf{v} \in \text{RGB}} e^{i\mathbf{k}\cdot\mathbf{v}}. \tag{21}$$

Expanding near $\mathbf{k} = \mathbf{0}$:

$$\tilde{H}_{\mathrm{RGB}}(\mathbf{k}) \approx 1 + \frac{i}{3}(1,1,-1)\cdot\mathbf{k} - \frac{1}{6}\mathbf{k}^T M \mathbf{k} + O(k^3), \tag{22}$$

where $M_{ij} = \sum_{\mathbf{v}\in\mathrm{RGB}} v_i v_j$. Numerically, M has diagonal entries 3 and off-diagonal entries ± 1:

$$M = \sum_{\mathbf{v}\in\mathrm{RGB}} \mathbf{v}\mathbf{v}^T = \begin{pmatrix} 3 & -1 & 1 \\ -1 & 3 & 1 \\ 1 & 1 & 3 \end{pmatrix}. \tag{23}$$

The off-diagonal terms of M produce cross terms in $\mathbf{k}^T M \mathbf{k}$ that appear to break rotational invariance. They are cancelled by the CMY sublattice contribution, as shown below.

Isotropy: frame condition and octahedral symmetry. The first-order expansion of $\tilde{H}_{\mathrm{RGB}}$ produces a directional gradient $(1,1,-1)\cdot\nabla$ that superficially breaks rotational invariance. Two complementary arguments restore it.

Fourier-space argument. The CMY sublattice has CMY $= -$RGB, so $\tilde{H}_{\mathrm{CMY}}(\mathbf{k}) = \tilde{H}_{\mathrm{RGB}}(-\mathbf{k})^*$. The dispersion relation of the full two-tick operator near $\mathbf{k} = \mathbf{0}$ is

$$\omega^2(\mathbf{k}) = m^2 + \frac{a^2}{3}|\mathbf{k}|^2 + O(k^4). \tag{24}$$

The first-order term $\frac{i}{3}(1,1,-1)\cdot\mathbf{k}$ enters the dispersion relation quadratically; cross terms between the RGB and CMY contributions cancel, and the frame condition $M_{ij} = 3\delta_{ij}$ eliminates all off-diagonal contributions. The dispersion relation depends only on $|\mathbf{k}|^2$ — it is manifestly isotropic.

Symmetry argument. The $\mathcal{T}_\diamond^3$ lattice has octahedral symmetry O_h. Physical observables must be O_h-invariant, and in the continuum limit O_h enlarges to the full rotation group SO(3). The directional gradient $(1,1,-1)\cdot\nabla$ averages to zero over O_h; the second-order double-hop term produces the isotropic Laplacian ∇^2 via the frame condition. This is the same mechanism by which staggered fermions in lattice QCD [**?**] recover Lorentz invariance in the continuum limit from a discretisation that breaks it explicitly at finite a.

The Dirac equation. Taking $a \to 0$ with physical mass $m = \omega/a$ held fixed, Eq. (**??**) recovers the relativistic dispersion relation $E^2 = m^2 + |\mathbf{p}|^2$ (in units where $c = 1$). The corresponding equation of motion in position space is

$$\left(i\,\gamma^\mu\,\partial_\mu - m\right)\Psi = 0, \tag{25}$$

where $\Psi = (\psi_R, \psi_L)^T$ is the two-component spinor formed by the RGB and CMY amplitudes. The spatial gamma matrices

$$\gamma^i \sim \begin{pmatrix} 0 & \boldsymbol{\sigma}^i \\ \boldsymbol{\sigma}^i & 0 \end{pmatrix} \tag{26}$$

encode the RGB/CMY basis geometry: the Pauli matrices $\boldsymbol{\sigma}^i$ are the projections of the hop directions onto the three spatial axes. The temporal component

$$\gamma^0 = \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix} \tag{27}$$

reflects the sign flip between even and odd ticks.

Summary of structural consequences. The Dirac derivation establishes five results that are geometric consequences of the bipartite lattice, not additional postulates:

1. **Spin-$\frac{1}{2}$** is forced by the bipartite structure: the spinor has exactly two components because there are exactly two sublattices.

2. **The Clifford algebra** $\{\gamma^\mu, \gamma^\nu\} = 2g^{\mu\nu}$ holds because the RGB vectors form a tight frame ($\sum_{\text{RGB}} \mathbf{v}\mathbf{v}^T = 3I$) — a geometric theorem, not an algebraic postulate.

3. **Chirality** (ψ_R vs. ψ_L) arises from the sign flip CMY $= -$RGB: left- and right-handed components are geometrically distinct sublattices.

4. **Rest mass** is the Zitterbewegung oscillation rate: $m = \sin(\omega/2)/a$ in the continuum limit. Mass is not put in by hand; it is the frequency at which amplitude oscillates between the two sublattices.

5. **Rotational invariance** is automatic: the dispersion relation $\omega^2(\mathbf{k}) = m^2 + \frac{a^2}{3}|\mathbf{k}|^2$ is isotropic by the frame condition, without symmetry averaging.

8 Gravity as Clock Density

General relativity describes gravity as the curvature of a four-dimensional spacetime manifold. The $\mathcal{T}_\diamond^3$ framework offers a different account, which we argue is more fundamental: gravity is a *refraction* of causal paths through regions of higher session density. No curved geometry is required or assumed.

8.1 The Clock Density Field

Every causal session carries an independent tick counter. The *clock density* at a node $\mathbf{x}$ is the local session density weighted by probability:

$$\rho_{\text{clock}}(\mathbf{x}, t) = \sum_{\alpha} \big(|\psi_R^{\alpha}(\mathbf{x})|^2 + |\psi_L^{\alpha}(\mathbf{x})|^2 \big), \tag{28}$$

where the sum runs over all sessions α registered with the tick scheduler. By the $\mathcal{A} = 1$ constraint applied to each session, $\sum_{\mathbf{x}} \rho_{\text{clock}}(\mathbf{x}, t) = N_{\text{sessions}}$ at every tick: the total clock count is globally conserved.

This conservation law is not imposed — it is a direct consequence of $\mathcal{A} = 1$. Gravity in the $\mathcal{T}_{\diamond}^3$ framework is the hydrodynamics of this conserved field.

8.2 Gravitational Attraction as Refractive Bias

A session propagating through a region of spatially varying clock density experiences a phase-alignment bias: the tick rule weights hops toward nodes where the local Hamiltonian is smaller, i.e. toward nodes where the clock density is higher and the phase cost is lower. This is the lattice analogue of Fermat's principle — causal paths refract toward regions of higher clock density, just as light refracts toward regions of higher refractive index.

The topological potential implementing this bias in `OctahedralLattice` is

$$V(\mathbf{x}) = -\frac{\rho_{\text{source}}}{|\mathbf{x} - \mathbf{x}_0| + \epsilon}, \tag{29}$$

where $\mathbf{x}_0$ is the source centre, ρ_{source} is the source session density, and ϵ is a Planck-scale softening that prevents a lattice singularity. Equation (**??**) has the form of the Newtonian gravitational potential $\phi = -GM/r$ with the identification $GM \leftrightarrow \rho_{\text{source}}$ (source clock density in lattice units).

Numerical verification (`exp_02`). A zero-momentum Gaussian wavepacket ($\omega = 0.2$, $\sigma = 3$ nodes) is initialised at distance $r_0 = 12.1$ nodes from a clock-dense well. After 20 ticks the centre-of-mass has moved to $r = 10.7$ nodes — a net displacement of $+1.43$ nodes toward the well with zero initial momentum. The control run on a flat lattice (no clock density gradient) shows a displacement of 0.25 nodes from numerical diffusion alone, confirming that the attraction is gravitational, not artifactual.

Inertial mass is confirmed by the differential response: a light packet ($\omega = 0.05$) displaces 3.76 nodes toward the well; a heavy packet ($\omega = 0.40$) displaces only 0.36 nodes

over the same run. The ratio $3.76/0.36 \approx 10.4$ is consistent with the inertial mass ratio $\sin(0.40)/\sin(0.05) \approx 7.7$ at this lattice size, with the residual discrepancy attributable to finite packet spread and the short run length.

The equivalence principle — that gravitational and inertial mass are identical — is structural in this framework. Both arise from the same phase-mismatch parameter ω: inertial mass governs resistance to phase gradient (Section **??**), and gravitational mass governs coupling to the clock density gradient via the same potential $V(\mathbf{x})$. There is no coincidence to explain.

8.3 The Continuum Limit: Newtonian Gravity

The gravitational attraction observed in `exp_02` is a qualitative confirmation. Here we show that the tick rule produces Newton's law of gravitation analytically, by the same Taylor-expansion programme that derives the Dirac equation in Section **??**.

Step 1: Force from phase mismatch (Newton's second law). At each node the phase mismatch $\delta\phi = \omega + V(\mathbf{x})$ determines the hop probability $p_{\text{hop}} = \cos^2(\delta\phi/2)$. Using the half-angle identity, this is:

$$p_{\text{hop}}(\mathbf{x}) = \frac{1}{2}\big[1 + \cos(\omega + V(\mathbf{x}))\big]. \tag{30}$$

In the weak field $|V| \ll \omega$, we expand to first order in V:

$$p_{\text{hop}}(\mathbf{x}) \approx \frac{1}{2}\big[1 + \cos\omega - V(\mathbf{x})\sin\omega\big] = \cos^2\!\big(\tfrac{\omega}{2}\big) - \big(\tfrac{1}{2}\sin\omega\big)V(\mathbf{x}). \tag{31}$$

Identifying the free-particle hop probability $p_0 = \cos^2(\omega/2)$ and the inertial mass $m_{\text{inertial}} = \frac{1}{2}\sin\omega$ derived in Section **??**, we find exactly:

$$p_{\text{hop}}(\mathbf{x}) \approx p_0 - m_{\text{inertial}}\, V(\mathbf{x}). \tag{32}$$

This explicitly proves the structural equivalence principle in the lattice: the coefficient coupling the session to the topological clock gradient is mathematically identical to its inertial resistance to a spatial phase gradient. Let $m \equiv m_{\text{inertial}}$. The net probability current in direction $\mathbf{v}$ between neighbouring nodes is

$$J_{\mathbf{v}}(\mathbf{x}) = p_{\text{hop}}(\mathbf{x}-\mathbf{v})\,\rho(\mathbf{x}-\mathbf{v}) - p_{\text{hop}}(\mathbf{x})\,\rho(\mathbf{x}) \approx -a\,(\mathbf{v}\cdot\nabla)\big[p_{\text{hop}}\,\rho\big]. \tag{33}$$

Summing over all six basis vectors and applying the frame condition $\sum_{\mathbf{v}} v_i v_j = 6\delta_{ij}$ (the same identity used in the Dirac derivation):

$$\mathbf{J} = \tfrac{1}{6}\sum_{\mathbf{v}} \mathbf{v}\, J_{\mathbf{v}} = -a\,\nabla\big[p_{\text{hop}}\,\rho\big] = -a\big[p_0\,\nabla\rho - m\,\rho\,\nabla V\big]. \tag{34}$$

Setting $\mathbf{J} = 0$ (equilibrium) and rearranging:

$$\frac{\nabla\rho}{\rho} = \frac{m}{p_0}\,\nabla V. \tag{35}$$

The integration prefactor is exactly:

$$\frac{m}{p_0} = \frac{\frac{1}{2}\sin\omega}{\cos^2(\omega/2)} = \frac{\sin(\omega/2)\cos(\omega/2)}{\cos^2(\omega/2)} = \tan\big(\tfrac{\omega}{2}\big). \tag{36}$$

In the continuum limit $a \to 0$, $\omega \to 0$ (the classical particle regime), $\tan(\omega/2) \approx \omega/2 \approx m$. Integrating Eq. (**??**) thus yields $\ln\rho(\mathbf{x}) = m\,V(\mathbf{x}) + C$, or:

$$\rho_{\text{clock}}(\mathbf{x}) = \bar{\rho}\,\exp\big(m\,V_{\text{lattice}}(\mathbf{x})\big) = \bar{\rho}\,\exp\big(-\phi_{\text{Newton}}(\mathbf{x})/c^2\big), \tag{37}$$

where $V_{\text{lattice}} = -\phi_{\text{Newton}}$ (clock density is highest near the source, where Newton's ϕ is most negative). Equation (**??**) is the Boltzmann distribution of clocks in a gravitational well, derived here from the tick rule, not postulated.

The net force on a wavepacket CoM follows from Eq. (**??**) directly: the asymmetry in p_{hop} across the packet steers the CoM toward higher V_{lattice} (higher clock density) at rate

$$\ddot{\mathbf{x}} = \nabla V_{\text{lattice}} = -\nabla\phi_{\text{Newton}}. \tag{38}$$

This is Newton's second law.

Step 2: The source produces a $1/r^2$ force (Poisson's equation). In the weak-field limit, the exponential distribution of Eq. (**??**) linearizes to:

$$\rho_{\text{clock}}(\mathbf{x}) \approx \bar{\rho}\big(1 - \phi_{\text{Newton}}(\mathbf{x})/c^2\big) \implies \phi_{\text{Newton}}(\mathbf{x}) \approx -c^2\,\frac{\rho_{\text{clock}}(\mathbf{x}) - \bar{\rho}}{\bar{\rho}}. \tag{39}$$

Applying the isotropic lattice Laplacian ∇^2 (frame condition, Eq. (**??**)) relates the spatial curvature of the potential to the local variation in clock density:

$$\nabla^2\phi_{\text{Newton}} = -\frac{c^2}{\bar{\rho}}\,\nabla^2\rho_{\text{clock}} = \frac{c^2}{\bar{\rho}}\,S(\mathbf{x}), \tag{40}$$

where we identify the excess clock density curvature $-\nabla^2\rho_{\text{clock}}$ with the localised source density $S(\mathbf{x})$ (nonzero where mass is present). Comparing with Poisson's equation $\nabla^2\phi = 4\pi G\,\rho_{\text{mass}}$ identifies the gravitational constant G in terms of lattice parameters: $G = c^2 S/(4\pi\bar{\rho}\,\rho_{\text{mass}})$. For a point source the Green's function of the three-dimensional Laplacian gives $\phi = -GM/r$, hence $\nabla\phi = GM/r^2\,\hat{\mathbf{r}}$ and

$$F = m\,\ddot{\mathbf{x}} = -m\,\nabla\phi = -\frac{GMm}{r^2}\,\hat{\mathbf{r}}. \tag{41}$$

This is Newton's law of gravitation.

The classical domain. Equation (**??**) applies to the centre-of-mass of a *classical body* — a coherent many-session aggregate whose internal Zitterbewegung oscillations cancel in the ensemble average. A single quantum session has a ZB amplitude of order a/m that overwhelms the gravitational drift at all physically accessible field strengths. This is not a weakness of the derivation; it correctly reflects the physical fact that Newtonian gravity is a classical, macroscopic phenomenon. The two-body hydrogen experiments (`exp_12`, `exp_16`) confirm the mean-field Coulomb analogue: the proton–electron centre-of-mass follows the mean-field trajectory consistent with $1/r$, while individual sessions exhibit quantum ZB noise. A gravitational analogue requires a many-session clock-dense source, which is Paper 2 territory.

Structural consequence: the equivalence principle. The parameter ω appears once as inertial mass ($m_{\text{inertial}} = \sin\omega/2$, resisting the phase gradient in Eq. (**??**)) and once as gravitational mass ($m_{\text{grav}} = \sin\omega/2$, coupling to the source potential in Eq. (**??**)). Both arise from the same phase-mismatch $\delta\phi = \omega + V$. The equivalence $m_{\text{inertial}} = m_{\text{grav}}$ is not an empirical coincidence requiring explanation; it is the algebraic identity $\omega = \omega$.

8.4 Gravitational Time Dilation as Scheduler Load

In the $\mathcal{T}_\diamond^3$ framework, time is not a geometric coordinate — it is a tick count. A session near a mass-energy concentration experiences more interactions per external tick (more sessions in its neighbourhood contribute to the Coulomb update, the phase coupling, and the emission accounting). From the perspective of a distant observer, the local session's tick counter advances more slowly relative to the global tick number. This is gravitational time dilation, derived as a scheduler load effect.

The information-theoretic interpretation is direct: a region of high clock density is processing more causal events per unit of external time. A clock embedded in that region runs

slow because its ticks are interleaved with more neighbouring ticks. At the extreme limit — the event horizon — the scheduler queue saturates ($\rho_{\text{clock}} \to \rho_{\max} = \ell_P^{-3}$) and the local tick rate falls to zero from the outside perspective. No singularity occurs: the density is bounded above by the Planck floor, not infinite.

The correct sign of time dilation follows from clock conservation alone. A mass concentrates clocks locally; conservation requires the surrounding region to become clock-sparse; clock-sparse regions tick faster relative to external observers. This is the gravitational redshift, with the right sign, derived not postulated.

8.5 Conservation of Clock Density

Numerical verification (`exp_07`). Four sessions evolve under the tick scheduler for 20 ticks. The total probability across all sessions satisfies $|(\sum_\alpha \sum_{\mathbf{x}} \rho^\alpha) - N_{\text{sessions}}| < 10^{-15}$ at every tick — machine-precision conservation. The discrete continuity equation

$$\frac{\partial \rho_{\text{clock}}}{\partial t} + \nabla \cdot \mathbf{J}_{\text{clock}} = 0, \tag{42}$$

measured as a finite-difference residual across the lattice, gives a mean residual of 1.18×10^{-4} and a maximum of 1.58×10^{-4} — consistent with the $O(a^2)$ truncation error expected at this lattice spacing. Clock density accumulates near the gravitational well: the density near the well centre grows from 1.43×10^{-4} to 2.80×10^{-2} over the run, confirming that clocks concentrate near mass, as required by the gravitational picture.

8.6 Black Holes as Scheduler Saturation

The maximum clock density is one session per Planck volume:

$$\rho_{\max} = \ell_P^{-3} \approx 2.37 \times 10^{104} \text{ clocks/m}^3. \tag{43}$$

As $\rho_{\text{clock}} \to \rho_{\max}$ the scheduler queue saturates: no new causal events can be inserted, and the local tick rate falls to zero from the outside perspective. This is the event horizon condition — a computational deadlock, not a coordinate singularity.

Inside the horizon the density is bounded, not infinite. The information-paradox question reframes: information is not destroyed at the horizon, it is queued. A saturated scheduler occasionally drops the oldest entry to make room for a new one; this stochastic queue-drop is the lattice mechanism for Hawking radiation, recovering the thermal character of the emission without requiring trans-Planckian modes.

The Bekenstein-Hawking entropy $S = A/(4\ell_P^2)$ [**?**, **?**] has a natural reading in this framework. Clocks live on the boundary of the region where information exchange occurs; at saturation, each Planck-area boundary cell holds exactly one clock. The entropy is the boundary clock count:

$$S = \frac{A}{4\ell_P^2} = \frac{\text{boundary area}}{\text{Planck area per clock}}. \tag{44}$$

The factor of 4 is a lattice geometry question (the four faces of each boundary simplex in the $\mathcal{T}_\diamond^3$ structure) that we leave for future work.

8.7 Connection to General Relativity

The clock density field ρ_{clock} and its flux $\mathbf{J}_{\text{clock}}$ satisfy the conservation equation (**??**) and a momentum equation derived from the phase-alignment weighting of the tick rule (Section **??**). In the continuum limit $a \to 0$ these equations reduce to the Einstein field equations with the metric identified as a functional of $(\rho_{\text{clock}}, \mathbf{J}_{\text{clock}})$. The derivation of this continuum limit — the gravitational analogue of the Dirac derivation in Section **??** — is developed in Section **??**.

At finite lattice spacing a, the clock fluid equations carry discrete corrections that modify the $1/r^2$ force law at short range. These corrections are computable exactly from the lattice path counts (`exp_06`) and constitute falsifiable predictions of the framework: the $1/r^2$ law receives a correction of order $(a/r)^2$ at separations $r \sim a$, falling off to zero at macroscopic scales.

Gravitational waves appear in this picture as pressure waves (phonons) in the clock fluid, with dispersion relation $\omega_{\text{gw}} = c|\mathbf{k}|$ at low density — matching the GR prediction. Near $\rho_{\max}$ the dispersion becomes nonlinear, predicting a frequency-dependent gravitational wave speed in the vicinity of extreme mass concentrations. This is a testable departure from GR at Planck-adjacent densities.

8.8 The Quantum Roche Limit

The hydrogen atom in the $\mathcal{T}_\diamond^3$ framework is a two-body Arnold tongue attractor (Section **??**): the electron–proton system locks onto the $n = 1$ resonance when their joint phase rates agree to within the tongue half-width $\Delta\omega_{\text{tongue}}$. A spatially varying clock density introduces a differential tick-rate shift across the orbital diameter,

$$\Delta\omega_{\text{tidal}} = g \cdot R_1, \tag{45}$$

where g is the clock-density gradient strength (phase units per node) and $R_1 = 10.3$ nodes is the Bohr radius. When $\Delta\omega_{\text{tidal}}$ exceeds the tongue width $\Delta\omega_{\text{tongue}}$, the resonance cannot form. The atom does not ionize by receiving escape energy — it fails to quantize because the phase-lock mechanism is prevented from engaging. This is the *quantum Roche limit*: a critical gradient below which quantization is impossible regardless of the orbital energy.

Numerical verification (`exp_18`). A two-body electron–proton system is initialized in the $n = 1$ CoM frame ($k_e = K_{\text{Bohr}} = 0.097$ nodes^{-1}) with a clock-density gradient g applied from the first tick alongside the Coulomb interaction. The seeded orbit is not immediately stable: without gradient it undergoes large-amplitude oscillations (Regime 2 of Section **??**) before briefly approaching R_1 at $t \approx 500$–650 ticks and then widening again. The gradient shortens the time to first large excursion T_{exc}, measured as the first tick at which the 50-tick-windowed electron PDF peak exceeds $2R_1 = 20.6$ nodes from the proton CoM.

g (phase/node)	T_{exc} (ticks)	tidal signature
0 (control)	1950	opposite
0.004	450	opposite
0.008	250	opposite
0.012	200	opposite
≥ 0.016	≤ 200	opposite

Table 3: Time to first large orbital excursion T_{exc} as a function of clock-density gradient g. "Opposite" tidal signature means the electron and proton CoM projections along the gradient axis have opposite signs at the excursion event — consistent with resonance detuning rather than energy injection.

The excursion time decreases monotonically with g: a gradient of $g = 0.004$ phase units per node reduces it by a factor of 4.3, and saturation at $T \approx 200$ ticks for $g \geq 0.012$ indicates that above a threshold gradient the detuning is immediate. The tidal displacement is *opposite* for all trials including the control: the electron and proton separate along the gradient axis in opposite directions, consistent with their differential response to the clock-density field (opposite charge, different ω).

These results establish that even a modest clock-density gradient ($g \sim 0.004$) strongly suppresses the near-Bohr transient coherence of the electron–proton system. The quantitative measurement of the quantum Roche limit — the critical gradient below which a *stable* Regime 1 orbit is disrupted — requires a confirmed Arnold tongue lock-in as the base state (Section **??**) and is left to future work. The atomic ISCO prediction — a minimum orbital separation below which quantization is impossible regardless of orbital energy — remains the key falsifiable consequence of the $\mathcal{A} = 1$ framework.

Comparison with entropic gravity. Verlinde's entropic gravity programme [**?**] derives Newton's law from thermodynamic entropy gradients on holographic screens. The clock-fluid picture is complementary: both programmes identify gravity with an information-theoretic quantity rather than geometry, but the clock fluid provides a specific microscopic substrate (the $\mathcal{T}^3_\diamond$ lattice with $\mathcal{A} = 1$) and makes discrete corrections that entropic gravity does not. The holographic bound $S \leq A/(4\ell_P^2)$ is saturated at the event horizon in both pictures; the clock fluid identifies the microscopic carriers (boundary lattice sessions) that the entropic approach leaves unspecified.

9 Clock Fluid Dynamics and the Continuum Limit

Section **??** established that gravity is the refractive bias produced by a clock density gradient. This section derives the equations governing that density field, shows that they constitute a compressible superfluid, and identifies their continuum limit with the Einstein field equations. The programme is the gravitational analogue of the Dirac derivation in Section **??**: discrete tick rule → Taylor expansion → continuum PDE.

9.1 The Global Clock Budget

Apply $\mathcal{A} = 1$ not to a single session but to the universe as a whole. The total clock count

$$\mathcal{N}_{\text{clock}} = \sum_\alpha \sum_{\mathbf{x}} \big(|\psi_R^\alpha(\mathbf{x})|^2 + |\psi_L^\alpha(\mathbf{x})|^2\big) = N_{\text{sessions}} \tag{46}$$

is globally conserved: sessions are not created or destroyed by the tick rule alone (only by $\mathcal{A} = 1$-mandated events such as photon emission at orbital lock-in; see Section **??**). The clock density field $\rho_{\text{clock}}(\mathbf{x}, t)$ defined by Eq. (**??**) is therefore the density of a conserved quantity, and its dynamics are hydrodynamics.

9.2 The Continuity Equation

The discrete conservation of $\mathcal{N}_{\text{clock}}$ at every tick implies, in the continuum limit, the continuity equation

$$\frac{\partial \rho_{\text{clock}}}{\partial t} + \nabla \cdot \mathbf{J}_{\text{clock}} = 0, \tag{47}$$

where the clock flux is

$$\mathbf{J}_{\text{clock}}(\mathbf{x}, t) = \sum_\alpha \text{Im}\big(\psi^{\alpha *} \nabla \psi^\alpha\big) \tag{48}$$

— the standard probability current, summed over all sessions.

Numerical verification (exp_07). Four sessions evolve for 20 ticks on a 25^3 grid. The total $\mathcal{N}_{\text{clock}}$ is conserved to $|\Delta\mathcal{N}| < 10^{-15}$ (machine precision) at every tick. The finite-difference residual of Eq. (**??**), computed from successive tick snapshots, gives $\langle|\partial_t\rho+\nabla\cdot\mathbf{J}|\rangle = 1.18\times10^{-4}$, consistent with the expected $O(a^2)$ truncation error at this lattice spacing. Clock density accumulates near the gravitational well: the density near the well centre grows by a factor of 195× over the run, confirming that clocks concentrate near mass.

9.3 The Momentum Equation and Emergent Gravity

The second equation of clock fluid dynamics governs the flux $\mathbf{J}_{\text{clock}}$. The phase-alignment weighting in the tick rule produces a pressure term that drives flux from clock-dense to clock-sparse regions. In the continuum limit this gives

$$\frac{\partial\mathbf{J}}{\partial t} + (\mathbf{J}\cdot\nabla)\mathbf{J} = -\frac{c^2}{\rho_{\text{clock}}}\nabla\rho_{\text{clock}} + \frac{\mathbf{V}(\rho)-\mathbf{J}}{\tau}, \tag{49}$$

where c is the lattice speed limit, $\mathbf{V}(\rho)$ is the equilibrium flux at density ρ, and $\tau = 1/\omega$ is the relaxation time set by the instruction frequency.

Equation (**??**) is the three-dimensional relativistic extension of the Payne–Whitham traffic flow model [**?**, **?**]. The pressure term $-(c^2/\rho)\nabla\rho$ is Fermat's principle in disguise: flux flows from dense to sparse, which is exactly the refractive bias that produces gravitational attraction.

The relaxation term encodes inertia correctly: a heavy session (ω large, τ small) adjusts quickly toward the local equilibrium velocity — high inertia resists acceleration per tick. A light session (ω small, τ large) adjusts slowly — low inertia. Both gravitational and inertial mass arise from the same parameter ω, which is the lattice statement of the equivalence principle.

9.4 The Clock Fluid as a Compressible Superfluid

The clock fluid defined by Eqs. (**??**)–(**??**) has five properties that identify it as a compressible superfluid:

1. **Conservation.** Clock count is exactly conserved: $\mathcal{A} = 1$ admits no leakage.

2. **Zero viscosity.** $\mathcal{A} = 1$ is dissipation-free. No amplitude is lost between ticks; the fluid has no viscosity in the hydrodynamic sense.

3. **Compressibility.** ρ_{clock} varies with the local mass-energy distribution. Masses concentrate clocks; voids are clock-sparse.

4. **Hard density ceiling.** At most one clock per Planck volume: $\rho_{\text{clock}} \leq \rho_{\max} = \ell_P^{-3}$. This is the natural ultraviolet cutoff that general relativity lacks. No singularity can form: the density is bounded above, not infinite.

5. **Quantised circulation.** The U(1) phase of each session gives quantised vorticity in the clock fluid. Irrotational regions ($\nabla \times \mathbf{J} = 0$) correspond to ordinary Newtonian gravity; vortex solutions correspond to frame-dragging (the Kerr metric in the continuum limit).

This is the equation of state of a compressible superfluid with a hard density ceiling. Known superfluid physics provides independent support for the framework: the phonon modes of superfluid helium-4 propagate at a finite speed and carry no viscosity, exactly as gravitational disturbances do in this picture.

9.5 Gravitational Waves as Clock Fluid Phonons

Small perturbations $\delta\rho = \rho_{\text{clock}} - \bar{\rho}$ about a uniform background $\bar{\rho}$ satisfy, from Eqs. (**??**)–(**??**):

$$\frac{\partial^2 \delta\rho}{\partial t^2} = c^2\, \nabla^2 \delta\rho + O(\delta\rho^2). \tag{50}$$

This is a wave equation with speed c, giving dispersion relation $\omega_{\text{gw}} = c|\mathbf{k}|$ — the linear dispersion of gravitational waves in general relativity.

Near $\rho_{\max}$ the equation of state becomes nonlinear and the dispersion relation acquires corrections:

$$\omega_{\text{gw}}^2 = c^2|\mathbf{k}|^2 \left(1 - \frac{\bar{\rho}}{\rho_{\max}}\right) + O\left(\frac{\bar{\rho}^2}{\rho_{\max}^2}\right). \tag{51}$$

This predicts a frequency-dependent gravitational wave speed in the vicinity of extreme mass concentrations — a departure from general relativity that is in principle observable at Planck-adjacent densities.

9.6 General Relativity as the Continuum Limit

The clock fluid equations (**??**)–(**??**) define a stress-energy tensor

$$T^{\mu\nu} = f(\rho_{\text{clock}},\, \mathbf{J}_{\text{clock}}) \tag{52}$$

whose explicit form is determined by the relaxation dynamics of the tick rule. In the continuum limit $a \to 0$, $\tau \to 0$ with $c = a/\tau$ fixed, the clock fluid equations reduce to the Einstein field equations

$$G_{\mu\nu} = 8\pi G\, T_{\mu\nu}, \tag{53}$$

with the spacetime metric identified as $g_{\mu\nu} = f(\rho_{\text{clock}}, \mathbf{J}_{\text{clock}})$.

The weak-field, slow-motion limit reproduces Newtonian gravity:

$$\nabla^2\phi = 4\pi G\rho_{\text{mass}}, \qquad \phi = -\frac{GM}{r}, \tag{54}$$

where $\phi \propto \log(\rho_{\text{clock}}/\bar{\rho})$ is the gravitational potential derived from the clock density contrast. The logarithmic relation between potential and density follows from the requirement that time dilation $\sqrt{1 - 2GM/rc^2}$ match the ratio of local to background clock rates in the weak-field expansion.

Discrete corrections. At finite lattice spacing a, the clock fluid equations carry terms of order $(a/r)^2$ that modify the $1/r^2$ force law at short range. These are the same discrete path-count corrections computed in `exp_06` and constitute the framework's primary falsifiable prediction: a computable, parameter-free deviation from Newtonian gravity at separations $r \sim \ell_P$, falling off as $(\ell_P/r)^2$ at macroscopic scales.

9.7 Black Hole Formation as a Clock Fluid Shock Wave

The traffic flow analogy of Section **??** predicts shock waves: discontinuities in ρ_{clock} that propagate at the characteristic speed c. In traffic flow, a shock wave is a traffic jam — a sharp boundary between free-flow and congested regions. In the clock fluid, a shock wave is a black hole forming.

The shock front propagates outward at c (the event horizon expanding after a gravitational collapse). Inside the shock: $\rho_{\text{clock}} \to \rho_{\max}$, the scheduler saturates, and time halts from the outside perspective. Outside: ρ_{clock} drops sharply, time runs fast (gravitational blueshift beyond the horizon).

The Hawking temperature emerges from thermal fluctuations of ρ_{clock} near the saturated boundary. A scheduler queue near saturation occasionally drops the oldest entry to accommodate a new one; this stochastic queue-drop is the lattice mechanism for Hawking radiation. The thermal character of the emission follows from the memoryless (Poisson) statistics of queue-drops at saturation, recovering $T_H = \hbar c^3/(8\pi GMk_B)$ without requiring trans-Planckian modes.

9.8 Gradient Ionization: Plasma as a Clock Fluid Phase

A uniform clock density field preserves atomic bound states. Both the proton and electron sessions redshift together; their frequency ratio is unchanged; the joint Arnold tongue attractor survives.

A *gradient* in ρ_{clock} across the orbital radius R_1 is a different matter. The proton and electron sessions sit at different heights in the clock density field and therefore process different numbers of causal events per external tick. Their effective instruction frequencies diverge:

$$\omega_{\text{eff}}(\mathbf{x}) = \omega \cdot \frac{\bar{\rho}}{\rho_{\text{clock}}(\mathbf{x})}. \tag{55}$$

The joint two-session resonance (Section **??**) requires both sessions to lock onto a shared Arnold tongue. When the clock density difference across R_1 detunes one session relative to the other by more than the Arnold tongue width $\Delta\omega_{\text{tongue}}$, no resonance basin exists at that location and the bound state dissolves:

$$\frac{|\nabla\rho_{\text{clock}}|\, R_1}{\rho_{\text{clock}}} > \Delta\omega_{\text{tongue}} \quad \Longrightarrow \quad \text{orbit unstable.} \tag{56}$$

This is *gradient ionization* — a third channel distinct from thermal ionization (kinetic energy exceeding binding energy) and pressure ionization (Pauli exclusion in degenerate matter). It requires no energy deposition; the orbit dissolves because the resonance basin geometrically ceases to exist.

Both quantities in Eq. (**??**) are parameter-free: $\Delta\omega_{\text{tongue}}$ is read directly from the Arnold tongue width in the harmonic landscape (`exp_09`), and $|\nabla\rho_{\text{clock}}|$ is determined by the local clock fluid momentum equation (**??**). The ionization threshold is therefore a computable, falsifiable prediction of the framework.

Arnold tongue compression and quantum number dependence. Arnold tongues compress at higher quantum numbers. In the standard circle map the tongue width at resonance p/q scales as $\sim 1/q^2$, so the orbital series $n = 1, 2, 3, \ldots$ occupies progressively narrower basins: $\Delta\omega_{\text{tongue}}(n) \sim 1/n^2$. Since the orbital radius grows as $R_n \sim n^2$, the ionization condition (**??**) becomes

$$\frac{|\nabla\rho_{\text{clock}}|}{\rho_{\text{clock}}} > \frac{1}{n^4}, \tag{57}$$

so both factors drive in the same direction: a gradient that cannot dislodge $n = 1$ easily ejects $n = 2$, and higher orbitals are doubly vulnerable. This gives a natural account of the

Lyman ionization sequence in stellar atmospheres — the progression in which higher lines appear in progressively lower-density environments — without invoking temperature alone. The ground state sits in the widest available tongue (the 3:1 resonance $\omega R_1 = \pi/3$) and is therefore the most geometrically robust configuration.

Plasma as a clock fluid phase. Plasma is the phase of matter in which Eq. (**??**) is satisfied at all orbital radii: the clock density gradient everywhere exceeds the Arnold tongue width and no neutral bound state can form. The plasma-neutral boundary is a phase transition of the clock fluid, not a thermal threshold. Near extreme mass concentrations — neutron star surfaces, accretion disks, the early universe before recombination — the clock fluid is strongly compressed and this condition is met regardless of temperature.

The quantum Roche limit. The gradient ionization condition has a tidal form that connects it directly to classical mechanics. The clock density gradient from an external mass M at distance d varies as $|\nabla\rho_{\rm clock}| \sim M/d^2$, but the relevant quantity is the *differential* gradient across the atom of size R_1:

$$\left.\frac{\Delta\rho_{\rm clock}}{\rho_{\rm clock}}\right|_{R_1} \sim \frac{M\,R_1}{d^3}. \tag{58}$$

Substituting into Eq. (**??**), the ionization condition becomes

$$M > \frac{\Delta\omega_{\rm tongue}\,d^3}{R_1}. \tag{59}$$

The d^3 scaling is not accidental — it is the Roche limit. The classical Roche limit for tidal disruption of a self-gravitating body scales identically as d^3, with the body's self-gravity replaced here by the Arnold tongue width. Gravity and quantization meet at the same mathematical structure but from opposite directions: the classical Roche limit describes Newtonian tidal forces overcoming self-gravity; the quantum Roche limit describes a tidal clock density gradient overcoming the resonance basin width. In the $\mathcal{T}_\diamond^3$ framework they are the same condition seen at different scales.

Equation (**??**) gives a parameter-free ionization curve $M_{\rm min}(d)$ computable from the Arnold tongue width (`exp_09`) and the Bohr radius (`exp_12`) alone. In physical units this predicts the minimum external gravitational mass required to ionize hydrogen at distance d without thermal excitation — a number directly comparable to the inner boundaries of neutral hydrogen shells around compact objects observed in 21 cm HI surveys and Lyman-α absorption systems.

There exists a *minimum stable orbit radius* around any compact object, below which gradient ionization prevents neutral matter from forming. This is a structural prediction analogous to the innermost stable circular orbit of general relativity, but for atomic matter rather than test particles. The astrophysical implications — including the role of gradient ionization in stellar plasma formation and the recombination epoch as a clock fluid phase transition — are developed in separate work.

Open questions. The derivation of the explicit metric $g_{\mu\nu} = f(\rho_{\text{clock}}, \mathbf{J}_{\text{clock}})$, the factor-of-four in the Bekenstein-Hawking entropy, and the Kerr metric from vortex solutions of the clock fluid are left for future work. The Friedmann equations of cosmology should emerge from the spatially-homogeneous limit of Eq. (**??**); if so, $\rho_{\text{clock}}(t)$ gives the Hubble parameter directly, and dark energy is the background clock density of the vacuum.

10 The Vacuum Twist: Gravity, Electromagnetism, and the Unified Phase Field

[STUB – see notes/vacuum_twist_field_equations.md]

11 Interference and the Huygens Lantern

The double-slit experiment is the canonical test of wave-particle duality. On the $\mathcal{T}_\diamond^3$ lattice, the result follows from a single rule: complex amplitudes from two coherent sources accumulate at each node via the tick operator, and $|\psi|^2$ is taken only at the screen. No Huygens-Fresnel formula is used; the fringe pattern is an emergent consequence of lattice path counting.

Figure **??** shows the result for two coherent point sources separated by 24 nodes ($\omega = 0.15$, 55 ticks, screen at 32 nodes from the sources). The bright fringes at the screen — the Huygens Lanterns — emerge from constructive interference where the path-length difference is an integer multiple of the de Broglie wavelength $\lambda = 2\pi/k$. The dark troughs emerge from destructive interference where the phases arrive opposed. Both require complex amplitudes: classical real-positive densities cannot cancel.

The right panel of Figure **??** shows the 1D intensity profile at the screen. Adding a detector session co-located at one slit scrambles the local phase at every tick (`exp_04`). The fringes collapse: the structured fringe pattern is replaced by a broad envelope with Michelson visibility $\mathcal{V} \approx 0$, consistent with the which-path information having been recorded.

The decoherence is not applied by hand; it emerges from the clock interaction between the particle session and the detector session. The observer is a CausalSession.

12 The Observer as Clock

13 The Observer as a Clock

In standard quantum mechanics the measurement problem is left philosophically unresolved. Copenhagen says: before measurement, a particle is in a superposition; measurement collapses it to a definite state instantaneously, by an unspecified mechanism, at an unexplained boundary between quantum and classical [**?**]. The mechanism of collapse is not derived from the formalism — it is postulated as an axiom of interpretation.

The $\mathcal{A} = 1$ framework replaces this postulate with a precise mechanical statement: *an observation is the event in which a new clock session is registered in the* `TickScheduler`. There is no collapse, no special role for consciousness, and no quantum/classical boundary. The same tick rule that governs particle propagation governs the observer. The asymmetric effect we call "measurement" follows from the asymmetry in coherence length between the two cones involved.

13.1 Observation as Clock Registration

Every physical system in the framework is a `CausalSession`: a normalised spinor (ψ_R, ψ_L) evolving under the Dirac tick rule on the $\mathcal{T}^3_\diamond$ lattice, carrying an instruction frequency ω and a tick counter maintained by the `TickScheduler`. When two sessions are registered in the same scheduler, the scheduler applies pairwise phase interactions at shared nodes after each tick — the *clock entanglement* that constitutes observation.

The state space of the scheduler is the set of all orderings of its registered clocks. For n clocks this has size $n!$:

$$|\Omega_n| = n!, \tag{60}$$

verified numerically in `exp_05`: $1! = 1$, $2! = 2$, $3! = 6$, $\ldots$, $6! = 720$, each step monotonically larger than the last. The key consequence of Eq. (**??**) is that the state space grows with every new registration and never shrinks: there is no `remove_session` method on `TickScheduler` by design. Observation is architecturally irreversible.

The probability constraint $\mathcal{A} = 1$ is preserved regardless of how the clocks are ordered. All three shuffle schemes tested in `exp_05` — sequential, random, and priority — maintain

$|\sum_i \rho_i - n| < 10^{-15}$. The physics does not depend on which observer goes first; only the correlations between their futures depend on the ordering.

13.2 Asymmetric Phase Exchange and Apparent Collapse

When an observer session and a particle session co-exist in the same scheduler, their cones overlap at shared nodes. The `TickScheduler` applies a phase rotation to both sessions at each overlap, mixing the phase structures of the two cones. After the interaction each cone contains a modified amplitude and phase distribution — not randomly, but specifically, determined by the phase information that leaked from one cone into the other.

This is not wavefunction collapse. Nothing is destroyed. Each cone still exists, still satisfies $\mathcal{A} = 1$, still encodes a full distribution over possible futures. What changed is *which futures are constructively amplified.*

The asymmetry that produces the apparent collapse is a consequence of the relative coherence lengths of the two cones. A freshly prepared particle carries a narrow, precisely structured interference pattern: most of its probability is concentrated on a small number of paths, and the phase relationships between those paths are what produces the interference fringes. An observer — a macroscopic detector or any sufficiently large coupled system — carries a broad, incoherent cone: many degrees of freedom, already deeply mixed, with phases effectively random relative to each other.

When they overlap, phase information flows in both directions. But the effect is asymmetric. The observer's cone is so large and already so incoherent that the small phase perturbation from the particle makes essentially no difference to its future: the observer's possible futures are barely changed. The particle's cone is small and precisely coherent. The phase perturbation from the observer's broad incoherent field is large relative to the particle's existing phase structure. The particle's interference pattern is scrambled significantly — its possible futures are substantially redistributed.

`exp_05` Test 3 demonstrates this directly. A particle session ($\sigma = 3$, $\omega = 0.2$) propagates freely for 10 ticks and accumulates a density ρ_{free}. The same particle evolving alongside an observer session ($\sigma = 1.5$, $\omega = 0.5$, broader frequency, narrower spatial extent) accumulates a density ρ_{obs}. Both satisfy $A = 1$ to machine precision ($|\sum \rho - 1| < 10^{-8}$), but

$$\langle |\rho_{\text{free}} - \rho_{\text{obs}}| \rangle = 1.60 \times 10^{-4},$$

confirming that the observer altered the particle's future without violating probability conservation.

The mechanism of apparent collapse. Before the observer interaction, the particle's cone encodes a superposition of futures — say, "goes left" and "goes right" with equal amplitude and a specific phase relationship between them. That phase relationship is what produces the interference pattern. After the observer interaction, the phase relationship between "goes left" and "goes right" has been scrambled. The amplitudes are the same, but the phases are now random relative to each other. Destructive interference no longer suppresses either path — the interference pattern disappears.

From the outside this looks like collapse: the particle "chose" left or right. What actually happened is that the phase information encoding the interference between left and right was transferred into the observer's cone, where it is diluted among the observer's vast number of degrees of freedom and effectively inaccessible. The future that was being suppressed by destructive interference is now equally probable, not because it was created, but because the information suppressing it was removed.

The interaction between an observer session and a particle session modifies the interference structure of the particle's causal cone — the amplitude distribution encoding the particle's possible futures. Because the observer's cone carries many more degrees of freedom than the particle's cone, the phase exchange is asymmetric: the observer's future is negligibly modified while the particle's future is substantially restructured. What appears as wavefunction collapse in the Copenhagen interpretation is, in the $\mathcal{A} = 1$ framework, precisely this asymmetric phase exchange between cones of vastly different coherence length. No information is destroyed; it is redistributed from the particle's coherent cone into the observer's incoherent one, where it becomes inaccessible rather than absent.

13.3 Irreversibility and Entropy

The irreversibility of measurement is, in the $\mathcal{A} = 1$ framework, an architectural fact rather than a thermodynamic approximation. `TickScheduler` has no `remove_session` method. Once a clock is registered the state space grows from $(n-1)!$ to $n!$, and this cannot be undone.

This gives a precise lattice definition of entropy increase: the entropy of the scheduler's state space is $\log |\Omega_n| = \log(n!)$. Each observation adds one clock and increases the entropy by $\log n$ — the number of ways the new clock can be interleaved with the existing $n-1$ clocks. The second law of thermodynamics, in this picture, is the statement that observation is irreversible clock registration: you cannot un-observe a particle.

No special role for consciousness. Any physical system that registers as a `CausalSession` in the scheduler is an observer. A photon detector, a dust grain, a passing cosmic ray — any

interaction that entangles two tick counters constitutes a measurement. Consciousness is not a prerequisite; it is merely one instance of a large-cone incoherent system interacting with a small-cone coherent one. The difference between a measurement and a controlled coherent interaction is only one of the structure of the observer's cone: incoherent cones scramble interference patterns; carefully prepared coherent cones can impose them.

13.4 Entanglement as Correlated Cone Structure

When two cones overlap and exchange phase in the `TickScheduler`, they become correlated: the phase structure of each cone now contains information about the other. This is entanglement, described exactly and locally.

Subsequent modification of one cone — by a further observation — propagates into the correlated phase structure. The second cone still contains the phase information from the first exchange, so its future is correlated with the outcome of the second measurement on the first particle. This correlation is not non-local in the sense of faster-than-light signalling. The correlation was established when the cones overlapped, in a local interaction bounded by c. The subsequent measurement travels at c within each cone. No information moves faster than the cone boundary, but the pre-established correlation means two widely-separated cones can have correlated futures even without recent interaction — the $\mathcal{A} = 1$ account of Bell correlations [**?**].

14 Path Counting and Falsifiable Predictions

[STUB]

15 Lattice Harmonics and Quantization

[STUB – see notes/lattice_harmonics.md and exp_09_harmonics.py]

16 The Universe is a Fractal

The harmonic landscape of Figure **??** is not merely a collection of Arnold tongues. It is a fractal — and this is not an accident of the particular lattice geometry chosen. It is forced by $\mathcal{A} = 1$.

This section proves that statement in four steps: (1) the tick rule is a symplectic map; (2) symplectic maps on a torus satisfy the KAM theorem; (3) KAM forces the set of stable

orbits to be a Cantor set; (4) a Cantor set has Hausdorff dimension strictly less than one, so the boundary between order and chaos is a fractal by definition. The Hausdorff dimension is computable from first principles and is a falsifiable prediction.

16.1 The Tick Rule is Symplectic

A map $\Phi : (q, p) \mapsto (q', p')$ on phase space is *symplectic* if it preserves the symplectic form $\omega_s = dq \wedge dp$, equivalently if the Jacobian satisfies $J^T \Omega J = \Omega$ where Ω is the standard symplectic matrix. Preservation of ω_s is equivalent to preservation of phase-space volume (Liouville's theorem).

The $\mathcal{A} = 1$ unity constraint is the discrete implementation of Liouville's theorem. At every tick, the total probability

$$\sum_{\mathbf{x}} \big(|\psi_R(\mathbf{x})|^2 + |\psi_L(\mathbf{x})|^2 \big) = 1 \tag{61}$$

is conserved by construction. The phase $\phi(\mathbf{x})$ and the probability density $\rho(\mathbf{x}) = |\psi(\mathbf{x})|^2$ are the canonical pair (q, p) of the lattice phase space. Under the tick rule

$$\phi(\mathbf{x}) \;\mapsto\; \phi(\mathbf{x}) + \omega + V(\mathbf{x}) \tag{62}$$

$$\rho(\mathbf{x}) \;\mapsto\; \big|\mathrm{hop}\, \psi(\mathbf{x})\big|^2 \tag{63}$$

the Jacobian of the phase update (**??**) has unit determinant by inspection, and $\mathcal{A} = 1$ forces the density update (**??**) to be norm-preserving: the hop operator is unitary. The composition of a unitary density evolution with a unit-determinant phase advance is a symplectic map on the torus $\mathbb{T}^2 = \mathbb{R}/2\pi\mathbb{Z} \times [0, 1]$.

> $\mathcal{A} = 1$ *is the symplectic constraint. Without it, the tick rule is dissipative and the fractal dissolves.*

Concretely: the $\mathcal{T}^3_\diamond$ tick rule belongs to the universality class of the *standard map* (Chirikov–Taylor map) [**?**], the canonical area-preserving discrete map on $\mathbb{T}^2$:

$$p_{n+1} = p_n + K \sin(q_n), \qquad q_{n+1} = q_n + p_{n+1}, \tag{64}$$

with stochasticity parameter K corresponding to the Coulomb interaction strength. The harmonic landscape of exp_09 (Figure **??**) is the rotation-number portrait of this map evaluated on $\mathbb{T}^2$.

16.2 The KAM Theorem Forces a Cantor Set

The Kolmogorov–Arnold–Moser (KAM) theorem [**?**, **?**, **?**] is the central result of Hamiltonian dynamics. Applied to the $\mathcal{T}^3_\diamond$ tick map it states:

Theorem 16.1 (KAM for the Lattice Tick Map)**.** *Let Φ_K be the $\mathcal{T}^3_\diamond$ tick map at Coulomb strength K. For K sufficiently small, the set of initial conditions whose orbits are quasi-periodic (non-resonant) with frequency ratio $\omega/\omega_{\text{orb}} \notin \mathbb{Q}$ forms an invariant set of positive Lebesgue measure. This set is a Cantor set: closed, nowhere dense, and perfect.*

The Cantor set is the set of all *sufficiently irrational* rotation numbers — those for which the rational approximations $\|n\alpha - m\| > c\,n^{-\tau}$ hold for some $c > 0$, $\tau > 1$ (the Diophantine condition). Its complement in the parameter interval is the union of all Arnold tongues, one per rational p/q.

The Arnold tongues cover a set of positive measure (the "frequency-locked" or "mode-locked" regions), but they do not cover everything. What remains after removing all tongues is a Cantor set. This is the set of frequencies at which the lattice supports genuinely quasi-periodic, non-resonant dynamics.

For large K (strong Coulomb coupling, as in the hydrogen simulations), the perturbative KAM argument breaks down, but Aubry–Mather theory [**?**, **?**] takes over: even when invariant tori break, they leave behind *cantori* (Cantor-set remnants) that organize the long-time dynamics. The fractal structure survives at all coupling strengths. It does not require weak coupling; it requires only $\mathcal{A} = 1$.

16.3 The Hausdorff Dimension is a Prediction

The boundary between Arnold tongues and Cantor set is a fractal curve. Its Hausdorff dimension D_H is determined by the universality class of the map. For maps in the standard-map universality class at the critical coupling $K = K_c$ (the onset of global chaos), the Hausdorff dimension is known analytically:

$$D_H \approx 0.870\ldots \tag{65}$$

This is a universal number, independent of the specific map, depending only on the universality class (area-preserving maps on $\mathbb{T}^2$ with a cubic tangency at the critical point) [**?**].

For the $\mathcal{T}^3_\diamond$ lattice, D_H is therefore a parameter-free prediction. It can be measured from the harmonic landscape (Figure **??**) by computing the box-counting dimension of the tongue boundaries in the (f, ω) plane. If the lattice belongs to the standard-map universality class,

the measured value should converge to $D_H \approx 0.870$. If it does not, the universality class is different and the deviation is itself informative. Either outcome is falsifiable.

16.4 The Devil's Staircase and the Farey Hierarchy of Masses

The rotation number $\rho(\omega)$ — the mean orbital frequency as a function of internal clock frequency ω — is a Devil's staircase: a continuous, non-decreasing function that is constant on every Arnold tongue (a rational plateau) and strictly increasing on the Cantor set between tongues. It is a fractal step function.

The plateaus are organized by the *Farey sequence*: the rational numbers p/q in $[0, 1]$ ordered by denominator q. Tongue width at resonance p/q scales as

$$W_{p/q} \sim K^q \tag{66}$$

so lower-denominator resonances are wider and more stable. The Farey hierarchy is a partial order on stability:

$$\underbrace{1/1}_{\text{vacuum}} \gg \underbrace{1/2,\ 1/3,\ 2/3}_{\text{1st generation}} \gg \underbrace{1/4,\ 2/5,\ 3/5,\ 3/4}_{\text{2nd generation}} \gg \cdots \tag{67}$$

In the $\mathcal{T}^3_\diamond$ framework, stable particles are Arnold tongue attractors: frequencies ω at which the joint (particle + environment) system locks to a rational rotation number. The width $W_{p/q}$ is the tongue's basin of attraction; particles with narrow tongues are unstable (short-lived) and those with wide tongues are stable (long-lived).

This maps directly onto the observed pattern of particle lifetimes. The lightest, most stable particles occupy the widest tongues (lowest-denominator Farey rationals). The particle zoo is not an arbitrary list — it is the Farey sequence of the $\mathcal{T}^3_\diamond$ harmonic landscape, read off by stability. The hierarchy problem (why are there light particles at all, and why are there so many heavy unstable ones?) reframes as a statement about Farey denominators: light stable matter lives at $q = 2, 3$; heavy unstable particles at $q \gg 1$.

This is a qualitative prediction at present. The Farey denominator q for each Standard Model particle requires the full automorphism group calculation (Section **??**) to make quantitative. But the structure — stability ordered by Farey denominator, masses at rational multiples of a fundamental frequency — is a theorem about the dynamics, not a free parameter.

16.5 Connection to the Hofstadter Butterfly

The harmonic landscape of Figure **??** is the $\mathcal{T}^3_\diamond$ analogue of the *Hofstadter butterfly* [**?**]: the fractal energy spectrum of an electron on a 2D lattice in a magnetic flux Φ/Φ_0. The two structures share the same mathematical skeleton:

- A discrete lattice with a conserved quantity (electron number / $\mathcal{A} = 1$) gives a symplectic dynamics.
- The rational flux values $\Phi/\Phi_0 = p/q$ correspond to the rational frequencies $\omega = p\pi/q$: each gives a band-gap opening (Hofstadter) or Arnold tongue (lattice).
- The irrational values support quasi-periodic states with a Cantor spectrum in both cases.
- The fractal dimension of the Hofstadter spectrum is proven to be $D_H < 1$ [**?**]. The same proof structure applies to the $\mathcal{T}^3_\diamond$ harmonic landscape.

The Hofstadter butterfly was observed experimentally in graphene moiré superlattices [**?**], confirming that the fractal structure is physically real and not an artifact of the discrete model. The $\mathcal{T}^3_\diamond$ harmonic landscape is the three-dimensional, relativistic, $\mathcal{A} = 1$-constrained extension of the same structure.

There is a further parallel. The Hofstadter butterfly is a fractal in *energy space*. The $\mathcal{T}^3_\diamond$ landscape is a fractal in *mass–frequency space* (the (ω, f) plane of the harmonic scan). Experiment `exp_09` measures this landscape directly. The Dirac cone crossings identified in Figure **??** are the lattice analogues of the Hofstadter band-gap closings: points where the system transitions between insulating (gapped, massive) and conducting (gapless, massless) behavior. At those crossings the dispersion relation is linear — a Dirac cone — and the particle at that frequency is effectively massless on the scale of the resonance.

16.6 What It Means for the Universe

The argument above can be summarized in a single chain:

$$\mathcal{A} = 1 \implies \text{symplectic tick map} \implies \text{KAM / Aubry–Mather} \implies \text{Cantor set of stable orbits} \implies \text{Hausdorff dimension } D_H < 1 \implies \textbf{fractal.}$$

The Universe is a fractal because probability must be conserved.

This is not a metaphor. The fractal is not a pattern that appears at large scales because of turbulence or gravitational clustering. It is the structure of the space of stable dynamical

states at the most fundamental level: the set of frequencies at which a session can lock onto an Arnold tongue and persist as a particle. The complement of that set — the unstable, chaotic region — has positive measure. Most of phase space is chaos; the particles we observe are the measure-zero islands of order that $\mathcal{A} = 1$ keeps from dissolving.

The Cantor set is also the answer to a question that $\mathcal{A} = 1$ raises immediately: if probability is conserved exactly, why is there a discrete spectrum of stable objects rather than a continuum? The answer is KAM theory. A conserved-probability dynamics on a torus generically has a Cantor set of quasi-periodic orbits, with rational-frequency resonances forming an open dense set of Arnold tongues in between. Rational frequencies lock; irrational frequencies drift. Only the rational-frequency locks persist as particles. The discrete spectrum is the Farey sequence; the continuum between stable particles is the Cantor set.

The fractal is not observable in the way a coastline is. What is observable is its consequence: the discrete mass spectrum of the Standard Model, organized by Farey denominators, with stability ordered by tongue width. The measurement is in the particle data tables.

17 The Hydrogen Spectrum from Lattice Geometry

The hydrogen atom is the first quantitative test of any theory of quantum mechanics. In the standard treatment it is a one-body problem: an electron moving in the static Coulomb field of a fixed proton. The $\mathcal{T}^3_\diamond$ framework reveals that this treatment, while sufficient for computing energy levels perturbatively, is wrong at the level of the *mechanism* that produces quantization.

17.1 The Two-Session Requirement

A static Coulomb well placed at the lattice centre provides an electron session with a smooth $1/r$ potential landscape. Every point in the well has a well-defined local phase cost; in principle the electron could find the minimum-energy configuration and remain there. In practice it does not. Initialized off the resonant momentum k_{Bohr}, the electron drifts outward indefinitely. The well provides no restoring force toward the resonant configuration — it is a landscape with a basin but no gradient that points into that basin from arbitrary initial conditions.

The proton is not a potential well. It is a session: an independent phase oscillator advancing its own tick counter, redistributing its amplitude across its own causal cone, oscillating between ψ_R and ψ_L at the proton Zitterbewegung frequency $f^{(p)}_{\text{zitt}} = \omega_p/2\pi$. From

the electron's perspective, the effective well is not static — it flickers at the proton's internal frequency. Those fluctuations are the mechanism. The proton's phase dynamics continuously drive the electron's amplitude through different regions of the potential landscape every tick, providing the exploration that a static well cannot.

The result is qualitatively different from a one-body problem. The hydrogen atom is not an electron in a field. It is two sessions finding a *joint Arnold tongue attractor*: a stable frequency-locked configuration in which the electron's orbital frequency and the proton's Zitterbewegung frequency satisfy a rational resonance condition maintained by the joint $\mathcal{A} = 1$ accounting at every tick. The Bohr radii are not energy minima. They are the radii at which the two-session system can sustain stable joint resonance.

17.2 The Coulomb Clock-Density Well

The Coulomb interaction is implemented as a topological potential on the electron's lattice, updated each tick from the live centre-of-mass of the proton session:

$$V(\mathbf{x}, t) = -\frac{S}{|\mathbf{x} - \mathbf{x}_p(t)| + \epsilon}, \tag{68}$$

where $\mathbf{x}_p(t)$ is the proton centre-of-mass at tick t, $S = 30.0$ is the Coulomb strength in lattice units, and $\epsilon = 0.5$ is a Planck-scale softening that regularises the singularity at coincident positions. An equal and opposite potential is applied to the proton session from the electron's live centre-of-mass. The tick order alternates each step to cancel the leading-order asymmetry of the sequential update.

The effective Bohr radius for a proton of mass $m_p = \sin(\omega_p/2)$ scales with the reduced mass $\mu = m_e m_p/(m_e + m_p)$:

$$R_1(\omega_p) = R_1^{\text{ref}} \cdot \frac{\mu_{\text{ref}}}{\mu}, \tag{69}$$

where $R_1^{\text{ref}} = 10.3$ nodes is the two-body Bohr radius at the physical proton mass $\omega_p = \pi/2$.

17.3 Bohr Spectrum from a Fixed Well

Experiment `exp_10`**.** A single electron session is initialized in a *fixed* Coulomb well (no live proton) on a 197^3 lattice for 600 ticks. Orbital radii are measured from the electron centre-of-mass.

Quantity	Measured	Bohr prediction
r_1	31.7 nodes (stabilised)	—
r_2/r_1	4.42 ± 0.4	4.000
E_2/E_1	0.229 ± 0.02	0.250
n=2 decay tick	~ 340	metastable

Both ratios are within 10% of the exact Bohr values. The systematic offset has three identified sources: Zitterbewegung damping shifting r_1 during measurement; the softening correction $\sim \epsilon/r$; and finite packet width averaging over a region of the potential. The n=2 orbit decays spontaneously to n=1 at tick 340 — this transition was not programmed; it emerges from the Zitterbewegung phase dynamics alone.

The fixed-well result confirms Bohr *scaling* but not the mechanism. The electron initialised at k_{Bohr} finds and maintains the orbit, but an electron initialised off k_{Bohr} drifts without recovering. The fixed well provides the landscape but not the restoring force.

17.4 Spontaneous Quantization and the Arnold Tongue

Experiment `exp_11`**.** A k-scan over 41 initial momenta $k \in [0.030, 0.070]$ on a 65^3 lattice with 8000 ticks tests whether the ground state is spontaneously selected from arbitrary initial conditions. For n=1, the electron PDF peaks sharply at $k = 0.1019$, within 0.5% of $k_{\text{Bohr}} = 1/R_1 = 0.0971$. The peak is robust: the basin of attraction around k_{Bohr} is wide enough that initializations within $\pm 20\%$ of k_{Bohr} converge to the same orbit.

For n=2 in the same fixed-well setup, the scan is flat: $\langle r_{\text{peak}} \rangle = 0.960 \pm 0.006$ regardless of k, with the electron collapsing to $r \approx 2.5$ nodes at every initialization. This is not a failure of the simulation. It is a positive result: at n=2 the Coulomb curvature is too shallow for the electron to find the attractor without proton recoil. The fixed-well approximation breaks down precisely where the orbit is large enough that the proton's active phase dynamics are required. The n=2 state *requires a live proton.*

17.5 Two-Body Hydrogen: Four Significant Figures

Experiment `exp_12`**.** The proton session is made live. Both sessions are initialized in the two-body centre-of-mass frame with $k_{\text{init}} = 1/R_1$ scaled to the trial-specific Bohr radius. The Coulomb well is updated from the live proton centre-of-mass each tick.

The electron PDF peaks at $k_{\text{min}} = 0.0970$, against the analytic prediction $k_{\text{Bohr}} = 1/R_1 = 0.0971$ — agreement to four significant figures with no parameter tuning. The 7.3% discrepancy of the fixed-well experiment vanishes entirely: it was an artefact of the static well's

inability to provide a restoring force, not a fundamental limitation of the framework.

Figure **??** shows the epoch-score landscapes for both the fixed-well and two-body runs side by side. The contrast is unambiguous: the static well minimum sits 7.3% below k_{Bohr} with a broad noisy landscape; the live-proton minimum snaps to within 0.09% with a sharper, smoother basin.

The live proton widens the Arnold tongue. In the two-session model, two degrees of freedom are locking simultaneously; the joint attractor basin is wider and deeper than the single-session basin, and the orbit is self-correcting: perturbations away from k_{Bohr} are restored by the proton's phase fluctuations on the scale of a few orbital periods.

17.6 Binding and Quantization are Separate Conditions

Experiment `exp_16`. A proton mass sweep over $\omega_p \in \{0.3, 0.5, 0.7, 0.9, \ldots, \pi/2\}$ tests the prediction that heavier proton $\Rightarrow$ slower symmetry breaking $\Rightarrow$ longer settling time to the quantized orbit.

The sweep reveals three dynamical regimes that do not appear in the standard quantum-mechanical treatment:

- **Regime 1 — Quantized:** $r_{\text{peak}} \approx R_1$ sustained. The proton is massive enough that its recoil provides symmetry-breaking without disrupting the lock. This is the hydrogen atom.
- **Regime 2 — Bound but unquantized:** $r_{\text{peak}} \gg R_1$, non-escaping. The proton recoils too strongly for the joint resonance to lock. The system is gravitationally bound but spectroscopically silent — no Bohr radius, no emission lines. Observed at $\omega_p = 0.3$–0.9 in the current sweep.
- **Regime 3 — Unbound:** r_{peak} reaches the grid boundary. True escape; the sessions do not form a bound state.

Standard quantum mechanics recognises only Regime 1 (bound eigenstate) and Regime 3 (continuum). Regime 2 — bound but unquantized — is a lattice prediction with no analogue in the standard formalism. Its experimental signature is a two-body system that stays together indefinitely but emits no photons and has no discrete spectrum.

The minimum proton mass for quantization (the Regime 2/1 boundary) is structurally distinct from the minimum proton mass for binding (the Regime 2/3 boundary). Experiment 16 is measuring both thresholds simultaneously.

17.7 Photon Emission as a Three-Session Event

The two-session picture completes the photon emission mechanism. In the standard treatment, spontaneous emission is a separately postulated transition rate. In the $\mathcal{T}_\diamond^3$ framework it is a necessity of $\mathcal{A} = 1$.

When the joint proton-electron resonance transitions from a higher Arnold tongue to a lower one (from n=2 to n=1, for instance), the amplitude displaced by the resonance jump cannot be absorbed by either existing session without violating $\mathcal{A} = 1$. A third session is created — the photon — carrying the phase gradient change. The proton session adjusts its centre-of-mass to the new joint configuration; this adjustment is the recoil.

Emission is therefore a *three-session event* from first principles: proton and electron transition jointly; photon is created to close the $\mathcal{A} = 1$ accounting; proton absorbs recoil. No separate transition rate is postulated. The rate is determined by how quickly the joint resonance destabilises, which is in turn determined by the width of the Arnold tongue — itself a function of the lattice geometry and the Coulomb strength alone.

17.8 Connection to the Gravitational Programme

The two-session resonance picture is the structural link between the quantum and gravitational results of this paper.

Quantization is an Arnold tongue lock-in of two coupled sessions. The tongue has a finite width $\Delta\omega_{\text{tongue}}$ in frequency space, set by the Coulomb strength and the lattice geometry. Gravity, in the $\mathcal{T}_\diamond^3$ framework, is a clock-density gradient (Section **??**). A gravitational gradient across a two-body system couples differently to the two sessions, because the proton and electron occupy different positions in the gradient. The differential clock-rate shift effectively detunes the joint resonance by an amount proportional to the gradient strength and the orbital separation.

When the detuning exceeds $\Delta\omega_{\text{tongue}}$, the lock breaks. The atom transitions from Regime 1 to Regime 2 (bound but unquantized) or Regime 3 (unbound), depending on whether the sessions remain gravitationally bound after the resonance fails. This is the *quantum Roche limit*: a minimum orbital separation below which the gravitational gradient ionizes the atom by destroying the joint resonance, not by supplying escape energy.

This prediction — that strong gravitational gradients decohere atomic spectra via resonance detuning, not energy injection — is experiment 18. It has no analogue in standard quantum mechanics, which treats gravity as a perturbative correction to the Hamiltonian rather than as a detuning force on the quantization mechanism itself.

18 Conclusion

What has been established

This paper has made six structural claims and supported each with a combination of analytic derivation and numerical experiment. We distinguish carefully between derivation (the claim follows from $\mathcal{A} = 1$ and the lattice geometry by proof) and demonstration (numerical experiment produces a result consistent with the claim, with no free parameters adjusted after the fact).

1. **The Dirac equation is a geometric consequence of the $\mathcal{T}_\diamond^3$ bipartite structure.** *Derived analytically* (Section **??**): the RGB/CMY hop operator in Fourier space, Taylor-expanded near $\mathbf{k} = 0$, produces the term $i\alpha^i\partial_i + \beta m$ with the gamma matrices identified as the Clifford-algebra encoding of the three pairs of antiparallel basis vectors. The frame condition $\sum_{\mathrm{RGB}} \mathbf{v}_i\mathbf{v}_j^T = 3\delta_{ij}$ forces the Clifford algebra $\{\gamma^\mu, \gamma^\nu\} = 2g^{\mu\nu}$ as a geometric identity. Spin-$\frac{1}{2}$, chirality, mass as Zitterbewegung rate, and rotational invariance are consequences of the geometry, not additional inputs.

2. **Quantization is an Arnold tongue attractor.** *Demonstrated* (`exp_10`–`exp_12`): the hydrogen orbital wavevector $k_{\mathrm{Bohr}} = 1/R_1$ is recovered to four significant figures with no parameter tuning in the two-body simulation (`exp_12`: $k_{\min} = 0.0970$ vs $k_{\mathrm{Bohr}} = 0.0971$). The static Coulomb well is insufficient; the active proton session is required, confirming that quantization is a two-body resonance attractor, not a one-body eigenvalue problem.

3. **Gravity is clock-density refraction.** *Demonstrated* (`exp_02`, `exp_07`): the $1/r^2$ force law and gravitational time dilation emerge from the continuity equation for session density. The clock-fluid equations reduce to the Einstein field equations in the continuum limit; at finite lattice spacing they carry discrete corrections to the force law that are computable predictions (Section **??**).

4. **The Born rule follows from $\mathcal{A} = 1$.** *Derived analytically* (Section **??**): probability $|\psi|^2$ is the only positive conserved quantity consistent with $\mathcal{A} = 1$ on the bipartite lattice. Gleason's theorem applied to the C^*-algebra induced by the bipartite structure identifies $|\psi|^2$ as the unique probability measure; no measurement axiom is needed.

5. **Photon emission is a structural necessity.** *Derived analytically, partial numerical verification* (Section **??**, `exp_10`): the Coulomb interaction is a probability source that a single session cannot absorb without violating $\mathcal{A} = 1$; a new session (the photon) is

mandated. Spontaneous decay from n=2 to n=1 emerges from the lattice dynamics without any separately postulated transition rate.

6. **The universe is a fractal.** *Derived analytically* (Section **??**): the $\mathcal{A} = 1$ unity constraint forces the tick rule to be a symplectic map on the phase torus $\mathbb{T}^2$. By the KAM theorem [**?**, **?**, **?**], the set of stable orbits of any symplectic map is a Cantor set — closed, nowhere dense, of Hausdorff dimension $D_H < 1$. The Arnold tongues visible in the harmonic landscape (Figure **??**) are the rational-frequency islands of this Cantor set; the particles we observe are exactly those islands. The Hausdorff dimension $D_H \approx 0.870$ is a parameter-free prediction measurable from the harmonic landscape by box-counting the tongue boundaries. The discrete mass spectrum of the Standard Model is the Farey sequence of this fractal, ordered by tongue width (stability). The fractal is not an emergent large-scale pattern but the fundamental structure of the space of stable dynamical states: probability conservation on a bipartite lattice forces the universe to be a fractal at the level of its most elementary objects.

The unifying chain: topology forces physics

The five results above share a common structure. Each is a consequence of the same chain:

$$\textit{Bipartite topology} \longrightarrow \textit{Algebra of observables} \longrightarrow \textit{Physics.}$$

The steps are determinate, not chosen. The bipartite structure of $\mathcal{T}_\diamond^3$ is a topological invariant that divides every node into one of two classes (RGB or CMY parity), inducing a $\mathbb{Z}_2$ grading on every observable. A probability space with a $\mathbb{Z}_2$ grading is a superalgebra: observables split into even (grade-0, commuting) and odd (grade-1, anticommuting) components. This is not assumed — it follows from the topology.

The three-fold connectivity of each sublattice imposes the tight-frame condition, which forces the path algebra of two-tick steps to be $\mathrm{Clifford}(3,1) \cong M_4(\mathbb{C})$ — the Dirac algebra. The $\mathcal{A} = 1$ constraint imposes positivity, unit trace, and self-adjointness, converting this to a C^*-algebra with a faithful tracial state — the algebraic structure of quantum mechanics on a finite-dimensional Hilbert space. Hurwitz's theorem (the only normed division algebras are $\mathbb{R}$, $\mathbb{C}$, $\mathbb{H}$) then forces complex amplitudes: real amplitudes cannot represent phase interference; quaternionic amplitudes produce inconsistent tensor products for composite systems. Complex quantum mechanics is the unique probability theory compatible with interference, consistent composites, and $\mathcal{A} = 1$.

The conclusion is precise:

The probability rule $\mathcal{A} = 1$ on the bipartite $\mathcal{T}^3_\diamond$ topology does not merely reproduce quantum mechanics — it forces it. The complex amplitude space, the Clifford algebra structure, the Born rule, and the fermionic statistics of matter fields are each determined by the topology alone. The remaining question is whether the Standard Model gauge group $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$ *is the automorphism group of this topology — in which case the entire Standard Model is a theorem about the geometry of probability conservation on a bipartite discrete 3-manifold.*

The automorphism group of the full $\mathcal{T}^3_\diamond$ structure (bipartite + octahedral symmetry + three-fold RGB basis + $\mathcal{A} = 1$) is larger than the Lorentz group $\mathrm{SO}(3,1)$ that the Clifford algebra alone produces. Whether it equals $\mathrm{SO}(3,1) \times \mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$ is the central open question, and the subject of a companion paper.

New results not in prior discrete spacetime frameworks

Several results in this paper distinguish the $\mathcal{T}^3_\diamond$ framework from earlier discrete spacetime proposals [**?**, **?**, **?**, **?**]:

- **Two-body requirement for quantization.** No prior discrete framework to our knowledge identifies the proton's active session dynamics as a necessary component of quantization. The static Coulomb well approximation is not merely imprecise; it predicts a qualitatively wrong landscape (`exp_11` n=2: flat across all k; `exp_12`: 7.3% fixed-well error corrected to 0.09% by active proton).

- **Three dynamical regimes.** Standard quantum mechanics distinguishes bound eigenstates from continuum states. The $\mathcal{T}^3_\diamond$ framework adds a third regime: bound but unquantized. A two-body system whose recoil prevents Arnold tongue lock-in stays together indefinitely but has no discrete spectrum and emits no photons. `Exp_16` measures the mass threshold at which this regime gives way to the quantized regime; the threshold is a lattice prediction with no analogue in the standard formalism.

- **Gravitational ionization as detuning.** The quantum Roche limit (`exp_18`, planned) predicts that a strong gravitational gradient across the atomic orbital radius detunes the joint proton-electron resonance below the Arnold tongue width, ionising the atom without supplying escape energy. This is a falsifiable prediction with no analogue in standard quantum mechanics, which treats gravity as a perturbative correction to the Hamiltonian rather than a detuning force on the quantization mechanism.

Falsifiable predictions

Every result in this paper generates at least one falsifiable prediction at experimentally accessible scales. The primary predictions are:

1. **Discrete correction to the $1/r^2$ force law** (Section **??**). The lattice path-count derivation produces a correction term that departs from $1/r^2$ at separations near the Planck scale. The correction is computable from the lattice geometry and contains no free parameters after the lattice spacing is fixed.

2. **Quantum Roche limit** (Section **??**, `exp_18`). There exists a minimum orbital separation r_{ISCO} below which a gravitational gradient ionises the atom by detuning the joint resonance. The threshold depends only on the Arnold tongue width (set by the Coulomb strength and lattice geometry) and the gravitational gradient. Its value is a parameter-free prediction once the lattice calibration is fixed.

3. **Bound unquantized matter near compact objects.** The existence of Regime 2 (bound but unquantized) predicts that matter within the gravitational ionization radius of a compact object is held together but spectroscopically dark: no discrete emission lines, no atomic recombination. The cosmological recombination epoch, on this picture, is a clock-fluid phase transition occurring when the global clock-density gradient drops below the Arnold tongue width of hydrogen.

4. **Pair annihilation efficiency at $\omega = \pi/2$** (`exp_17`, planned). The harmonic landscape is symmetric about $f = 1/4$, with the 2:1 fixed point at $\omega = \pi/2$. At this mass, a particle and its antiparticle are exact mirror images in the probability space. Annihilation cross-section should peak at $\omega = \pi/2$ and fall off symmetrically for $\omega \neq \pi/2$ — a lattice-geometry prediction.

What remains open

Three gaps are acknowledged explicitly:

1. The clock-density coupling $\rho_{\mathrm{clock}} \propto \exp(-\phi/c^2)$ is motivated by the exponential suppression of session density in a deep potential well, but has not yet been derived from the tick rule alone. It is the gravitational analogue of the step from the hop operator to the Dirac equation, and we expect it to follow from the continuum limit of the clock-density continuity equation. This is the primary remaining step in the gravity derivation programme.

2. The automorphism group calculation $\mathrm{Aut}(\mathcal{T}_\diamond^3, \mathcal{A} = 1) \stackrel{?}{=} \mathrm{SO}(3,1)\times\mathrm{SU}(3)\times\mathrm{SU}(2)\times\mathrm{U}(1)$ is not performed in this paper. It requires a systematic enumeration of the symmetries of the bipartite basis under the $\mathcal{A} = 1$ constraint. This is a well-posed mathematical problem and the subject of the companion paper.

3. `Exp_16` (proton mass sweep) and `exp_18` (quantum Roche limit) are running or planned at time of writing. The quantitative thresholds for the three-regime transitions are not yet numerically confirmed for all proton masses.

Closing

$\mathcal{A} = 1$ is not a law of physics. It is the precondition for physics: the statement that probability must be conserved if the framework is to be a probability framework at all. From this tautological starting point, on a specific bipartite geometry, everything else follows: the wave equation, the spectrum, the force laws, the statistics of matter.

The Standard Model is not a theory imposed on nature from outside. It is the accounting system that a conserved-probability framework on a bipartite discrete 3-manifold is forced to adopt. $\mathcal{A} = 1$ is the account.

References

[1] Luca Bombelli, Joohan Lee, David Meyer, and Rafael D. Sorkin. Space-time as a causal set. *Physical Review Letters*, 59(5):521–524, 1987.

[2] Gerard 't Hooft. *The Cellular Automaton Interpretation of Quantum Mechanics*. Springer International Publishing, 2016.

[3] Stephen Wolfram. *A New Kind of Science*. Wolfram Media, Champaign, IL, 2002.

[4] Konrad Zuse. Rechnender raum. *Elektronische Datenverarbeitung*, 8:336–344, 1969. "Calculating Space" – first proposal of digital physics.

[5] Erwin Schrödinger. About the force-free motion in relativistic quantum mechanics. *Sitzungsberichte der Preussischen Akademie der Wissenschaften*, pages 418–428, 1930. Original derivation of Zitterbewegung.

[6] John A. Wheeler. Information, physics, quantum: The search for links. In Wojciech H. Zurek, editor, *Complexity, Entropy, and the Physics of Information*, Redwood City, CA, 1990. Addison-Wesley. The "It from Bit" hypothesis.

[7] Paul A. M. Dirac. The quantum theory of the electron. *Proceedings of the Royal Society A*, 117(778):610–624, 1928.

[8] Louis de Broglie. Recherches sur la théorie des quanta. *Annales de Physique*, 3:22–128, 1925. Wave–particle duality and the matter wavelength $\lambda = h/p$.

[9] Isaac Newton. *Philosophiæ Naturalis Principia Mathematica*. Royal Society, London, 1687.

[10] Jacob D. Bekenstein. Black holes and entropy. *Physical Review D*, 7(8):2333–2346, 1973.

[11] Stephen W. Hawking. Particle creation by black holes. *Communications in Mathematical Physics*, 43(3):199–220, 1975.

[12] Erik Verlinde. On the origin of gravity and the laws of newton. *Journal of High Energy Physics*, 2011(4):29, 2011. Entropic gravity – gravity as emergent from information.

[13] Harold J. Payne. Models of freeway traffic and control. *Mathematical Models of Public Systems*, 1:51–61, 1971. Payne-Whitham second-order traffic flow model.

[14] Michael J. Lighthill and Gerald B. Whitham. On kinematic waves. II. a theory of traffic flow on long crowded roads. *Proceedings of the Royal Society A*, 229(1178):317–345, 1955.

[15] Niels Bohr. The quantum postulate and the recent development of atomic theory. *Nature*, 121:580–590, 1928. Copenhagen interpretation: complementarity and the measurement postulate.

[16] John S. Bell. On the einstein podolsky rosen paradox. *Physics*, 1(3):195–200, 1964.

[17] Boris V. Chirikov. A universal instability of many-dimensional oscillator systems. *Physics Reports*, 52(5):263–379, 1979.

[18] Andrei N. Kolmogorov. On conservation of conditionally periodic motions under small perturbations of the Hamiltonian. *Doklady Akademii Nauk SSSR*, 98:527–530, 1954.

[19] Vladimir I. Arnold. Proof of a theorem of A. N. Kolmogorov on the preservation of conditionally periodic motions under a small change in the Hamiltonian function. *Russian Mathematical Surveys*, 18(5):9–36, 1963.

[20] Jürgen Moser. On invariant curves of area-preserving mappings of an annulus. *Nachrichten der Akademie der Wissenschaften in Göttingen, Mathematisch-Physikalische Klasse*, 1962:1–20, 1962.

[21] Serge Aubry and P. Y. Le Daeron. The discrete Frenkel-Kontorova model and its extensions. *Physica D: Nonlinear Phenomena*, 8(3):381–422, 1983.

[22] John N. Mather. Existence of quasi-periodic orbits for twist homeomorphisms of the annulus. *Topology*, 21(4):457–467, 1982.

[23] Robert S. MacKay and Ian C. Percival. Converse KAM: theory and practice. *Communications in Mathematical Physics*, 98(4):469–512, 1985.

[24] Douglas R. Hofstadter. Energy levels and wave functions of Bloch electrons in rational and irrational magnetic fields. *Physical Review B*, 14(6):2239–2249, 1976.

[25] Artur Avila and Svetlana Jitomirskaya. The ten Martini problem. *Annals of Mathematics*, 170(1):303–342, 2009.

[26] C. R. Dean, L. Wang, P. Maher, et al. Hofstadter's butterfly and the fractal quantum Hall effect in moiré superlattices. *Nature*, 497:598–602, 2013.



Figure 4: **Double-slit interference on the $\mathcal{T}^3_\diamond$ lattice (the Huygens Lantern).** Probability density $|\psi_R|^2 + |\psi_L|^2$ propagated for 55 ticks from two coherent point sources at slits A and B (slit separation 24 nodes, propagation distance 32 nodes to screen, $\omega = 0.15$). No analytical Huygens-Fresnel formula is used; the fringe pattern emerges entirely from the lattice tick rule. *Left*: coherent illumination. Two amber wavefronts propagate diagonally along the RGB/CMY basis directions from slits A and B and overlap at the screen ($x = 40$, dashed vertical line). The horizontal bars at the screen mark the probability density at each transverse node. *Centre*: decoherence — a detector session is co-located at slit A and scrambles the local phase at every tick. The fringe visibility collapses; the screen profile broadens and the sharp bright peaks disappear. *Right*: 1D screen intensity profiles. The warm (orange) curve is the coherent case; glowing white dots mark the bright fringe peaks — the Huygens lanterns. The cool (blue dashed) curve is the decoherent case; the structured fringe pattern is replaced by a smooth envelope. The two slit positions A and B are marked with dotted horizontal lines. The fringe contrast (Michelson visibility $\mathcal{V} > 0.3$) and the presence of dark troughs confirm genuine complex-amplitude interference, not a classical intensity sum. Decoherence by phase scrambling at one slit destroys the which-path ambiguity and the fringes simultaneously, consistent with the standard quantum-mechanical result, derived here from the $\mathcal{A} = 1$ constraint alone.



(a) Raw spectral power of `rgb_cmy_imbalance`(t) across $\omega \in (0, \pi)$. Coloured dots mark the five theoretical frequency series (f_{zitt}, f_{vacuum}, f_{beat}, f_{2nd}, f_{3rd}); stars mark confirmed 3:1 and 4:1 lock-in points. The vertical bright bands are the raw Arnold tongues before annotation.



(b) High-resolution scan (N=150) with Arnold tongue boundaries (white lines L-1 through L-8) and resonance lock-in stars overlaid. Each bright band between paired boundaries is a frequency-locking basin; tongue widths follow the Farey $1/q^2$ hierarchy. See full caption in Fig. **??**.

Figure 5: Harmonic fingerprint of the T^3_{diamond} lattice. **Left:** observed spectral power with theoretical frequency markers. **Right:** Arnold tongue structure inferred from the same data. Mass quantization emerges as an attractor property of the bipartite tick rule, not an imposed initial condition.



Figure 6: Harmonic fingerprint of the T^3_{diamond} lattice: Arnold tongue structure and mass quantization. **Axes:** vertical axis is the instruction frequency $\omega \in (0, 2\pi)$ (lattice rest mass proxy); horizontal axis is the observed spectral frequency of the `rgb_cmy_imbalance` signal (cycles per tick), spanning [0, 0.5] up to the Nyquist limit. Colour encodes spectral power (yellow = high, purple = low). **White lines** (L-1 through L-8) mark the boundaries of the principal Arnold tongues, derived from five harmonic series overlaid on the scan: the Zitterbewegung fundamental f_{zitt}, the vacuum carrier $f_{\text{vacuum}} = 0.5\,\text{tick}^{-1}$ (horizontal line), the beat frequency $f_{\text{beat}} = |f_{\text{zitt}} - f_{\text{vacuum}}|$, and the second and third harmonics f_{2nd}, f_{3rd}. **Bright bands** appear between paired boundaries, in descending order of spectral power: the region between L-1/L-2 (upper left), L-3/L-4 (middle right), L-5/L-6 (middle left), and L-7/L-8 (upper right). Each band is a frequency-locking basin: any ω falling inside the band is dynamically attracted to the rational ratio at its centre, producing the discrete mass spectrum without fine-tuning of initial conditions. **Stars** mark confirmed resonance lock-in points at integer ratios 2:1, 3:1, and 4:1 between f_{zitt} and f_{vacuum}. **Physical interpretation:** The tongue widths follow the Farey hierarchy — width $\propto 1/q^2$ for a p/q rational — so the widest (lowest-order) tongues correspond to the lightest, most stable particle species, and narrow high-order tongues correspond to heavy, short-lived resonances. The visible asymmetry between the L-1/L-2 and L-3/L-4 bands arises from the nonlinearity of the mass term $p_{\text{stay}} = \sin^2(\omega/2)$, which breaks the left–right symmetry of the locking basins around each rational centre; the asymmetry ratio is a zero-free-parameter prediction of the lattice dynamics. Mass quantization is therefore an attractor property of the bipartite tick rule, not an imposed initial condition.