

Asteroids Progress Report

July 2021

Jack Dinsmore

Asteroid Dynamics

- **Ingredients**

- Linear equation of motion

$$\ddot{\mathbf{r}} = -\frac{\mu}{r^2}$$

- Angular equation of motion

$$\dot{\omega} = I^{-1}(\tau - \omega \times (I\omega))$$

- Torque
 - Moment of inertia

Asteroid Dynamics

- **Parameters:** K_{lm} control both shape and density distribution
 - Chosen so that they relate easily to torque
 - Moment of inertia derived from K_{lm} as well.
- **Initial conditions**
 - Asteroid starts on hyperbolic trajectory
 - Initially spinning around lowest energy principal axis

Asteroid Dynamics

- The torque is given by
- MOI is diagonal:

$$\tau = -\frac{1}{2} \sum_{lm} (-1)^l \mathcal{J}_{lm} \sum_{l'm'} S_{l+l', m+m'}(D) \left[(i\hat{x} + \hat{y})(l' - m' + 1) \mathcal{K}_{l', m'-1}^* + (i\hat{x} - \hat{y})(l' + m' + 1) \mathcal{K}_{l', m'+1}^* + 2im'\hat{z} \mathcal{K}_{l'm'}^* \right].$$

$$I_{xx} = \frac{2}{3} \mathcal{K}_{20} - 2\mathcal{K}_{2,-2} - 2\mathcal{K}_{22} + \frac{2}{5} \mathcal{K}_{00}$$

$$I_{yy} = \frac{2}{3} \mathcal{K}_{20} + 2\mathcal{K}_{2,-2} + 2\mathcal{K}_{22} + \frac{2}{5} \mathcal{K}_{00}$$

$$I_{zz} = -\frac{4}{3} \mathcal{K}_{20} + \frac{2}{5} \mathcal{K}_{00}$$


Asteroid Dynamics

- To force a diagonal MOI, only the real part of K_{22} and K_{20} are nonzero.
- We may choose $K_{1m} = 0$
- K_{00} is the mass

Data

- We collect spin data separated in time by “cadences.”
 - E.g., 2 min to increase data, 1 hr to increase speed, etc.

Example Data



Fit Method


- With an MCMC, find the best K_{lm} s to maximize Gaussian likelihood:

$$\ln \mathcal{L} = - \sum_{\text{data}} \frac{(y_i - y_i^*)^2}{\sigma_i^2}$$



where error is simulated by rotating each spin vector by a small amount in a random direction.

Current status

- Fits are running, but I'm still generating some to better understand them.
- Degeneracy sometimes appears with certain choices of parameters



More fit examples



Density distribution model


- Suppose we can extract K_{lm} from spin data.
- Suppose we can extract a shape parameterization from light curve data
- What is the density distribution?

Density distribution model

- Idea:
 - 1) Separate the shape model into N different sections
 - 2) Give each section its own density
 - 3) Find densities that give the fitted K_{lm} values
- Use $N = \text{number of } K_{lm}\text{'s}$ so that the math is as simple as N volume calculations and a matrix inversion!

Density distribution model

- How do you choose the N sections?
 - 1) Find a sphere that encloses the asteroid
 - 2) Fill the sphere with N points
 - 3) Repopulate all points outside the asteroid model until they're all inside
 - 4) Form Voronoi cells based on the points



Density distribution model

- Pros:
 - Easy to execute
 - Possibly more information can be gained by making many selections of the N cells and combining
 - Non-degenerate
 - Not too complicated
- Cons
 - A shape model is needed
 - Cannot react to uncertainties in shape model
 - Does not capture very much density information
 - Cells may have strange shape

Conclusion

- Done:
 - Simulation of asteroid has been made
 - Fits of asteroid parameters have been started
 - Way to extract asteroid density from parameters has been proposed
- To do:
 - Fit higher order parameters
 - Implement asteroid density extraction
 - Fit to light curve data?