

Mosaics

Jack Hanke

July 21, 2025

1 THIS IS NEW

$$\begin{array}{cccc}
 \begin{array}{c} 0 - 0 - 0 \\ | \quad | \quad | \\ 0 - 0 - 0 \\ | \quad | \quad | \\ 0 - 0 - 0 \\ | \quad | \quad | \\ 0 - 0 - 0 \\ | \quad | \quad | \\ 0 - 0 - 0 \end{array} &
 \begin{array}{c} 0 - 0 - 0 \\ | \quad | \quad | \\ 0 - 0 - 0 \\ | \quad | \quad | \\ 0 - 0 - 0 \\ | \quad | \quad | \\ 0 - 1 - 0 \\ | \quad | \quad | \\ 0 - 0 - 0 \end{array} &
 \begin{array}{c} 0 - 0 - 0 \\ | \quad | \quad | \\ 0 - 0 - 0 \\ | \quad | \quad | \\ 0 - 1 - 0 \\ | \quad | \quad | \\ 0 - 0 - 0 \\ | \quad | \quad | \\ 0 - 0 - 0 \end{array} &
 \begin{array}{c} 0 - 0 - 0 \\ | \quad | \quad | \\ 0 - 0 - 0 \\ | \quad | \quad | \\ 0 - 1 - 0 \\ | \quad | \quad | \\ 0 - 1 - 0 \\ | \quad | \quad | \\ 0 - 0 - 0 \end{array}
 \end{array}$$

11^8

11^4

11^4

11^2

$$\begin{array}{cccc}
 \begin{array}{c} 0 - 0 - 0 \\ | \quad | \quad | \\ 0 - 1 - 0 \\ | \quad | \quad | \\ 0 - 0 - 0 \\ | \quad | \quad | \\ 0 - 0 - 0 \\ | \quad | \quad | \\ 0 - 0 - 0 \end{array} &
 \begin{array}{c} 0 - 0 - 0 \\ | \quad | \quad | \\ 0 - 1 - 0 \\ | \quad | \quad | \\ 0 - 0 - 0 \\ | \quad | \quad | \\ 0 - 1 - 0 \\ | \quad | \quad | \\ 0 - 0 - 0 \end{array} &
 \begin{array}{c} 0 - 0 - 0 \\ | \quad | \quad | \\ 0 - 1 - 0 \\ | \quad | \quad | \\ 0 - 1 - 0 \\ | \quad | \quad | \\ 0 - 0 - 0 \\ | \quad | \quad | \\ 0 - 0 - 0 \end{array} &
 \begin{array}{c} 0 - 0 - 0 \\ | \quad | \quad | \\ 0 - 1 - 0 \\ | \quad | \quad | \\ 0 - 1 - 0 \\ | \quad | \quad | \\ 0 - 1 - 0 \\ | \quad | \quad | \\ 0 - 0 - 0 \end{array}
 \end{array}$$

11^4

1

11^2

1

We first re-prove the enumeration of $k_{n,m}$ from Theorem ?? using our general method for enumerating mosaic systems, and show it immediately gives an analogous result for $p_{n,m}$. We state the enumeration of $p_{n,m}$ for $m, n \geq 2$ in Theorem 1.1. First we define $A(m) \in \mathbb{Z}^{2^{m-1} \times 2^{m-1}}$ for integers

$m \geq 2$, which we use to enumerate $p_{m,n}$. To be clear, throughout the paper we index the rows and columns of matrices starting at 0.

Definition 1.1. Let $A(2) = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$. We recursively define $A(k+1)$ given $A(k)$. Begin by writing $A(k) = \begin{bmatrix} A_{0,0} & A_{0,1} \\ A_{1,0} & A_{1,1} \end{bmatrix}$, where the block matrices $A_{i,j}$ are square block matrices of size $2^{k-2} \times 2^{k-2}$. We then define

$$A(k+1) = \begin{bmatrix} A_{0,0} & A_{0,0} & A_{0,1} & A_{0,1} \\ A_{0,0} & A_{0,0} & 0A_{0,1} & A_{0,1} \\ A_{1,0} & 0A_{1,0} & A_{1,1} & A_{1,1} \\ A_{1,0} & A_{1,0} & A_{1,1} & A_{1,1} \end{bmatrix},$$

Construct $A(m)$ by starting with $k = 2$ and recursing until $k = m$.

Theorem 1.1. *The number of polygon mosaics $p_{m,n}$ is the $(0,0)$ entry of $A(m)^n$.*

We then introduce messy knot mosaics—a variant of knot mosaics— in Section ?? and enumerate them.

2 Proof of Theorem 1.1

Proof. For a given mosaic, label the vertices of the tiles as follows. If the vertex is surrounded by an even number of polygons, label it 0. If the vertex is surrounded by an odd number of polygons, label it 1. To make a *vertex labeling* we also remove the dotted lines from all tiles in the mosaic. Using the mosaic from Figure ??, we show both the labeling of the vertices, and then the removal of the dotted lines.

These two cell labelings aren't associated with any of the cells T_1, \dots, T_7 , and so cannot correspond with a polygon mosaic. We seek a recursive solution to enumerate the number of vertex labelings for an $m \times n$ grid with these conditions: the labeling has a boundary of all 0's and does not contain the above cell labelings.

Our solution accomplishes this by building $m \times n$ vertex labelings for fixed m using vertex labelings of $m \times 1$ columns of cells. These columns have vertex labelings such that the top and bottom edge all labeled 0. One of these columns for $m = 3$ is below.

$$\begin{array}{c}
 0 \text{ --- } 0 \\
 | \quad | \\
 0 \text{ --- } 0 \\
 | \quad | \\
 0 \text{ --- } 1 \\
 | \quad | \\
 0 \text{ --- } 0
 \end{array}$$

It is useful to define an index for vertex labelings of these $m \times 1$ columns. To do this, consider reading a column of vertex labelings from bottom to top, ignoring the first and last 0's. If we interpret these sequences for the left and right column as binary numbers b_{left} and b_{right} , the index in base ten is the pair (b_{left}, b_{right}) . For example, for the above $m = 3$ column, we have sequences 00 for the left column and 10 for the right column, so the index is $(0, 2)$.

Next consider a column with index $(0, b_1)$ for some $b_1 \in [0, 2^{m-1} - 1]$. For reasons we will see later, let's call this the *starting column*. Now consider appending a column with index (b_1, b_2) for some $b_2 \in [0, 2^{m-1} - 1]$. For example, below is a diagram for appending our starting column $(0, 2)$ with column $(2, 3)$.

$$\begin{array}{ccc}
 \begin{array}{c} 0 \text{ --- } 0 \\ | \quad | \\ 0 \text{ --- } 0 \\ | \quad | \\ 0 \text{ --- } 1 \\ | \quad | \\ 0 \text{ --- } 0 \end{array} & + & \begin{array}{c} 0 \text{ --- } 0 \\ | \quad | \\ 0 \text{ --- } 1 \\ | \quad | \\ 1 \text{ --- } 1 \\ | \quad | \\ 0 \text{ --- } 0 \end{array} \rightarrow \begin{array}{c} 0 \text{ --- } 0 \text{ --- } 0 \\ | \quad | \quad | \\ 0 \text{ --- } 0 \text{ --- } 1 \\ | \quad | \quad | \\ 0 \text{ --- } 1 \text{ --- } 1 \\ | \quad | \quad | \\ 0 \text{ --- } 0 \text{ --- } 0 \end{array}
 \end{array}$$

Notice that in the above example we have created a vertex labeling for an $m \times 2$ grid of cells, in which the left-most vertex column labels are all 0 (left index of 0). Also note that we did not create either illegal cell labelings with column $(2, 3)$. However, we could append the column $(2, 1)$, which would create an illegal cell labeling.

Finally, if we further appended a column with label $(b_2, 0)$, we would have created a vertex labeling with a boundary of all 0's. For this reason, let's call a column with index $(b, 0)$ for $b \in [0, 2^{m-1} - 1]$ an *ending column*. This vertex labeling would correspond with 1 polygon mosaic if the middle column had index $(2, 3)$, but not if the middle column had index $(2, 1)$.

This motivates the creation of a matrix $A(m)$ to every column index (i, j) , where $A(m)_{i,j} = 1$ if the column labeling corresponds with a legal polygon mosaic, and 0 otherwise. For example, the $A(3)$ matrix corresponds with the following labeled columns.

$$\begin{array}{cccc}
\begin{array}{c} 0 \text{ --- } 0 \\ | \quad | \\ 0 \text{ --- } 0 \\ | \quad | \\ 0 \text{ --- } 0 \\ | \quad | \\ 0 \text{ --- } 0 \end{array} &
\begin{array}{c} 0 \text{ --- } 0 \\ | \quad | \\ 0 \text{ --- } 1 \\ | \quad | \\ 0 \text{ --- } 0 \\ | \quad | \\ 0 \text{ --- } 0 \end{array} &
\begin{array}{c} 0 \text{ --- } 0 \\ | \quad | \\ 0 \text{ --- } 0 \\ | \quad | \\ 0 \text{ --- } 1 \\ | \quad | \\ 0 \text{ --- } 0 \end{array} &
\begin{array}{c} 0 \text{ --- } 0 \\ | \quad | \\ 0 \text{ --- } 1 \\ | \quad | \\ 0 \text{ --- } 1 \\ | \quad | \\ 0 \text{ --- } 0 \end{array} \\
\\
\begin{array}{c} 0 \text{ --- } 0 \\ | \quad | \\ 1 \text{ --- } 0 \\ | \quad | \\ 0 \text{ --- } 0 \\ | \quad | \\ 0 \text{ --- } 0 \end{array} &
\begin{array}{c} 0 \text{ --- } 0 \\ | \quad | \\ 1 \text{ --- } 1 \\ | \quad | \\ 0 \text{ --- } 0 \\ | \quad | \\ 0 \text{ --- } 0 \end{array} &
\begin{array}{c} 0 \text{ --- } 0 \\ | \quad | \\ 1 \text{ --- } 0 \\ | \quad | \\ 0 \text{ --- } 1 \\ | \quad | \\ 0 \text{ --- } 0 \end{array} &
\begin{array}{c} 0 \text{ --- } 0 \\ | \quad | \\ 1 \text{ --- } 1 \\ | \quad | \\ 0 \text{ --- } 1 \\ | \quad | \\ 0 \text{ --- } 0 \end{array} \\
\\
\begin{array}{c} 0 \text{ --- } 0 \\ | \quad | \\ 0 \text{ --- } 0 \\ | \quad | \\ 1 \text{ --- } 0 \\ | \quad | \\ 0 \text{ --- } 0 \end{array} &
\begin{array}{c} 0 \text{ --- } 0 \\ | \quad | \\ 0 \text{ --- } 1 \\ | \quad | \\ 1 \text{ --- } 0 \\ | \quad | \\ 0 \text{ --- } 0 \end{array} &
\begin{array}{c} 0 \text{ --- } 0 \\ | \quad | \\ 0 \text{ --- } 0 \\ | \quad | \\ 1 \text{ --- } 1 \\ | \quad | \\ 0 \text{ --- } 0 \end{array} &
\begin{array}{c} 0 \text{ --- } 0 \\ | \quad | \\ 0 \text{ --- } 1 \\ | \quad | \\ 1 \text{ --- } 1 \\ | \quad | \\ 0 \text{ --- } 0 \end{array} \\
\\
\begin{array}{c} 0 \text{ --- } 0 \\ | \quad | \\ 1 \text{ --- } 0 \\ | \quad | \\ 1 \text{ --- } 0 \\ | \quad | \\ 0 \text{ --- } 0 \end{array} &
\begin{array}{c} 0 \text{ --- } 0 \\ | \quad | \\ 1 \text{ --- } 1 \\ | \quad | \\ 1 \text{ --- } 0 \\ | \quad | \\ 0 \text{ --- } 0 \end{array} &
\begin{array}{c} 0 \text{ --- } 0 \\ | \quad | \\ 1 \text{ --- } 0 \\ | \quad | \\ 1 \text{ --- } 1 \\ | \quad | \\ 0 \text{ --- } 0 \end{array} &
\begin{array}{c} 0 \text{ --- } 0 \\ | \quad | \\ 1 \text{ --- } 1 \\ | \quad | \\ 1 \text{ --- } 1 \\ | \quad | \\ 0 \text{ --- } 0 \end{array}
\end{array} \rightarrow \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 0 & 1 \\ 1 & 0 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{bmatrix}$$

$A(3)$

$A(m)$ has the property that the 0-th row represents all starting columns, and the 0-th column represents all ending columns. Even more importantly, notice that $A(m)_{i,j}^2$ represents the number of $m \times 2$ grids with left-most index i and right-most index j that correspond with a legal polygon mosaic. In general, $(A(m)^n)_{i,j}$ represents this quantity for an $m \times n$ grid of cells, and so if we know $A(m)$ for some m , then $(A(m)^n)_{0,0} = p_{m,n}$.

The final component of the proof is constructing $A(m)$ for any m . Begin by calculating $A(2)$ by identifying the number of legal polygon mosaics that correspond with each vertex coloring index $\{(0,0), (0,1), (1,0), (1,1)\}$, like so.

$$\begin{array}{cc}
 \begin{array}{c} 0 \text{ --- } 0 \\ | \quad | \\ 0 \text{ --- } 0 \\ | \quad | \\ 0 \text{ --- } 0 \end{array} & \begin{array}{c} 0 \text{ --- } 0 \\ | \quad | \\ 0 \text{ --- } 1 \\ | \quad | \\ 0 \text{ --- } 0 \end{array} \\
 \\
 \begin{array}{c} 0 \text{ --- } 0 \\ | \quad | \\ 1 \text{ --- } 0 \\ | \quad | \\ 0 \text{ --- } 0 \end{array} & \begin{array}{c} 0 \text{ --- } 0 \\ | \quad | \\ 1 \text{ --- } 1 \\ | \quad | \\ 0 \text{ --- } 0 \end{array}
 \end{array} \rightarrow \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$$

Next consider an arbitrary value $A(k)_{i,j}$ for any $k \geq 2$. This value is 1 if the $k \times 1$ column with index (i,j) can be part of a polygon mosaic, and 0 otherwise. We can determine that specific values of $A(k+1)$ are multiples of $A(k)_{i,j}$ by considering the following operation on an arbitrary column with index (i,j) . Copy the column four times and replace the top two 0's of each column with the bottom row of labels from one of the four cell labelings below.

$$\begin{array}{cccc}
 \begin{array}{c} 0 \text{ --- } 0 \\ | \quad | \\ 0 \text{ --- } 0 \end{array} & \begin{array}{c} 0 \text{ --- } 0 \\ | \quad | \\ 0 \text{ --- } 1 \end{array} & \begin{array}{c} 0 \text{ --- } 0 \\ | \quad | \\ 1 \text{ --- } 0 \end{array} & \begin{array}{c} 0 \text{ --- } 0 \\ | \quad | \\ 1 \text{ --- } 1 \end{array}
 \end{array}$$

For example, for the $m = 2$ column with index $(0,1)$, this operation looks like the following.

$$\begin{array}{c}
\begin{array}{ccc}
0 & \text{---} & 0 \\
| & & | \\
0 & \text{---} & 1 \\
| & & | \\
0 & \text{---} & 0
\end{array} \\
A(2)_{0,1}
\end{array}
\rightarrow
\begin{array}{cc}
\begin{array}{ccc}
0 & \text{---} & 0 \\
| & & | \\
0 & \text{---} & 0 \\
| & & | \\
0 & \text{---} & 1 \\
| & & | \\
0 & \text{---} & 0
\end{array}
&
\begin{array}{ccc}
0 & \text{---} & 0 \\
| & & | \\
0 & \text{---} & 1 \\
| & & | \\
0 & \text{---} & 1 \\
| & & | \\
0 & \text{---} & 0
\end{array}
\end{array}
\rightarrow
\begin{bmatrix}
A(3)_{0,2} & A(3)_{0,3} \\
A(3)_{1,2} & A(3)_{1,3}
\end{bmatrix}$$

This operation results in 4 new columns that are represented in $A(k+1)$. In our example, specifically we get the following.

$$\begin{bmatrix}
A(3)_{0,2} & A(3)_{0,3} \\
A(3)_{1,2} & A(3)_{1,3}
\end{bmatrix}
=
\begin{bmatrix}
1A(2)_{i,j} & 1A(2)_{i,j} \\
0A(2)_{i,j} & 1A(2)_{i,j}
\end{bmatrix},$$

Critically, this transformation *only* changes the identity of the top two tiles. This implies that the same value coefficients computed by comparing $A(2)$ and $A(3)$ can be used for any $m \times 1$ column, as long as both column indices (i, j) are congruent mod 2. Furthermore, if one writes $A(k)$ as the block matrix

$$A(k) = \begin{bmatrix} A_{0,0} & A_{0,1} \\ A_{1,0} & A_{1,1} \end{bmatrix},$$

where $A_{\hat{i},\hat{j}} \in \mathbb{R}^{2^{k-2} \times 2^{k-2}}$, then all column's represented in $A_{\hat{i},\hat{j}}$ have indices $(i, j) \equiv (\hat{i}, \hat{j}) \pmod{2}$. This allows us to write that in general, if $A(k) = \begin{bmatrix} A_{0,0} & A_{0,1} \\ A_{1,0} & A_{1,1} \end{bmatrix}$, then

$$\begin{bmatrix}
V_{0,0}A_{0,0} & V_{0,1}A_{0,0} & V_{0,2}A_{0,1} & V_{0,3}A_{0,1} \\
V_{1,0}A_{0,0} & V_{1,1}A_{0,0} & V_{1,2}A_{0,1} & V_{1,3}A_{0,1} \\
V_{2,0}A_{1,0} & V_{2,1}A_{1,0} & V_{2,2}A_{1,1} & V_{2,3}A_{1,1} \\
V_{3,0}A_{1,0} & V_{3,1}A_{1,0} & V_{3,2}A_{1,1} & V_{3,3}A_{1,1}
\end{bmatrix}. \tag{1}$$

where $V \in \mathbb{R}^{4 \times 4}$ can be found after directly computing $A(2)$ and $A(3)$, then solving the following equation.

$$\begin{bmatrix} A(3)_{0,0} & A(3)_{0,1} & A(3)_{0,2} & A(3)_{0,3} \\ A(3)_{1,0} & A(3)_{1,1} & A(3)_{1,2} & A(3)_{1,3} \\ A(3)_{2,0} & A(3)_{2,1} & A(3)_{2,2} & A(3)_{2,3} \\ A(3)_{3,0} & A(3)_{3,1} & A(3)_{3,2} & A(3)_{3,3} \end{bmatrix} = \begin{bmatrix} V_{0,0}A(2)_{0,0} & V_{0,1}A(2)_{0,0} & V_{0,2}A(2)_{0,1} & V_{0,3}A(2)_{0,1} \\ V_{1,0}A(2)_{0,0} & V_{1,1}A(2)_{0,0} & V_{1,2}A(2)_{0,1} & V_{1,3}A(2)_{0,1} \\ V_{2,0}A(2)_{1,0} & V_{2,1}A(2)_{1,0} & V_{2,2}A(2)_{1,1} & V_{2,3}A(2)_{1,1} \\ V_{3,0}A(2)_{1,0} & V_{3,1}A(2)_{1,0} & V_{3,2}A(2)_{1,1} & V_{3,3}A(2)_{1,1} \end{bmatrix}$$





This is solved by

$$V = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 0 & 1 \\ 1 & 0 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{bmatrix}$$

which completes the proof. □

The method detailed in Theorem 1.1 generalizes to other tile sets, which we tabularize in Section ?? without proof. Interestingly, the method not only generalizes to other tile sets, but can also be augmented to enumerate the more complicated “messy” polygon mosaics.

As demonstrated in the proof of Theorem ??, the enumeration of both polygon mosaics and mosaics that do not contain a polygon share the same structure, and only differ by the identity of the matrices $A(2), V$. We summarize these matrices for various collections of tiles below.

Tile Set	Polygon Mosaics	Messy Polygon Mosaics
	$\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 0 & 1 \\ 1 & 0 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{bmatrix}$	$\begin{bmatrix} 7^2 & 1 \\ -1 & 1 \end{bmatrix}, \begin{bmatrix} 7 & \frac{1}{7} & 7 \\ -\frac{1}{7} & 1 & 0 \\ 7 & 0 & 7 \\ 1 & -1 & 1 \end{bmatrix} \begin{matrix} 1 \\ 1 \\ 1 \\ 7 \end{matrix}$
	TODO	$\begin{bmatrix} 6^2 & 1 \\ -1 & 1 \end{bmatrix}, \begin{bmatrix} 6 & \frac{1}{6} & 6 \\ -\frac{1}{6} & 1 & 0 \\ 6 & 0 & 6 \\ 1 & -1 & 1 \end{bmatrix} \begin{matrix} 1 \\ 1 \\ 1 \\ 6 \end{matrix}$
	$\begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}, \text{TODO}$	TODO
	TODO	TODO

Theorem 1. Let the probability that an (m, n) -mosaic does not contain a SAP be denoted $p_{m,n} = \frac{t_{m,n}}{7^{nm}}$. Then the growth rate of the main diagonal $p_{n,n}$ has

$$\gamma = \lim_{n \rightarrow \infty} \frac{p_{n+1,n+1} p_{n-1,n-1}}{p_{n,n}^2} = ?$$

3 Introduction


Consider the following 6 unit squares with markings on them.



Call these squares *tiles*.


Definition 3.1. An (n, m) -mosaic is a rectangular grid made up of tiles.

Example 3.1. An example of a $(7, 5)$ -mosaic:




Clearly there are 6^{nm} possible mosaics. Which of these mosaics contain self-avoiding polygons?

Example 3.2. An example of a $(7, 5)$ -mosaic with a self-avoiding polygon:



Let $t_{n,m}$ be the number of mosaics that have at least one self avoiding polygon (SAP). From the fact that the smallest SAP is



we have that $t_{n,1} = t_{1,m} = 0$, and $t_{2,2} = 1$. What else can be said?

Theorem 2. *Setting $m = 2$ gives*

$$T_2(x) = \sum_{n \geq 2} t_{n,2} x^n = \frac{x^2}{(1 - 36x)(1 - 37x + 37x^2)}. \quad (2)$$

This can be solved for $n \geq 2$ to give

$$t_{n,2} = 6^{2n} - \frac{1}{\beta - \alpha} ((36\beta - 35)\beta^{-n+1} - (36\alpha - 35)\alpha^{-n+1}) \quad (3)$$

where $\alpha = \frac{1}{2} + \frac{1}{2}\sqrt{\frac{33}{37}}$ and $\beta = \frac{1}{2} - \frac{1}{2}\sqrt{\frac{33}{37}}$.

Proof. We prove that $t_{n,2}$ has

$$t_{n,2} = 36t_{n-1,2} + \sum_{i=2}^n (6^{2(n-i)} - t_{n-i,2}).$$

Split $t_{n,2}$ into S_n and S_n^c . S_n contains the mosaics that have just 1 SAP that contains the left-most two cells. This means S_n^c contains all mosaics that contain multiple SAPs and mosaics that contain only 1 SAP, but that does not contain the two left-most cells.

The subset S_n can be split further by the length of each SAP i .




Figure 1: Members of S_n of lengths $i = 2$ and $i = 3$

As S_n counts the number of mosaics that only contain 1 SAP, the blank space in Figure 3 must have no SAPs. The number of mosaics that have no SAPs is $6^{2(n-i)} - t_{n-i,2}$. As a SAP's width can range from 2 to n , we have $|S_n| = \sum_{i=2}^n (6^{2(n-i)} - t_{n-i,2})$.

Now consider S_n^c . The mosaics that belong to this set can be represented by the following diagram,




Figure 2: Representation of S_n^c

where the red dot in the left most cells indicate any marking. For this paper, we will refer to a cell that can have any marking as an *open*. We can conclude that $|S_n^c| = 6^2 t_{n-1,2}$. Combining S_n and S_n^c gives the recurrence relation. Standard techniques then give the generating function and formula. \square

Theorem 3. *Setting $m = 3$ gives*

$$T_3(x) = \sum_{n \geq 2} t_{n,3} x^n = \frac{(73 - 414x)x^2}{(1 - 216x)(1 - 228x + 2699x^2 - 7758x^3)} \quad (4)$$

Proof. For $t_{n,3}$ we directly compute the generating function $T_3(x) = \sum_{n \geq 2} t_{n,3} x^n$ using the following recurrence relation

$$t_{3,n} = 6^3 t_{n-1,3} + \sum_{i=2}^n (6^{3(n-i)} - t_{n-i,3}) f_i,$$

where f_i is the number of mosaics in an $i \times 3$ grid that contain just one SAP that has cells in the left-most column. We similarly split $t_{n,3}$ into S_n and S_n^c . Here let S_n be the set that contains the mosaics that have just 1 SAP that has cells in the left-most column. Therefore S_n^c contains all mosaics that contain multiple SAPs and mosaics that contain 1 SAP that does not have cells in the left-most column.

Similarly for the $n = 2$ case, S_n^c can be easily enumerated.





Figure 3: Representation of S_n^c

It is clear to see that $|S_n^c| = 6^3 t_{n-1,3}$, and so

$$\sum_{n \geq 2} |S_n^c| x^n = 6^3 x T(x).$$

To enumerate S_n , as in the $n = 2$ case, the SAP starts in the first column and ends at column i , after which there are no SAPs. This allows us to conclude that $|S_n| = \sum_{i=2}^n (6^{3(n-i)} - t_{n-i,3}) f_i$, where f_i is the number of ways the cells to the left of and including column i can contain 1 SAP that includes the left-most column. To find the identity of f_i , we study the 4 cases below.



It is easy to see that $a_2 = 36$ and $a_3 = 216$. As i increases, one can see that the enumeration of a_i is related to smaller values of a_i and b_i , more

specifically

$$a_i = 6a_{i-1} + 6^2b_{i-2}.$$

We find similar relations with the other 3 cases, namely

$$b_n = 6b_{n-1} + 6^2d_{n-2}$$

where $b_2 = 0$ and $b_3 = 6$

$$c_n = 6c_{n-1} + 6^2b_{n-2}$$

where $c_2 = 0$ and $c_3 = 0$

$$d_n = 6d_{n-1} + 6^2b_{n-2}$$

where $d_2 = 1$ and $d_3 = 6$.

Combining these 4 cases, and accounting for the appropriate symmetries, we arrive at

$$f_i = 2a_i + 4b_i + 2c_i + d_i.$$

Solving this series of recurrence relations using generating functions gives

$$F(x) = \sum_{i \geq 2} f_i x^i = \frac{73 - 414x}{1 - 12x + 43x^2}$$

This allows us to write

$$\sum_{n \geq 2} |S_n| x^n = \left(\frac{1}{1 - 6^3 x} - T(x) \right) F(x).$$

Combining these two generating functions and simplifying gives the result.

□

4 Full Solution

TODO writeup, credit Farstar31!

$$M(2) = \begin{bmatrix} 36 & 1 \\ -1 & 1 \end{bmatrix}$$

For $h \geq 2$, then if

$$M(h) = \begin{bmatrix} M_1 & M_2 \\ M_3 & M_4 \end{bmatrix}$$

then

$$M(h+1) = \begin{bmatrix} 6M_1 & 6M_2 & \frac{1}{6}M_1 & 1M_2 \\ 6M_3 & 6M_4 & 0M_3 & 1M_4 \\ -\frac{1}{6}M_1 & 0M_2 & \frac{1}{6}M_1 & 1M_2 \\ 1M_3 & 1M_4 & -1M_3 & 6M_4 \end{bmatrix}$$

where M_i is a sub-matrix (or possible a scalar) of the block matrix M .