

Jack Wright Project Proposal

Jack Wright

10/26/2020

Libraries

```
library(tidyverse)

## -- Attaching packages -----

## v ggplot2 3.3.2      v purrr   0.3.4
## v tibble  3.0.3      v dplyr  1.0.2
## v tidyr   1.1.2      v stringr 1.4.0
## v readr   1.4.0      v forcats 0.5.0

## -- Conflicts -----
## x dplyr::filter() masks stats::filter()
## x dplyr::lag()     masks stats::lag()
```

Data preparation

```
# load data from fivethirtyeight

url<-"https://projects.fivethirtyeight.com/nfl-api/nfl_elo_latest.csv"

dat<-read.csv(url)

# i will subset the columns that I need for the analysis

dat<-dat%>%
  select(elo1_pre,elo2_pre,score1,score2)

#add the elo difference column

dat<-dat%>%
  mutate(elo_diff=elo1_pre-elo2_pre)

## add score difference column (if the elo_diff is positive, then score1-score2, if negative score2-score1)

dat<-dat%>%
  mutate(score_diff=
```

```

        ifelse(elo_diff>=0,score1-score2,score2-score1)
    )

#now i can take the absolute value of the elo_difference

dat$elo_diff<-abs(dat$elo_diff)

```

Research Question

I want to look at the NFL elo data from 2019 and find out if the difference in elo (a rating system adopted from chess) and the difference in final score are correlated.

Cases and collection

there are 269 cases, each is a game played during the 2019 season in the NFL.

I am pulling this information in from “fivethirtyeight,” so I will be relying on their elo calculations.

link:

fivethirtyeight elo

Type of study

this is an observational study

Data source

This data is collected and the elo rank is created by fivethirtyeight.com, and is available at the above link

Response

The response variable will be a numerical difference in score at the end of the game.

Explanatory

The explanatory variable will be the numerical difference in elo before the game was played.

Relevant summary statistics

The summary statistics will be correlation coefficient and the R^2 of the data.

```


ggplot(data=dat, aes(x=elo_diff, y=score_diff))+
  geom_point()+
  stat_smooth(method = "lm", se=FALSE)

```

```
## 'geom_smooth()' using formula 'y ~ x'
```

```
## Warning: Removed 165 rows containing non-finite values (stat_smooth).
```

```
## Warning: Removed 165 rows containing missing values (geom_point).
```



```
m1<-lm(score_diff~ elo_diff, data=dat)
summary(m1)
```

```
##
## Call:
## lm(formula = score_diff ~ elo_diff, data = dat)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -33.533  -6.782   0.074   8.621  27.099
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)  2.27969    2.28982   0.996  0.322
## elo_diff     0.02895    0.01902   1.522  0.131
##
## Residual standard error: 12.52 on 102 degrees of freedom
## (165 observations deleted due to missingness)
## Multiple R-squared:  0.02221,    Adjusted R-squared:  0.01262
## F-statistic: 2.316 on 1 and 102 DF,  p-value: 0.1311
```