Lecture Notes: From Basic Algebra to Quantum Exchange, Matter Types, and Anyons

May 2026

Abstract

These notes build a conceptual and mathematical bridge from elementary algebraic ideas—commutativity, isotropy, idempotence, and projection—to the quantum-mechanical exchange rules for identical particles. The central result is that, in ordinary three-dimensional quantum mechanics, exchanging two identical particles twice gives the identity operation. This implies an exchange eigenvalue satisfying $p^2 = 1$, hence $p = +1$ for bosons or $p = -1$ for fermions. The notes then explain how two-dimensional topology enlarges the possibilities, giving anyons whose exchange can carry fractional or matrix-valued braid information.

Contents

1 Basic Algebraic Language **3**
- 1.1 Operations and Composition . . . 3
- 1.2 Abelian: Order Does Not Matter . . . 3
- 1.3 Isotropic: No Preferred Direction . . . 3
- 1.4 Idempotent: Once Is Enough . . . 4
- 1.5 Idempotent Versus Self-Inverse . . . 4

2 Projection Operators **4**
- 2.1 Eigenvalues of an Idempotent . . . 5

3 Quantum States and Identical Particles **5**

4 The Exchange Operator **6**

5 Deriving $p^2 = 1$ **6**

6 Bosons: Symmetric Exchange **6**
- 6.1 Physical Intuition . . . 7

7 Fermions: Antisymmetric Exchange **7**
- 7.1 The Pauli Exclusion Principle . . . 7

8 Projection onto Bosonic and Fermionic Subspaces **7**

9 Spin and Statistics **8**

10 Matter and Forces as Exchange Behaviors **8**
10.1 Fermions Create Separateness 8
10.2 Bosons Create Coherence 9
10.3 A Useful Intuition 9

11 Why Three Dimensions Give Only Bosons and Fermions **9**

12 The Two-Dimensional Loophole: Anyons **9**
12.1 Bosons and Fermions as Special Cases 9

13 Abelian Anyons **10**

14 Non-Abelian Anyons **10**

15 Experimental Status of Anyons **11**

16 Smoothly Interpolating Between "Stuff" and "Forces" **11**

17 Conceptual Summary **11**

18 Suggested Further Reading **12**

1 Basic Algebraic Language

Before discussing quantum particles, it is useful to define a few algebraic concepts that describe how operations behave when repeated, reordered, or viewed from different directions.

1.1 Operations and Composition

An *operation* is a rule that transforms an object. If A and B are operations, their composition AB means:

$$AB(x) = A(B(x)).$$

That is, B acts first and A acts second. Many important physical ideas reduce to questions about how operations compose.

1.2 Abelian: Order Does Not Matter

Definition 1.1 (Abelian or commutative)**.** *A set of operations is called* Abelian *if every pair of operations commutes:*

$$AB = BA.$$

Addition of ordinary numbers is Abelian:

$$2 + 3 = 3 + 2.$$

Multiplication of ordinary numbers is also Abelian:

$$4 \cdot 5 = 5 \cdot 4.$$

Matrix multiplication, however, is usually not Abelian. In general,

$$AB \neq BA.$$

This distinction becomes crucial for non-Abelian anyons, where the order of braids changes the final quantum state.

1.3 Isotropic: No Preferred Direction

Definition 1.2 (Isotropic)**.** *A system is* isotropic *if its properties are the same in every direction. Equivalently, it is invariant under rotations.*

For example, an ideal gas at rest is approximately isotropic: its pressure does not depend on whether one measures upward, sideways, or diagonally. Wood is anisotropic: it is easier to split along the grain than across it.

In mathematical terms, if R is a rotation and Q is some property of the system, isotropy means

$$R(Q) = Q.$$

In quantum mechanics, isotropy is related to rotational symmetry and angular momentum conservation.

1.4 Idempotent: Once Is Enough

Definition 1.3 (Idempotent)**.** *An operation P is* idempotent *if applying it twice is the same as applying it once:*

$$P^2 = P.$$

The intuitive meaning is: after the first application, the system is already in the stable output subspace, so repeating the operation changes nothing.

[A simple idempotent matrix] Consider

$$P = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}.$$

Then

$$P^2 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} = P.$$

This matrix projects a vector onto the x-axis:

$$P \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \\ 0 \end{pmatrix}.$$

Once the y component has been removed, projecting again does nothing.

[A non-diagonal integer idempotent matrix] Another 2×2 integer idempotent matrix is

$$P = \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix}.$$

Indeed,

$$P^2 = \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix} = P.$$

1.5 Idempotent Versus Self-Inverse

It is important not to confuse two different equations:

$$P^2 = P$$

and

$$P^2 = \mathbb{I}.$$

The first says that P is idempotent. Repeating the operation stabilizes.

The second says that P is self-inverse, or an *involution*. Repeating the operation undoes it.

A projection is idempotent. A swap is self-inverse.

2 Projection Operators

Projection operators are the most important idempotents in quantum mechanics.

Definition 2.1 (Projection operator)**.** *A linear operator P is a projection if*

$$P^2 = P.$$

If it is also Hermitian,

$$P^\dagger = P,$$

then it is an orthogonal projection.

Projection operators answer questions of the form: "Is the state inside this subspace?" Once a state has been projected into the subspace, applying the same projection again gives no further change.

2.1 Eigenvalues of an Idempotent

Suppose P is idempotent and v is an eigenvector:

$$Pv = \lambda v.$$

Then applying P again gives

$$P^2 v = P(\lambda v) = \lambda P v = \lambda^2 v.$$

But $P^2 = P$, so

$$P^2 v = Pv = \lambda v.$$

Therefore

$$\lambda^2 v = \lambda v.$$

For nonzero v,

$$\lambda^2 = \lambda,$$

so

$$\lambda(\lambda - 1) = 0.$$

Thus the only eigenvalues are

$$\lambda = 0 \quad \text{or} \quad \lambda = 1.$$

This is exactly what a projection does: it completely keeps some directions and completely erases others.

3 Quantum States and Identical Particles

In quantum mechanics, a physical state is represented by a vector in a Hilbert space, or by a wavefunction in a suitable representation. For a single particle with coordinate x, one may write

$$\psi(x).$$

For two particles, one writes

$$\psi(x_1, x_2).$$

If the two particles are distinguishable, then label 1 and label 2 may correspond to physically different objects. But if the particles are identical, such as two electrons, the labels are not physical names. They are bookkeeping devices.

The physical principle is:

> Exchanging two identical particles cannot change any observable fact about the system.

This is the starting point of quantum exchange statistics.

4 The Exchange Operator

For two identical particles, define the exchange operator X by

$$(X\psi)(x_1, x_2) = \psi(x_2, x_1).$$

It swaps the particle labels.

Now apply the exchange operation twice:

$$X^2\psi(x_1, x_2) = X\psi(x_2, x_1) = \psi(x_1, x_2).$$

Therefore

$$X^2 = \mathbb{I}.$$

This is not idempotence. Exchange is not a projection. It is an involution.

5 Deriving $p^2 = 1$

Suppose a two-particle state is an eigenstate of exchange:

$$X\psi = p\psi.$$

Apply X again:

$$X^2\psi = X(p\psi) = pX\psi = p^2\psi.$$

But $X^2 = \mathbb{I}$, so

$$X^2\psi = \psi.$$

Hence

$$p^2\psi = \psi.$$

For a nonzero state,

$$p^2 = 1.$$

Therefore

$$p = +1 \quad \text{or} \quad p = -1.$$

This is the elementary algebraic origin of the two familiar exchange classes in ordinary three-dimensional quantum mechanics.

6 Bosons: Symmetric Exchange

If

$$p = +1,$$

then

$$X\psi = +\psi.$$

Equivalently,

$$\psi(x_1, x_2) = \psi(x_2, x_1).$$

Such a state is symmetric under exchange.

Particles whose allowed many-particle states are symmetric are called *bosons*. Examples include photons, gluons, the Higgs boson, and composite particles with integer total spin such as helium-4 atoms.

6.1 Physical Intuition

Bosons can occupy the same quantum state. Their wavefunctions reinforce rather than exclude one another. This makes bosons naturally associated with coherence, fields, and collective behavior.

Examples of bosonic collective behavior include:

- laser light,
- Bose-Einstein condensates,
- superfluid helium-4,
- macroscopic electromagnetic fields.

This is one reason bosons often appear as force carriers in quantum field theory: many bosons can pile into the same field mode, producing smooth macroscopic fields.

7 Fermions: Antisymmetric Exchange

If

$$p = -1,$$

then

$$X\psi = -\psi.$$

Equivalently,

$$\psi(x_1, x_2) = -\psi(x_2, x_1).$$

Such a state is antisymmetric under exchange.

Particles whose allowed many-particle states are antisymmetric are called *fermions*. Examples include electrons, quarks, neutrinos, protons, and neutrons.

7.1 The Pauli Exclusion Principle

Suppose two identical fermions try to occupy the same state. In coordinate language, set $x_1 = x_2 = x$. Antisymmetry gives

$$\psi(x, x) = -\psi(x, x).$$

Therefore

$$2\psi(x, x) = 0,$$

so

$$\psi(x, x) = 0.$$

Thus two identical fermions cannot occupy exactly the same quantum state.

This is the Pauli exclusion principle. It is responsible for atomic shell structure, chemistry, the stability of ordinary matter, and degeneracy pressure in white dwarfs and neutron stars.

8 Projection onto Bosonic and Fermionic Subspaces

Although the exchange operator X is not idempotent, one can build idempotent projection operators from it.

Define the symmetric projection operator

$$S = \frac{1 + X}{2}.$$

Then

$$S^2 = \left(\frac{1+X}{2}\right)^2 = \frac{1+2X+X^2}{4}.$$

Using $X^2 = 1$,

$$S^2 = \frac{1+2X+1}{4} = \frac{2+2X}{4} = \frac{1+X}{2} = S.$$

So S is idempotent.

Similarly define the antisymmetric projection operator

$$A = \frac{1-X}{2}.$$

Then

$$A^2 = A.$$

These operators carve the two-particle Hilbert space into bosonic and fermionic sectors:

$$\mathcal{H} = \mathcal{H}_{\text{sym}} \oplus \mathcal{H}_{\text{antisym}}.$$

9 Spin and Statistics

Experimentally and theoretically, particles obey the spin-statistics connection:

$$\text{integer spin} \Rightarrow \text{boson},$$

$$\text{half-integer spin} \Rightarrow \text{fermion}.$$

The full spin-statistics theorem is not a theorem of elementary nonrelativistic quantum mechanics alone. Its standard derivation requires relativistic quantum field theory, especially locality, Lorentz symmetry, and positivity of energy.

A useful summary is:

Spin	Exchange statistics	Examples
$0, 1, 2, \ldots$	bosonic	Higgs, photon, graviton candidate
$\frac{1}{2}, \frac{3}{2}, \ldots$	fermionic	electron, quark, neutrino

10 Matter and Forces as Exchange Behaviors

The ordinary language of physics often says:

- fermions are matter,
- bosons are force carriers.

This is a useful first approximation but not a fundamental divide.

Both matter particles and force particles are excitations of quantum fields. What differs is how their states transform under exchange and rotation.

10.1 Fermions Create Separateness

Fermionic antisymmetry creates exclusion. This produces stable structure. Atoms do not collapse because electrons cannot all fall into the same state. Solid matter has rigidity because fermionic states resist compression.

In this sense, fermions create the individuality and impenetrability associated with "stuff."

10.2 Bosons Create Coherence

Bosons can accumulate in the same state. They naturally form coherent collective modes. A classical electromagnetic wave can be interpreted as a coherent state containing many photons.

In this sense, bosons create the smooth collective behavior associated with "forces."

10.3 A Useful Intuition

One compact way to phrase the distinction is:

Type	Exchange rule	Intuition
Boson	$\psi \mapsto +\psi$	overlap, coherence, collective field behavior
Fermion	$\psi \mapsto -\psi$	exclusion, individuality, stable matter

This is not an absolute metaphysical distinction. Rather, it is a structural distinction in how quantum states organize identity.

11 Why Three Dimensions Give Only Bosons and Fermions

The derivation $p^2 = 1$ relies on treating exchange as an element of the permutation group. In three spatial dimensions, exchanging two identical particles twice can be continuously untwisted back to doing nothing. The topology of configuration space enforces the familiar rule.

For n identical particles in three dimensions, the relevant exchange structure is the permutation group S_n. Its simplest one-dimensional representations are:

- the trivial representation, giving bosons;
- the sign representation, giving fermions.

The crucial point is that in three dimensions, particle paths can move around one another with enough room to untangle.

12 The Two-Dimensional Loophole: Anyons

In two spatial dimensions, particle exchange has richer topology. Paths can wind around one another, and those windings cannot always be smoothly undone. The relevant mathematical object is no longer merely the permutation group S_n but the braid group B_n.

For two-dimensional quasiparticles, an exchange can produce

$$\psi \mapsto e^{i\theta}\psi,$$

where θ need not be only 0 or π.

12.1 Bosons and Fermions as Special Cases

Bosons correspond to

$$\theta = 0,$$

so

$$e^{i\theta} = 1.$$

Fermions correspond to

$$\theta = \pi,$$

so

$$e^{i\theta} = -1.$$

Anyons allow intermediate phases:

$$0 < \theta < \pi,$$

or other fractional values modulo 2π.

Thus anyons interpolate between bosonic and fermionic exchange behavior.

13 Abelian Anyons

Definition 13.1 (Abelian anyon)**.** *An Abelian anyon is a two-dimensional quasiparticle whose exchange multiplies the wavefunction by a phase:*

$$\psi \mapsto e^{i\theta}\psi.$$

Because phases are ordinary complex numbers, their multiplication commutes:

$$e^{i\theta_1}e^{i\theta_2} = e^{i\theta_2}e^{i\theta_1}.$$

Therefore the exchange operations are Abelian.

The physical intuition is that Abelian anyons remember braiding only through an accumulated phase.

14 Non-Abelian Anyons

Definition 14.1 (Non-Abelian anyon)**.** *A non-Abelian anyon is a quasiparticle whose exchange acts not just by a phase but by a matrix on a degenerate quantum state space:*

$$\Psi \mapsto U\Psi.$$

Different exchanges may fail to commute:

$$U_1U_2 \neq U_2U_1.$$

Here, the order of braids matters. The system stores information in the collective topological state of many anyons. This is why non-Abelian anyons are important for topological quantum computing: quantum information can be encoded nonlocally and manipulated by braiding.

A concise intuition is:

Type	Exchange action	Memory of motion
Boson	$+1$	none
Fermion	-1	sign memory
Abelian anyon	$e^{i\theta}$	fractional phase memory
Non-Abelian anyon	matrix U	order-dependent braid memory

15 Experimental Status of Anyons

Anyons are not fundamental particles in the Standard Model. They are emergent quasiparticles in effectively two-dimensional many-body systems.

The primary experimental setting is the fractional quantum Hall effect, where electrons confined to two dimensions at low temperature and high magnetic field form topologically ordered fluids. The elementary excitations can carry fractional charge and fractional exchange statistics.

Important experimental milestones include:

- interferometric observations of Abelian anyonic braiding statistics in fractional quantum Hall devices around 2020;
- extensions to additional fractional quantum Hall states in subsequent interferometry experiments;
- active experimental and simulated studies of non-Abelian anyons, including Majorana zero-mode platforms, even-denominator fractional quantum Hall systems, and quantum-processor simulations of anyon braiding.

A cautious summary is:

Object	Status
Bosons and fermions	firmly established fundamental exchange classes in 3D
Abelian anyons	strong experimental evidence in 2D condensed matter systems
Non-Abelian anyons	promising evidence and simulations; direct, broadly settled braiding remains an active fr

16 Smoothly Interpolating Between "Stuff" and "Forces"

The language of "stuff" and "forces" is classical. Quantum theory replaces this with fields, excitations, exchange rules, and topology.

Fermions behave like stuff because antisymmetry creates exclusion and structure. Bosons behave like forces because symmetry permits coherence and macroscopic field-like behavior. Anyons reveal that this division is not absolute: exchange behavior can carry fractional or topological memory.

One intuitive spectrum is:

Exchange type	Identity behavior	Physical flavor
Bosonic	fluid identity	coherence, collective modes, fields
Anyonic	braided identity	topological memory, relational structure
Fermionic	rigid identity	exclusion, matter, stability

The deepest lesson is that particle type is not merely about what something is made of. It is about how the quantum state transforms under allowed motions in configuration space.

17 Conceptual Summary

The logical chain is:

1. Identical particle labels are not physically observable.
2. Exchanging two identical particles must preserve all observable physics.

3. In ordinary three-dimensional quantum mechanics, exchange twice gives the identity:

$$X^2 = \mathbb{I}.$$

4. If a state has exchange eigenvalue p, then

$$p^2 = 1.$$

5. Therefore

$$p = +1 \quad \text{or} \quad p = -1.$$

6. $p = +1$ gives symmetric bosonic states.
7. $p = -1$ gives antisymmetric fermionic states.
8. Symmetrizers and antisymmetrizers are idempotent projections:

$$S = \frac{1+X}{2}, \qquad A = \frac{1-X}{2}.$$

9. In two dimensions, the topology of exchange paths changes from permutations to braids.
10. This permits anyons, whose exchange can produce fractional phases or noncommuting matrix transformations.

18 Suggested Further Reading

References

[1] F. Wilczek, "Quantum Mechanics of Fractional-Spin Particles," *Physical Review Letters* 49, 957 (1982).

[2] R. B. Laughlin, "Anomalous Quantum Hall Effect: An Incompressible Quantum Fluid with Fractionally Charged Excitations," *Physical Review Letters* 50, 1395 (1983).

[3] J. Nakamura, S. Liang, G. C. Gardner, and M. J. Manfra, "Direct observation of anyonic braiding statistics," *Nature Physics* 16, 931–936 (2020).

[4] H. Bartolomei et al., "Fractional statistics in anyon collisions," *Science* 368, 173–177 (2020).

[5] C. Nayak, S. H. Simon, A. Stern, M. Freedman, and S. Das Sarma, "Non-Abelian anyons and topological quantum computation," *Reviews of Modern Physics* 80, 1083 (2008).

[6] S. Xu et al., "Non-Abelian braiding of Fibonacci anyons with a superconducting quantum processor," *Nature Physics* 20, 1469–1475 (2024).