

Supplemental Material for “Interband Pairing Signatures in the Superfluid Response of Magic-Angle Twisted Bilayer Graphene”

Jian Zhou¹

¹*JZ Institute of Science, Hong Kong, China**

(Dated: June 9, 2026)

TABLE I. Default parameters used for the mechanism scans.

quantity	value
twist angle	$\theta = 1.05^\circ$
Dirac velocity	$v_F = 2.135 \text{ eV \AA}$
AA tunneling	$w_0 = 87.2 \text{ meV}$
AB tunneling	$w_1 = 109.0 \text{ meV}$
reciprocal cutoff	$N_{\text{shell}} = 3$
retained bands	$n_{\text{keep}} = 6$
momentum grid	$nk = 7$ for dense maps, $nk = 9$ for key-point validation
gap scale	$\Delta_0 = 1 \text{ meV}$
velocity finite difference	$dk = 10^{-6} \text{ \AA}^{-1}$

I. SCOPE OF THE SUPPLEMENTAL MATERIAL

This Supplemental Material documents the executable numerical workflow behind the PRB manuscript. It is organized as a reproducibility and self-audit record: the Bistritzer-MacDonald (BM) model conventions [1], the orbital-projected pairing construction, the finite-band response formula, the figure and table provenance, and the historical PRB benchmark reconstruction are recorded separately.

The main text makes a conservative mechanism claim: an orbital-projected interband-pairing direction produces a reproducible, normalization-conditioned change in the normalized finite-band superfluid response. The present Supplemental Material does not promote the raw response units to final experimental stiffness values. This choice is motivated by the distinction between robust quantum-geometric mechanism diagnostics [2–5] and direct quantitative comparison with recent stiffness experiments [6, 7].

* jack@jzis.org

II. BM MODEL AND NUMERICAL PARAMETERS

The normal-state calculation uses the single-valley BM continuum Hamiltonian with orbital basis

$$(tA, tB, bA, bB), \quad (1)$$

where t, b label layers and A, B label sublattices. Unless explicitly stated otherwise, the production mechanism scans use the parameters in Table I.

The moire unit-cell area and reciprocal scale used in the unit audit are

$$A_M = 1.5606 \times 10^4 \text{ \AA}^2, \quad G_M = 0.054047 \text{ \AA}^{-1}. \quad (2)$$

The raw response conversion used in the audit is

$$D[\text{eV \AA}^2] = \frac{D_{\text{raw}}[\text{meV \AA}^2]}{1000}, \quad (3)$$

with all remaining spin, valley, or convention factors kept explicit rather than absorbed into a hidden prefactor.

III. ORBITAL-PROJECTED PAIRING CONSTRUCTION

The pairing direction is specified first in the layer-sublattice orbital basis,

$$\Delta_{\text{orb}}(\eta) = \Delta_0 \mathcal{N}_\eta (M_0 + \eta M_1), \quad (4)$$

with

$$M_0 = \tau_0 \sigma_0, \quad M_1 = \tau_x \sigma_x. \quad (5)$$

The two normalization controls used in the manuscript are

$$\text{fixed Frobenius:} \quad \|M_0 + \eta M_1\|_F = \|M_0\|_F, \quad (6)$$

$$\text{fixed } \Delta_0 : \quad \mathcal{N}_\eta = 1. \quad (7)$$

The band-basis pairing matrix is

$$\Delta_{\text{band}}(\mathbf{k}) = U^\dagger(\mathbf{k}) \Delta_{\text{orb}} U^*(-\mathbf{k}). \quad (8)$$

The working valley convention in the present diagnostic is the time-reversal sewn proxy

$$U_-(-\mathbf{k}) = U_+^*(\mathbf{k}). \quad (9)$$

This convention is useful for isolating an orbital-defined interband-pairing signature, but the readiness audit identifies a fully documented two-valley implementation as a submission-critical improvement.

The projected interband-pairing weight is

$$W_{\text{inter}} = \frac{\sum_{n \neq m} |\Delta_{nm}|^2}{\sum_{n,m} |\Delta_{nm}|^2}. \quad (10)$$

For the dense $nk = 7$, $n_{\text{keep}} = 6$ scan, $W_{\text{inter}} \simeq 0.108$ at $\eta = 1$.

IV. FINITE-BAND RESPONSE AND CHANNEL SPLIT

For a retained band subspace, the finite-band BdG Hamiltonian is

$$H_{\text{BdG}}(\mathbf{k}) = \begin{pmatrix} \varepsilon(\mathbf{k}) - \mu & \Delta_{\text{band}}(\mathbf{k}) \\ \Delta_{\text{band}}^\dagger(\mathbf{k}) & -[\varepsilon(\mathbf{k}) - \mu] \end{pmatrix}. \quad (11)$$

The zero-temperature static current-current response is evaluated as

$$\Lambda_{\alpha\beta} = \sum_{a \neq b} \frac{\langle a | J_\alpha | b \rangle \langle b | J_\beta | a \rangle (f_a - f_b)}{E_b - E_a}. \quad (12)$$

The band-basis velocity is split into diagonal and off-diagonal parts,

$$v_\alpha^{\text{band}} = v_\alpha^{\text{intra}} + v_\alpha^{\text{inter}}. \quad (13)$$

This produces conventional, geometric, and cross-channel response diagnostics. The primary normalized response map uses

$$\delta D_{\text{iso}}(\mu, \eta) = \frac{D_{\text{iso}}(\mu, \eta)}{D_{\text{iso}}(\mu, 0)} - 1, \quad D_{\text{iso}} = \frac{D_{xx} + D_{yy}}{2}. \quad (14)$$

V. FIGURE AND TABLE PROVENANCE

Table II summarizes the current figure and table provenance. The full machine-readable record is maintained in `notes/figure_table_provenance_2026-06-09.md`.

The dense response data are reproduced by

```
python3 scripts/run_mu_response_scan.py \
--nk 7 --n-shell 3 --n-keep 6 \
```

TABLE II. Manuscript artifact provenance. CSV values used in the manuscript tables are checked by `scripts/verify_manuscript_tables.py`.

artifact	data source	script	check
Frobenius heatmap	dense scan CSV	heatmap plotter	figure file exists
fixed- Δ_0 heatmap	dense scan CSV	heatmap plotter	figure file exists
dense response table	eta-summary CSV	eta summarizer	table verifier
$nk = 9$ validation table	key-point summary CSV	eta summarizer	table verifier
PRB audit table	reconstruction CSVs	audit scripts	table verifier

```
--mus -5 -4 -3 -2 -1 0 1 2 3 4 5 \
--etas 0 0.25 0.5 0.75 1 \
--output data/processed/mu_eta_response_scan_nk7_nkeep6.csv
```

and summarized by

```
python3 scripts/summarize_eta_effect.py \
data/processed/mu_eta_response_scan_nk7_nkeep6.csv \
--output data/processed/mu_eta_response_scan_nk7_nkeep6_eta1_summary.csv
```

The table verifier is

```
python3 scripts/verify_manuscript_tables.py
```

VI. KEY-POINT MOMENTUM-GRID CONVERGENCE

The dense maps in the main text use $nk = 7$, while selected key points were checked at $nk = 9$ and $nk = 11$. The $nk = 11$ scan uses

```
python3 scripts/run_mu_response_scan.py \
--nk 11 --n-shell 3 --n-keep 6 \
--mus -4 0 2 4 --etas 0 0.5 1 \
--output data/processed/mu_response_scan_nk11_nkeep6_keypoints.csv
```

and is summarized with `scripts/summarize_eta_effect.py`.

TABLE III. Key-point momentum-grid check for $D_{\text{iso}}(\eta = 1)/D_{\text{iso}}(\eta = 0) - 1$.

mode, μ (meV)	$nk = 9$	$nk = 11$	difference
fixed Δ_0 , -4	-0.0068	-0.0061	+0.0007
fixed Δ_0 , 0	+0.0103	+0.0080	-0.0023
fixed Δ_0 , 2	-0.0029	+0.0118	+0.0147
fixed Δ_0 , 4	-0.0017	-0.0003	+0.0014
fixed Frobenius, -4	+0.0604	+0.0617	+0.0012
fixed Frobenius, 0	+0.0863	+0.0796	-0.0067
fixed Frobenius, 2	+0.0643	+0.0927	+0.0285
fixed Frobenius, 4	+0.0605	+0.0608	+0.0002

Table III compares the normalized $\eta = 1$ response relative to $\eta = 0$. The fixed-Frobenius response remains positive at all four key chemical potentials. The fixed- Δ_0 response remains small and sign-sensitive, consistent with the main-text interpretation that the interband-pairing signal is normalization-conditioned.

The largest pointwise change in the fixed-Frobenius sector occurs at $\mu = 2$ meV, where the response increases by about 2.85 percentage points between $nk = 9$ and $nk = 11$. Therefore the current manuscript claims sign and order-of-magnitude stability of the normalized mechanism signal, not a strict continuum-limit extrapolation of the tabulated values.

VII. HISTORICAL PRB BENCHMARK RECONSTRUCTION

The production interband-pairing scan and the historical PRB table reconstruction are kept separate. The best simple unified reconstruction route for the old uniform- s benchmark is

$$D_{\text{conv}} = 2D_{\text{intra},\tau_z}, \quad D_{\text{geom}} = D_{\text{inter},\tau_z}. \quad (15)$$

For $n_{\text{keep}} = 6$, this gives $D_{\text{total}} = 126.66 \text{ eV } \text{\AA}^2$ versus the historical target $129.30 \text{ eV } \text{\AA}^2$, and $D_{\text{geom}} = 75.22 \text{ eV } \text{\AA}^2$ versus $75.30 \text{ eV } \text{\AA}^2$.

The old $n_{\text{keep}} = 2$ endpoint is more convention-sensitive. The closest audited reconstruction combines a Gamma-centered mesh, central-pair band selection, an all- τ_z current convention, a factor of two in the intraband term, and a full curvature-like conventional

correction. This gives

$$D_{\text{total}} = 66.18 \text{ eV } \text{\AA}^2 \quad \text{versus} \quad 67.50 \text{ eV } \text{\AA}^2. \quad (16)$$

Because this full-curvature correction does not work uniformly at $n_{\text{keep}} = 6$, the reconstruction is treated as a historical audit rather than a production convention.

VIII. CURRENT LIMITATIONS AND NEXT VALIDATION GATES

The current submission-critical gaps are:

1. a systematic continuum-grid extrapolation beyond the present $nk = 9$ and $nk = 11$ key-point checks;
2. a fully documented two-valley sewing convention or an explicitly limited diagnostic-model framing;
3. a final absolute stiffness convention before any direct experimental comparison;
4. a calibrated filling variable replacing the retained-band proxy ν_{proxy} .

These limitations do not invalidate the normalized-response mechanism claim, but they prevent the current draft from being treated as a final PRB submission without further convergence and convention work.

ACKNOWLEDGMENTS

The author acknowledges use of open-source scientific computing tools including NumPy and Matplotlib.

DECLARATIONS

Funding: Not applicable.

Conflict of interest: The author declares no conflict of interest.

Data availability: All data and code supporting this work are available from the corresponding author upon reasonable request.

Ethics approval: Not applicable.

- [1] R. Bistritzer and A. H. MacDonald, Moiré bands in twisted double-layer graphene, [Proceedings of the National Academy of Sciences](#) **108**, 12233 (2011).
- [2] S. Peotta and P. Törmä, Superfluidity in topologically nontrivial flat bands, [Nature Communications](#) **6**, 8944 (2015).
- [3] X. Hu, T. Hyart, D. I. Pikulin, and E. Rossi, Geometric and conventional contribution to the superfluid weight in twisted bilayer graphene, [Physical Review Letters](#) **123**, 237002 (2019).
- [4] A. Julku, T. J. Peltonen, L. Liang, T. T. Heikkilä, and P. Törmä, Superfluid weight and Berezinskii-Kosterlitz-Thouless transition temperature of twisted bilayer graphene, [Physical Review B](#) **101**, 060505 (2020).
- [5] F. Xie, Z. Song, B. Lian, and B. A. Bernevig, Topology-bounded superfluid weight in twisted bilayer graphene, [Physical Review Letters](#) **124**, 167002 (2020).
- [6] H. Tian, X. Gao, Y. Zhang, *et al.*, Evidence for Dirac flat band superconductivity enabled by quantum geometry, [Nature](#) **614**, 440 (2023).
- [7] M. Tanaka, J. I.-J. Wang, T. H. Dinh, *et al.*, Superfluid stiffness of magic-angle twisted bilayer graphene, [Nature](#) **638**, 99 (2025).