Un colorifies produce due tipi de colorants C. Cz utilitants tre preparati P<sub>1</sub>, P<sub>2</sub>, P<sub>3</sub>. Tol preparati dievano emen aquestati e subisciono un processo de volfmarore prime de enere cisti. Le quertite (in litri) de prepunti de hore de equistare per prolum un boro de colorante de coscerno des due tiji;

	Cı	62	
PI	1	1	
PZ	l	2	
P3	<u> </u>	1	

Opr mer le prontrè de pront che 1911ons eur equisión à le pronte

P1 P2 P3
750 1000 400

 $mox 7C_1 + 10C_2$   $CL + C_2 \le 750$ 0001 > 226 t C2 4 400 ×1 1×27,0

_ ,	b	C	CZ	S	52	83	
3	Q	-7	-10	0	0	6	
5,	710	J	1	1	0	0	
ړک	1000	1	2	0	1	0	
Sz	400	0	1	J	0	)	
						6 0 0	

	þ	C	CZ	SI	5 ي	83	
3	5250	Q	13	1	0	6	
4	tso	J	1	1	0	O	B3 -7 R3-
٢,	250	0			1	0	Bf-7 61 + 76
Sz	400	0	1	J	0	)	

	b	C	CZ	S <sub>1</sub>	Si	53	
Z	6000	0	0	4	<u> </u>	6	
C1	500	J	0	S	- (	O	
Cz	250	0			1		
53	150	0	0		-1	)	

B4-7B4-B3
B2-7B3-R2
B1-7B1+3B3

C1 = 800 C2 = 280 Sz = 180 S1,52=0 E= 6000