

immunosuppression. Nevertheless, as with other vaccinations, the benefit of immune protection, most probably, outweighs the risk of relapse.

1. Kervella D, Jacquemont A, Chapelet-Debout A, et al. Minimal change disease relapse following SARS-CoV2 mRNA vaccine. *Kidney Int.* 2021;100:457–458.
2. Banerjee S, Dissanayake PV, Abeyagunawardena AS. Vaccinations in children on immunosuppressive medications for renal disease. *Pediatr Nephrol.* 2016;31:1437–1448.

Nora Schwotzer¹, Sébastien Kissling¹ and Fadi Fakhouri¹

¹Service of Nephrology and Hypertension, Department of Medicine, Lausanne University Hospital, Lausanne, Switzerland

Correspondence: Nora Schwotzer, Service of Nephrology and Hypertension, Department of Medicine, Lausanne University Hospital, Lausanne, Switzerland. E-mail: nora.schwotzer@chuv.ch

Kidney International (2021) **100**, 458–459; <https://doi.org/10.1016/j.kint.2021.05.006>

Copyright © 2021, International Society of Nephrology. Published by Elsevier Inc. All rights reserved.

Relapse of minimal change disease following the AstraZeneca COVID-19 vaccine



To the editor: Anecdotal reports linking minimal change disease (MCD) to vaccinations possibly due to immune dysregulation,¹ including influenza vaccine,² pneumococcal,³ meningococcal C vaccines,⁴ and BNT162b2 coronavirus disease 2019 (COVID-19) vaccine (Pfizer-BioNTech)^{5,6} have been published. We report 2 cases of biopsy-proven MCD relapsing within 2 days of receiving an AstraZeneca COVID-19 vaccine.

A 30-year-old man had received 1 g of rituximab in August 2020, having experienced annual relapses on tacrolimus. His prednisolone had been weaned to 1 mg/day by January and discontinued altogether by February 2021. Two days after his COVID-19 vaccine, he developed a headache and frothy urine. Urine protein-to-creatinine ratio 1 week later was 213 mg/mmol; albumin was preserved at 47 g/l; creatinine was stable at 82 µmol/l. At that time, lymphocyte subsets showed complete B-cell depletion; CD19 was 0.00. He did not seek medical attention until 2 months after receiving the vaccine when his urine protein-to-creatinine ratio was 142 mg/mmol. Repeat lymphocyte subsets then revealed B-cell return; CD19 was 0.06. Complete remission was achieved with 10 days of starting prednisolone 20 mg daily.

A 40-year-old woman was maintained on prednisolone 5 mg daily and tacrolimus (Adoport); trough level was 4.6 µg/l before vaccination. One day after receiving her first COVID-19 vaccine, she developed a headache, frothy urine, and ankle swelling. After 1 week, her general practitioner recorded 3+ dipstick proteinuria. Unfortunately, no laboratory samples were sent. Prednisolone was increased to 30 mg daily, and

complete remission was achieved within 2 weeks. Creatinine was unchanged at 105 µmol/l.

The association with various vaccines has been described, occurring between 4 days to several weeks later.^{1,5,6,7} The timing of COVID-19 vaccination and the very early development of relapse of MCD in our cases raises questions as to the mechanisms involved. At 2 days after vaccination, one would assume the vaccine triggered a more generalized cytokine-mediated response.⁷ Others have postulated that symptoms after 4 days represent a rapid T cell-mediated response to viral mRNA.^{2,5,6}

We administered the second dose of a different COVID vaccine, and neither patient suffered an adverse effect. However, both patients were taking 15 mg prednisolone daily at the time. This may prove a useful strategy in similar cases.

We await further reports to evaluate the true incidence.

1. Clajus C, Spiegel J, Brocker V, et al. Minimal change nephrotic syndrome in an 82 year old patient following a tetanus-diphtheria-poliomyelitis-vaccination. *BMC Nephrol.* 2009;10:21.
2. Kielstein JT, Termuhlen L, Sohn J. Minimal change nephrotic syndrome in a 65-year-old patient following influenza vaccination. *Clin Nephrol.* 2000;54:246–248.
3. Kikuchi Y, Imakiire T, Hyodo T, et al. Minimal change nephrotic syndrome, lymphadenopathy and hyperimmunoglobulinemia after immunization with a pneumococcal vaccine. *Clin Nephrol.* 2002;58:68–72.
4. Abeyagunawardena A, Goldblatt D, Andrews N, Trompeter R. Risk of relapse after meningococcal C conjugate vaccine in nephrotic syndrome. *Lancet.* 2003;362:449–450.
5. Lebedev L, Sapojnikov M, Wechsler A, et al. Minimal change disease following the Pfizer-BioNTech COVID-19 vaccine. *Am J Kidney Dis.* 2021;78:142–145.
6. D'Agati VD, Kudose S, Bombardieri AS, et al. Minimal change disease and acute kidney injury following the Pfizer-BioNTech COVID-19 vaccine. *Kidney Int.* 2021;100:461–463.
7. Sette A, Crotty S. Adaptive immunity to SARS-CoV-2 and COVID-19. *Cell.* 2021;184:861–880.

Clare Morlidge¹, Sally El-Kateb², Praveen Jeevaratnam² and Barbara Thompson²

¹Pharmacy Department, Lister Hospital, Stevenage, Hertfordshire, UK; and

²Renal Department, Lister Hospital, Stevenage, Hertfordshire, UK

Correspondence: Clare Morlidge, Pharmacy Department, East and North Hertfordshire NHS Trust, Coreys Mill Lane, Stevenage, Hertfordshire, SG1 4AB, UK. E-mail: claremorlidge@nhs.net

Kidney International (2021) **100**, 459; <https://doi.org/10.1016/j.kint.2021.06.005>

Copyright © 2021, International Society of Nephrology. Published by Elsevier Inc. All rights reserved.

Post-vaccinal minimal change disease



To the editor: Previous reports have described the onset of minimal change disease after the administration of certain vaccines.¹

Recently a 61-year-old woman was admitted to our hospital 8 days after her first coronavirus disease 2019 (COVID-19) vaccination (BioNTech/Pfizer SARS-CoV-2 COM-IRNATY) because of edema and weight gain (6 kg). Medical