Math 475 – Project 1 Bobcat Population

Jackson Brienen

October 8 2025

1 Introduction

In this paper we discuss how we can model bobcat populations under a variety of environmental conditions and external manipulations (hunting and introduction). While there are a variety of modeling techniques that can be used for populations we focus on using a basic exponential model. With this come a variety of assumptions:

- 1. There is no carrying capacity/ maximum population.
- 2. The growth rate of the population is always constant, i.e. the growth rate in the first year will be the same as the hundredth year, and thousandth year, and so on.
- 3. We assume, in most cases, that fractional populations are valid. While one half of a bobcat does not usually make sense, for the sake simplicity we will allow it to be the case unless we explicitly say we use integer based populations.
- 4. All births and deaths happen at the end of a year/ start of a new year.

This paper will be divided into three main topics. First, the analysis of the basic model under three different growth rates in both the long and short term. Second, we will how the model changes when hunting is introduced and devising a strategy to meet a stable population with hunting. Third and last, we will analyze how the model changes when external introduction is used along with devising a strategy to meet a stable population with introduction.

2 Basic Growth

This section discusses the basic growth model which we use throughout this paper. We start with a very basic recursive function P(t) which takes our previous population P(t-1) and adds the annual change to the population rP(t-1), where r is some growth rate. This simplifies to Equation 1.

$$P(t) = (r+1)P(t-1)$$
 (1)

Previously we mentioned we were using an exponential model, yet this recursive formula does not look exponential. While it might not be immediately obvious, we can solve the recurrence for its closed form expression.

$$P(t) = (r+1)P(t-1) P(t-1) = (r+1)P(t-2)$$

$$P(t) = (r+1)((r+1)P(t-2)) P(t-2) = (r+1)P(t-3)$$

$$P(t) = (r+1)((r+1)((r+1)P(t-3))) P(t-3) = (r+1)P(t-4)$$

$$\vdots$$

$$P(t) = (r+1)^t P(0)$$

In this closed form we can see clearly that this is in fact an exponential model. We can see this exponential nature by graphing our model. There are three growth rates based on environmental conditions, 1.676% in the best case, 0.549% in moderate cases, and -4.5% in the worst case, we will look at these when the starting population is 100.




Figure 1: Bobcat populations in the short (30 year) and long (70 year) term. Using Equation 1 with P(0) = 100, and three different growth rates, r, being 1.676%, 0.549%, and -4.5%.

While in Figure 1a it is difficult to see our exponential behavior in all cases except r = -0.045, it becomes obvious in Figure 1b. In the case where r = -0.045, we know our function takes the form $P(t) = (1 - 0.045)^t P(0)$, which means 0.955 will be multiplied together t times, to get our population for any given t. Since 0.955 < 1, our population will be decreasing, as we can see in both figures. We also know that, while population will continue to get smaller as $t \to \infty$, it will never reach zero. This behavior is seen in Figure 1b as we see the line become asymptotic to zero.

For r = 0.01676 (in blue) and r = 0.00549 (in orange) we see similar behavior, since in both cases r + 1 > 1 we know the population will always be increasing. We know that there is nothing in our model to stop this growth, so as $t \to \infty$ $P(t) \to \infty$, but at different rates depending on r. This is highlighted in the figures as both lines go upwards in similar manners, but at different rates.

This model appears to only have one point of stability, when P(t) = 0, and we only approach this stable point with negative r values. We will discuss different methods of identifying stable, or fixed, points later in this paper.

¹github.com/JacksonBrienen – Graphs github.com/JacksonBrienen – Code for Graphs

- 3 Hunting
- 4 Population Introduction