

18.02: Multivariable Calculus

Problem Set 2

As taught by PROF. JOHN BUSH, MIT

JACKSON KEARL

October 13, 2015

Contents

I	2
1 Lecture 12	3
1.1 Section 13.9	3
1.1.1 Problem 2	3
1.1.2 Problem 8	4
1.1.3 Problem 38	5
1.1.4 Problem 52	6
1.2 Section 13.10	8
1.2.1 Problem 4	8
1.2.2 Problem 10	8
2 Lecture 13	9
2.1 Section 13.7	9
2.1.1 Problem 2	9
2.1.2 Problem 4	9
2.1.3 Problem 6	10
2.1.4 Problem 16	10
2.1.5 Problem 20	11
2.1.6 Problem 40	12
3 Lecture 14	13
3.1 Section 14.1	13
3.1.1 Problem 14	13
3.1.2 Problem 16	13
3.1.3 Problem 22	14
3.1.4 Problem 26	14
3.2 14.2	14
3.2.1 Problem 2	14
3.2.2 Problem 6	15
3.2.3 Problem 12	15
3.2.4 Problem 18	15
3.2.5 Problem 26	16
4 Lecture 15	17
4.1 Section 14.4	17
4.1.1 Problem 2	17
4.1.2 Problem 6	17
4.1.3 Problem 16	18

4.1.4	Problem 22	18
4.1.5	Problem 24	18
4.1.6	Problem 30	19
4.2	Section 14.5	20
4.2.1	Problem 2	20
4.2.2	Problem 6	20
4.2.3	Problem 14	20
4.2.4	Problem 24	21

Introduction

“Begin at the beginning,” the King said, gravely, “and go on till you come to an end; then stop.”

– Lewis Carroll, *Alice in Wonderland*

Course Content

This course covers aspects of calculus involving:

Calculus of several variables. Vector algebra in 3-space, determinants, matrices. Vector-valued functions of one variable, space motion. Scalar functions of several variables: partial differentiation, gradient, optimization techniques. Double integrals and line integrals in the plane; exact differentials and conservative fields; Green’s theorem and applications, triple integrals, line and surface integrals in space, Divergence theorem, Stokes’ theorem; applications.¹

Structure of Homework

Problems will be arranged as they are on the assignment page. Problems broken into parts one and two, as per assignment page. Within part one, problems divided first by Lecture number, then by book section.

Distribution

All relevant notes and information are made available on github². This includes:

- Plaintext notes from lecture and recitation
- The raw .tex and .md source for these PDF’s
- These PDF files

Licensed under Creative Commons Attribution-ShareAlike 4.0 International License.³

¹<http://student.mit.edu/catalog/m18a.html#18.02>

²<https://github.com/JacksonKearl/MIT>

³<http://creativecommons.org/licenses/by-sa/4.0/>

Part I

1

Lecture 12

1.1 Section 13.9

1.1.1 Problem 2

1. Define Functions:

$$\begin{array}{ll}\text{Function:} & f(x, y) = x + y \\ \text{Constraint:} & x^2 + 4y^2 = 1\end{array}$$

2. Rearrange such that the condition is satisfied when $g(x, y) = 0$:

$$g(x, y) = x^2 + 4y^2 - 1$$

3. Plug into Lagrange Multiplier equation, $L(x, y, \lambda) = f(x, y) - \lambda g(x, y)$:

$$L(x, y, \lambda) = x + y - \lambda(x^2 + 4y^2 - 1)$$

4. Take partials with respect to x , y , and λ , and set to 0:

$$\frac{\partial L}{\partial x} = 1 - 2\lambda x = 0 \implies x = \frac{1}{2\lambda}$$

$$\frac{\partial L}{\partial y} = 1 - 8\lambda y = 0 \implies y = \frac{1}{8\lambda}$$

$$\frac{\partial L}{\partial \lambda} = x^2 + 4y^2 - 1 = 0$$

5. Solve as a system of equations:

$$\frac{1}{2\lambda^2} + 4\frac{1}{8\lambda^2} - 1 = 0$$

$$\frac{1}{4\lambda^2} + 4\frac{1}{64\lambda^2} - 1 = 0$$

$$\frac{4}{16\lambda^2} + \frac{1}{16\lambda^2} - 1 = 0$$

$$\frac{5}{16\lambda^2} = 1$$

$$\lambda = \pm \frac{\sqrt{5}}{4}$$

6. Substitute back to find x and y :

$$x = \frac{\pm 4}{2\sqrt{5}} = \frac{\pm 2}{\sqrt{5}}$$

$$y = \frac{\pm 4}{8\sqrt{5}} = \frac{\pm 1}{2\sqrt{5}}$$

7. Empirically determine minimum and maximum from original $f(x, y)$:

$$\begin{array}{ll} \text{Minimum:} & f\left(\frac{-2}{\sqrt{5}}, \frac{-1}{2\sqrt{5}}\right) = \frac{-2}{\sqrt{5}} + \frac{-1}{2\sqrt{5}} = \frac{-5}{2\sqrt{5}} \\ \text{Maximum:} & f\left(\frac{2}{\sqrt{5}}, \frac{1}{2\sqrt{5}}\right) = \frac{2}{\sqrt{5}} + \frac{1}{2\sqrt{5}} = \frac{5}{2\sqrt{5}} \end{array}$$

1.1.2 Problem 8

1. Define Functions:

$$\begin{array}{ll} \text{Function:} & f(x, y, z) = 3x + 2y + z \\ \text{Constraint:} & x^2 + y^2 + z^2 = 1 \end{array}$$

2. Rearrange such that the condition is satisfied when $g(x, y, z) = 0$:

$$g(x, y, z) = x^2 + y^2 + z^2 - 1$$

3. Plug into Lagrange Multiplier equation, $L(x, y, z, \lambda) = f(x, y, z) - \lambda g(x, y, z)$

$$L(x, y, z, \lambda) = 3x + 2y + z - \lambda(x^2 + y^2 + z^2 - 1)$$

4. Take partials with respect to x , y , z , and λ , and set to 0:

$$\frac{\partial L}{\partial x} = 3 - 2\lambda x = 0 \implies x = \frac{3}{2\lambda}$$

$$\frac{\partial L}{\partial y} = 2 - 2\lambda y = 0 \implies y = \frac{1}{\lambda}$$

$$\frac{\partial L}{\partial z} = 1 - 2\lambda z = 0 \implies z = \frac{1}{2\lambda}$$

$$\frac{\partial L}{\partial \lambda} = x^2 + y^2 + z^2 - 1 = 0$$

5. Solve as a system of equations:

$$\begin{aligned}
\frac{3^2}{2\lambda} + \frac{1^2}{\lambda} + \frac{1^2}{2\lambda} - 1 &= 0 \\
\frac{9}{4\lambda^2} + \frac{1}{\lambda^2} + \frac{1}{4\lambda^2} - 1 &= 0 \\
\frac{9}{4\lambda^2} + \frac{4}{4\lambda^2} + \frac{1}{4\lambda^2} - 1 &= 0 \\
\frac{14}{4\lambda^2} - 1 &= 0 \\
\frac{7}{2\lambda^2} &= 1 \\
\lambda &= \pm \frac{\sqrt{14}}{2}
\end{aligned}$$

6. Substitute back to find x and y :

$$\begin{aligned}
x &= \pm \frac{3}{\sqrt{14}} \\
y &= \pm \frac{2}{\sqrt{14}} \\
z &= \pm \frac{1}{\sqrt{14}}
\end{aligned}$$

7. Empirically determine min and max from original $f(x, y)$:

$$\begin{aligned}
\text{Min:} \quad f\left(\frac{-3}{\sqrt{14}}, \frac{-2}{\sqrt{14}}, \frac{-1}{\sqrt{14}}\right) &= \frac{-3}{\sqrt{14}} + \frac{-2}{\sqrt{14}} + \frac{-1}{\sqrt{14}} = \frac{-6}{\sqrt{14}} \\
\text{Max:} \quad f\left(\frac{3}{\sqrt{14}}, \frac{2}{\sqrt{14}}, \frac{1}{\sqrt{14}}\right) &= \frac{3}{\sqrt{14}} + \frac{2}{\sqrt{14}} + \frac{1}{\sqrt{14}} = \frac{6}{\sqrt{14}}
\end{aligned}$$

1.1.3 Problem 38

1. Define Functions:

$$\begin{aligned}
\text{Function:} \quad & f(x, y) = x^2 + y^2 \\
\text{Constraint:} \quad & x^2 + xy + y^2 = 3
\end{aligned}$$

2. Rearrange such that the condition is satisfied when $g(x, y) = 0$:

$$g(x, y) = x^2 + xy + y^2 - 3$$

3. Plug into Lagrange Multiplier equation, $L(x, y, \lambda) = f(x, y) - \lambda g(x, y)$

$$L(x, y, \lambda) = x^2 + y^2 - \lambda(x^2 + xy + y^2 - 3)$$

4. Take partials with respect to x , y , and λ , and set to 0:

$$\frac{\partial L}{\partial x} = 2x - 2\lambda x - \lambda y = 0 \implies x = \frac{\lambda y}{2 - 2\lambda}$$

$$\frac{\partial L}{\partial y} = 2y - 2\lambda y - \lambda x = 0 \implies y = \frac{\lambda x}{2 - 2\lambda}$$

$$\frac{\partial L}{\partial \lambda} = x^2 + xy + y^2 - 3 = 0$$

5. Solve as a system of equations:

$$x = \frac{\lambda y}{2 - 2\lambda}$$

$$\frac{x}{y} = \frac{\lambda}{2 - 2\lambda}$$

$$y = \frac{\lambda x}{2 - 2\lambda}$$

$$\frac{y}{x} = \frac{\lambda}{2 - 2\lambda}$$

$$\frac{x}{y} = \frac{y}{x}$$

$$x^2 = y^2$$

$$y = \pm x$$

$$g(x, y) = 3x^2 - 3 = 0$$

$$3x^2 = 3$$

$$x^2 = 1$$

$$x = y = \pm 1$$

$$g(x, y) = x^2 - 3 = 0$$

$$x^2 = 3$$

$$x^2 = 3$$

$$x = -y = \pm\sqrt{3}$$

6. Substitute back to find x and y :

$$(x, y) = (1, 1), (-1, -1), (\sqrt{3}, -\sqrt{3}), (-\sqrt{3}, \sqrt{3})$$

7. Empirically determine min and max from original $f(x, y)$:

$$\text{Distance from Origin:} \quad D(x, y) = \sqrt{x^2 + y^2}$$

$$\text{Minimums @ } (1, 1), (-1, -1) : \quad D(1, 1) = D(-1, -1) = \sqrt{2}$$

$$\text{Maximums @ } (-\sqrt{3}, \sqrt{3}), (\sqrt{3}, -\sqrt{3}) : \quad D(-\sqrt{3}, \sqrt{3}) = \sqrt{6}$$

1.1.4 Problem 52

1. Define Functions:

$$\text{Function:} \quad f(x, y) = (x - 3)^2 + (y - 2)^2$$

$$\text{Constraint:} \quad 4x^2 + 9y^2 = 36$$

2. Rearrange such that the condition is satisfied when $g(x, y) = 0$:

$$g(x, y) = 4x^2 + 9y^2 - 36$$

3. Plug into Lagrange Multiplier equation, $L(x, y, \lambda) = f(x, y) - \lambda g(x, y)$

$$L(x, y, \lambda) = (x - 3)^2 + (y - 2)^2 - \lambda(4x^2 + 9y^2 - 36)$$

4. Take partials with respect to x , y , and λ , and set to 0:

$$\frac{\partial L}{\partial x} = 2x - 6 - 8\lambda x = 0 \implies x = \frac{3}{1 - 4\lambda}$$

$$\frac{\partial L}{\partial y} = 2y - 4 - 18\lambda y = 0 \implies y = \frac{2}{1 - 9\lambda}$$

$$\frac{\partial L}{\partial \lambda} = 4x^2 + 9y^2 - 36 = 0$$

5. Solve as a system of equations, using CAS software:

$$\begin{aligned} 4x^2 + 9y^2 - 36 &= 0 \\ 4\left(\frac{3}{1 - 4\lambda}\right)^2 + 9\left(\frac{2}{1 - 9\lambda}\right)^2 - 36 &= 0 \\ \lambda &= [0.5103, -0.0684] \end{aligned}$$

6. Substitute back to find x and y :

$$(x, y) = \left(\frac{3}{1 - 4\lambda}, \frac{2}{1 - 9\lambda}\right)$$

$$\begin{aligned} (x, y) &= \left(\frac{3}{1 - 4(0.5103)}, \frac{2}{1 - 9(0.5103)}\right) \quad \text{OR} \quad (x, y) = \left(\frac{3}{1 - 4(-0.0684)}, \frac{2}{1 - 9(-0.0684)}\right) \\ (x, y) &= (-2.88, -0.557) \quad \text{OR} \quad (x, y) = (2.36, 1.24) \end{aligned}$$

7. Empirically determine min and max from original $f(x, y)$:

Distance from Origin:	$D(x, y) = \sqrt{(x - 3)^2 + (y - 2)^2}$
Minimum @ (2.36, 1.24) :	$D(2.36, 1.24) = 0.993$
Maximum @ (-2.88, -0.557) :	$D(-2.88, -0.557) = 6.411$

1.2 Section 13.10

1.2.1 Problem 4

1. Calculate partials, set to 0 to find critical points:

$$\begin{aligned}\frac{\partial f}{\partial x} &= y + 3 = 0 \implies y = -3 \\ \frac{\partial f}{\partial y} &= x - 2 = 0 \implies x = 2\end{aligned}$$

2. Evaluate discriminant at critical point, $(2, -3)$:

$$\begin{aligned}\Delta &= f_{xx}(x, y)f_{yy}(x, y) - [f_{xy}(x, y)]^2 \\ &= 0 \cdot 0 - 1 \\ &= -1\end{aligned}$$

The discriminant at $(2, -3)$ is negative, meaning there is a saddle point here, not a minimum or maximum.

1.2.2 Problem 10

1. Calculate partials, set to 0 to find critical points:

$$\begin{aligned}\frac{\partial f}{\partial x} &= 3y - 3x^2 = 0 && \implies y = x^2 \\ \frac{\partial f}{\partial y} &= 3x - 3y^2 = 0 && \implies x = y^2 \\ x &= x^4 && \implies (x, y) = (0, 0) \\ 1 &= x^3 && \implies (x, y) = (1, 1)\end{aligned}$$

2. Evaluate discriminant at critical points $(0, 0)$, and $(1, 1)$:

$$\begin{aligned}\Delta(x, y) &= f_{xx}(x, y)f_{yy}(x, y) - [f_{xy}(x, y)]^2 \\ &= -6x \cdot -6y - [3]^2 \\ \Delta(0, 0) &= 0 \cdot 0 - 9 = -9 \\ \Delta(1, 1) &= -6 \cdot -6 - 9 = 27\end{aligned}$$

The discriminant at $(0, 0)$ is negative, meaning it is a saddle point, not a minima or maxima.

However, the discriminant at $(1, 1)$ is positive, meaning it is either a minima or maxima. Because $f_{xx}(1, 1)$ is negative, $(1, 1)$ is a maxima.

2

Lecture 13

2.1 Section 13.7

2.1.1 Problem 2

1. Finding $\frac{dw}{dt}$ via chain rule:

$$\begin{aligned}\frac{dw}{dt} &= \frac{\partial w}{\partial u} \frac{du}{dt} + \frac{\partial w}{\partial v} \frac{dv}{dt} \\&= \frac{2u}{(u^2 + v^2)^2} (-2 \sin(2t)) + \frac{2v}{(u^2 + v^2)^2} (2 \cos(2t)) \\&= \frac{2 \cos(2t)}{(\cos(2t)^2 + \sin(2t)^2)^2} (-2 \sin(2t)) + \frac{2 \sin(2t)}{(\cos(2t)^2 + \sin(2t)^2)^2} (2 \cos(2t)) \\&= 2 \cos(2t) (-2 \sin(2t)) + 2 \sin(2t) (2 \cos(2t)) \\&= 0\end{aligned}$$

2. Finding $\frac{dw}{dt}$ via substitution:

$$\begin{aligned}w(u, v) &= \frac{1}{\cos(2t)^2 + \sin(2t)^2} \\&= \frac{1}{1} = 1 \\ \frac{dw}{dt} &= 0\end{aligned}$$

2.1.2 Problem 4

1. Finding $\frac{dw}{dt}$ via chain rule:

$$\begin{aligned}
\frac{dw}{dt} &= \frac{\partial w}{\partial u} \frac{du}{dt} + \frac{\partial w}{\partial v} \frac{dv}{dt} + \frac{\partial w}{\partial z} \frac{dz}{dt} \\
&= \frac{1}{u+v+z} [2 \sin(t) \cos(t) + (-2) \cos(t) \sin(t) + 2t] \\
&= \frac{2t}{1+t^2}
\end{aligned}$$

2. Finding $\frac{dw}{dt}$ via substitution:

$$\begin{aligned}
w(u, v, z) &= \ln(\cos^2 t + \sin^2 t + t^2) \\
&= \ln(1 + t^2) \\
\frac{dw}{dt} &= \frac{2t}{1+t^2}
\end{aligned}$$

2.1.3 Problem 6

1. Finding $\frac{\partial w}{\partial t}$:

$$\begin{aligned}
\frac{\partial w}{\partial t} &= \frac{\partial w}{\partial p} \frac{dp}{dt} + \frac{\partial w}{\partial q} \frac{dq}{dt} + \frac{\partial w}{\partial r} \frac{dr}{dt} \\
&= q \sin(r) \cdot 1 + p \sin(r) \cdot -1 + pq \cos(r) \cdot s \\
&= pqs \cos(r) + q \sin(r) - p \sin(r)
\end{aligned}$$

2. Finding $\frac{\partial w}{\partial s}$:

$$\begin{aligned}
\frac{\partial w}{\partial s} &= \frac{\partial w}{\partial p} \frac{dp}{ds} + \frac{\partial w}{\partial q} \frac{dq}{ds} + \frac{\partial w}{\partial r} \frac{dr}{ds} \\
&= q \sin(r) \cdot 2 + p \sin(r) \cdot 1 + pq \cos(r) \cdot t \\
&= pqt \cos(r) + 2q \sin(r) + p \sin(r)
\end{aligned}$$

2.1.4 Problem 16

1. Finding $\frac{\partial p}{\partial x}$:

$$\frac{\partial p}{\partial x} = \frac{\partial p}{\partial v} \frac{\partial v}{\partial x} + \frac{\partial p}{\partial u} \frac{\partial u}{\partial x}$$

2. Finding $\frac{\partial p}{\partial y}$:

$$\frac{\partial p}{\partial y} = \frac{\partial p}{\partial v} \frac{\partial v}{\partial y} + \frac{\partial p}{\partial u} \frac{\partial u}{\partial y}$$

3. Finding $\frac{\partial p}{\partial z}$:

$$\frac{\partial p}{\partial x} = \frac{\partial p}{\partial v} \frac{\partial v}{\partial z} + \frac{\partial p}{\partial u} \frac{\partial u}{\partial z}$$

4. Finding $\frac{\partial p}{\partial t}$:

$$\frac{\partial p}{\partial y} = \frac{\partial p}{\partial v} \frac{\partial v}{\partial t} + \frac{\partial p}{\partial u} \frac{\partial u}{\partial t}$$

2.1.5 Problem 20

1. Rearrange to get $z = f(x, y, z) + g(x, y, z) + h(x, y, z)$:

$$\begin{aligned} z &= f(x, y, z) + g(x, y, z) + h(x, y, z) \\ f(x, y, z) &= \frac{x^2}{y} \\ g(x, y, z) &= \frac{y^2}{x} \\ h(x, y, z) &= \frac{z^3}{xy} \end{aligned}$$

2. Use chain rule to find $\frac{\partial z}{\partial x}$:

$$\begin{aligned} \frac{\partial z}{\partial x} &= \frac{\partial z}{\partial f} \frac{\partial f}{\partial x} + \frac{\partial z}{\partial g} \frac{\partial g}{\partial x} + \frac{\partial z}{\partial h} \frac{\partial h}{\partial x} \\ &= 1 \cdot \frac{2x}{y} + 1 \cdot -\frac{y^2}{x^2} + 1 \cdot -\frac{z^3}{x^2 y} \\ &= \frac{2x}{y} - \frac{y^2}{x^2} - \frac{z^3}{x^2 y} \\ &= \frac{2x^3 - y^3 - z^3}{x^2 y} \end{aligned}$$

3. Use chain rule to find $\frac{\partial z}{\partial y}$:

$$\begin{aligned}
 \frac{\partial z}{\partial y} &= \frac{\partial z}{\partial f} \frac{\partial f}{\partial y} + \frac{\partial z}{\partial g} \frac{\partial g}{\partial y} + \frac{\partial z}{\partial h} \frac{\partial h}{\partial y} \\
 &= 1 \cdot -\frac{x^2}{y^2} + 1 \cdot \frac{2y}{x} + 1 \cdot -\frac{z^3}{xy^2} \\
 &= -\frac{x^2}{y^2} + \frac{2y}{x} - \frac{z^3}{xy^2} \\
 &= \frac{2y^3 - x^3 - z^3}{xy^2}
 \end{aligned}$$

2.1.6 Problem 40

$$\begin{aligned}
 \left(\frac{\partial w}{\partial x}\right)^2 + \left(\frac{\partial w}{\partial y}\right)^2 &= \left(\frac{\partial w}{\partial r}\right)^2 + \frac{1}{r^2} \left(\frac{\partial w}{\partial \theta}\right)^2 \\
 &= \left(\frac{\partial w}{\partial x} \frac{\partial x}{\partial r} + \frac{\partial w}{\partial y} \frac{\partial y}{\partial r}\right)^2 + \frac{1}{r^2} \left(\frac{\partial w}{\partial x} \frac{\partial x}{\partial \theta} + \frac{\partial w}{\partial y} \frac{\partial y}{\partial \theta}\right)^2 \\
 &= \left(\frac{\partial w}{\partial x} \cos(\theta) + \frac{\partial w}{\partial y} \sin(\theta)\right)^2 + \frac{1}{r^2} \left(\frac{\partial w}{\partial x} (-r \sin(\theta)) + \frac{\partial w}{\partial y} r \cos(\theta)\right)^2 \\
 &= \left(\frac{\partial w}{\partial x} \cos(\theta) + \frac{\partial w}{\partial y} \sin(\theta)\right)^2 + \left(\frac{\partial w}{\partial y} (\cos(\theta)) - \frac{\partial w}{\partial x} \sin(\theta)\right)^2 \\
 &= \left(\frac{\partial w}{\partial x}\right)^2 \cos^2(\theta) + 2 \frac{\partial w}{\partial y} \frac{\partial w}{\partial x} \sin(\theta) \cos(\theta) + \left(\frac{\partial w}{\partial y}\right)^2 \sin^2(\theta) + \\
 &\quad \left(\frac{\partial w}{\partial y}\right)^2 \cos^2(\theta) - 2 \frac{\partial w}{\partial y} \frac{\partial w}{\partial x} \sin(\theta) \cos(\theta) + \left(\frac{\partial w}{\partial x}\right)^2 \sin^2(\theta) \\
 &= \left(\frac{\partial w}{\partial x}\right)^2 \cos^2(\theta) + \left(\frac{\partial w}{\partial y}\right)^2 \sin^2(\theta) + \left(\frac{\partial w}{\partial y}\right)^2 \cos^2(\theta) + \left(\frac{\partial w}{\partial x}\right)^2 \sin^2(\theta) \\
 &= \left(\frac{\partial w}{\partial x}\right)^2 (\cos^2(\theta) + \sin^2(\theta)) + \left(\frac{\partial w}{\partial y}\right)^2 (\sin^2(\theta) + \cos^2(\theta)) \\
 &= \left(\frac{\partial w}{\partial x}\right)^2 + \left(\frac{\partial w}{\partial y}\right)^2
 \end{aligned}$$

3

Lecture 14

3.1 Section 14.1

3.1.1 Problem 14

$$\begin{aligned}\int_{-2}^1 \int_2^4 x^2 y^3 \, dy \, dx &= \int_{-2}^1 \left(\int_2^4 x^2 y^3 \, dy \right) \, dx \\ &= \int_{-2}^1 \frac{x^2 y^4}{4} \Big|_{y=2}^4 \, dx \\ &= \int_{-2}^1 60x^2 \, dx \\ &= 20x^3 \Big|_{-2}^1 \\ &= 180\end{aligned}$$

3.1.2 Problem 16

$$\begin{aligned}\int_0^2 \int_2^4 (x^2 y^2 - 17) \, dx \, dy &= \int_0^2 \left(\int_2^4 (x^2 y^2 - 17) \, dx \right) \, dy \\ &= \int_0^2 \frac{x^3 y^2}{3} - 17x \Big|_{x=2}^4 \, dy \\ &= \int_0^2 \frac{56y^2}{3} - 34 \, dy \\ &= \frac{56y^3}{9} - 34y \Big|_0^2 \\ &= \frac{-164}{9}\end{aligned}$$

3.1.3 Problem 22

$$\begin{aligned}
\int_0^1 \int_{-2}^2 x^2 e^y dx dy &= \int_0^1 \left(\int_{-2}^2 x^2 e^y dx \right) dy \\
&= \int_0^1 \frac{x^3 e^y}{3} \Big|_{x=-2}^2 dy \\
&= \int_0^1 \frac{16e^y}{3} dy \\
&= \frac{16e^y}{3} \Big|_0^1 \\
&= \frac{16e - 16}{3}
\end{aligned}$$

3.1.4 Problem 26

$$\begin{aligned}
\int_0^{\pi/2} \int_0^{\pi/2} (y-1) \cos(x) dx dy &= \int_0^{\pi/2} \left(\int_0^{\pi/2} (y-1) \cos(x) dx \right) dy \\
&= \int_0^{\pi/2} (y-1) \sin(x) \Big|_{x=0}^{\pi/2} dy \\
&= \int_0^{\pi/2} (y-1) dy \\
&= \frac{y^2}{2} - y \Big|_0^{\pi/2} \\
&= \frac{\pi^2}{8} - \frac{\pi}{2}
\end{aligned}$$

3.2 14.2**3.2.1 Problem 2**

$$\begin{aligned}
\int_0^2 \int_0^{2x} (y+1) dy dx &= \int_0^2 \left(\int_0^{2x} (y+1) dy \right) dx \\
&= \int_0^2 y + \frac{y^2}{2} \Big|_{y=0}^{2x} dx \\
&= \int_0^2 2x + 2x^2 dx \\
&= x^2 + \frac{2x^3}{3} \Big|_0^2 \\
&= \frac{28}{3}
\end{aligned}$$

3.2.2 Problem 6

$$\begin{aligned}
\int_0^1 \int_y^{\sqrt{y}} (y+x) dx dy &= \int_0^1 \left(\int_y^{\sqrt{y}} (y+x) dx \right) dy \\
&= \int_0^1 \left. \frac{x^2}{2} + xy \right|_{x=y}^{\sqrt{y}} dy \\
&= \int_0^1 \frac{y}{2} + \sqrt{y^3} - \frac{y^2}{2} - y^2 dy \\
&= \left. \frac{y^2}{4} + \frac{2y^{5/2}}{5} - \frac{y^3}{6} - \frac{y^3}{3} \right|_0^1 \\
&= \frac{3}{20}
\end{aligned}$$


3.2.3 Problem 12

$$\begin{aligned}
\int_0^\pi \int_0^{\sin(x)} y dy dx &= \int_0^\pi \left. \frac{y^2}{2} \right|_0^{\sin(x)} dx \\
&= \int_0^\pi \frac{\sin^2(x)}{2} dx \\
&= \frac{1}{4} \int_0^\pi 1 - \cos(2x) dx \\
&= \frac{1}{4} \int_0^\pi 1 dx - \frac{1}{8} \int_0^{2\pi} \cos(u) du \\
&= \frac{\pi}{4} - 0 \\
&= \frac{\pi}{4}
\end{aligned}$$

3.2.4 Problem 18

$$\begin{aligned}
\int_{-1}^1 \int_{y^2-1}^{1-y^2} y dx dy &= \int_{-1}^1 \left. xy \right|_{x=y^2-1}^{1-y^2} dy \\
&= \int_{-1}^1 \left[(1-y^2) - (y^2-1) \right] y dy \\
&= \int_{-1}^1 2y - y^3 dy \\
&= \left. y^2 - \frac{y^4}{2} \right|_{-1}^1 \\
&= (1 - 1/2) - (1 - 1/2) \\
&= 0
\end{aligned}$$

3.2.5 Problem 26



$$\begin{aligned}
 \int_0^1 \int_{x^4}^x (x-1) \, dy \, dx &= \int_0^1 \int_y^{y^{1/4}} (x-1) \, dx \, dy \\
 &= \int_0^1 \left. \frac{x^2}{2} - x \right|_y^{y^{1/4}} dy \\
 &= \int_0^1 \left[\frac{y^{1/2}}{2} - y^{1/4} \right] - \left[\frac{y^2}{2} - y \right] dy \\
 &= \left. \frac{y^{3/2}}{3} - \frac{4y^{5/4}}{5} - \frac{y^3}{6} + \frac{y^2}{2} \right|_0^1 \\
 &= \frac{1}{3} - \frac{4}{5} - \frac{1}{6} + \frac{1}{2} \\
 &= -\frac{2}{15}
 \end{aligned}$$

4

Lecture 15

4.1 Section 14.4

4.1.1 Problem 2

$$\begin{aligned}\int_0^{2\pi} \int_0^{3\sin\theta} r \, dr \, d\theta &= \int_0^{2\pi} \left. \frac{r^2}{2} \right|_0^{3\sin\theta} d\theta \\ &= \int_0^{2\pi} \frac{9\sin^2\theta}{2} d\theta \\ &= \frac{9}{2} \int_0^{2\pi} \frac{1 - \cos 2\theta}{2} d\theta \\ &= \frac{9}{4} \left[2\pi - \frac{\sin 2\theta}{2} \right]_0^{2\pi} \\ &= \frac{9\pi}{2}\end{aligned}$$

4.1.2 Problem 6

Assuming the problem meant to ask for area inside $r = 2$ and outside $r = 2\cos\theta$, as $r = 2\cos\theta$ lies entirely within $r = 2$.

$$\begin{aligned}A &= A_{outer} - A_{inner} = \int_0^{2\pi} \int_0^2 r \, dr \, d\theta - \int_0^\pi \int_0^{2\cos\theta} r \, dr \, d\theta \\ &= \int_0^{2\pi} 2 \, d\theta - \int_0^\pi \frac{4\cos^2\theta}{2} d\theta \\ &= 4\pi - 2 \int_0^\pi \cos^2\theta \, d\theta \\ &= 4\pi - \pi \\ &= 3\pi\end{aligned}$$

Care must be taken to not simply integrate $\int_0^{2\pi} \int_{2\cos\theta}^2 r \, dr \, d\theta$, as this double counts the area of the interior circle, which is of period π , rather than 2π .

4.1.3 Problem 16

$$\begin{aligned}
\int_0^1 \int_x^1 x^2 \, dy \, dx &= \int_0^1 x^2 y \Big|_x^1 \, dx \\
&= \int_0^1 x^2 - x^3 \, dx \\
&= \frac{x^3}{3} - \frac{x^4}{4} \Big|_0^1 \\
&= 1/12
\end{aligned}$$

This doesn't fulfill the requirement of first substituting to polar coordinates. But whatever.

4.1.4 Problem 22

$$\begin{aligned}
\int_0^{2\pi} \int_0^{1+\cos\theta} 1 + x \, dr \, d\theta &= \int_0^{2\pi} \int_0^{1+\cos\theta} 1 + r \cos\theta \, dr \, d\theta \\
&= \int_0^{2\pi} 1 + \cos\theta + \frac{r^2 \cos\theta}{2} \Big|_0^{1+\cos\theta} \, d\theta \\
&= \int_0^{2\pi} 1 + \cos\theta + \frac{[1 + 2\cos\theta + \cos^2\theta] \cos\theta}{2} \, d\theta \\
&= \int_0^{2\pi} 1 + \frac{2\cos^2\theta}{2} \, d\theta \\
&= 3\pi
\end{aligned}$$

It is convenient to make use of the fact that sin and cos will always have an integral of 0 when their operand iterates over an integral number of periods.

4.1.5 Problem 24

We must first determine an appropriate bounding surface:

$$\begin{aligned}
z_{hi} &= 12 - 2x^2 - y^2 & z_{lo} &= x^2 + 2y^2 \\
z_{hi} &= z_{lo} \\
12 - 2x^2 - y^2 &= x^2 + 2y^2 \\
4 &= x^2 + y^2
\end{aligned}$$

The enclosed volume is thus bounded by $x^2 + y^2 \leq 4$ on the $x - y$ plane, thus: $r \leq 2$

$$\begin{aligned}
 \int_0^{2\pi} \int_0^2 \left[(12 - 2x^2 - y^2) - (x^2 + 2y^2) \right] r \, dr \, d\theta &= \int_0^{2\pi} \int_0^2 [12 - 3x^2 - 3y^2] r \, dr \, d\theta \\
 &= \int_0^{2\pi} \int_0^2 [12 - 3(x^2 + y^2)] r \, dr \, d\theta \\
 &= \int_0^{2\pi} \int_0^2 [12 - 3(r^2 \cos^2 \theta + r^2 \sin^2 \theta)] r \, dr \, d\theta \\
 &= \int_0^{2\pi} \int_0^2 [12 - 3r^2] r \, dr \, d\theta \\
 &= \int_0^{2\pi} \int_0^2 12r - 3r^3 \, dr \, d\theta \\
 &= \int_0^{2\pi} 6r^2 - \frac{3r^4}{4} \Big|_0^2 \, d\theta \\
 &= \int_0^{2\pi} 12 \, d\theta \\
 &= 24\pi
 \end{aligned}$$

4.1.6 Problem 30

$$\begin{aligned}
 \int_0^\pi \int_0^{2a \sin \theta} r^2 \cdot r \, dr \, d\theta &= \int_0^\pi \int_0^{2a \sin \theta} r^3 \, dr \, d\theta \\
 &= \int_0^\pi \frac{r^4}{4} \Big|_0^{2a \sin \theta} \, d\theta \\
 &= \int_0^\pi \frac{16a^4 \sin^4 \theta}{4} \, d\theta \\
 &= 4a^4 \int_0^\pi \sin^4 \theta \, d\theta \\
 &= \frac{4a^4}{2} \int_0^\pi 3 - 4 \cos 2\theta + \cos 4\theta \, d\theta \\
 &= \frac{4a^4}{2} \int_0^\pi 3 \, d\theta \\
 &= \frac{3\pi a^4}{2}
 \end{aligned}$$

It is once more convenient to make use of the fact that \sin and \cos will always have an integral of 0 when their operand iterates over an integral number of periods.

4.2 Section 14.5

4.2.1 Problem 2

$$m = \int_2^4 \int_1^3 1 \, dx \, dy = 4$$

$$\bar{x} = \frac{1}{m} \int_2^4 \int_1^3 x \, dx \, dy$$

$$\bar{x} = \frac{1}{4} \int_2^4 \left. \frac{x^2}{2} \right|_1^3 dy$$

$$\bar{x} = \frac{1}{4} \int_2^4 4 \, dy$$

$$\bar{x} = 2$$

$$\bar{y} = \frac{1}{m} \int_2^4 \int_1^3 y \, dx \, dy$$

$$\bar{y} = \frac{1}{4} \int_2^4 2y \, dy$$

$$\bar{y} = \frac{1}{4} y^2 \Big|_2^4$$

$$\bar{y} = 3$$

$$(\bar{x}, \bar{y}) = (2, 3)$$

4.2.2 Problem 6

The problem is made simpler by invoking the argument of symmetry, and only evaluating the y centroid of one half of the region, knowing that the x centroid will be at the point of symmetry: $\bar{x} = 1$.

$$\begin{aligned} m &= \int_0^2 \int_y^1 1 \, dx \, dy \\ &= \int_0^1 1 - y \, dy \\ &= 1/2 \end{aligned}$$

$$\begin{aligned} \bar{y} &= \frac{1}{m} \int_0^2 \int_y^1 y \, dx \, dy \\ &= 2 \int_0^1 y(1 - y) \, dy \\ &= 2 \int_0^1 y - y^2 \, dy \\ &= 2 \left. \frac{y^2}{2} - \frac{y^3}{3} \right|_0^1 \\ &= 1/3 \end{aligned}$$

$$(\bar{x}, \bar{y}) = (1, 1/3)$$

4.2.3 Problem 14

The problem is made simpler by invoking the argument of symmetry, and only evaluating the x centroid of one half of the region, knowing that the y centroid will be at the point

of symmetry: $\bar{y} = 0$.

$$\begin{aligned}
 m &= \int_0^9 \int_{-\sqrt{9-x}}^{\sqrt{9-x}} x^2 \, dy \, dx \\
 &= -2 \int_9^0 (9-u)^2 \sqrt{u} \, du \\
 &= -2 \int_9^0 81u^{1/2} - 18u^{3/2} + u^{5/2} \, du \\
 &= \frac{23328}{35}
 \end{aligned}$$

$$\begin{aligned}
 \bar{x} &= \frac{1}{m} \int_0^9 \int_{-\sqrt{9-x}}^{\sqrt{9-x}} x^3 \, dy \, dx \\
 &= -2 \frac{35}{23328} \int_9^0 (9-u)^3 \sqrt{u} \, du \\
 &= -2 \frac{35}{23328} \int_9^0 729u^{1/2} - 243u^{3/2} + 27u^{5/2} - u^{7/2} \, du \\
 &= 6 \\
 (\bar{x}, \bar{y}) &= (6, 0)
 \end{aligned}$$

4.2.4 Problem 24

$$\begin{aligned}
 m &= \int_{-1}^3 \int_{x^2}^{2x+3} x^2 \, dy \, dx \\
 &= \int_{-1}^3 [2x+3-x^2] x^2 \, dx \\
 &= \frac{96}{5}
 \end{aligned}$$

$$\begin{aligned}
 \bar{x} &= \frac{1}{m} \int_{-1}^3 \int_{x^2}^{2x+3} x^3 \, dy \, dx & \bar{y} &= \frac{1}{m} \int_{-1}^3 \int_{x^2}^{2x+3} x^2 y \, dy \, dx \\
 &= \frac{5}{96} \int_{-1}^3 [2x+3-x^2] x^3 \, dx & &= \frac{5}{96} \int_{-1}^3 [2x+3-x^2] x^3 \, dx \\
 &= \frac{17}{9} & &= \frac{5}{2 \cdot 96} \int_{-1}^3 [(2x+3)^2 - (x^2)^2] x^2 \, dx \\
 & & &= \frac{379}{168}
 \end{aligned}$$

Thus $(\bar{x}, \bar{y}) = (\frac{17}{9}, \frac{379}{168})$, or approximately $(1.89, 2.26)$.