

- 11 Suppose that Cauchy's estimate is an equality. Let's create a disk around the origin of radius  $r > 0$ . We know that the maximum is either on a boundary or at a zero. Using Cauchy's estimate we get

$$|f^{(0)}(0)| = |f(0)| = M(0!)/r^0 = M$$

This means that the origin is a maximum and thus a zero. So the maximum of the entire function must be 0. Thus  $f(z) = 0$

- 12 For the domain  $D$  we have a fixed center  $z_0$  and a radius  $r$ .  $\partial D$  can be written as the path  $\partial D(t) = z_0 + re^{it}$  with  $t \in [0, 2\pi]$ . If we integrate  $f(\partial D(t))$  we get the sum of all  $f(z)$  along the path so to get the average we can divide by the length of the path giving us:

$$\frac{1}{2\pi} \int_0^{2\pi} f(\partial D(t)) dt = \frac{1}{2\pi} \int_0^{2\pi} f(z_0 + re^{it}) dt$$

We can then do a substitution with  $z = z_0 + re^{it}$  and  $dz = ire^{it} dt$ .

$$\begin{aligned} \frac{1}{2\pi} \int_0^{2\pi} \frac{f(z)}{ire^{it}} dt &= \frac{1}{2i\pi} \int_0^{2\pi} \frac{f(z)}{re^{it}} dz \\ &= \frac{1}{2i\pi} \int_{\partial D} \frac{f(z)}{z - z_0} dz \end{aligned}$$

by Cauchy's integral formula we can see

$$\frac{1}{2i\pi} \int_{\partial D} \frac{f(z)}{z - z_0} dz = f(z_0)$$

- 13 Suppose we have a domain  $D$  with radius  $r > 0$ . The maximum would be  $|f(z)| \leq Kr^c$ . Then using Cauchy's estimate for  $n = c$ :

$$|f^{(c)}(z)| \leq \frac{Kr^c(c)!}{r^c} = Kc!$$

This means that  $f^{(c)}(z) = w$  and  $w$  is constant. Let's find the antiderivative of  $f^{(c)}(z)$ ,  $c$  times:

$$\begin{aligned} f^{(c-1)}(z) &= wz + l_0 (l \text{ is a constant of integer}) \\ f^{(c-2)}(z_0) &= \frac{wz^2}{2!} + l_0z + l_1 \\ f^{(c-(c-1))}(z_0) &= \frac{wz^{c-1}}{(c-1)!} + l_0 \frac{z^{c-2}}{(c-2)!} + l_1 \frac{z^{c-3}}{(c-3)!} + \dots \\ f(z_0) &= \frac{wz^c}{(c)!} + l_0 \frac{z^{c-1}}{(c-1)!} + l_1 \frac{z^{c-2}}{(c-2)!} + \dots \end{aligned}$$

This gives us a polynomial of degree  $\leq c$

14 Let's consider the Taylor expansion of  $f, g$  centered at 0

$$f(z) = \sum_{n=0}^{\infty} \frac{f^{(n)}(0)z^n}{n!}$$

$$g(z) = \sum_{n=0}^{\infty} \frac{g^{(n)}(0)z^n}{n!}$$

We also know that

$$f^{(n)}(0) = \frac{n!}{2i\pi} \left| \int_{C_r} \frac{f(z)}{(z - z_0)^{n+1}} dz \right|$$

if we choose a small enough  $r$  that would mean

$$\frac{n!}{2i\pi} \left| \int_{C_r} \frac{f(z)}{(z - z_0)^{n+1}} dz \right| = \frac{n!}{2i\pi} \left| \int_{C_r} \frac{g(z)}{(z - z_0)^{n+1}} dz \right|$$

Thus:

$$\begin{aligned} f^{(n)}(0) &= g^{(n)}(0) \\ \implies f(z) &= \sum_{n=0}^{\infty} \frac{f^{(n)}(0)z^n}{n!} \\ &= \sum_{n=0}^{\infty} \frac{g^{(n)}(0)z^n}{n!} \\ &= g(z) \end{aligned}$$

15