

# CS70 HW2

February 22, 2021

## 1 Hit or Miss?

- (a) Incorrect. What the proof proves is that for all positive integer  $n \in \mathbb{R}, n^2 \geq n$ , but the claim is to prove for any positive number  $\in \mathbb{R}$ .

For a counter-example, when  $n = 0.5, n^2 = 0.25, n = 0.5$ , thus  $n^2 < n$ .

- (b) Correct.

- (c) Incorrect. From the reference to the Well Ordering Principle, the first counter-example is when  $n = 0$ , where  $0 < a, b \leq n \implies a = 1, b = 0$ , and we don't know whether  $2 * 1 = 0$  or not.

## 2 A Coin Game

Proof by Strong Induction on the total coin number  $N$ .

*Base Case:* when  $N = 2$ , which can only split into 1 and 1. Thus  $1 * 1 = \frac{2*(2-1)}{2} = 1$ .

*Induction Hypothesis:* Assume total score will be  $\frac{n*(n-1)}{2}$  when  $2 \leq n \leq k$  no matter how to split;

*Induction Step:* We must show when  $n = k + 1$ , the equation still holds. Say we split the stack into  $a$  and  $b$  parts where  $a + b = k + 1, a \in \mathbb{R}, b \in \mathbb{R}$ . Use induction hypothesis on these two parts, we gain the equation

$$\frac{a * (a - 1)}{2} + \frac{b * (b - 1)}{2} + a * b = \frac{(a + b)^2 - (a + b)}{2} = \frac{(k + 1) * k}{2}$$

and since  $a \in \mathbb{R}, b \in \mathbb{R}$ , so proof is done.

## 3 Grid Induction

Claim: Pacman needs  $i + j$  steps to reach  $(0, 0)$ .

Proof by induction on the sum of its coordinates  $N = i + j$ .

*Base Case:* when  $N = 0 \implies i = 0, j = 0$ , which takes no step to reach  $(0, 0)$ ;

*Induction Hypothesis:* Assume pacman at  $(i, j)$  needs exactly  $N = i + j = k$  steps to reach origin.

*Induction Step:* when  $N = k + 1$ , there are two cases:

(a) Pacman walks one step down if allowed. Thus the new sum  $N' = k$ , by induction hypothesis, the total step is  $1 + k$ .

(b) Pacman walks one step left if allowed. Thus the new sum  $N' = k$ , by induction hypothesis, the total step is  $1 + k$ .

In both cases, the total steps needed are  $k + 1$ , so the proof is done.

## 4 Stable Marriage

|     | day 1 |         | day 2 |         | day 3 |      | day 4 |      | day 5 |   |
|-----|-------|---------|-------|---------|-------|------|-------|------|-------|---|
| (a) | women | men     | women |         | women |      | women |      | women |   |
|     | 1     | A, B, C | 1     | A       | 1     | A, D | 1     | D    | 1     | D |
|     | 2     |         | 2     |         | 2     | C    | 2     | A, C | 2     | A |
|     | 3     | D       | 3     | D, B, C | 3     | B    | 3     | B    | 3     | B |
|     | 4     |         | 4     |         | 4     |      | 4     |      | 4     | C |

So the final result is  $\{(D, 1), (A, 2), (B, 3), (C, 4)\}$ ;

(b) The proof consists of two parts : First is to prove it's stable and section is to prove it puts out a male-optimal pairing.

(a) Stability.

Say there is a rough couple in the result produced by this Algorithm, namely in  $(W, M^*)$  and  $(W-, M)$ ,  $W$  and  $M$  are rough couple.

By the definition of rough couple, for  $M$ :  $W > W-$ ; for  $W$ :  $M > M^*$ . Since  $M$  proposes, he proposes to  $W$  earlier than  $W-$ . But  $W$  ends up with  $M^*$ , so for  $W$ :  $M^* > M$ , which contradicts the earlier fact.

So it's stable.

(b) Male-optimal pairing.

Proof similar in the Notebook. Proof by Well Ordering Principle: first introduce a first man  $M$  who is rejected by his optimal woman  $W$ , ...

And it's Male-optimal pairing.

In short, proving stability and optimality needn't the precise day to propose. So the output remains the same.

## 5 Optimal Partners

Proof by Contradiction.

Assume both  $M$  and  $M^*$  's optimal woman are  $W$ . And we have two sets where  $M^*$  and  $M$  end up with his optimal women  $W$ :

Set  $T : (M, W), (M^*, W^*)$ ;

Set  $S : (M, W^-), (M^*, W)$ ;

And for women  $W$ , there are two cases:

- (a) If woman  $W$  likes  $M^*$  more than  $M$ . Then in the Set  $S$ ,  $(M^*, W)$  is a rough couple.
- (b) If woman  $W$  likes  $M$  more than  $M^*$ . Then in the Set  $T$ ,  $(M, W)$  is a rough couple.

In both cases, there exists a contradiction.

So no two men can have the same optimal partner.

## 6 Examples or It's Impossible

- (a) Possible.

| men | preferences | women | preferences |
|-----|-------------|-------|-------------|
| A   | 1, 2, 3     | 1     | A, B, C     |
| B   | 2, 1, 3     | 2     | B, A, C     |
| C   | 3, 2, 1     | 3     | C, B, A     |

- (b) Possible.

| men | preferences | women | preferences |
|-----|-------------|-------|-------------|
| A   | 2, 1, 3     | 1     | A, B, C     |
| B   | 1, 2, 3     | 2     | B, C, A     |
| C   | 2, 3, 1     | 3     | C, A, B     |

- (c) Possible.

| men | preferences | men | preferences |
|-----|-------------|-----|-------------|
| A   | 1, 2, 3     | 1   | C, B, A     |
| B   | 2, 1, 3     | 2   | C, A, B     |
| C   | 3, 2, 1     | 3   | A, B, C     |

- (d) Possible.

But I can't think of an instance.

**Impossible.**

The proof is quite beautiful written by a student. I will write it our tomorrow to see if I have really understood it.

---

*Proof.* Proof by contradiction.

Say every man pairs with his last choice, namely  $\{(M_1, W_1), (M_2, W_2), \dots, (M_n, W_n)\}$ . Then assume the algorithm ends at the  $K^{th}$  day, on that day  $M_1$  proposes to  $W_1$ .

And on  $K - 1^{th}$ ,  $M_1$  proposes to another woman  $W^*$  and is rejected, since  $W^*$  has already someone  $M^*$  on her string.

Having these observation,

1.  $M_1$  proposes to  $W_1$  at the last day  $\implies W_1$  has nobody on her string and never was proposed by any man before.
2.  $W^*$  has  $M^*$  on her string and  $W^*$  is  $M^*$  last choice  $\implies M^*$  must propose to  $W_1$  before.

Contradiction. Thus the proof is done.

□

(e) Impossible.

Say a man  $M$  ends up with his last choice woman  $W$ . Next we wanna prove that on  $W$ 's list,  $M$  is her first choice.

Let  $M$  pairs up with other  $n - 1$  women, say  $W_1, W_2, \dots, W_{n-1}$  whose responding men are  $M_1, M_2, \dots, M_{n-1}$ .

$M$  like  $W_1, W_2, \dots, W_{n-1}$  more than  $W$ , and since they are not rough couple, which implies  $W_1, W_2, \dots, W_{n-1}$  like their  $M_1, M_2, \dots, M_{n-1}$  more than  $M$ .

Besides  $M$  is on  $W_1, W_2, \dots, W_{n-1}$  second place, so  $M_1, M_2, \dots, M_{n-1}$  are all  $W_1, W_2, \dots, W_{n-1}$  first choice.

Thus the remaining couple, for  $W$ , her first choice is not determined.

---

All the above is not related.

Answer should be Possible.

| men | preferences |
|-----|-------------|
| A   | 2, 3, 1     |
| B   | 2, 1, 3     |
| C   | 3, 2, 1     |

| men | preferences |
|-----|-------------|
| 1   | C, A, B     |
| 2   | B, A, C     |
| 3   | C, A, B     |