

# The Hecke algebra action on Morava $E$ -theory of height 2

YIFEI ZHU

Given a one-dimensional formal group of height 2, let  $E$  be the Morava  $E$ -theory spectrum associated to its universal deformation over the Lubin-Tate ring. By computing with moduli spaces of elliptic curves, we give an explication for an algebra of Hecke operators acting on  $E$ -cohomology. This leads to a vanishing result for Rezk’s logarithmic cohomology operation on the units of  $E$ . It identifies a family of elements in the kernel with meromorphic modular forms whose Serre derivative is zero. Our calculation finds a connection to logarithms of modular units. In particular, we work out an action of Hecke operators on certain “logarithmic”  $q$ -series, in the sense of Knopp and Mason, that agrees with our vanishing result and extends the classical Hecke action on modular forms.

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
1.1	Acknowledgements . . . . .	5
1.2	Conventions . . . . .	5
1.3	Correction to two cited formulas . . . . .	6
<b>2</b>	<b>From rings of modular forms to homotopy groups of Morava <math>E</math>-theories</b>	<b>6</b>
2.1	Models for an $E$ -theory of height 2 . . . . .	7
2.2	Homotopy groups of an $E$ -theory at height 2 . . . . .	9
<b>3</b>	<b>From classical to topological Hecke operators</b>	<b>12</b>
3.1	Isogenies and power operations . . . . .	12

3.2	Hecke operators . . . . .	17
3.3	Ando’s coordinate . . . . .	20
<b>4</b>	<b>Kernels of logarithmic cohomology operations at height 2</b>	<b>22</b>
4.1	Connecting to Hecke operators . . . . .	22
4.2	Detecting the kernels . . . . .	26
<b>5</b>	<b>Extending the action of Hecke operators onto logarithmic <math>q</math>-series</b>	<b>30</b>
<b>6</b>	<b>Topological Hecke operators in terms of individual power operations</b>	<b>34</b>
	<b>References</b>	<b>42</b>

## 1 Introduction

In the context of elliptic cohomology, Hecke operators have been studied as cohomology operations by Baker [[Baker1989](#), [Baker1990](#)], Ando [[Ando1995](#)], and Ganter [[Ganter2006](#), [Ganter2009](#), [Ganter2013a](#), [Ganter2013b](#)], among others. The various cohomology theories each have as coefficients a ring of modular forms of a particular type—modular forms on  $\mathrm{SL}_2(\mathbb{Z})$  with no condition at the cusp,  $p$ -adic modular forms, modular forms with level structure—or it is a closely related ring, e.g., a completion of the former at an ideal. These cohomology theories are topological realizations of the domain that Hecke operators act on.

The notion of an elliptic spectrum [[Ando-Hopkins-Strickland2001](#), Definition 1.2] and the theory of multiplicative ring spectra, notably the theorem of Goerss, Hopkins, and Miller [[Goerss-Hopkins2004](#), Corollary 7.6], enable studying this action of Hecke operators via power operations. These operations arise from the multiplicative ring structure on a spectrum, and they capture all the algebraic structure that naturally adheres to the cohomology theory represented by the spectrum [[Brunner-May-McClure-Steinberger1986](#), [Rezk2006a](#)].

In particular, Morava  $E$ -theories are a family of cohomology theories whose power operations are better understood thanks to the work of Ando, Hopkins, Strickland, and

Rezk [Ando-Hopkins-Strickland2004, Rezk2009, Rezk2014]. This family is organized by heights  $n$  and primes  $p$ . Here the theory of formal groups plays a key role: each  $E$ -theory spectrum  $E$  corresponds to a specific one-dimensional commutative formal group, whose finite flat subgroups, in turn, correspond precisely to power operations on  $E$ -cohomology. At height  $n = 2$ , formal groups of elliptic curves provide concrete models for  $E$ -theories. The associated power operations can then be computed from moduli spaces that parametrize isogenies between elliptic curves.

In this paper, we study the action of Hecke operators on Morava  $E$ -theories at height 2, based on the explicit calculations of power operations in [Rezk2008, Zhu2014]. This is motivated by the interpretation—in terms of Hecke operators—for Rezk’s logarithmic operations on  $E$ -cohomology at an arbitrary height (see [Rezk2006b, 1.12]). At height 2, these “logarithms” are critical in the work of Ando, Hopkins, and Rezk on rigidification of the string-bordism elliptic genus, or more precisely, on  $E_\infty$ -string orientations of the spectrum of topological modular forms [Ando-Hopkins-Rezk2010]. For the space of such orientations, its set of components can be detected by elements in the kernel of a logarithm. Explicitly, these elements are identified with Eisenstein series  $\mathcal{E}_k$ , which are eigenforms of Hecke operators [Ando-Hopkins-Rezk2010, Theorem 12.3] (cf. [Sprang2014, Theorem 40] and [Wilson2015, Theorem 0.4]).

In Section 4, we give a different account of certain elements in the kernel of a logarithm at height 2: they are meromorphic modular forms with vanishing Serre derivative, including modular forms whose zeros and poles are located only at the cusps, such as the modular discriminant  $\Delta$ . This is stated as follows.

**Theorem 1.1** (Proposition 2.9 and Theorem 4.13) *Let  $E$  be a Morava  $E$ -theory spectrum of height 2 at the prime  $p$ , and let  $N > 3$  be any integer prime to  $p$ .*

- (i) *There is a ring homomorphism  $\beta_N^{(p)}: \mathrm{MF}(\Gamma_1(N)) \rightarrow E^0$ , where  $\mathrm{MF}(\Gamma_1(N))$  is the graded ring of weakly holomorphic modular forms on  $\Gamma_1(N)$ .*
- (ii) *Given  $f \in \mathrm{MF}(\Gamma_1(N))^\times$  with trivial Nebentypus, if its Serre derivative is zero, then the element  $\beta_N^{(p)}(f)$  is contained in the kernel of the logarithmic operation  $\ell_{2,p}: (E^0)^\times \rightarrow E^0$ .*

The discrepancy between the aforementioned two sets of elements—Eisenstein series as opposed to the discriminant—results from the different domains of the logarithms. The former is the group of units in the zeroth cohomology of an even-dimensional sphere, while the latter is with respect to the infinite-dimensional complex projective space  $\mathbb{CP}^\infty$ , that is, the logarithms are defined on  $E^0(S^{2k})^\times$  and  $E^0(\mathbb{CP}^\infty)^\times$  respectively.

The finiteness of  $E^0(S^{2k})$  as a module over  $E^0$  leads to a simple formula for the logarithm. Specifically, the logarithm can be written as a combination of Hecke operators acting on  $\log(1+f)$ , where  $f$  is a generator of the truncated polynomial ring  $E^0(S^{2k})$ . Note that the formal power series expansion of  $\log(1+f)$  simply equals  $f$ , because  $f^2 = 0$  (see [Ando-Hopkins-Rezk2010, Proposition 4.8 and Example 4.9]).

In the latter case of  $E^0(\mathbb{CP}^\infty)$ , we calculate instead with  $\log(g)$  for units  $g$  in  $E^0$ . Through a generator  $u$  of the formal power series ring  $E^0(\mathbb{CP}^\infty)$ , certain  $g$  can be represented by meromorphic modular forms (cf. Proposition 2.9). Without nilpotency of  $g$  as in the case studied by Ando, Hopkins, and Rezk, a different set of tools from number theory is applied. In particular, it came as a surprise to the author that the formula of Rezk for the logarithms, which arose from a purely homotopy-theoretic construction, resembled the *logarithms of ratios of Siegel functions* studied by Katz in [Katz1976, Section 10.1] (see Remark 4.7). Indeed, Katz’s approach to those logarithms inspired a final step in our proof of Theorem 4.13.

Section 5 provides a “feedback” to number theory. As mentioned above, various types of modular forms have been associated with elliptic cohomology theories. In our calculation of the logarithms on  $E$ -cohomology, the occurrence of  $\log(g)$  indicates that Morava  $E$ -theories at height 2 witness a larger class of functions on elliptic curves than modular forms. In fact, for meromorphic modular forms  $g$  in a  $q$ -expansion,  $\log(g)$  are *logarithmic  $q$ -series* studied by Knopp and Mason [Knopp-Mason2011]. A paradigm is  $\log(\Delta)$ , which we discuss in Example 5.1 and Remark 5.2. Motivated by the logarithmic operations from homotopy theory, we sketch how the domain of Hecke operators may extend to include certain logarithmic  $q$ -series. This is Definition 5.9, from which the statement for the case  $f = \Delta$  in Theorem 4.13 falls out.

This interplay via Hecke operators between homotopy theory and number theory is bookended by the foundational material in Sections 2–3 and further discussions in Section 6 on Hecke operators as additive power operations.

Crucial to our understanding of the action of Hecke operators are the explicit computations for power operations. Building on the prime-2 case in [Rezk2008] and the prime-3 case in [Zhu2014], we present in Section 3.1 a general recipe for computing power operations on Morava  $E$ -theories at height 2 for all primes, and make some calculations at the prime 5 available as a working example throughout the paper (Examples 2.6, 3.5, 3.19, 4.8, 6.1, and 6.7). The explicit formulas show that the “topological” Hecke operators—constructed directly from power operations as in [Ando1995, Proposition 3.6.2]—do not agree with the classical Hecke operators acting on modular forms (Re-

mark 3.18). This explicitation also shows that a topological Hecke operator may not commute with all additive power operations on  $E$ -cohomology, which contradicts a claim in the literature known to experts.

**Theorem 1.2** (Theorem 6.8) *The center of the Dyer-Lashof algebra for a Morava  $E$ -theory  $E$  at height 2 differs from an algebra of Hecke operators acting on  $E$ -cohomology.*

Finally, Section 3.3 discusses an analog of Ando's criterion for  $H_\infty$ - $MU$ -orientations at chromatic level 2, which we hope is of independent interest.

## 1.1 Acknowledgements

I thank Andrew Baker, Paul Goerss, Tyler Lawson, Charles Rezk, and Joel Specter for helpful discussions.

I thank my friends Tzu-Yu Liu and Meng Yu, and their parents, for the hospitality I received during me writing this paper in their home in California.

## 1.2 Conventions

Throughout the paper,  $p$  will denote a prime, and  $N$  an integer prime to  $p$  with  $N > 3$ .

Let  $\mathrm{MF}(\Gamma_1(N))$  be the graded ring of modular forms of level  $\Gamma_1(N)$  over  $\mathbb{Z}[1/N]$  with no condition at the cusps (cf. [Katz1973, Section 1.2]). Complex-analytically, this corresponds to the graded complex vector space of weakly holomorphic modular forms on  $\Gamma_1(N)$  (cf. [Ono2004, Definition 1.12]). For simplicity, we will refer to these as *modular forms of level  $\Gamma_1(N)$* , or *modular forms* if the congruence subgroup  $\Gamma_1(N)$  is clear from the context.

By a *meromorphic modular form*, we mean a complex-valued modular form that is meromorphic both at the cusps and over the upper half of the complex plane [Ono2004, Definition 1.8].

We use a list of symbols.

$\mathbb{F}_{p^n}$	a field with $p^n$ elements
$\mathbb{Z}_p$	the ring of $p$ -adic integers
$\mathbb{Q}_p$	the field of $p$ -adic numbers
$\bar{k}$	a separable closure of a field $k$
$\mathbb{W}(k)$	the ring of $p$ -typical Witt vectors over a field $k$ of characteristic $p$
$\Sigma_m$	the symmetric group on $m$ letters
$R^\times$	the multiplicative group of invertible elements in a ring $R$
$R_I^\wedge$	the completion of a ring $R$ with respect to an ideal $I$
$R((x))$	the ring of formal Laurent series over a ring $R$ in a variable $x$ write $\sum_{n>-\infty} a_n x^n$ for a general element in this ring
$C/R$ or $C_R$	a scheme $C$ over a ring $R$
$[m]$	the multiplication-by- $m$ map on a group scheme
$C[m]$	the kernel of $[m]$ on a group scheme $C$
$\widehat{C}$	the formal completion of an elliptic curve $C$ at the identity
$q$	$e^{2\pi iz}$ with $z$ a complex variable
$\sigma_s(m)$	$\sum_{\substack{d m \\ d \geq 1}} d^s$ for any positive integer $m$ , with $s \in \mathbb{Z}$

### 1.3 Correction to two cited formulas

For the reader's convenience, we list the following typographical corrections.

- [Katz1973, (1.11.0.4)] The last line should start with  $\ell^{-k}$  instead of  $\ell^{-1}$ .
- [Milnor-Stasheff1974, Problem 16-A] The left-hand side of Girard's formula should read  $(-1)^n s_n / n$ . The summation on the right-hand side is over  $i_1 + 2i_2 + \cdots + ki_k = n$ .

## 2 From rings of modular forms to homotopy groups of Morava $E$ -theories

In this section, we explain how every modular form of level  $\Gamma_1(N)$  gives an element in the zeroth coefficient ring of a Morava  $E$ -theory at height 2 and prime  $p$ . The connection comes from finding explicit models for the  $E$ -theory.

## 2.1 Models for an $E$ -theory of height 2

Morava  $E$ -theory spectra can be viewed as topological realizations of Lubin-Tate rings. Specifically, given a formal group  $\mathbb{G}_0$  of height  $n < \infty$  over a perfect field  $k$  of characteristic  $p$ , the associated Morava  $E$ -theory (of height  $n$  at the prime  $p$ ) is a complex-oriented cohomology theory  $E$  whose formal group  $\mathrm{Spf} E^0 \mathbb{C}P^\infty$  is a universal deformation of  $\mathbb{G}_0$  over the Lubin-Tate ring

$$\mathbb{W}(k)[[u_1, \dots, u_{n-1}]] \cong E^0$$

(see [Lubin-Tate1966, Section 3] and [Goerss-Hopkins2004, Section 7]). Such a universal deformation is unique up to unique isomorphism. For height  $n = 2$ , via the Serre-Tate theorem [Lubin-Serre-Tate1964] (cf. [Katz-Mazur1985, Theorem 2.9.1]), this universal deformation of a formal group can be obtained from a universal deformation of a supersingular elliptic curve. We construct the latter as the universal family of elliptic curves equipped with a level- $\Gamma_1(N)$  structure.

Write  $\mathcal{P}_N$  for the representable moduli problem of smooth elliptic curves over  $\mathbb{Z}[1/N]$  with a choice of a point  $P_0$  of exact order  $N$  and a nonvanishing one-form  $\omega$ . The following two universal families represent  $\mathcal{P}_N$  for  $N = 4$  and  $N = 5$  respectively, and they suffice to model  $E$ -theories of height 2 at all primes.

**Example 2.1** ([Zhu2014, Proposition 2.1]) The moduli problem  $\mathcal{P}_4$  is represented by the curve

$$\mathcal{C}_4: y^2 + Axy + ABy = x^3 + Bx^2$$

over the graded ring

$$S_4 := \mathbb{Z}[1/4][A, B, \Delta^{-1}]$$

where  $|A| = 1$ ,  $|B| = 2$ , and  $\Delta = A^2 B^4 (A^2 - 16B)$ . The chosen point is  $P_0 = (0, 0)$  and the chosen one-form is  $\omega = du$  with  $u = x/y$ .

**Example 2.2** ([Behrens-Ormsby2015, Corollary 1.1.10]) The moduli problem  $\mathcal{P}_5$  is represented by

$$\mathcal{C}_5: y^2 + Axy + B^2(A - B)y = x^3 + B(A - B)x^2$$

over the graded ring

$$S_5 := \mathbb{Z}[1/5][A, B, \Delta^{-1}]$$

where  $|A| = |B| = 1$  and  $\Delta = B^5(B - A)^5(A^2 + 9AB - 11B^2)$ . Again the chosen point is  $P_0 = (0, 0)$  and the chosen one-form is  $\omega = du$  with  $u = x/y$ . Moreover, writing  $\zeta := e^{2\pi i/5}$ , we have  $\Delta = B^5(B - A)^5(A - (5\zeta^4 + 5\zeta - 2)B)(A + (5\zeta^4 + 5\zeta + 7)B)$ .

**Remark 2.3** The moduli problem  $\mathcal{P}_3$  is also representable [Mahowald-Rezk2009, Proposition 3.2], but  $[\Gamma_1(3)]$  is not (cf. [Katz-Mazur1985, Corollary 2.7.4]). The beginning of [Behrens-Ormsby2015, Section 1] explains the relationship between these two moduli problems. For the prime 2, compare [Behrens-Ormsby2015, Corollary 1.1.11] and [Lawson-Naumann2014, Section 3.1] (see also Proposition 2.5 below).

From each of the above examples, restricting  $\mathcal{C}_N$  over a closed point in the supersingular locus at  $p$ , we get a supersingular elliptic curve  $C_0$  over  $\overline{\mathbb{F}}_p$ . By the Serre-Tate theorem, the formal completion  $\widehat{\mathcal{C}}_N$  of  $\mathcal{C}_N$  at the identity then gives a universal deformation of the formal group  $\widehat{C}_0$ . Here  $\widehat{C}_0/\overline{\mathbb{F}}_p$  is of height 2, and it models  $\mathbb{G}_0/k$  for the  $E$ -theory we begin with.

**Remark 2.4** For a fixed  $E$ -theory, the various models each involve a choice of  $N$  for the  $\mathcal{P}_N$ -structures, and a choice of a supersingular elliptic curve over  $\overline{\mathbb{F}}_p$  equipped with a  $\mathcal{P}_N$ -structure. Over a separably closed field of characteristic  $p$ , any two formal group laws of the same height are isomorphic [Lazard1955, Théorème IV]. In view of this and the universal property in the Lubin-Tate theorem [Lubin-Tate1966, Theorem 3.1], we see that up to isomorphism and extension of scalars, the  $E$ -theory is independent of these choices.

Topologically, there is a  $K(2)$ -localization that corresponds to the above completion along the supersingular locus at  $p$ , as we now describe.

The graded rings  $S_N$  in Examples 2.1 and 2.2 can be identified as  $\mathrm{MF}(\Gamma_1(N))$ . Their topological realizations are the periodic spectra  $\mathrm{TMF}(\Gamma_1(N))$  of topological modular forms of level  $\Gamma_1(N)$  (see [Mahowald-Rezk2009, Section 2] and cf. [Hill-Lawson2016]). Following the convention that elements in algebraic degree  $k$  lie in topological degree  $2k$ , we have

$$\pi_* \mathrm{TMF}(\Gamma_1(4)) \cong \mathbb{Z}[1/4][A, B, \Delta^{-1}]$$

with  $|A| = 2$ ,  $|B| = 4$ , and  $\Delta = A^2 B^4 (A^2 - 16B)$ , and

$$\pi_* \mathrm{TMF}(\Gamma_1(5)) \cong \mathbb{Z}[1/5][A, B, \Delta^{-1}]$$

with  $|A| = |B| = 2$  and  $\Delta = B^5 (B - A)^5 (A^2 + 9AB - 11B^2)$ .

**Proposition 2.5** (cf. [Behrens-Ormsby-Stapleton-Stojanoska2015, Section 3.5] and [Ormsby2012, Corollary on page 20]) *Let  $K(2)$  be the Morava  $K$ -theory spectrum at height 2 and prime  $p$ , with  $\pi_* K(2) \cong \overline{\mathbb{F}}_p[v_2^{\pm 1}]$ , where  $|v_2| = 2(p^2 - 1)$ . Let  $N > 3$  be an integer prime to  $p$ . Denote by  $L_{K(2)}\mathrm{TMF}(\Gamma_1(N))$  the Bousfield localization of*



$\mathrm{TMF}(\Gamma_1(N))$  with respect to  $K(2)$ . Given a Morava  $E$ -theory spectrum  $E$  of height 2 at the prime  $p$ , there is a noncanonical isomorphism

$$L_{K(2)}\mathrm{TMF}(\Gamma_1(N)) \cong \underbrace{E \times \cdots \times E}_{m \text{ copies}}$$

of  $E_\infty$ -ring spectra, where  $m$  is the number of isomorphism classes of supersingular elliptic curves over  $\overline{\mathbb{F}}_p$  equipped with a level- $\Gamma_1(N)$  structure.

**Proof** Let  $C_0$  be a supersingular elliptic curve over  $\overline{\mathbb{F}}_p$  equipped with a level  $\Gamma_1(N)$ -structure. Its formal group  $\widehat{C}_0/\overline{\mathbb{F}}_p$  gives a model for the  $E$ -theory. By the Goerss-Hopkins-Miller theorem [Goerss-Hopkins2004, Corollary 7.6], the spectrum  $E$  admits an action of the automorphism group  $\mathrm{Aut}(C_0/\overline{\mathbb{F}}_p)$ .

Let  $G$  be the subgroup of  $\mathrm{Aut}(C_0/\overline{\mathbb{F}}_p)$  consisting of automorphisms that preserve the  $\Gamma_1(N)$ -structure on  $C_0$ . In view of Remark 2.4, we then obtain  $L_{K(2)}\mathrm{TMF}(\Gamma_1(N))$  as taking homotopy fixed points for the action of  $G$  on  $E$ , one such copy for each closed point in the supersingular locus at  $p$ . By [Katz-Mazur1985, Corollary 2.7.4], the moduli problem  $[\Gamma_1(N)]$  is rigid when  $N > 3$ , and thus  $G$  is trivial. Again by the Goerss-Hopkins-Miller theorem, we get the stated isomorphism as one between  $E_\infty$ -ring spectra.  $\square$

## 2.2 Homotopy groups of an $E$ -theory at height 2

Thanks to the explicit models for  $E$ -theories, the global-to-local relationship between  $\mathrm{TMF}(\Gamma_1(N))$  and  $E$  in Proposition 2.5 can be spelled out on homotopy groups. The next example illustrates a passage from  $\pi_*\mathrm{TMF}(\Gamma_1(N))$  to

$$E_* \cong \mathbb{W}(\overline{\mathbb{F}}_p)[[u_1]][u^{\pm 1}]$$

where we extend the scalars to  $\mathbb{W}(\overline{\mathbb{F}}_p)$ . The *deformation* parameter  $u_1$  in degree 0 comes from a Hasse invariant. The 2-periodic unit  $u$  in degree  $-2$  corresponds to a local uniformizer at the identity of the universal elliptic curve  $\mathcal{C}_N$ .

**Example 2.6** Let  $p = 5$  and  $N = 4$ .

By [Silverman2009, V.4.1a], the Hasse invariant of  $\mathcal{C}_4$  at the prime 5 equals

$$A^4 - A^2B + B^2 = (A^2 + 2(1 + \eta)B)(A^2 + 2(1 - \eta)B)$$

in  $\overline{\mathbb{F}}_5[A, B]$  with  $\eta^2 = 2$ . Fix this  $\eta$ , and fix one of its lifts to  $\mathbb{W}(\overline{\mathbb{F}}_5)$ , also denoted by  $\eta$  as an abuse of notation. For a reason that will become clear in (3.6), we choose an integral lift of this Hasse invariant given by

$$H := A^4 - 16A^2B + 26B^2 = (A^2 + 2(-4 + \eta)B)(A^2 + 2(-4 - \eta)B) \in S_4[\eta]$$

with  $\eta^2 = 38/4$ . The supersingular locus of  $\mathcal{C}_4$  at 5 is then the closed subscheme of  $\text{Proj } S_4$  cut out by the ideal  $(5, H)$ . Over  $\overline{\mathbb{F}}_5$ , it consists of two closed points in view of the factorization above. Setting  $A = \eta$  and  $B = 1 - \eta$ , we get a supersingular curve  $C_0$  over  $\mathbb{F}_{25}$ .


Since  $\Delta = A^2B^4(A^2 - 16B)$  gets inverted in  $S_4$ , the scheme  $\text{Proj } S_4$  is affine, and is contained in the affine open chart

$$\text{Proj } \mathbb{Z}[1/4][A, B][A^{-1}]$$

for the weighted projective space  $\text{Proj } \mathbb{Z}[1/4][A, B]$ . There is another affine chart

$$\text{Proj } \mathbb{Z}[1/4][A, B][u]$$

with  $u^2 = B^{-1}$  that is étale over  $\text{Proj } \mathbb{Z}[1/4][A, B]$ . The supersingular points are contained in both charts, as illustrated in the following diagram.



We now pass to the homotopy groups of the corresponding  $E$ -theory spectrum  $E$  by a procedure of *dehomogenization*. Define elements

$$\begin{aligned} a &:= uA \\ u_1 &:= u^2(A^2 + 2(-4 + \eta)B) = a^2 - 8 + 2\eta \\ h &:= u^4H = a^4 - 16a^2 + 26 = (a^2 - 8 + 2\eta)(a^2 - 8 - 2\eta) \\ \delta &:= u^{12}\Delta = h - 26 \end{aligned}$$

Following the convention that elements in algebraic degree  $k$  lie in topological degree  $2k$ , we then have

$$E_* \cong \mathbb{W}(\overline{\mathbb{F}}_5)[[u_1]][u^{\pm 1}]$$

with  $|u_1| = 0$  and  $|u| = -2$ . In particular, by Hensel's lemma, both  $a$  and  $\delta$  are contained in  $(E_0)^\times$ . Moreover,  $u$  corresponds to a local uniformizer at the identity of  $\mathcal{C}_4$  via a chosen isomorphism  $\mathrm{Spf} E^0 \mathbb{C}\mathbb{P}^\infty \cong \widehat{\mathcal{C}}_4$  of formal groups over

$$\mathrm{Spf} E^0 \cong \mathrm{Spf} \left( S_4[\eta]_{(5, A^2+2(-4+\eta)B)}^\wedge \otimes_{\mathbb{W}(\mathbb{F}_5)} \mathbb{W}(\overline{\mathbb{F}}_5) \right)$$

(cf. [Ando-Hopkins-Strickland2001, Definition 1.2]).

**Remark 2.7** As an abuse of notation, in Examples 2.1 and 2.2 we have written the local uniformizer

$$u = \frac{x}{y}$$

where  $x$  and  $y$  are the affine coordinates in the Weierstrass equation for  $\mathcal{C}_N$ . The resulting algebraic degree  $-1$  of  $u$  matches the topological degree  $-2$  of  $u$  in  $E_*$ .

To summarize, we arrive at the following.

**Definition 2.8** Given a Morava  $E$ -theory  $E$  of height 2 at the prime  $p$ , a  $\mathcal{P}_N$ -model for  $E$  consists of the data:

- a supersingular elliptic curve  $C_0/\overline{\mathbb{F}}_p$
- a universal deformation  $\mathcal{C}_N/S_N$  of  $C_0$
- a coordinate<sup>1</sup>  $u$  on  $\widehat{\mathcal{C}}_N$ , that is, a local uniformizer at the identity of  $\mathcal{C}_N$
- upon extension of scalars to  $\mathbb{W}(\overline{\mathbb{F}}_p)$ , an isomorphism

$$\mathrm{Spf} (S_N)_{I_0}^\wedge \cong \mathrm{Spf} E^0$$

where  $I_0$  corresponds to the special fiber  $C_0/\overline{\mathbb{F}}_p$

- an isomorphism

$$\widehat{\mathcal{C}}_N \cong \mathrm{Spf} E^0 \mathbb{C}\mathbb{P}^\infty$$

of formal groups under which  $u$  corresponds to a unit  $u \in \pi_{-2}E$ , that is, a strict isomorphism of formal group laws

Recall that elements in  $\mathrm{MF}(\Gamma_1(N))$  are functions  $f$  on elliptic curves  $C/R$  equipped with a  $\mathcal{P}_N$ -structure  $(P_0, \omega)$ . Each value  $f(C/R, P_0, \omega) \in R$  depends only on the  $R$ -isomorphism class of the triple  $(C/R, P_0, \omega)$ , and is subject to a modular transformation property that encodes the weight of  $f$ ; moreover, its formation commutes with arbitrary base change (see, e.g., [Katz1973, Section 1.2]).

---

<sup>1</sup>Cf. [Ando2000, Definition 1.4].

**Proposition 2.9** *Let  $E$  be a Morava  $E$ -theory of height 2 at the prime  $p$ . There is a noncanonical ring homomorphism*

$$\beta_N^{(p)}: \mathrm{MF}(\Gamma_1(N)) \rightarrow E^0$$

**Proof** Choose a  $\mathcal{P}_N$ -model for the  $E$ -theory. As in Example 2.6, via dehomogenization a modular form  $f$  of weight  $k$  maps to  $u^k \cdot f(\mathcal{C}_N, P_0, du) \in E^0$ .  $\square$

We will drop the indices in “ $\beta_N^{(p)}$ ” when there is no ambiguity.

### 3 From classical to topological Hecke operators

Morava  $E$ -theory spectra are  $E_\infty$ -ring spectra [Goerss-Hopkins2004, Corollary 7.6], and thus they come equipped with power operations. The work of Ando, Hopkins, and Strickland [Ando-Hopkins-Strickland2004] (cf. [Rezk2009, Theorem B]) gives a correspondence between power operations on an  $E$ -theory and deformations of Frobenius isogenies of formal groups.

At height 2, the Serre-Tate theorem [Lubin-Serre-Tate1964] (cf. [Katz-Mazur1985, Theorem 2.9.1]) gives a second correspondence between isogenies of formal groups and isogenies of elliptic curves, in terms of their deformation theory.

Via these two bridges, the classical action of Hecke operators on modular forms then corresponds to an action of “topological” Hecke operators on  $E$ -cohomology [Rezk2006b, Section 14]. The purpose of this section is to give an explicit comparison between these two actions. In particular, we will explain why the ring homomorphism  $\beta$  in Proposition 2.9 *fails* to become a map of modules over a Hecke algebra.

#### 3.1 Isogenies and power operations

Let  $\mathcal{C}_N$  over  $S_N$  be the universal curve in Section 2.1 for the moduli problem  $\mathcal{P}_N$ . Denote by  $\mathcal{G}_N^{(p)}$  the universal example of a degree- $p$  subgroup of  $\mathcal{C}_N$ . It is defined over an extension ring  $S_N^{(p)}$  that is free of rank  $p + 1$  as an  $S_N$ -module [Katz-Mazur1985, Theorem 6.6.1]. Explicitly,

$$S_N^{(p)} \cong S_N[\kappa]/(W(\kappa))$$

where  $\kappa$  is a generator that satisfies  $W(\kappa) = 0$  for a monic polynomial  $W$  of degree  $p + 1$ . The roots  $\kappa_0, \kappa_1, \dots, \kappa_p$  of  $W$  each correspond to a degree- $p$  subgroup of  $\mathcal{C}_N$ . In particular, let  $\kappa_0$  correspond to the subgroup whose formal completion over an ordinary point is the unique degree- $p$  subgroup of the formal group  $\widehat{\mathcal{C}}_N$ .

We write

$$\Psi_N^{(p)}: \mathcal{C}_N \rightarrow \mathcal{C}_N/\mathcal{G}_N^{(p)}$$

for the universal degree- $p$  isogeny over  $S_N^{(p)}$ , and write  $\mathcal{C}_N^{(p)}$  for the quotient curve  $\mathcal{C}_N/\mathcal{G}_N^{(p)}$ . Following Lubin [Lubin1967, proof of Theorem 1.4], we construct  $\Psi_N^{(p)}$  as a *deformation of Frobenius*: over any closed point in the supersingular locus at  $p$ ,  $\Psi_N^{(p)}$  restricts as the  $p$ -power Frobenius isogeny on the corresponding supersingular curve.

**Construction 3.1** (cf. [Lubin1967, proof of Theorem 1.4] and [Katz-Mazur1985, Section 7.7]) Let  $P$  be any point on  $\mathcal{C}_N$ . Let  $u \in S_N$  be a local uniformizer at the identity  $O$  of  $\mathcal{C}_N$ , so that  $|u| = -1$  tautologically.

- (i) Define  $\Psi_N^{(p)}: \mathcal{C}_N \rightarrow \mathcal{C}_N^{(p)}$  by the formula

$$u(\Psi_N^{(p)}(P)) := \prod_{Q \in \mathcal{G}_N^{(p)}} u(P - Q)$$

where the  $u$  on the left-hand side, by an abuse of notation, denotes the local coordinate on  $\mathcal{C}_N^{(p)}$  induced by  $\Psi_N^{(p)}$  as given on the right-hand side (cf. [Ando2000, Section 4.3]).

- (ii) Define  $\kappa \in S_N^{(p)}$  in degree  $-p + 1$  as

$$\kappa := \prod_{Q \in \mathcal{G}_N^{(p)} \setminus \{O\}} u(Q)$$

We verify that the isogeny  $\Psi_N^{(p)}$  has kernel precisely the subgroup  $\mathcal{G}_N^{(p)}$ . Moreover, it is a deformation of Frobenius, since at a supersingular point the  $p$ -divisible group is formal so that  $Q = O$  for all  $Q \in \mathcal{G}_N^{(p)}$ .

**Remark 3.2** The element  $\kappa$  in Construction 3.1 (ii) gives a *norm* parameter for the moduli problem  $[\Gamma_0(p)]$  as an “open arithmetic surface,” the other parameter being a deformation parameter (see [Katz-Mazur1985, Section 7.7]). As  $u$  is a local coordinate near the identity  $O$ , we note that by the above construction the cotangent map  $(\Psi_N^{(p)})^*$  at  $O$  sends  $du$  to  $\kappa du$ .

**Remark 3.3** Both  $\Psi_N^{(p)}$  and  $\kappa$  as defined above depend on a choice of  $u$ .

Via completion at a supersingular point, we see in Section 2.2 that the ring  $S_N \cong \text{MF}(\Gamma_1(N))$  representing  $\mathcal{P}_N$  is locally realized in homotopy theory as  $E^0$ . Here, given the ring  $S_N^{(p)}$  representing the simultaneous moduli problem  $(\mathcal{P}_N, [\Gamma_0(p)])$ , Strickland's theorem [Strickland1998, Theorem 1.1] identifies the completion of  $S_N^{(p)}$  at the supersingular point as  $E^0(B\Sigma_p)/I$ , where  $I$  is a transfer ideal. In particular,  $E^0(B\Sigma_p)/I$  is free over  $E^0$  of rank  $p+1$ , isomorphic to  $E^0[\alpha]/(w(\alpha))$  for a monic polynomial  $w$  of degree  $p+1$ .

The universal degree- $p$  isogeny  $\Psi_N^{(p)}: \mathcal{C}_N \rightarrow \mathcal{C}_N^{(p)}$  is constructed above as a deformation of Frobenius. By [Rezk2009, Theorem B] it then corresponds to an *(additive) total power operation*

$$(3.4) \quad \psi^p: A^0 \rightarrow A^0(B\Sigma_p)/J$$

where  $A$  is any  $K(2)$ -local commutative  $E$ -algebra, and  $J$  is the corresponding transfer ideal. Since  $E^0(B\Sigma_p)$  is free over  $E^0$  of finite rank [Strickland1998, Theorem 3.2], we have  $J \cong A^0 \otimes_{E^0} I$  and

$$A^0(B\Sigma_p)/J \cong (A^0 \otimes_{E^0} E^0(B\Sigma_p))/J \cong A^0 \otimes_{E^0} (E^0(B\Sigma_p)/I) \cong A^0[\alpha]/(w(\alpha))$$

Note that up to isomorphism,  $\psi^p$  is independent of the choice of a  $\mathcal{P}_N$ -model (see Remark 2.4). Moreover, taking quotient by the transfer ideal makes this operation additive and hence a local ring homomorphism.

**Example 3.5** We continue Example 2.6 with  $p = 5$  and  $N = 4$ .

Let  $u = x/y$  as in Example 2.1, and let  $v = 1/y$ . The universal isogeny  $\Psi_4^{(5)}: \mathcal{C}_4 \rightarrow \mathcal{C}_4^{(5)}$  in Construction 3.1 is defined over the graded ring  $S_4^{(5)} \cong S_4[\kappa]/(W(\kappa))$ , where  $|\kappa| = -4$  and

$$(3.6) \quad W(\kappa) = \kappa^6 - \frac{10}{B^2}\kappa^5 + \frac{35}{B^4}\kappa^4 - \frac{60}{B^6}\kappa^3 + \frac{55}{B^8}\kappa^2 - \frac{H}{B^{12}}\kappa + \frac{5}{B^{12}}$$

This polynomial is computed from the division polynomial  $\psi_5$  in [Silverman2009, Exercise 3.7] for the curve  $\mathcal{C}_4$ . We first deduce from  $\psi_5$  identities satisfied by the  $uv$ -coordinates of a universal example  $Q \in \mathcal{G}_4^{(5)} \setminus \{O\}$  as in [Zhu2014, proof of Proposition 2.2]. We then compute an explicit formula for

$$\kappa = u(Q) \cdot u(-Q) \cdot u(2Q) \cdot u(-2Q)$$

using methods analogous to [Silverman2009, III.2.3]. Finally we solve for a monic degree-6 equation satisfied by  $\kappa$  and obtain the polynomial  $W$ . (We do not compute an equation for  $\mathcal{C}_4^{(5)}$  as in [Zhu2014, Proposition 2.3].)

Passing to the corresponding power operation, we write

$$(3.7) \quad \alpha := u^{-4} \kappa_0$$

(see Remark 2.7) where  $\kappa_0$  corresponds to the subgroup of  $\mathcal{C}_4$  whose formal completion over an ordinary point is the unique degree-5 subgroup of  $\widehat{\mathcal{C}}_4$ . The total power operation  $\psi^5: E^0 \rightarrow E^0(B\Sigma_5)/I$  then lands in

$$E^0(B\Sigma_5)/I \cong \mathbb{W}(\overline{\mathbb{F}}_5)[[u_1, \alpha]]/(w(\alpha))$$

where  $w(\alpha) = \alpha^6 - 10\alpha^5 + 35\alpha^4 - 60\alpha^3 + 55\alpha^2 - h\alpha + 5$ . We can now compute the effect of  $\psi^5$  on  $h$  as follows.

Consider a second universal degree-5 isogeny

$$\widetilde{\Psi}_4^{(5)}: \mathcal{C}_4^{(5)} \rightarrow \mathcal{C}_4^{(5)}/\widetilde{\mathcal{G}}_4^{(5)}$$

where  $\widetilde{\mathcal{G}}_4^{(5)} = \mathcal{C}_4[5]/\mathcal{G}_4^{(5)}$ . It is defined the same way as in Construction 3.1, with a parameter  $\widetilde{\kappa} \in S_4^{(5)}$  in degree  $-20$ . Over  $S_4^{(5)}$ , the assignment

$$\left((\mathcal{C}_4, P_0, du), \mathcal{G}_4^{(5)}\right) \mapsto \left((\mathcal{C}_4^{(5)}, \Psi_4^{(5)}(P_0), du), \widetilde{\mathcal{G}}_4^{(5)}\right)$$

is an involution on the moduli problem  $(\mathcal{P}_4, [\Gamma_0(5)])$  (cf. [Katz-Mazur1985, 11.3.1]). In particular, by rigidity, we have an identity

$$\widetilde{\Psi}_4^{(5)} \circ \Psi_4^{(5)} = \tau \circ [5]$$

that lifts  $\text{Frob}^2 = [5]$  over the supersingular point, where  $\text{Frob}$  is the 5-power Frobenius isogeny, and  $\tau: \mathcal{C}_4/\mathcal{C}_4[5] \rightarrow \mathcal{C}_4^{(5)}/\widetilde{\mathcal{G}}_4^{(5)}$  is the *canonical* isomorphism. Thus in view of Remark 3.2 and  $\kappa_0 \kappa_1 \cdots \kappa_5 = 5/B^{12}$  from (3.6) we obtain a relation

$$\widetilde{\kappa} \cdot \kappa = \frac{5}{B^{12}}$$

in  $S_4^{(5)}$  (cf. [Zhu2014, proof of Corollary 3.2]).

Correspondingly, there is an involution

$$(h, \alpha) \mapsto (\widetilde{h}, \widetilde{\alpha})$$

on  $E^0(B\Sigma_5)/I$ , coming from the Atkin-Lehner involution of modular forms on  $\Gamma_0(5)$  (cf. [Atkin-Lehner1970, Lemmas 7–10]). In particular, the relation

$$(3.8) \quad \alpha^6 - 10\alpha^5 + 35\alpha^4 - 60\alpha^3 + 55\alpha^2 - h\alpha + 5 = 0$$

has an analog

$$(3.9) \quad \widetilde{\alpha}^6 - 10\widetilde{\alpha}^5 + 35\widetilde{\alpha}^4 - 60\widetilde{\alpha}^3 + 55\widetilde{\alpha}^2 - \widetilde{h}\widetilde{\alpha} + 5 = 0$$

and we have

$$(3.10) \quad \tilde{\alpha} \cdot \alpha = 5$$

Based on these, we compute that

$$\begin{aligned}
 \psi^5(h) &= \tilde{h} \\
 &= \tilde{\alpha}^5 - 10\tilde{\alpha}^4 + 35\tilde{\alpha}^3 - 60\tilde{\alpha}^2 + 55\tilde{\alpha} + \alpha && \text{by (3.9) and (3.10)} \\
 &= (-\alpha^5 + 10\alpha^4 - 35\alpha^3 + 60\alpha^2 - 55\alpha + h)^5 \\
 &\quad - 10(-\alpha^5 + 10\alpha^4 - 35\alpha^3 + 60\alpha^2 - 55\alpha + h)^4 \\
 &\quad + 35(-\alpha^5 + 10\alpha^4 - 35\alpha^3 + 60\alpha^2 - 55\alpha + h)^3 \\
 &\quad - 60(-\alpha^5 + 10\alpha^4 - 35\alpha^3 + 60\alpha^2 - 55\alpha + h)^2 \\
 &\quad + 55(-\alpha^5 + 10\alpha^4 - 35\alpha^3 + 60\alpha^2 - 55\alpha + h) \\
 &\quad + \alpha && \text{by (3.10) and (3.8)} \\
 (3.11) \quad &= h^5 - 10h^4 - 1065h^3 + 12690h^2 + 168930h \\
 &\quad - 1462250 + (-55h^4 + 850h^3 + 39575h^2 \\
 &\quad - 608700h - 1113524)\alpha + (60h^4 - 775h^3 \\
 &\quad - 45400h^2 + 593900h + 2008800)\alpha^2 + (-35h^4 \\
 &\quad + 400h^3 + 27125h^2 - 320900h - 1418300)\alpha^3 \\
 &\quad + (10h^4 - 105h^3 - 7850h^2 + 86975h \\
 &\quad + 445850)\alpha^4 + (-h^4 + 10h^3 + 790h^2 - 8440h \\
 &\quad - 46680)\alpha^5 && \text{by (3.8)}
 \end{aligned}$$

We also have  $\psi^5(c) = Fc$  for  $c \in \mathbb{W}(\overline{\mathbb{F}}_5)$ , where  $F$  is the Frobenius automorphism. We would like to compute  $\psi^5(u_1)$  as well. As  $\psi^5$  is a local ring homomorphism, a formula for  $\psi^5(u_1)$  determines  $\psi^5(x)$  for all  $x \in E^0 \cong \mathbb{W}(\overline{\mathbb{F}}_5)[[u_1]]$ . We will address this using a modified presentation for  $E^0(B\Sigma_5)/I$  in a subsequent paper.

The previous example illustrates a general recipe for computing power operations on a Morava  $E$ -theory at height 2 and prime  $p$ , with a model based on the moduli problem  $(\mathcal{P}_N, [\Gamma_0(p)])$ . Crucial in this computation is an explicit expression for

$$(3.12) \quad w(\alpha) = \alpha^{p+1} + w_p\alpha^p + \cdots + w_1\alpha + w_0 \in E^0[\alpha]$$

Computing  $w(\alpha)$  appears to be essentially computing a “modular equation” (see, e.g., [Milne2012, II.6] and [Zureick-Brown2010], and compare the *canonical modular polynomials* in Magma’s Modular Polynomial Databases).



### 3.2 Hecke operators

We first describe the classical action of Hecke operators on modular forms in terms of isogenies between elliptic curves. The  $p$ 'th Hecke operator  $T_p$  that acts on  $\mathrm{MF}(\Gamma_1(N))$  can be built from universal isogenies as follows.

**Construction 3.13** (cf. [Katz1973, (1.11.0.2)]) Let the notation be as in Section 3.1, with the subscripts “ $N$ ” suppressed. Given any  $f \in \mathrm{MF}(\Gamma_1(N))$  of weight  $k \geq 1$ ,  $T_p f \in \mathrm{MF}(\Gamma_1(N))$  is a modular form such that

$$(3.14) \quad T_p f(\mathcal{C}_S, P_0, du) := \frac{1}{p} \sum_{i=0}^p \kappa_i^k \cdot f(\mathcal{C}_{S^{(p)}}/\mathcal{G}_i^{(p)}, \Psi_i^{(p)}(P_0), du)$$

where each  $\mathcal{G}_i^{(p)}$  denotes a degree- $p$  subgroup of  $\mathcal{C}$  over  $S^{(p)}$ , and  $\Psi_i^{(p)}$  is the quotient map with kernel  $\mathcal{G}_i^{(p)}$  as in Construction 3.1.

**Remark 3.15** The terms  $\kappa_i^k$  appear in the above formula so that  $T_p$  is independent of the choice of a basis for the cotangent space. Explicitly, we have

$$(\Psi_i^{(p)})^* du = \kappa_i \cdot du \quad \text{and} \quad (\Psi_i^{(p)})^* (\check{\Psi}_i^{(p)})^* du = p \cdot du$$

where each  $\check{\Psi}_i^{(p)}: \mathcal{C}/\mathcal{G}_i^{(p)} \rightarrow \mathcal{C}$  is a dual isogeny, and  $(\check{\Psi}_i^{(p)})^* du$  is the choice in [Katz1973] for a nonvanishing one-form on a quotient curve (cf. Remark 3.2 and the discussion before (1.11.0.0) in [Katz1973, Section 1.11]). Thus we rewrite (3.14) as

$$\begin{aligned} & \frac{1}{p} \sum_{i=0}^p \kappa_i^k \cdot f(\mathcal{C}_{S^{(p)}}/\mathcal{G}_i^{(p)}, \Psi_i^{(p)}(P_0), du) \\ &= \frac{1}{p} \sum_{i=0}^p \kappa_i^k \cdot f(\mathcal{C}_{S^{(p)}}/\mathcal{G}_i^{(p)}, \Psi_i^{(p)}(P_0), (\kappa_i/p)(\check{\Psi}_i^{(p)})^* du) \\ &= \frac{1}{p} \sum_{i=0}^p p^k \cdot f(\mathcal{C}_{S^{(p)}}/\mathcal{G}_i^{(p)}, \Psi_i^{(p)}(P_0), (\check{\Psi}_i^{(p)})^* du) \end{aligned}$$

and this agrees with [Katz1973, (1.11.0.2)].

**Construction 3.16** (cf. [Rezk2006b, 1.12]) There is a *topological* Hecke operator  $t_p: E^0 \rightarrow p^{-1}E^0$  defined by

$$(3.17) \quad t_p(x) := \frac{1}{p} \sum_{i=0}^p \psi_i^p(x)$$

where  $\psi_i^p$  denotes the power operation  $\psi^p: E^0 \rightarrow E^0(B\Sigma_p)/I \cong E^0[\alpha]/(w(\alpha))$  with the parameter  $\alpha$  replaced by  $\alpha_i = u^{-p+1}\kappa_i$  (cf. (3.7)).

Since the parameters  $\alpha_i$  are the roots of  $w(\alpha) \in E^0[\alpha]$ ,  $t_p$  indeed lands in  $p^{-1}E^0$ .

**Remark 3.18** Given the ring homomorphism  $\beta$  in Proposition 2.9, the topological Hecke operator above does not coincide with the classical one on modular forms. Comparing (3.17) to (3.14), note that there are no terms  $\alpha_i^k$  corresponding to  $\kappa_i^k$ .<sup>2</sup> This is related to the fact that  $\beta$  is not injective: if we include  $\alpha_i^k$  in the definition, for each  $x \in E^0$  we need to determine a unique value of its “weight”  $k$  so that  $t_p$  is well-defined.

Note that the map  $\beta$  is not surjective either, so an element in  $E^0$  may not come from any modular form. On the other hand, the total power operation  $\psi^p$  (and hence  $t_p$ ) is defined on the entire  $E^0$ .

**Example 3.19** Again, let  $p = 5$  and  $N = 4$ . Recall from Example 2.6 that we have

$$\beta(\Delta) = \delta = h - 26$$

In view of (3.8), we then compute from (3.11) that

$$\begin{aligned} t_5(\delta) &= \frac{1}{5} \sum_{i=0}^5 \psi_i^5(\delta) \\ &= \frac{1}{5} \sum_{i=0}^5 (\psi_i^5(h) - 26) \\ &= \frac{1}{5} (h^5 - 10h^4 - 1340h^3 + 18440h^2 + 267430h - 3178396) \\ &= \frac{1}{5} (h^4 + 16h^3 - 924h^2 - 5584h + 122246) \cdot \delta \end{aligned}$$

Given that the modular form  $\Delta$  is of weight 12, we define and compute

$$\begin{aligned} \check{t}_5(\delta) &:= \frac{1}{5} \sum_{i=0}^5 \alpha_i^{12} \cdot \psi_i^5(\delta) \\ &= 4830h - 125580 \\ &= \tau(5) \cdot \delta \end{aligned}$$

where  $\tau: \mathbb{N} \rightarrow \mathbb{Z}$  is the Ramanujan tau-function, with  $\tau(5) = 4830$ . This recovers the action of  $T_5$  on  $\Delta$ .

---

<sup>2</sup>Ando originally defines power operations that include such factors for a version of  $\mathbb{Z}$ -graded elliptic cohomology theory due to Landweber, Ravenel, and Stong, but he drops the factors when defining Hecke operators (cf. [Ando1992, (6.3.16) and Theorem 6.5.2]).

More generally, in the theory of automorphic forms on  $\Gamma_1(N) \subset \mathrm{SL}_n(\mathbb{Z})$ , the above Hecke operator  $T_p$  can be renamed as  $T_{1,p}$ . It belongs to a family of operators  $T_{i,p}$ ,  $1 \leq i \leq n$  which generate the  $p$ -primary Hecke algebra (see, e.g., [Shimura1994, Theorems 3.20 and 3.35]).

For the case  $n = 2$ , the other Hecke operator  $T_{2,p}$  arises from the isogeny of multiplication by  $p$ , whose kernel is the degree- $p^2$  subgroup consisting of the  $p$ -torsion points. Explicitly (again with the subscripts “ $N$ ” suppressed), given any  $f \in \mathrm{MF}(\Gamma_1(N))$  of weight  $k \geq 2$ , there exists  $\lambda \in S^\times$  of degree  $-p^2 + 1$  such that

$$\begin{aligned}
 T_{2,p} f(\mathcal{C}, P_0, du) &:= \frac{1}{p^2} \left( (p\lambda)^k \cdot f(\mathcal{C}/\mathcal{C}[p], [p]P_0, du) \right) \\
 (3.20) \quad &= p^{k-2} \cdot f(\mathcal{C}/\mathcal{C}[p], [p]P_0, \lambda^{-1} du) \\
 &= p^{k-2} \cdot f(\mathcal{C}, [p]P_0, du) \quad \text{via } \mathcal{C}/\mathcal{C}[p] \xrightarrow{\sim} \mathcal{C}
 \end{aligned}$$

For example, when  $N = 4$  with  $p = 3$  or  $p = 5$ , we have  $\lambda = B^{(1-p^2)/2}$  (cf. Example 2.1). Via the canonical  $S^{(p)}$ -isomorphism

$$(3.21) \quad \mathcal{C}/\mathcal{C}[p] \cong \frac{\mathcal{C}/\mathcal{G}^{(p)}}{\mathcal{C}[p]/\mathcal{G}^{(p)}} =: \frac{\mathcal{C}^{(p)}}{\widetilde{\mathcal{G}}^{(p)}}$$

the quotient curve  $\mathcal{C}/\mathcal{C}[p]$  can be identified with the target in the composite

$$\mathcal{C} \xrightarrow{\Psi^{(p)}} \mathcal{C}^{(p)} \xrightarrow{\widetilde{\Psi}^{(p)}} \mathcal{C}^{(p)}/\widetilde{\mathcal{G}}^{(p)}$$

of deformations of Frobenius isogenies.

Correspondingly, given any  $K(2)$ -local commutative  $E$ -algebra  $A$ , there is a composite  $\phi$  of total power operations  $\psi^p \circ \psi^p: A^0 \rightarrow A^0$  (cf. [Zhu2014, (26)]). In view of (3.4), note that  $\phi$  lands in  $A^0$ , because up to the isomorphism (3.21) the composite  $\widetilde{\Psi}^{(p)} \circ \Psi^{(p)}$  is an endomorphism on  $\mathcal{C}$  over  $S$ . In particular, on  $\mathbb{W}(\mathbb{F}_{p^2})[[u_1]] \subset E^0$ , the operation  $\phi$  becomes the identity map,<sup>3</sup> which is a manifest of the Atkin-Lehner involution (cf. Example 3.5).

Taking  $A = E$ , we define a *topological* Hecke operator  $t_{2,p}: E^0 \rightarrow p^{-1}E^0$  by

$$(3.22) \quad t_{2,p}(x) := p^{-2}\phi(x)$$

(we will discuss the general case on  $A^0$  in Section 6). Rewrite  $t_{1,p} := t_p$ . The ring  $\mathbb{Z}[t_{1,p}, t_{2,p}]$  then acts on  $p^{-1}E^0$ . As we see in Remark 3.18, along the map  $\beta: \mathrm{MF}(\Gamma_1(N)) \rightarrow E^0$ , this action is not compatible with the action of the Hecke algebra  $\mathbb{Z}[T_{1,p}, T_{2,p}]$  on modular forms. However, in certain instances, the two actions

<sup>3</sup>This is also true on  $\mathbb{W}(\mathbb{F}_{p^2})[[u_1, \alpha]]/(w(\alpha)) \subset E^0(B\Sigma_p)/I$ . Again, cf. [Zhu2014, (26)].

do interact well. Section 4 will be about this connection. Below we prepare a technicality.

### 3.3 Ando's coordinate

In [Ando1995], Ando constructs power operations for a family of Landweber-exact cohomology theories  $E_n$ , using power operations in the complex cobordism  $MU$ . For a fixed prime  $p$ , each  $E_n$  is a Morava  $E$ -theory whose formal group is a universal deformation of the Honda formal group  $\Phi$  of height  $n$  over  $\mathbb{F}_p$ . This universal deformation is defined over the Lubin-Tate ring  $\mathbb{Z}_p[[u_1, \dots, u_{n-1}]]$ . In particular, for each  $\star$ -isomorphism class  $F$  of such a universal deformation, Ando constructs a unique coordinate  $x$  on  $F$  that lifts a particular coordinate on  $\Phi$  and satisfies

$$(3.23) \quad f_p^x(t) = [p]_{F_x}(t)$$

The left-hand side is the formal power series of a degree- $p^n$  *Lubin isogeny*, constructed with the coordinate  $x$ , for the subgroup of  $p$ -torsions on  $F$ . The right-hand side is the formal power series of multiplication by  $p$  under the group law  $F_x$ . See [Ando1995, Theorems 2.5.7 and 2.6.4]. Ando shows that such coordinates give rise to precisely those orientations for  $E_n$  which intertwine the power operations on  $MU$  and  $E_n$  [Ando1995, Theorem 4.1.1].

In our context of elliptic cohomology, let  $E$  be a Morava  $E$ -theory of height 2 at  $p$ . Choose a  $\mathcal{P}_N$ -model for  $E$  in the sense of Definition 2.8. Given the supersingular elliptic curve  $C_0$  over  $\overline{\mathbb{F}}_p$ , there exists a supersingular curve  $C'_0$  over  $\mathbb{F}_{p^2}$ , isomorphic to  $C_0$  over  $\overline{\mathbb{F}}_p$ , such that its  $p^2$ -power Frobenius endomorphism  $\text{Frob}^2$  satisfies

$$(3.24) \quad \text{Frob}^2 = \begin{cases} [-p] & \text{if } p = 2 \\ [p] & \text{if } p \neq 2 \end{cases}$$

[Poonen2010] (cf. [Baker et al. 2005, Lemma 3.21], [Zhu2014, Remark 3.3], and [Rezk2012, 3.8]). For a reason that will become clear in (3.30), replace  $C_0/\overline{\mathbb{F}}_p$  by  $C'_0/\mathbb{F}_{p^2}$ . Correspondingly, replace  $\mathcal{C}_N$  up to an isomorphism as needed above so that its restriction over a particular supersingular point at  $p$  is this  $C_0$ .

With notation as in (3.14), Construction 3.1 extends to give a degree- $p^2$  isogeny  $\Psi^{(p^2)}: \mathcal{C} \rightarrow \mathcal{C}/\mathcal{C}[p]$  such that

$$u(\Psi^{(p^2)}(P)) = \prod_{Q \in \mathcal{C}[p]} u(P - Q) = \prod_{i=0}^{p-1} u(\Psi_i^{(p)}(P))$$

(again, with an abuse of the notation  $u$ ). This Lubin isogeny is defined over  $S$ , as the parameters  $\kappa_i$  for  $\Psi_i^{(p)}$  are the roots of the polynomial  $W(\kappa) \in S[\kappa]$ . Over the supersingular point, we have  $\Psi^{(p^2)} = \text{Frob}^2$ . By rigidity, this lifts to

$$\sigma \circ \Psi^{(p^2)} = [p]$$

over  $S$  for some *noncanonical* isomorphism  $\sigma$  that depends on the choice of  $u$ . In particular, comparing the induced cotangent maps at the identity, we get

$$(3.25) \quad s \cdot \alpha_0 \alpha_1 \cdots \alpha_p = p$$

for some degree-0 unit  $s$  in  $S$ .

We next show that there exists a coordinate  $u_A$  on  $\widehat{C}$  for which  $s = 1$  in (3.25), hence analogous to Ando's coordinate characterized by (3.23). In fact, they are related via restriction to the cusps.

Over a punctured formal neighborhood of each cusp of the compactified moduli scheme for  $\mathcal{P}_N$ , the universal curve  $\mathcal{C}_N$  is isomorphic to the Tate curve  $\text{Tate}(q^N)$  with the level- $\Gamma_1(N)$  structure corresponding to that cusp. Over  $\mathbb{Z}[1/N][[q]]$ , the formal group of  $\text{Tate}(q^N)$  is canonically isomorphic to the multiplicative formal group  $\widehat{\mathbb{G}}_m$ . After base change to  $\mathbb{Z}_p$ , there is a unique coordinate  $u_A$  of Ando's on this  $\star$ -isomorphism class of universal deformations for  $\Phi$ , the Honda formal group of height 1 over  $\mathbb{F}_p$  (cf. [Ando1995, Example 2.7]). In particular, the Lubin isogeny  $\Psi_0^{(p)}$  on  $\text{Tate}(q^N)$  in this coordinate gives

$$(3.26) \quad \alpha_0 = p$$

for the induced cotangent map at the identity.

In view of Remark 3.15, we then have

$$(3.27) \quad \alpha_1 = \cdots = \alpha_p = (-1)^{p+1}$$

by construction of the degree- $p$  isogenies on  $\text{Tate}(q^N)$  in [Katz1973, Section 1.11] (see particularly the first new paragraph on page Ka-23). In fact, (3.24) lifts to guarantee that, when  $p$  is odd, the dual isogenies used by Katz in his construction are the Lubin isogenies as deformations of Frobenius. When  $p = 2$ , they differ by a sign.

In this particular coordinate  $u_A$ , (3.25) now becomes

$$(3.28) \quad \alpha_0 \alpha_1 \cdots \alpha_p = p$$

near the cusps. Thus, the modular unit  $s$  in (3.25) must be constant. It equals 1 by continuity. In other words, the above identity holds over a formal neighborhood of the supersingular point.

**Definition 3.29** Given a  $\mathcal{P}_N$ -model for an  $E$ -theory in Definition 2.8, we call it a *preferred* one if in addition we have the following.

- The supersingular curve  $C_0$  satisfies (3.24).
- With the chosen coordinate  $u$  on  $\widehat{C}_N$ , the identity (3.28) holds over a formal neighborhood of the supersingular point.

Given a preferred model, the constant term in (3.12) satisfies

$$(3.30) \quad w_0 = (-1)^{p+1} p = \widetilde{\alpha} \cdot \alpha$$

as a result of (3.28) and (3.24).<sup>4</sup> Our model for  $p = 5$  and  $N = 4$  in Examples 2.6 and 3.5 is preferred, but the one for  $p = 3$  and  $N = 4$  in [Zhu2014] is not. We will choose preferred models for  $E$ -theories at height 2 through the rest of this paper.

## 4 Kernels of logarithmic cohomology operations at height 2

In Section 3.2, we give an explication for an algebra of topological Hecke operators acting on  $E^0$ , where  $E$  is a Morava  $E$ -theory of height 2. Based on this, we now use a number-theoretic calculation to understand a topological construction. This is Rezk's construction of logarithmic cohomology operations via Bousfield-Kuhn functors [Rezk2006b]. We first recall a formula of Rezk for computing these operations, which leads to a connection with Hecke operators.

### 4.1 Connecting to Hecke operators

Given any  $E_\infty$ -ring spectrum  $R$ , let  $\mathrm{gl}_1 R$  be the spectrum of units of  $R$ . Rezk constructs a family of operations that naturally acts on  $\mathrm{gl}_1 R$  [Rezk2006b, Definition 3.6]. Specifically, given a positive integer  $n$  and a prime  $p$ , let  $L_{K(n)}$  denote localization with respect to the  $n$ 'th Morava  $K$ -theory at  $p$ , and let  $\Phi_n$  be the corresponding Bousfield-Kuhn functor from the category of based topological spaces to the category of spectra. Consider the composite

$$(4.1) \quad \mathrm{gl}_1 R \rightarrow L_{K(n)} \mathrm{gl}_1 R \simeq \Phi_n \Omega^\infty \mathrm{gl}_1 R \xrightarrow{\sim} \Phi_n \Omega^\infty R \simeq L_{K(n)} R$$

---

<sup>4</sup>The right equality depends only on the curve  $C_0$ , not on the choice of coordinate.

Note that  $\Omega^\infty \mathrm{gl}_1 R$  and  $\Omega^\infty R$  have weakly equivalent basepoint components, but the standard inclusion  $\Omega^\infty \mathrm{gl}_1 R \hookrightarrow \Omega^\infty R$  is not basepoint-preserving. The equivalence  $\Phi_n \Omega^\infty \mathrm{gl}_1 R \rightarrow \Phi_n \Omega^\infty R$  thus involves a “basepoint shift” (see [Rezk2006b, 3.4]).

Let  $E$  be a Morava  $E$ -theory of height  $n$  at the prime  $p$ . With  $R = E$ , applying  $\pi_0(-)$  to (4.1), we then obtain the logarithmic operation

$$\ell_{n,p}: (E^0)^\times \rightarrow E^0$$

which is a homomorphism from a multiplicative group to an additive group. More generally, let  $X$  be a space. Taking  $R$  to be the spectrum of functions from  $\Sigma_+^\infty X$  to  $E$ , we obtain the operation  $\ell_{n,p}$  that acts on  $E^0(X)^\times$ , naturally in both  $X$  and  $E$ .

Rezk gives a formula for this operation [Rezk2006b, Theorem 1.11]. In particular, for any  $x \in (E^0)^\times$ ,

$$(4.2) \quad \ell_{n,p}(x) = \frac{1}{p} \log(1 + pM(x))$$

where  $M: (E^0)^\times \rightarrow E^0$  is a cohomology operation that can be expressed in terms of power operations  $\psi_A$  associated to certain subgroups  $A$  of  $(\mathbb{Q}_p/\mathbb{Z}_p)^n$ . Explicitly,

$$1 + pM(x) = \prod_{j=0}^n \prod_{\substack{A \subset (\mathbb{Q}_p/\mathbb{Z}_p)^n[p] \\ \#A = p^j}} \psi_A(x)^{(-1)^j p^{j(j-1)(j-2)/2}}$$

Now let the  $E$ -theory be of height  $n = 2$ . In the presence of a model for  $E$ , the operations  $\psi_A$  above coincide with the power operations in Section 3. In particular,

$$(4.3) \quad \ell_{2,p}(x) = \frac{1}{p} \log \left( x^p \cdot \frac{1}{\psi_0^p(x) \cdots \psi_p^p(x)} \cdot \phi(x) \right)$$

Note that  $\ell_{2,p}(x) = 0$  for any  $x \in \mathbb{Z}_p \cap (E^0)^\times$ .

Suppose that  $x = \beta(f)$  for some modular form  $f$  of weight  $k$  (see Proposition 2.9). We next write (4.3) in terms of Hecke operators. We note the following.

- Recall from Definition 3.29 that the parameters  $\alpha_i$  in the operations  $\psi_i^p$  satisfy

$$\alpha_0 \cdots \alpha_p = p$$

upon we choose a preferred  $\mathcal{P}_N$ -model for the  $E$ -theory.

- Recall that  $\psi_i^p$ ,  $\phi$  and  $\beta$  are ring homomorphisms. Multiplications by the terms  $\kappa_i^k$  in (3.14) and by  $p^k$  in (3.20) can be viewed as ring homomorphisms

as well. More precisely, we allow the exponent  $k$  to vary, according to the weight of the modular form that immediately follows. We will denote these ring homomorphisms by  $\kappa_i^\bullet$  and  $p^\bullet$ . The topological analog of  $\kappa_i^\bullet$  on homotopy groups is  $\alpha_i^\bullet$ . In particular, given as a formal power series, the usual logarithm “log” commutes with these continuous ring homomorphisms.

In Section 3.2, we give a comparison between classical and topological Hecke operators, particularly as illustrated in Example 3.19. In view of this comparison and the two observations above, we can now rewrite (4.3) as follows (cf. [Rezk2006b, 1.12]). Given the element  $x = \beta(f)$ ,

$$\begin{aligned}
 \ell_{2,p}(x) &= \frac{1}{p} \log \frac{x^p \cdot p^\bullet \phi(x)}{\prod_{i=0}^p \alpha_i^\bullet \psi_i^p(x)} \\
 (4.4) \quad &= \left( 1 - \frac{1}{p} \sum_{i=0}^p \alpha_i^\bullet \psi_i^p + p \cdot \frac{1}{p^2} p^\bullet \phi \right) \log x \\
 &= \beta \left( (1 - T_{1,p} + p \cdot T_{2,p}) \log f \right)
 \end{aligned}$$

Following Rezk [Rezk2006b, 1.12], we write

$$(4.5) \quad F_X := 1 - T_{1,p} \cdot X + p T_{2,p} \cdot X^2 \in \mathbb{Z}[T_{1,p}, T_{2,p}][X]$$

In particular, with  $X = 1$ , (4.4) becomes

$$(4.6) \quad \ell_{2,p}(x) = \beta(F_1(\log f))$$

**Remark 4.7** Let  $x = \beta(f) \in (E^0)^\times$  for some unit  $f$  in  $\mathrm{MF}(\Gamma_1(N))$  of weight  $k$  (cf. [Kubert-Lang1981]). The formula (4.3) expresses  $p\ell_{2,p}(x)$  as the logarithm of a ratio: the numerator, via  $\beta$ , corresponds to a modular form of weight  $pk + p^2k$ , and the denominator corresponds to one having weight  $(p+1)pk = pk + p^2k$ . Thus  $p\ell_{2,p}(x)$  corresponds to a  $p$ -adic modular form of weight 0 in view of (4.2) (cf. [Katz1976, Section 10.1]). We note the similarity between the logarithm in [Katz1976, 10.2.7] and the one following Theorem 1.9 in [Rezk2006b].

**Example 4.8** We revisit the case  $p = 5$  with the preferred model for the  $E$ -theory given by the moduli problem  $\mathcal{P}_4$ . Consider  $\delta = \beta(\Delta) \in (E^0)^\times$ . As in Example 3.19,



we calculate from (3.11) and (3.8) that

$$\begin{aligned}\ell_{2,5}(\delta) &= \frac{1}{5} \log \left( \delta^5 \cdot \frac{1}{\psi_0^p(\delta) \cdots \psi_p^p(\delta)} \cdot \phi(\delta) \right) \\ &= \frac{1}{5} \log \left( \delta^5 \cdot \frac{1}{\delta^6} \cdot \delta \right) \\ &= \frac{1}{5} \log 1 \\ &= 0\end{aligned}$$

Compare this to the prime-2 case with a preferred  $\mathcal{P}_3$ -model from [Mahowald-Rezk2009, Proposition 3.2]. Mahowald and Rezk show that  $\mathcal{P}_3$  is represented by

$$y^2 + Axy + By = x^3$$

over  $\mathbb{Z}[1/3][A, B, \Delta^{-1}]$  with  $|A| = 1$ ,  $|B| = 3$ , and  $\Delta = B^3(A^3 - 27B)$ . In [Rezk2008, 2.8], Rezk computes that

$$\ell_{2,2}(\beta_3^{(2)}(\Delta)) = \frac{1}{2} \log(-1) = 0$$

We interpret the “log” here as the 2-adic logarithm (see, e.g., [Koblitz1984, §IV.1] and cf. [Katz1976, 10.2.16]). Again, the modular form  $\Delta$  produces an element in the kernel of a logarithmic operation.

Moreover, since  $\Delta = B^3(A^3 - 27B)$  in this case, we have

$$\beta_3^{(2)}(\Delta) = (a - 3)(a^2 + 3a + 9)$$

with  $E^0 \cong \mathbb{W}(\overline{\mathbb{F}}_2)[[a]]$  (cf. [Rezk2008, Section 4]). Using Rezk’s formula for the total power operation on  $E^0$ , we compute that

$$\ell_{2,2}(a - 3) = \frac{1}{2} \log(-1) = 0$$

$$\ell_{2,2}(a^2 + 3a + 9) = \frac{1}{2} \log 1 = 0$$

Thus, as  $\ell_{2,2}$  is a group homomorphism, the formal power series

$$\frac{1}{a - 3} = - \sum_{i=0}^{\infty} \left( \sum_{j=0}^{\infty} (-2)^j \right)^{i+1} a^i \in \mathbb{W}(\overline{\mathbb{F}}_2)[[a]]^\times$$

is also contained in the kernel of  $\ell_{2,2}$ .

The above turn out to be instances of a general vanishing result for the logarithms that we discuss next.

## 4.2 Detecting the kernels

In this section, via the formula (4.4) for a logarithmic operation in terms of Hecke operators, we detect a family of elements contained in the kernel of this operation. It includes those elements computed in Example 4.8.

We first review some preliminaries about differential structures on rings of modular forms (see, e.g., [Zagier2008, §5] and [Ono2004, Section 2.3]). In connection to Morava  $E$ -theories, we have been considering the  $p$ -local behavior of integral modular forms of level  $\Gamma_1(N)$ , with  $p$  not dividing  $N$  (which embed into the ring of  $p$ -adic modular forms). Thus for our purpose these modular forms can equivalently be viewed as defined over  $\mathbb{C}$ . Henceforth we will freely move between the algebraic and analytic perspectives.

Recall that there is a differential operator  $D$  that acts on meromorphic elliptic functions over  $\mathbb{C}$ . Specifically, any meromorphic modular form  $f$  has a  $q$ -expansion

$$f(z) = \sum_{j > -\infty} a_j q^j$$

with  $a_j \in \mathbb{C}$ , where  $q = e^{2\pi iz}$  as usual. We then have

$$(4.9) \quad Df := \frac{1}{2\pi i} \frac{df}{dz} = q \frac{df}{dq}$$

If the  $q$ -expansion of  $f$  has coefficients in  $\mathbb{Z}$ , so does the  $q$ -series for  $Df$ .

In general, the function  $Df$  is no longer modular. There is another derivation  $\vartheta$  that preserves modularity. If  $f$  has weight  $k$ , its *Serre derivative* is defined by

$$(4.10) \quad \vartheta f := Df - \frac{k}{12} \mathcal{E}_2 \cdot f$$

where  $\mathcal{E}_2(z) = 1 - 24 \sum_{j=1}^{\infty} \sigma_1(j) q^j$  is the quasimodular Eisenstein series of weight 2. This  $\vartheta f$  is a meromorphic modular form of weight  $k + 2$  [Ono2004, Proposition 2.11]. Cf. [Serre1973, Théorème 5(a)] for the action of  $D$  on  $p$ -adic modular forms.

**Example 4.11** Consider the modular discriminant  $\Delta$ , a cusp form of weight 12. The product expansion

$$\Delta = q \prod_{n=1}^{\infty} (1 - q^n)^{24}$$

implies that

$$\begin{aligned}
 \log \Delta &= \log q + 24 \sum_{n=1}^{\infty} \log(1 - q^n) \\
 (4.12) \quad &= \log q + 24 \sum_{n=1}^{\infty} \sum_{k=1}^{\infty} (-1)^{k-1} \frac{(-q^n)^k}{k} \\
 &= \log q - 24 \sum_{m=1}^{\infty} \sigma_{-1}(m) q^m
 \end{aligned}$$

Thus  $D \log \Delta = \mathcal{E}_2$ , and hence  $\vartheta \Delta = 0$ .

**Theorem 4.13** *Let  $E$  be a Morava  $E$ -theory of height 2 at the prime  $p$ , and  $N > 3$  be any integer prime to  $p$ . Choose any preferred  $\mathcal{P}_N$ -model for  $E$  in the sense of Definition 3.29. Let*

$$\beta: \mathbf{MF}(\Gamma_1(N)) \rightarrow E^0$$

*be the ring homomorphism in Proposition 2.9 built from this model. Let*

$$\ell_{2,p}: (E^0)^\times \rightarrow E^0$$

*be Rezk's logarithmic cohomology operation. Suppose that  $f \in \mathbf{MF}(\Gamma_1(N))^\times$  has trivial Nebentypus character. If the Serre derivative  $\vartheta f = 0$ , then  $\beta(f)$  is contained in the kernel of  $\ell_{2,p}$ .*

Our proof consists of two parts. In the first part (Lemma 4.14 below), we show that  $\ell_{2,p}(\beta(f))$  is constant, i.e., it is the image of a constant modular form under  $\beta$ . This is based on an interplay between the differential structures and the action of Hecke operators on modular forms. Looking at  $q$ -expansions, we then show in the second part that this constant equals zero. It boils down to an analysis of the behavior of Tate curves under isogenies.

**Lemma 4.14** *Given any  $f \in \mathbf{MF}(\Gamma_1(N))^\times$  such that  $\vartheta f = 0$ , the elliptic function  $F_1(\log f)$  is constant, where  $F_1 = 1 - T_{1,p} + pT_{2,p}$  is the operator defined in (4.5).*

**Proof** Suppose that  $f$  is of weight  $k$ . By (4.10), since  $\vartheta f = 0$ , we have

$$D \log f = \frac{Df}{f} = \frac{k\mathcal{E}_2}{12}$$

Note that  $\mathcal{E}_2$  is a weight-2 eigenform for every Hecke operator  $T_m = T_{1,m}$ ,  $m \geq 1$  with eigenvalue  $\sigma_1(m)$ .

By comparing the effects on  $q$ -expansions, we have

$$(4.15) \quad D \circ T_{i,p} = \frac{1}{p^i} \cdot T_{i,p} \circ D$$

for  $i = 1$  and  $i = 2$ . We then compute that

$$\begin{aligned} D(F_1(\log f)) &= D((1 - T_{1,p} + pT_{2,p}) \log f) \\ &= \left(1 - \frac{1}{p} \cdot T_{1,p} + \frac{1}{p^2} \cdot pT_{2,p}\right) (D \log f) \\ &= \left(1 - \frac{1}{p} \cdot T_{1,p} + \frac{1}{p^2} \cdot pT_{2,p}\right) \frac{k\mathcal{E}_2}{12} \\ &= \left(1 - \frac{1}{p} \cdot (1+p) + \frac{1}{p^2} \cdot p \left(\frac{1}{p^2} \cdot p^2\right)\right) \frac{k\mathcal{E}_2}{12} \\ &= 0 \end{aligned}$$

Thus, as a function of the complex variable  $z$ ,  $F_1(\log f)$  is constant.  $\square$

**Proof of Theorem 4.13** Let  $x := \beta(f)$ . Since  $\beta$  is a ring homomorphism,  $x \in (E^0)^\times$ . Recall from (4.4) that

$$(4.16) \quad \ell_{2,p}(x) = \frac{1}{p} \log \frac{x^p \cdot p^\bullet \phi(x)}{\prod_{i=0}^p \alpha_i^\bullet \psi_i^p(x)} = \beta(F_1(\log f))$$

By Lemma 4.14, the above equals an element in  $\mathbb{W}(\overline{\mathbb{F}}_p)^\times$ , which we denote by  $c_f$ .

Suppose that  $f$  has weight  $k$ . Via the correspondence between power operations and deformations of Frobenius in [Rezk2009, Theorem B], the ratio of values of power operations on  $x$  in (4.16) equals a ratio of values of  $f$  on the corresponding universal elliptic curves. Explicitly, with notation as in (3.14) and (3.20), we have

$$(4.17) \quad \frac{x^p \cdot p^\bullet \phi(x)}{\prod_{i=0}^p \alpha_i^\bullet \psi_i^p(x)} = \frac{f(\mathcal{C}_S, P_0, du)^p \cdot p^k f(\mathcal{C}_S, [p]P_0, du)}{\prod_{i=0}^p \kappa_i^k f(\mathcal{C}_{S^{(p)}}/\mathcal{G}_i^{(p)}, \Psi_i^{(p)}(P_0), du)}$$

To determine  $c_f$ , we need only inspect the constant term in a  $q$ -expansion for the right-hand side.

Over a punctured formal neighborhood of each cusp, the universal curve  $\mathcal{C}$  is isomorphic to the Tate curve  $\text{Tate}(q^N)$  with the level- $\Gamma_1(N)$  structure corresponding to that cusp. The universal degree- $p$  isogeny on  $\text{Tate}(q^N)$  is defined over the ring  $\mathbb{Z}[1/pN, \zeta_p][\langle q^{1/p} \rangle]$ , where  $\zeta_p$  is a primitive  $p$ 'th root of unity (see [Katz1973, Sections 1.2, 1.4, and 1.11]). In particular, the  $(p+1)$  subgroups of order  $p$  are

$$G_0^{(p)}, \text{ generated by } \zeta_p, \quad \text{and} \quad G_i^{(p)}, 1 \leq i \leq p, \text{ generated by } (\zeta_p^i q^{1/p})^N$$

Let  $\sum_{j=m}^{\infty} a_j q^j$  be a  $q$ -expansion of  $f$ . We now compare as follows the lowest powers of  $q$  at the denominator and the numerator of (4.17).

By [Katz1973, (1.11.0.3) and (1.11.0.4)] (cf. Section 1.3) and in view of Remark 3.15, we have at the denominator a leading term as the product of

$$p^k a_m (q^p)^m \quad \text{and} \quad p^k \left( ((-1)^{p+1} p)^{-k} a_m (\zeta_p^i q^{1/p})^m \right)$$

where  $i$  runs from 1 to  $p$ .<sup>5</sup> Note that since the Nebentypus character of  $f$  is trivial, the coefficient  $a_m$  is independent of where the level- $\Gamma_1(N)$  structure goes under each degree- $p$  isogeny.

At the numerator, for the first factor, we have a leading term  $(a_m q^m)^p$ . For the second factor, we have a leading term  $p^k a_m q^m$ , again under the assumption that  $f$  has trivial Nebentypus character.

Combining these, we see that the ratio has a leading constant term

$$\begin{aligned} \frac{(a_m q^m)^p \cdot p^k a_m q^m}{p^k a_m (q^p)^m \cdot \prod_{i=1}^p p^k \left( ((-1)^{p+1} p)^{-k} a_m (\zeta_p^i q^{1/p})^m \right)} &= \zeta_p^{-m(1+p)p/2} \\ &= \begin{cases} (-1)^m & \text{if } p = 2 \\ 1 & \text{if } p \neq 2 \end{cases} \end{aligned}$$

Thus applying the  $p$ -adic logarithm (cf. Example 4.8) we get  $c_f = 0$ .  $\square$

**Remark 4.18** Let  $f$  be any nonzero meromorphic modular form on  $\mathrm{SL}_2(\mathbb{Z})$ . Bruinier, Kohnen, and Ono give an explicit formula for  $\vartheta f$  as a multiple of  $-f$  by a certain function  $f_{\Theta}$  [Bruinier-Kohnen-Ono2004, Theorem 1]. The function  $f_{\Theta}$  encodes a sequence of modular functions  $j_m(z)$ ,  $m \geq 0$  defined by applying Hecke operators to the usual  $j$ -invariant, whose values are of arithmetic and combinatorial significance.

In particular, the formula of Bruinier, Kohnen, and Ono immediately shows that a nonzero meromorphic modular form  $f$  has vanishing Serre derivative precisely when its zeros and poles are located only at the cusp (cf. [Dumas-Royer2014, Proposition 6]). The functions in Example 4.8, and in fact any unit in  $\mathrm{MF}(\Gamma_1(N))$ , all have the latter property, though they are associated to  $\Gamma_1(N)$  instead of  $\mathrm{SL}_2(\mathbb{Z})$ .

Their theorem has been generalized by Ahlgren to  $\Gamma_0(p)$  with  $p \in \{2, 3, 5, 7, 13\}$  [Ahlgren2003, Theorem 2] and further by Choi to  $\Gamma_0(n)$  for any  $n$  that is square-free

<sup>5</sup>Comparing these to (3.26) and (3.27), we see  $\alpha_0 = p$  in the first expression and  $\alpha_i = p \cdot ((-1)^{p+1} p)^{-1} = (-1)^{p+1}$  in the second expression.

[Choi2006, Theorem 3.4]. In view of the assumption on Nebentypus character in Theorem 4.13, we note that modular forms of level  $\Gamma_1(N)$  with trivial Nebentypus character are precisely those of level  $\Gamma_0(N)$ .

## 5 Extending the action of Hecke operators onto logarithmic $q$ -series

The purpose of this section is to give an account of elliptic functions of the form  $\log f$ , where  $f$  is a meromorphic modular form. As we have seen in Section 4, such functions arise in computations for logarithmic cohomology operations on Morava  $E$ -theories at height 2, e.g., in (4.4).

**Example 5.1** Consider the function  $\log \Delta$  in Example 4.11. In the final form of (4.12), the second summand is a convergent  $q$ -series. We may call it an *Eisenstein series of weight 0*, by analogy to  $q$ -expansions for the usual Eisenstein series of higher weight (also cf. the real analytic Eisenstein series of weight 0 discussed in [Funke2007, Sections 3.3 and 4.1]).

The first summand,  $\log q$ , never shows up in the  $q$ -expansion of a meromorphic modular form. It is indeed this term that we will address, given the importance of  $\Delta$  in the context of logarithmic operations (see Theorem 4.13). Specifically, with motivations from homotopy theory, we aim to extend the classical action of Hecke operators on modular forms to incorporate series such as (4.12). We will then apply the extended action in Example 5.11.

**Remark 5.2** Given a cusp form  $f = \sum_{n=1}^{\infty} a_n q^n$  of weight  $k$ , its Eichler integral  $\tilde{f} = \sum_{n=1}^{\infty} n^{-k+1} a_n q^n$  is a mock modular form of weight  $2 - k$  (see the end of [Zagier2009, § 6]). Recall from Example 4.11 that  $D \log \Delta = \mathcal{E}_2$ . Since  $D^{k-1} \tilde{f} = f$ , we may then view  $\log \Delta$  as a generalized Eichler integral, “generalized” in that  $\mathcal{E}_2$  is not a cusp form. We may even approach proving Lemma 4.14 from this viewpoint.

In [Knopp-Mason2011], given a representation  $\rho: \mathrm{SL}_2(\mathbb{Z}) \rightarrow \mathrm{GL}_n(\mathbb{C})$ , Knopp and Mason consider  $n$ -dimensional vector-valued modular forms associated to  $\rho$ . In particular, they show in [Knopp-Mason2011, Theorem 2.2] that the components of certain vector-valued modular forms are functions

$$(5.3) \quad f(z) = \sum_{j=0}^t (\log q)^j h_j(z)$$

where  $t \geq 0$  is an integer, and each  $h_j$  is a convergent  $q$ -series with at worst real exponents (cf. [Knopp-Mason2011, (7), (13), and Sections 3.2–3.3]). They remark that  $q$ -expansions of this form occur in logarithmic conformal field theory (e.g., cf. [Zhu1996, (5.3.9)] and [Dong-Li-Mason2000, (6.12)]). The one in (4.12) gives another example. Following Knopp and Mason, we call the series in (5.3) a *logarithmic  $q$ -series*.

**Proposition 5.4** *The action of the Hecke operator  $T_p$  on modular forms of level  $\Gamma_0(N)$  extends so that*

$$T_p(\log q) = (p^{-1} + p^{-2}) \log q$$

**Proof** Consider modular forms in  $\text{MF}(\Gamma_1(N))$  with trivial Nebentypus character, i.e., those on  $\Gamma_0(N)$ . We follow the modular description in [Katz1973, Section 1.11] for Hecke operators in the presence of the Tate curve  $\text{Tate}(q^N)$  over  $\mathbb{Z}[1/pN, \zeta_p][[q^{1/p}]]$ , where  $\zeta_p$  is a primitive  $p$ 'th root of unity.

Write  $\mathcal{F} := \log q$ . Let  $\omega_{\text{can}}$  be the canonical differential on  $\text{Tate}(q^N)$ . Let  $\mathcal{M}_N := \text{Proj } S_N$  be the scheme over  $\mathbb{Z}[1/N]$  representing the moduli problem  $\mathcal{P}_N$  (see Examples 2.1, 2.2, and 2.6). Denote by  $\underline{\omega} := \text{pr}_* \Omega_{\mathcal{C}_N/\mathcal{M}_N}^1$  the pushforward along the structure morphism  $\text{pr}: \mathcal{C}_N \rightarrow \mathcal{M}_N$  of the relative cotangent sheaf  $\Omega_{\mathcal{C}_N/\mathcal{M}_N}^1$ .

By [Katz-Mazur1985, Theorem 10.13.11], the isomorphism

$$\underline{\omega}^{\otimes 2} \xrightarrow{\sim} \Omega_{\mathcal{M}_N/\mathbb{Z}[1/N]}^1$$

on  $\mathcal{M}_N$  extends to an isomorphism

$$\underline{\omega}^{\otimes 2} \xrightarrow{\sim} \Omega_{\mathcal{M}_N/\mathbb{Z}[1/N]}^1(\log \text{cusps})$$

on the compactification  $\overline{\mathcal{M}}_N$ , where the target is the invertible sheaf of one-forms with at worst simple poles along the cusps. In particular, over the cusps,  $\omega_{\text{can}}^2$  corresponds to  $N \cdot d\mathcal{F}$  under this isomorphism (cf. [Katz1973, Section 1.5]). Therefore, as  $\mathcal{F} = \log q = 2\pi iz$  is linear in  $z$ , we have

$$(5.5) \quad \mathcal{F}(\text{Tate}(q^N), P_0, p \cdot \omega_{\text{can}}) = p^2 \cdot \mathcal{F}(\text{Tate}(q^N), P_0, \omega_{\text{can}})$$

By [Katz1973, (1.11.0.3) and (1.11.0.4)] (cf. Section 1.3), given the assumption of

trivial Nebentypus character, we then calculate that

$$\begin{aligned}
 T_p(\log q) &= \frac{1}{p} \cdot p^k \left( \log(q^p) + \sum_{i=1}^p p^{-k} \log(\zeta_p^i q^{1/p}) \right) \\
 &= p^{k-1} \left( p \log q + p^{-k} \sum_{i=1}^p (\log \zeta_p^i + p^{-1} \log q) \right) \\
 &= p^{k-1} \left( p \log q + p^{-k} \sum_{i=1}^p p^{-1} \log q \right) \\
 &= p^{k-1} (p + p^{-k}) \log q
 \end{aligned}
 \tag{5.6}$$

where we interpret “log” as the  $p$ -adic logarithm so that  $\log \zeta_p^i = 0$  (cf. Example 4.8). Setting  $k = -2$  in view of (5.5), we obtain the stated identity.  $\square$

**Remark 5.7** More generally, if we view  $\log q$  as generalizing modular forms in  $\text{MF}(\Gamma_1(N))$  with Nebentypus character  $\chi$ , we compute as in (5.6) and obtain

$$T_p(\log q) = (p^{-1} + \chi(p)p^{-2}) \log q$$

For the rest of this section, we focus on the case when  $\chi$  is trivial (cf. Theorem 4.13).

Let  $K$  be a number field. In [Bruinier-Ono2003], Bruinier and Ono study meromorphic modular forms  $g$  on  $\text{SL}_2(\mathbb{Z})$  with  $q$ -expansion

$$g(z) = q^m \left( 1 + \sum_{n=1}^{\infty} a_n q^n \right) \tag{5.8}$$

where  $m \in \mathbb{Z}$  and  $a_n \in \mathcal{O}_K$  (the ring of integers in  $K$ ). They show that if  $g$  satisfies a certain condition with respect to a prime  $p$ , its logarithmic derivative  $D \log(g)$  is a  $p$ -adic modular form of weight 2 [Bruinier-Ono2003, Theorem 1] (the differential operator  $D$  is defined in (4.9)).

This theorem has been generalized to meromorphic modular forms on  $\Gamma_0(p)$  for  $p \geq 5$  [Getz2004, Theorem 4]. Examples include  $\mathcal{E}_{p-1}$  at each  $p \geq 5$  and, for all  $p$ , meromorphic modular forms whose zeros and poles are located only at the cusps (cf. [Bruinier-Ono2003, Definition 3.1]). In particular, when  $g = \Delta$ , we have  $D \log \Delta = \mathcal{E}_2$  (cf. the discussion before [Serre1973, Théorème 5]).

Given any  $g$  as in (5.8), note that  $\log(g)$  is a logarithmic  $q$ -series. The following definition is based on Proposition 5.4 (specifically (5.5) and (5.6)) and the theorem of Bruinier and Ono above.



**Definition 5.9**

- (i) Given any integer  $j \geq 0$ , define the *weight* of  $(\log q)^j$  to be  $-2j$ .
- (ii) Let  $g$  be a meromorphic modular form such that  $D \log(g)$  is a  $p$ -adic modular form of weight 2 for all  $p$ . Define the *weight* of  $\log(g)$  to be 0.
- (iii) Given any prime  $p$  and any logarithmic  $q$ -series

$$f = \sum_{j=0}^t (\log q)^j h_j$$

of weight  $k$ , define  $T_p f$  as follows. For each  $j$ , if

$$h_j = \sum_{m > -\infty} a_m q^m$$

(the index  $m$  and the coefficients  $a_m$  depend on  $j$ ), define

$$T_p((\log q)^j h_j) := (\log q)^j \sum_{m > -\infty} b_m q^m$$

where the coefficients

$$b_m = p^{j+k-1} a_{m/p} + p^{-j} a_{pm}$$

with the convention that  $a_{m/p} = 0$  unless  $p|m$ . We then define

$$T_p f := \sum_{j=0}^t T_p((\log q)^j h_j)$$

**Remark 5.10**

- (i) The definitions of weight above are compatible with the action of the differential operator  $D$ . Specifically, we have

$$D \log q = q \frac{d}{dq} (\log q) = 1$$

Thus applying  $D$  to  $\log q$  increases the weight by 2, which agrees with [Serre1973, Théorème 5 (a)]. More generally, for  $j \geq 0$ , since  $D(\log q)^j = j(\log q)^{j-1}$ , the same compatibility holds.

- (ii) The definition for  $T_p((\log q)^j h_j)$  extends [Katz1973, Formula 1.11.1] by a computation analogous to (5.6) under the assumption of trivial Nebentypus character

(see Remark 5.7). Moreover, the following identities for operators acting on modular forms extend to the series  $(\log q)^j h_j$ :

$$\begin{aligned} D \circ T_p &= \frac{1}{p} \cdot T_p \circ D && \text{cf. (4.15)} \\ T_\ell \circ T_p &= T_p \circ T_\ell && \text{for primes } \ell \text{ and } p \end{aligned}$$

(assuming that Hecke operators preserve weight). We can also define the Hecke operators  $T_m$  acting on  $(\log q)^j h_j$  for any positive integer  $m$  as in the remark after [Serre1973, Théorème 4].

**Example 5.11** We now return to Example 5.1. By Definition 5.9(ii),  $\log \Delta$  is a logarithmic  $q$ -series of weight 0. In view of (4.12), we then compute by Definition 5.9(iii) that

$$T_p(\log \Delta) = \sigma_{-1}(p) \log \Delta$$

Thus in (4.6) we have

$$F_1(\log \Delta) = (1 - \sigma_{-1}(p) + p^{-1}) \log \Delta = 0$$

This gives a second proof of Theorem 4.13 in the case  $f = \Delta$ .

## 6 Topological Hecke operators in terms of individual power operations

Constructed from total power operations, the topological Hecke operators in (3.17) and (3.22) can be defined more generally on  $E^0 X$  for any space  $X$ . In this section, we examine their role in the algebra of additive  $E$ -cohomology operations, by expressing them in terms of generators of this algebra.

Let  $E$  be a Morava  $E$ -theory of height  $n$  at the prime  $p$ . There is an algebraic theory—in the sense of Lawvere [Lawvere1963]—constructed from the extended power functors on the category of  $E$ -modules (cf. [Rezk2006a, Section 9] and [Rezk2009, Section 4]). It describes all homotopy operations on  $K(n)$ -local commutative  $E$ -algebras.

Rezk shows that under a certain congruence condition, a model for this algebraic theory amounts to a  $p$ -torsion-free graded commutative algebra over an associative ring  $\Gamma$  [Rezk2009, Theorem A]. Following Rezk, we call  $\Gamma$  the *Dyer-Lashof algebra* of the  $E$ -theory. It is constructed in [Rezk2009, 6.2] as a direct sum of the  $E^0$ -linear duals of the rings  $E^0(B\Sigma_{p^k})/I$ , for all  $k \geq 0$  (cf. (3.4)).

**Example 6.1** A Dyer-Lashof algebra  $\Gamma$  at height 2 and prime 5 is a graded twisted bialgebra over  $E_0 \cong \mathbb{W}(\overline{\mathbb{F}}_5)[[u_1]]$  with generators  $Q_i$ ,  $0 \leq i \leq 5$ . In the absence of a formula for  $\psi^5(u_1)$ , using those for  $\tilde{h}$  and  $\tilde{\alpha}$  in (3.11), we compute as in [Zhu2014, Proposition 3.6] and obtain *some* relations for  $\Gamma$ . Note that the generators  $Q_i$ , as well as relations among them, depend on the choice of  $\alpha$ .

- “Twists” in terms of *commutation relations*:

$$Q_i c = (Fc)Q_i \text{ for } c \in \mathbb{W}(\overline{\mathbb{F}}_5) \text{ and all } i, \text{ with } F \text{ the Frobenius automorphism}$$

$$\begin{aligned} Q_0 h &= (h^5 - 10h^4 - 1065h^3 + 12690h^2 + 168930h - 1462250)Q_0 + (5h^4 \\ &\quad - 50h^3 - 3950h^2 + 42200h + 233400)Q_1 + (25h^3 - 250h^2 - 12875h \\ &\quad + 104750)Q_2 + (125h^2 - 1250h - 30000)Q_3 + (625h - 6250)Q_4 \\ &\quad + 3120Q_5 \end{aligned}$$

$$\begin{aligned} Q_1 h &= (-55h^4 + 850h^3 + 39575h^2 - 608700h - 1113524)Q_0 + (-275h^3 \\ &\quad + 4250h^2 + 122250h - 1462250)Q_1 + (-1375h^2 + 21250h \\ &\quad + 233400)Q_2 + (-6875h + 104750)Q_3 - 30000Q_4 + (h - 6250)Q_5 \end{aligned}$$

$$\begin{aligned} Q_2 h &= (60h^4 - 775h^3 - 45400h^2 + 593900h + 2008800)Q_0 + (300h^3 \\ &\quad - 3875h^2 - 144500h + 1453876)Q_1 + (1500h^2 - 19375h \\ &\quad - 310000)Q_2 + (7500h - 96600)Q_3 + 36000Q_4 + 4320Q_5 \end{aligned}$$

$$\begin{aligned} Q_3 h &= (-35h^4 + 400h^3 + 27125h^2 - 320900h - 1418300)Q_0 + (-175h^3 \\ &\quad + 2000h^2 + 87500h - 792000)Q_1 + (-875h^2 + 10000h + 196876)Q_2 \\ &\quad + (-4375h + 50000)Q_3 - 21600Q_4 - 1440Q_5 \end{aligned}$$

$$\begin{aligned} Q_4 h &= (10h^4 - 105h^3 - 7850h^2 + 86975h + 445850)Q_0 + (50h^3 - 525h^2 \\ &\quad - 25500h + 215500)Q_1 + (250h^2 - 2625h - 58750)Q_2 + (1250h \\ &\quad - 13124)Q_3 + 6250Q_4 + 240Q_5 \end{aligned}$$

$$\begin{aligned} Q_5 h &= (-h^4 + 10h^3 + 790h^2 - 8440h - 46680)Q_0 + (-5h^3 + 50h^2 + 2575h \\ &\quad - 20950)Q_1 + (-25h^2 + 250h + 6000)Q_2 + (-125h + 1250)Q_3 \\ &\quad - 624Q_4 + 10Q_5 \end{aligned}$$

- A product structure subject to *Adem relations*:

$$\begin{aligned} Q_1 Q_0 &= 55Q_0 Q_1 + (55h - 300)Q_0 Q_2 + (55h^2 - 300h - 14250)Q_0 Q_3 + (55h^3 \\ &\quad - 300h^2 - 29375h + 163750)Q_0 Q_4 + (55h^4 - 300h^3 - 44500h^2 \\ &\quad + 328750h + 3228750)Q_0 Q_5 + 275Q_1 Q_2 + (275h - 1500)Q_1 Q_3 \\ &\quad + (275h^2 - 1500h - 71250)Q_1 Q_4 + (275h^3 - 1500h^2 - 146875h \\ &\quad + 818750)Q_1 Q_5 - 5Q_2 Q_1 + 1375Q_2 Q_3 + (1375h - 7500)Q_2 Q_4 \end{aligned}$$

$$\begin{aligned}
& + (1375h^2 - 7500h - 356250)Q_2Q_5 - 25Q_3Q_2 + 6875Q_3Q_4 \\
& + (6875h - 37500)Q_3Q_5 - 125Q_4Q_3 + 34375Q_4Q_5 - 625Q_5Q_4 \\
Q_2Q_0 = & -60Q_0Q_1 + (-60h + 175)Q_0Q_2 + (-60h^2 + 175h + 16250)Q_0Q_3 \\
& + (-60h^3 + 175h^2 + 32750h - 138000)Q_0Q_4 + (-60h^4 + 175h^3 \\
& + 49250h^2 - 276125h - 3943750)Q_0Q_5 - 300Q_1Q_2 + (-300h \\
& + 875)Q_1Q_3 + (-300h^2 + 875h + 81250)Q_1Q_4 + (-300h^3 + 875h^2 \\
& + 163750h - 690000)Q_1Q_5 - 1500Q_2Q_3 + (-1500h + 4375)Q_2Q_4 \\
& + (-1500h^2 + 4375h + 406250)Q_2Q_5 - 5Q_3Q_1 - 7500Q_3Q_4 \\
& + (-7500h + 21875)Q_3Q_5 - 25Q_4Q_2 - 37500Q_4Q_5 - 125Q_5Q_3 \\
Q_3Q_0 = & 35Q_0Q_1 + (35h - 50)Q_0Q_2 + (35h^2 - 50h - 9600)Q_0Q_3 + (35h^3 \\
& - 50h^2 - 19225h + 66250)Q_0Q_4 + (35h^4 - 50h^3 - 28850h^2 \\
& + 132500h + 2411875)Q_0Q_5 + 175Q_1Q_2 + (175h - 250)Q_1Q_3 \\
& + (175h^2 - 250h - 48000)Q_1Q_4 + (175h^3 - 250h^2 - 96125h \\
& + 331250)Q_1Q_5 + 875Q_2Q_3 + (875h - 1250)Q_2Q_4 + (875h^2 \\
& - 1250h - 240000)Q_2Q_5 + 4375Q_3Q_4 + (4375h - 6250)Q_3Q_5 \\
& - 5Q_4Q_1 + 21875Q_4Q_5 - 25Q_5Q_2 \\
Q_4Q_0 = & -10Q_0Q_1 + (-10h + 5)Q_0Q_2 + (-10h^2 + 5h + 2750)Q_0Q_3 \\
& + (-10h^3 + 5h^2 + 5500h - 16375)Q_0Q_4 + (-10h^4 + 5h^3 + 8250h^2 \\
& - 32750h - 705000)Q_0Q_5 - 50Q_1Q_2 + (-50h + 25)Q_1Q_3 + (-50h^2 \\
& + 25h + 13750)Q_1Q_4 + (-50h^3 + 25h^2 + 27500h - 81875)Q_1Q_5 \\
& - 250Q_2Q_3 + (-250h + 125)Q_2Q_4 + (-250h^2 + 125h \\
& + 68750)Q_2Q_5 - 1250Q_3Q_4 + (-1250h + 625)Q_3Q_5 - 6250Q_4Q_5 \\
& - 5Q_5Q_1 \\
Q_5Q_0 = & Q_0Q_1 + hQ_0Q_2 + (h^2 - 275)Q_0Q_3 + (h^3 - 550h + 1500)Q_0Q_4 + (h^4 \\
& - 825h^2 + 3000h + 71250)Q_0Q_5 + 5Q_1Q_2 + 5hQ_1Q_3 + (5h^2 \\
& - 1375)Q_1Q_4 + (5h^3 - 2750h + 7500)Q_1Q_5 + 25Q_2Q_3 + 25hQ_2Q_4 \\
& + (25h^2 - 6875)Q_2Q_5 + 125Q_3Q_4 + 125hQ_3Q_5 + 625Q_4Q_5
\end{aligned}$$

- A coproduct structure given by *Cartan formulas*:

$$\begin{aligned}
Q_0(xy) = & Q_0(x)Q_0(y) - 5(Q_1(x)Q_5(y) + Q_2(x)Q_4(y) + Q_3(x)Q_3(y) \\
& + Q_4(x)Q_2(y) + Q_5(x)Q_1(y)) - 50(Q_2(x)Q_5(y) + Q_3(x)Q_4(y) \\
& + Q_4(x)Q_3(y) + Q_5(x)Q_2(y)) - 325(Q_3(x)Q_5(y) + Q_4(x)Q_4(y) \\
& + Q_5(x)Q_3(y)) - 1800(Q_4(x)Q_5(y) + Q_5(x)Q_4(y))
\end{aligned}$$

$$\begin{aligned}
& - 9350Q_5(x)Q_5(y) \\
Q_1(xy) &= (Q_0(x)Q_1(y) + Q_1(x)Q_0(y)) + h(Q_1(x)Q_5(y) + Q_2(x)Q_4(y) \\
& + Q_3(x)Q_3(y) + Q_4(x)Q_2(y) + Q_5(x)Q_1(y)) + (10h - 5)(Q_2(x)Q_5(y) \\
& + Q_3(x)Q_4(y) + Q_4(x)Q_3(y) + Q_5(x)Q_2(y)) + (65h \\
& - 50)(Q_3(x)Q_5(y) + Q_4(x)Q_4(y) + Q_5(x)Q_3(y)) + (360h \\
& - 325)(Q_4(x)Q_5(y) + Q_5(x)Q_4(y)) + (1870h - 1800)Q_5(x)Q_5(y) \\
Q_2(xy) &= (Q_0(x)Q_2(y) + Q_1(x)Q_1(y) + Q_2(x)Q_0(y)) - 55(Q_1(x)Q_5(y) \\
& + Q_2(x)Q_4(y) + Q_3(x)Q_3(y) + Q_4(x)Q_2(y) + Q_5(x)Q_1(y)) + (h \\
& - 550)(Q_2(x)Q_5(y) + Q_3(x)Q_4(y) + Q_4(x)Q_3(y) + Q_5(x)Q_2(y)) \\
& + (10h - 3580)(Q_3(x)Q_5(y) + Q_4(x)Q_4(y) + Q_5(x)Q_3(y)) + (65h \\
& - 19850)(Q_4(x)Q_5(y) + Q_5(x)Q_4(y)) + (360h - 103175)Q_5(x)Q_5(y) \\
Q_3(xy) &= (Q_0(x)Q_3(y) + Q_1(x)Q_2(y) + Q_2(x)Q_1(y) + Q_3(x)Q_0(y)) \\
& + 60(Q_1(x)Q_5(y) + Q_2(x)Q_4(y) + Q_3(x)Q_3(y) + Q_4(x)Q_2(y) \\
& + Q_5(x)Q_1(y)) + 545(Q_2(x)Q_5(y) + Q_3(x)Q_4(y) + Q_4(x)Q_3(y) \\
& + Q_5(x)Q_2(y)) + (h + 3350)(Q_3(x)Q_5(y) + Q_4(x)Q_4(y) \\
& + Q_5(x)Q_3(y)) + (10h + 18020)(Q_4(x)Q_5(y) + Q_5(x)Q_4(y)) + (65h \\
& + 92350)Q_5(x)Q_5(y) \\
Q_4(xy) &= (Q_0(x)Q_4(y) + Q_1(x)Q_3(y) + Q_2(x)Q_2(y) + Q_3(x)Q_1(y) \\
& + Q_4(x)Q_0(y)) - 35(Q_1(x)Q_5(y) + Q_2(x)Q_4(y) + Q_3(x)Q_3(y) \\
& + Q_4(x)Q_2(y) + Q_5(x)Q_1(y)) - 290(Q_2(x)Q_5(y) + Q_3(x)Q_4(y) \\
& + Q_4(x)Q_3(y) + Q_5(x)Q_2(y)) - 1730(Q_3(x)Q_5(y) + Q_4(x)Q_4(y) \\
& + Q_5(x)Q_3(y)) + (h - 9250)(Q_4(x)Q_5(y) + Q_5(x)Q_4(y)) + (10h \\
& - 47430)Q_5(x)Q_5(y) \\
Q_5(xy) &= (Q_0(x)Q_5(y) + Q_1(x)Q_4(y) + Q_2(x)Q_3(y) + Q_3(x)Q_2(y) \\
& + Q_4(x)Q_1(y) + Q_5(x)Q_0(y)) + 10(Q_1(x)Q_5(y) + Q_2(x)Q_4(y) \\
& + Q_3(x)Q_3(y) + Q_4(x)Q_2(y) + Q_5(x)Q_1(y)) + 65(Q_2(x)Q_5(y) \\
& + Q_3(x)Q_4(y) + Q_4(x)Q_3(y) + Q_5(x)Q_2(y)) + 360(Q_3(x)Q_5(y) \\
& + Q_4(x)Q_4(y) + Q_5(x)Q_3(y)) + 1870(Q_4(x)Q_5(y) + Q_5(x)Q_4(y)) \\
& + (h + 9450)Q_5(x)Q_5(y)
\end{aligned}$$

- Subject to the above commutation and Adem relations,  $\Gamma$  becomes a free left module over  $E_0$ . A basis consists of monomials in the generators  $Q_i$  which are in the form  $Q_0^m Q_{i_1} \cdots Q_{i_n}$ , with  $m \geq 0$ ,  $n \geq 0$  ( $n = 0$  corresponds to  $Q_0^m$ ), and

$1 \leq i_k \leq 5$ . The grading on  $\Gamma$  refers to the sum of the exponents in each of these monomials.

Let  $E$  be a Morava  $E$ -theory of height  $n$  at the prime  $p$ . Given any  $K(n)$ -local commutative  $E$ -algebra  $A$ , the elements  $Q_i$  above are examples of additive power operations having  $A^0$  as both domain and range. Such operations arise from the total power operation

$$(6.2) \quad \psi^p: A^0 \rightarrow A^0(B\Sigma_p)/J \cong A^0[\alpha]/(w(\alpha))$$

where  $J$  is a transfer ideal, and  $w(\alpha) \in E^0[\alpha]$  is a monic polynomial of degree  $r := 1 + p + p^2 + \cdots + p^{n-1}$  (cf. (3.4)). We define *individual power operations*  $Q_i: A^0 \rightarrow A^0$  by the formula

$$(6.3) \quad \psi^p(x) = \sum_{i=0}^{r-1} Q_i(x) \alpha^i$$

In particular, given a space  $X$ , each  $Q_i$  acts on  $E^0X$  if we take  $A$  to be the spectrum of functions from  $\Sigma_+^\infty X$  to  $E$ . These individual power operations  $Q_i$  generate the Dyer-Lashof algebra of the  $E$ -theory (cf. [Zhu2014, proof of Theorem 3.10]).

**Proposition 6.4** *Let  $E$  be a Morava  $E$ -theory of height  $n = 2$  at the prime  $p$ . Choose any preferred  $\mathcal{P}_N$ -model for  $E$  in Definition 3.29. Given any  $K(2)$ -local commutative  $E$ -algebra  $A$ , let  $\psi^p$  be the total power operation in (6.2) with the polynomial*

$$(6.5) \quad w(\alpha) = w_{p+1}\alpha^{p+1} + \cdots + w_1\alpha + w_0 \quad w_{p+1} = 1$$

*associated to this model, and let  $Q_i$ ,  $0 \leq i \leq p$  be the corresponding individual power operations in (6.3). For  $\mu = 1$  and  $\mu = 2$ , let*

$$t_{\mu,p}: A^0 \rightarrow p^{-1}A^0$$

*be the topological Hecke operators that are defined as in (3.17) and (3.22) from the above total power operation  $\psi^p$  on  $A^0$ . Then the following identities hold.*

$$t_{1,p} = \frac{1}{p} \sum_{i=0}^p c_i Q_i \quad \text{and} \quad t_{2,p} = \frac{1}{p^2} \sum_{j=0}^p \sum_{i=0}^j w_0^i d_{j-i} Q_i Q_j$$

where

$$c_i = \begin{cases} p+1 & i=0 \\ -\sum_{k=0}^{i-1} w_{p+1+k-i} c_k + (p+1-i) w_{p+1-i} & 1 \leq i \leq p \end{cases}$$

$$d_\tau = \begin{cases} 1 & \tau=0 \\ -\sum_{k=0}^{\tau-1} w_0^{\tau-k-1} w_{\tau-k} d_k & 1 \leq \tau \leq p \end{cases}$$

Moreover, for  $i \geq 1$  and  $\tau \geq 1$ ,

$$c_i = i \sum_{\substack{m_1+2m_2+\dots+vm_v=i \\ m_s \geq 1}} (-1)^{m_1+\dots+m_v} \frac{(m_1+\dots+m_v-1)!}{m_1! \dots m_v!} w_{p+1-1}^{m_1} \dots w_{p+1-v}^{m_v}$$

$$d_\tau = \sum_{n=0}^{\tau-1} (-1)^{\tau-n} w_0^n \sum_{\substack{m_1+\dots+m_{\tau-n}=\tau \\ 1 \leq m_s \leq m_{s+1} \leq p}} w_{m_1} \dots w_{m_{\tau-n}}$$

**Proof** By definitions (3.17) and (6.3),

$$\begin{aligned} t_{1,p}(x) &= \frac{1}{p} \sum_{j=0}^p \psi_j^p(x) \\ &= \frac{1}{p} \sum_{j=0}^p \sum_{i=0}^p Q_i(x) \alpha_j^i \\ &= \frac{1}{p} \sum_{i=0}^p \left( \sum_{j=0}^p \alpha_j^i \right) Q_i(x) \end{aligned}$$

Since the parameters  $\alpha_j$  are the roots of  $w(\alpha)$  in (6.5), the formulas for  $c_i = \sum_{j=0}^p \alpha_j^i$  then follow from Newton's and Girard's formulas relating power sums and elementary symmetric functions (see, e.g., [Milnor-Stasheff1974, Problem 16-A] and Section 1.3).

For  $t_{2,p}(x)$ , we first write by (3.22) that

$$\begin{aligned} t_{2,p}(x) &= \frac{1}{p^2} \psi^p(\psi^p(x)) \\ &= \frac{1}{p^2} \psi^p \sum_{j=0}^p Q_j(x) \alpha^j \\ &= \frac{1}{p^2} \sum_{j=0}^p \psi^p(Q_j(x)) \psi^p(\alpha)^j \\ &= \frac{1}{p^2} \sum_{j=0}^p \left( \sum_{i=0}^p Q_i Q_j(x) \alpha^i \right) \tilde{\alpha}^j \end{aligned}$$

Since the target of  $t_{2,p}$  is  $p^{-1}A^0$ , the above identity should simplify to contain neither  $\alpha$  nor  $\tilde{\alpha}$ . Thus in view of (3.30), we rewrite

$$t_{2,p}(x) = \frac{1}{p^2} \sum_{j=0}^p \sum_{i=0}^j w_0^i Q_i Q_j(x) \tilde{\alpha}^{j-i}$$

For  $0 \leq \tau \leq p$ , we next express each  $\tilde{\alpha}^\tau$  as a polynomial in  $\alpha$  of degree at most  $p$  with coefficients in  $E^0$ , and verify that the constant term of this polynomial is  $d_\tau$  as stated in the proposition.

The case  $\tau = 0$  is clear. For  $1 \leq \tau \leq p$ , we have

$$\begin{aligned}
 \tilde{\alpha}^\tau &= \left( \frac{w_0}{\alpha} \right)^\tau && \text{by (3.30)} \\
 &= \frac{w_0^{\tau-1}(-w_{p+1}\alpha^{p+1} - \dots - w_1\alpha)}{\alpha^\tau} && \text{by (6.5)} \\
 &= w_0^{\tau-1}(-w_{p+1}\alpha^{p+1-\tau} - \dots - w_{\tau+1}\alpha) \\
 &\quad - w_0^{\tau-1}w_\tau - \frac{w_0^{\tau-1}w_{\tau-1}}{\alpha} - \dots - \frac{w_0^{\tau-1}w_1}{\alpha^{\tau-1}} \\
 &= w_0^{\tau-1}(-w_{p+1}\alpha^{p+1-\tau} - \dots - w_{\tau+1}\alpha) \\
 &\quad - w_0^{\tau-1}w_\tau - w_0^{\tau-2}w_{\tau-1}\tilde{\alpha} - \dots - w_1\tilde{\alpha}^{\tau-1} && \text{by (3.30)}
 \end{aligned}$$

and thus

$$\begin{aligned}
 d_\tau &= -w_0^{\tau-1}w_\tau - w_0^{\tau-2}w_{\tau-1}d_1 - \dots - w_1d_{\tau-1} \\
 &= -\sum_{k=0}^{\tau-1} w_0^{\tau-k-1}w_{\tau-k}d_k
 \end{aligned}$$

This gives the first identity for  $d_\tau$  stated in the proposition. From this relation, we show the second stated identity for  $d_\tau$  by induction on  $\tau$ . The base case  $\tau = 1$ , with  $d_1 = -w_1$ , can be checked directly. For  $\tau \geq 2$ , we first rewrite

$$d_\tau = -w_0^{\tau-1}w_\tau - \sum_{r=1}^{\tau-1} w_0^{r-1}w_r d_{\tau-r}$$

Expanding  $d_{\tau-r}$  for  $1 \leq r \leq \tau-1$  by the induction hypothesis, we then arrange  $\sum_{r=1}^{\tau-1} w_0^{r-1}w_r d_{\tau-r}$  above into a sum indexed by  $n$  with  $0 \leq n \leq \tau-2$ , where we collect terms that contain exactly an  $n$ -power of  $w_0$ :

$$\begin{aligned}
 d_\tau &= -w_0^{\tau-1}w_\tau - \sum_{n=0}^{\tau-2} \left( \sum_{r=1}^{n+1} w_0^{r-1}w_r (-1)^{\tau-r-(n-r+1)} w_0^{n-r+1} \right. \\
 &\quad \left. \sum_{\substack{m_{1,r} + \dots + m_{\tau-r-(n-r+1),r} = \tau-r \\ 1 \leq m_{s,r} \leq m_{s+1,r} \leq p}} w_{m_{1,r}} \cdots w_{m_{\tau-r-(n-r+1),r}} \right)
 \end{aligned}$$



$$\begin{aligned}
&= -w_0^{\tau-1}w_\tau - \sum_{n=0}^{\tau-2} (-1)^{\tau-n-1} w_0^n \sum_{r=1}^{n+1} w_r \sum_{\substack{m_{1,r}+\dots+m_{\tau-n-1,r}=\tau-r \\ 1 \leq m_{s,r} \leq m_{s+1,r} \leq p}} w_{m_{1,r}} \cdots w_{m_{\tau-n-1,r}} \\
&= -w_0^{\tau-1}w_\tau + \sum_{n=0}^{\tau-2} (-1)^{\tau-n} w_0^n \sum_{\substack{m_1+\dots+m_{\tau-n}=\tau \\ 1 \leq m_s \leq m_{s+1} \leq p}} w_{m_1} \cdots w_{m_{\tau-n}} \\
&= \sum_{n=0}^{\tau-1} (-1)^{\tau-n} w_0^n \sum_{\substack{m_1+\dots+m_{\tau-n}=\tau \\ 1 \leq m_s \leq m_{s+1} \leq p}} w_{m_1} \cdots w_{m_{\tau-n}}
\end{aligned}$$

□

**Remark 6.6** In the proof above, the method for computing  $t_{2,p}$  applies more generally. It enables us to find formulas for Adem relations—one for each  $Q_i Q_0$ ,  $1 \leq i \leq p$ —in terms of the coefficients  $w_j$  of  $w(\alpha)$  (see Example 6.1 and cf. [Zhu2014, proof of Proposition 3.6 (iv)]). We can also express Cartan formulas using these coefficients. However, commutation relations are determined by both the terms  $w_j$  and  $\psi^p(w_j)$ .

**Example 6.7** For  $p = 5$ , by Proposition 6.4, we compute from (3.8) that

$$\begin{aligned}
t_{1,5} &= \frac{6}{5}Q_0 + 2Q_1 + 6Q_2 + 26Q_3 + 126Q_4 + (h + 600)Q_5 \\
t_{2,5} &= \frac{1}{25}(Q_0Q_0 + hQ_0Q_1 + (h^2 - 275)Q_0Q_2 + (h^3 - 550h + 1500)Q_0Q_3 \\
&\quad + (h^4 - 825h^2 + 3000h + 71250)Q_0Q_4 + (h^5 - 1100h^3 + 4500h^2 \\
&\quad + 218125h - 818750)Q_0Q_5 + 5Q_1Q_1 + 5hQ_1Q_2 + (5h^2 - 1375)Q_1Q_3 \\
&\quad + (5h^3 - 2750h + 7500)Q_1Q_4 + (5h^4 - 4125h^2 + 15000h \\
&\quad + 356250)Q_1Q_5 + 25Q_2Q_2 + 25hQ_2Q_3 + (25h^2 - 6875)Q_2Q_4 + (25h^3 \\
&\quad - 13750h + 37500)Q_2Q_5 + 125Q_3Q_3 + 125hQ_3Q_4 + (125h^2 \\
&\quad - 34375)Q_3Q_5 + 625Q_4Q_4 + 625hQ_4Q_5 + 3125Q_5Q_5)
\end{aligned}$$

**Theorem 6.8** Let  $E$  be a Morava  $E$ -theory of height 2 at the prime  $p$ , and let  $\Gamma$  be its Dyer-Lashof algebra. Define  $\tilde{t}_{\mu,p} := p^\mu t_{\mu,p}$  for  $\mu = 1$  and  $\mu = 2$ , where  $t_{\mu,p}$  are the topological Hecke operators in Proposition 6.4. Then  $\tilde{t}_{2,p}$  lies in the center of  $\Gamma$  but  $\tilde{t}_{1,p}$  does not.

**Proof** By Proposition 6.4, both  $\tilde{t}_{1,p}$  and  $\tilde{t}_{2,p}$  are contained in  $\Gamma$ . In fact,  $\tilde{t}_{2,p} = \psi^p \circ \psi^p$ , and is thus a ring homomorphism. Note that  $\tilde{t}_{2,p}(\alpha) = \alpha$  as a result of the relation

(3.30). Since  $\psi^p = Q_0 + Q_1\alpha + \cdots + Q_p\alpha^p$ , we can then write the three-fold composite  $\psi^p \circ \psi^p \circ \psi^p$  in two ways:

$$\begin{aligned}\tilde{t}_{2,p}(Q_0 + Q_1\alpha + \cdots + Q_p\alpha^p) &= \tilde{t}_{2,p}Q_0 + (\tilde{t}_{2,p}Q_1)(\tilde{t}_{2,p}\alpha) + \cdots + (\tilde{t}_{2,p}Q_p)(\tilde{t}_{2,p}\alpha)^p \\ &= \tilde{t}_{2,p}Q_0 + (\tilde{t}_{2,p}Q_1)\alpha + \cdots + (\tilde{t}_{2,p}Q_p)\alpha^p \\ (Q_0 + Q_1\alpha + \cdots + Q_p\alpha^p)\tilde{t}_{2,p} &= Q_0\tilde{t}_{2,p} + (Q_1\tilde{t}_{2,p})\alpha + \cdots + (Q_p\tilde{t}_{2,p})\alpha^p\end{aligned}$$

For each  $0 \leq i \leq p$ , comparing the coefficients for  $\alpha^i$ , we see that  $\tilde{t}_{2,p}$  commutes with  $Q_i$ . Thus  $\tilde{t}_{2,p}$  lies in the center of  $\Gamma$ .

It remains to show that the other operation  $\tilde{t}_{1,p}$  is not central in  $\Gamma$ . By means analogous to the proof of Proposition 6.4, we find that the term  $Q_0Q_1$  has coefficient  $w_2$  in the Adem relation for  $Q_1Q_0$  (see Remark 6.6). Near the cusps, since  $p|w_i$  for  $2 \leq i \leq p$  according to (3.26) and (3.27), we then compute by Proposition 6.4 that

$$\begin{aligned}Q_1\tilde{t}_{1,p} &= Q_1((p+1)Q_0 - w_pQ_1 + c_2Q_2 + \cdots + c_pQ_p) \\ &\equiv Q_1Q_0 + Q_1(c_2Q_2 + \cdots + c_pQ_p) \pmod{p}\end{aligned}$$

Thus the reduction of  $Q_1\tilde{t}_{1,p}$  modulo  $p$  does not contain  $Q_0Q_1$ . On the other hand, the term  $Q_0Q_1$  in

$$\tilde{t}_{1,p}Q_1 = ((p+1)Q_0 - w_pQ_1 + c_2Q_2 + \cdots + c_pQ_p)Q_1$$

has coefficient  $p+1$ . Therefore  $\tilde{t}_{1,p}Q_1 \neq Q_1\tilde{t}_{1,p}$ .  $\square$

## References

- [Ahlgren2003] Scott Ahlgren, *The theta-operator and the divisors of modular forms on genus zero subgroups*, Math. Res. Lett. **10** (2003), no. 6, 787–798. [MR2024734\(2004m:11059\)](#)
- [Ando1992] Matthew Ando, *Operations in complex-oriented cohomology theories related to subgroups of formal groups*, ProQuest LLC, Ann Arbor, MI, 1992, Thesis (Ph.D.)–Massachusetts Institute of Technology. [MR2716371](#)
- [Ando1995] Matthew Ando, *Isogenies of formal group laws and power operations in the cohomology theories  $E_n$* , Duke Math. J. **79** (1995), no. 2, 423–485. [MR1344767\(97a:55006\)](#)
- [Ando2000] Matthew Ando, *Power operations in elliptic cohomology and representations of loop groups*, Trans. Amer. Math. Soc. **352** (2000), no. 12, 5619–5666. [MR1637129](#)
- [Ando-Hopkins-Rezk2010] Matthew Ando, Michael J. Hopkins, and Charles Rezk, *Multiplicative orientations of  $KO$ -theory and the spectrum of topological modular forms*, available at <http://www.math.uiuc.edu/~mando/papers/koandtmf.pdf>.

- [Ando-Hopkins-Strickland2001] M. Ando, M. J. Hopkins, and N. P. Strickland, *Elliptic spectra, the Witten genus and the theorem of the cube*, Invent. Math. **146** (2001), no. 3, 595–687. [MR1869850\(2002g:55009\)](#)
- [Ando-Hopkins-Strickland2004] Matthew Ando, Michael J. Hopkins, and Neil P. Strickland, *The sigma orientation is an  $H_\infty$  map*, Amer. J. Math. **126** (2004), no. 2, 247–334. [MR2045503\(2005d:55009\)](#)
- [Atkin-Lehner1970] A. O. L. Atkin and J. Lehner, *Hecke operators on  $\Gamma_0(m)$* , Math. Ann. **185** (1970), 134–160. [MR0268123\(42 #3022\)](#)
- [Baker1989] Andrew Baker, *Elliptic cohomology,  $p$ -adic modular forms and Atkin’s operator  $U_p$* , Algebraic topology (Evanston, IL, 1988), Contemp. Math., vol. 96, Amer. Math. Soc., Providence, RI, 1989, pp. 33–38. [MR1022672\(90i:55007\)](#)
- [Baker1990] Andrew Baker, *Hecke operators as operations in elliptic cohomology*, J. Pure Appl. Algebra **63** (1990), no. 1, 1–11. [MR1037690\(91m:55005\)](#)
- [Baker et al. 2005] Matthew H. Baker, Enrique González-Jiménez, Josep González, and Bjorn Poonen, *Finiteness results for modular curves of genus at least 2*, Amer. J. Math. **127** (2005), no. 6, 1325–1387. [MR2183527\(2006i:11065\)](#)
- [Behrens-Ormsby2015] Mark Behrens and Kyle Ormsby, *On the homotopy of  $Q(3)$  and  $Q(5)$  at the prime 2*. [arXiv:1211.0076](#)
- [Behrens-Ormsby-Stapleton-Stojanoska2015] M. Behrens, K. Ormsby, N. Stapleton, and V. Stojanoska, *On the ring of cooperations for 2-primary connective topological modular forms*. [arXiv:1501.01050](#)
- [Bruinier-Kohnen-Ono2004] Jan H. Bruinier, Winfried Kohnen, and Ken Ono, *The arithmetic of the values of modular functions and the divisors of modular forms*, Compos. Math. **140** (2004), no. 3, 552–566. [MR2041768\(2005h:11083\)](#)
- [Bruinier-Ono2003] Jan H. Bruinier and Ken Ono, *The arithmetic of Borcherds’ exponents*, Math. Ann. **327** (2003), no. 2, 293–303. [MR2015071\(2005b:11055\)](#)
- [Bruner-May-McClure-Steinberger1986] R. R. Bruner, J. P. May, J. E. McClure, and M. Steinberger,  *$H_\infty$  ring spectra and their applications*, Lecture Notes in Mathematics, vol. 1176, Springer-Verlag, Berlin, 1986. [MR836132\(88e:55001\)](#)
- [Choi2006] D. Choi, *On values of a modular form on  $\Gamma_0(N)$* , Acta Arith. **121** (2006), no. 4, 299–311. [MR2224397\(2006m:11051\)](#)
- [Dong-Li-Mason2000] Chongying Dong, Haisheng Li, and Geoffrey Mason, *Modular-invariance of trace functions in orbifold theory and generalized Moonshine*, Comm. Math. Phys. **214** (2000), no. 1, 1–56. [MR1794264\(2001k:17043\)](#)
- [Dumas-Royer2014] François Dumas and Emmanuel Royer, *Poisson structures and star products on quasimodular forms*, Algebra Number Theory **8** (2014), no. 5, 1127–1149. [MR3263138](#)

- [Funke2007] Jens Funke, *CM points and weight 3/2 modular forms*, Analytic number theory, Clay Math. Proc., vol. 7, Amer. Math. Soc., Providence, RI, 2007, pp. 107–127. [MR2362196\(2008j:11054\)](#)
- [Ganter2006] Nora Ganter, *Orbifold genera, product formulas and power operations*, Adv. Math. **205** (2006), no. 1, 84–133. [MR2254309](#)
- [Ganter2009] Nora Ganter, *Hecke operators in equivariant elliptic cohomology and generalized Moonshine*, Groups and symmetries, CRM Proc. Lecture Notes, vol. 47, Amer. Math. Soc., Providence, RI, 2009, pp. 173–209. [MR2500561\(2011j:55012\)](#)
- [Ganter2013a] Nora Ganter, *Power operations in orbifold Tate  $K$ -theory*, Homology Homotopy Appl. **15** (2013), no. 1, 313–342. [MR3079210](#)
- [Ganter2013b] Nora Ganter, *Stringy power operations in Tate  $K$ -theory*. [arXiv:math/0701565](#)
- [Getz2004] Jayce Getz, *Classical and  $p$ -adic modular forms arising from the Borcherds exponents of other modular forms*, available at [http://www.math.harvard.edu/theses/senior/getz/getz\\_senior\\_thesis.pdf](http://www.math.harvard.edu/theses/senior/getz/getz_senior_thesis.pdf).
- [Goerss-Hopkins2004] P. G. Goerss and M. J. Hopkins, *Moduli spaces of commutative ring spectra*, Structured ring spectra, London Math. Soc. Lecture Note Ser., vol. 315, Cambridge Univ. Press, Cambridge, 2004, pp. 151–200. [MR2125040\(2006b:55010\)](#)
- [Hill-Lawson2016] Michael Hill and Tyler Lawson, *Topological modular forms with level structure*, Invent. Math. **203** (2016), no. 2, 359–416. [MR3455154](#)
- [Katz1973] Nicholas M. Katz,  *$p$ -adic properties of modular schemes and modular forms*, Modular functions of one variable, III (Proc. Internat. Summer School, Univ. Antwerp, Antwerp, 1972), Springer, Berlin, 1973, pp. 69–190. Lecture Notes in Mathematics, Vol. 350. [MR0447119\(56 #5434\)](#)
- [Katz1976] Nicholas M. Katz,  *$p$ -adic interpolation of real analytic Eisenstein series*, Ann. of Math. (2) **104** (1976), no. 3, 459–571. [MR0506271\(58 #22071\)](#)
- [Katz-Mazur1985] Nicholas M. Katz and Barry Mazur, *Arithmetic moduli of elliptic curves*, Annals of Mathematics Studies, vol. 108, Princeton University Press, Princeton, NJ, 1985. [MR772569\(86i:11024\)](#)
- [Knopp-Mason2011] Marvin Knopp and Geoffrey Mason, *Logarithmic vector-valued modular forms*, Acta Arith. **147** (2011), no. 3, 261–262. [MR2773205\(2012c:11094\)](#)
- [Koblitz1984] Neal Koblitz,  *$p$ -adic numbers,  $p$ -adic analysis, and zeta-functions*, second ed., Graduate Texts in Mathematics, vol. 58, Springer-Verlag, New York, 1984. [MR754003\(86c:11086\)](#)
- [Kubert-Lang1981] Daniel S. Kubert and Serge Lang, *Modular units*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Science], vol. 244, Springer-Verlag, New York-Berlin, 1981. [MR648603\(84h:12009\)](#)
- [Lawson-Naumann2014] Tyler Lawson and Niko Naumann, *Strictly commutative realizations of diagrams over the Steenrod algebra and topological modular forms at the prime 2*, Int. Math. Res. Not. IMRN (2014), no. 10, 2773–2813. [MR3214285](#)

- [Lawvere1963] F. William Lawvere, *Functorial semantics of algebraic theories*, Proc. Nat. Acad. Sci. U.S.A. **50** (1963), 869–872. [MR0158921\(28 #2143\)](#)
- [Lazard1955] Michel Lazard, *Sur les groupes de Lie formels à un paramètre*, Bull. Soc. Math. France **83** (1955), 251–274. [MR0073925\(17,508e\)](#)
- [Lubin1967] Jonathan Lubin, *Finite subgroups and isogenies of one-parameter formal Lie groups*, Ann. of Math. (2) **85** (1967), 296–302. [MR0209287\(35 #189\)](#)
- [Lubin-Serre-Tate1964] J. Lubin, J.-P. Serre, and J. Tate, *Elliptic curves and formal groups*, available at <http://www.ma.utexas.edu/users/voloch/lst.html>.
- [Lubin-Tate1966] Jonathan Lubin and John Tate, *Formal moduli for one-parameter formal Lie groups*, Bull. Soc. Math. France **94** (1966), 49–59. [MR0238854\(39 #214\)](#)
- [Mahowald-Rezk2009] Mark Mahowald and Charles Rezk, *Topological modular forms of level 3*, Pure Appl. Math. Q. **5** (2009), no. 2, Special Issue: In honor of Friedrich Hirzebruch. Part 1, 853–872. [MR2508904\(2010g:55010\)](#)
- [Milne2012] J.S. Milne, *Modular functions and modular forms*, available at <http://www.jmilne.org/math/CourseNotes/MF.pdf>.
- [Milnor-Stasheff1974] John W. Milnor and James D. Stasheff, *Characteristic classes*, Princeton University Press, Princeton, N. J.; University of Tokyo Press, Tokyo, 1974, Annals of Mathematics Studies, No. 76. [MR0440554\(55 #13428\)](#)
- [Ono2004] Ken Ono, *The web of modularity: arithmetic of the coefficients of modular forms and  $q$ -series*, CBMS Regional Conference Series in Mathematics, vol. 102, Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 2004. [MR2020489\(2005c:11053\)](#)
- [Ormsby2012] Kyle M. Ormsby, *Tate normal form and level resolutions of the  $K(2)$ -local sphere*, available at <https://dl.dropboxusercontent.com/u/18335215/TateNormalLevelResolutions.pdf>.
- [Poonen2010] Bjorn Poonen, *Supersingular elliptic curves and their “functorial” structure over  $\mathbb{F}_{p^2}$* , MathOverflow, <http://mathoverflow.net/a/19013> (version: 2010-03-22).
- [Rezk2006a] Charles Rezk, *Lectures on power operations*, available at <http://www.math.uiuc.edu/~rezk/power-operation-lectures.dvi>.
- [Rezk2006b] Charles Rezk, *The units of a ring spectrum and a logarithmic cohomology operation*, J. Amer. Math. Soc. **19** (2006), no. 4, 969–1014. [MR2219307\(2007h:55006\)](#)
- [Rezk2008] Charles Rezk, *Power operations for Morava  $E$ -theory of height 2 at the prime 2*, [arXiv:0812.1320](#)
- [Rezk2009] Charles Rezk, *The congruence criterion for power operations in Morava  $E$ -theory*, Homology, Homotopy Appl. **11** (2009), no. 2, 327–379. [MR2591924\(2011e:55021\)](#)
- [Rezk2012] Charles Rezk, *Modular isogeny complexes*, Algebr. Geom. Topol. **12** (2012), no. 3, 1373–1403. [MR2966690](#)

- [Rezk2014] Charles Rezk, *Isogenies, power operations, and homotopy theory*, available at <http://www.math.uiuc.edu/~rezk/rezk-icm-talk-posted.pdf>.
- [Serre1973] Jean-Pierre Serre, *Formes modulaires et fonctions zêta  $p$ -adiques*, Modular functions of one variable, III (Proc. Internat. Summer School, Univ. Antwerp, 1972), Springer, Berlin, 1973, pp. 191–268. Lecture Notes in Math., Vol. 350. [MR0404145\(53#7949a\)](#)
- [Shimura1994] Goro Shimura, *Introduction to the arithmetic theory of automorphic functions*, Publications of the Mathematical Society of Japan, vol. 11, Princeton University Press, Princeton, NJ, 1994, Reprint of the 1971 original, Kanô Memorial Lectures, 1. [MR1291394\(95e:11048\)](#)
- [Silverman2009] Joseph H. Silverman, *The arithmetic of elliptic curves*, second ed., Graduate Texts in Mathematics, vol. 106, Springer, Dordrecht, 2009. [MR2514094\(2010i:11005\)](#)
- [Sprang2014] Johannes Sprang,  *$p$ -adic interpolation and multiplicative orientations of  $KO$  and  $tmf$  (with an appendix by Niko Naumann)*. [arXiv:1409.5314](#)
- [Strickland1998] N. P. Strickland, *Morava  $E$ -theory of symmetric groups*, Topology **37** (1998), no. 4, 757–779. [MR1607736\(99e:55008\)](#)
- [Wilson2015] Dylan Wilson, *Orientations and topological modular forms with level structure*. [arXiv:1507.05116](#)
- [Zagier2008] Don Zagier, *Elliptic modular forms and their applications*, The 1-2-3 of modular forms, Universitext, Springer, Berlin, 2008, pp. 1–103. [MR2409678\(2010b:11047\)](#)
- [Zagier2009] Don Zagier, *Ramanujan’s mock theta functions and their applications [d’après Zwegers and Bringmann-Ono]*, Astérisque (2009), no. 326, Exp. No. 986, vii–viii, 143–164 (2010), Séminaire Bourbaki. Vol. 2007/2008. [MR2605321\(2011h:11049\)](#)
- [Zhu1996] Yongchang Zhu, *Modular invariance of characters of vertex operator algebras*, J. Amer. Math. Soc. **9** (1996), no. 1, 237–302. [MR1317233\(96c:17042\)](#)
- [Zhu2014] Yifei Zhu, *The power operation structure on Morava  $E$ -theory of height 2 at the prime 3*, Algebr. Geom. Topol. **14** (2014), no. 2, 953–977. [MR3160608](#)
- [Zureick-Brown2010] David Zureick-Brown, *Where can I find a comprehensive list of equations for small genus modular curves?*, MathOverflow, <http://mathoverflow.net/q/14054> (version: 2010-02-03).