

MR662118 (83i:55019) 55Q40

Mahowald, Mark

The image of J in the EHP sequence.

Ann. of Math. (2) **116** (1982), no. 1, 65–112.

A more apt title for this very important paper might be “ v_1 -periodic unstable homotopy and the J -spectrum”. The classical EHP sequence and image of J may be an underlying motivation, but seem somewhat peripheral to the theorems proved.

The v_1 -periodicity arose in BP-theory [H. R. Miller, D. C. Ravenel and W. S. Wilson, same journal (2) **106** (1977), no. 3, 469–516; [MR0458423](#)]; in this paper it is given an elegant unstable homotopy-theoretic description. Let Y^n denote the 4-cell complex $\Sigma^{n-6}\mathbf{C}P^2 \wedge \mathbf{R}P^2$, and define $\pi_n(\ ; Y) = [Y^n, \]$, a homotopy functor on topological spaces or spectra. There is a map $v_1: Y^{n+2} \rightarrow Y^n$, such that v_1 acts on $\pi_*(\ ; Y)$. By localizing with respect to v_1 , a theory $V^{-1}\pi_*(\ ; Y)$ is obtained, which is effectively the v_1 -periodic homotopy of a space. The reviewer and the author [Amer. J. Math. **103** (1981), no. 4, 615–659; [MR0623131](#)] discussed the relationship between v_1 -periodic Y -homotopy and ordinary homotopy.

One of the main theorems (1.3) is that the Snaith map $\Omega^{2n}S^{2n+1} \rightarrow \Omega^\infty\Sigma^{\infty+1}P^{2n}$ induces an isomorphism in $V^{-1}\pi_*(\ ; Y)$. Note that the domain is unstable homotopy, while the target is effectively stable homotopy. Both are shown to have precisely four v_1 -periodic families.

Let J denote the fibre of $\psi^3 - 1: bo \rightarrow \Sigma^4 bsp$, where these spectra are localized at 2. It has been known since Adams’ work on $J(X)$ that $\pi_*(J)$ is closely related to the classical image of J . A new discussion of this relationship is given in Section 8. Unfortunately, Theorem 8.4 is marred by serious misprints: The first sentence should say $\mathcal{O}(j) = 3, 2, 5$ if $j \equiv 0, 1, 3 \pmod{8}$. The second sentence need not be included.

The close relationship between J and v_1 -periodicity was already noted [the author and the reviewer, op. cit.]. This relationship is expanded in Theorem 6.2: If X is a finite complex, then $\Omega^\infty\Sigma^\infty X \rightarrow X \wedge J$ induces an isomorphism in $V^{-1}\pi_*(\ ; Y)$. The special case $X = P^{2n}$ is given as 1.4, the statement of which is somewhat misleading in suggesting that it deals with $\pi_*(P^{2n})$ rather than $\pi_*(\Omega^\infty\Sigma^\infty P^{2n})$. When combined with 1.3, the remarkable result that the v_1 -periodic unstable Y -homotopy of odd spheres is isomorphic to that of the easily calculated (stable) $(J \wedge Y)$ -homology of projective spaces is obtained.

Another main result (1.0) is in some sense a limiting version of 1.3, but has the added feature of comparing the homotopy theory $V^{-1}\pi_*(\ ; Y)$ with the homology theory $V^{-1}(J \wedge Y)_*(\)$. It states that there is an isomorphism $V^{-1}\pi_*(\Omega^\infty S^\infty; Y) \rightarrow V^{-1}(J \wedge Y)_*(\mathbf{R}P^\infty)$. Several warnings to the reader are in order here. First, 1.1, of which 1.0 is a corollary, is misleading in that its blanks are not to be filled in with the same spaces. Second, the Y which appears in the homology functor $(J \wedge Y)_*$ is really Y^3 , i.e. its bottom cell has dimension zero, which is different from what one would expect if thinking of the smash product of the spectra J and Y . Third, underlying the switch of Y from the domain (in $\pi_*(\ ; Y)$) to the target (in $(J \wedge Y)_*(\)$) is an application of S -duality and the fact that Y is self-dual, ideas which are not mentioned in the paper.

The fourth main result tells the extent to which $s_*: \pi_i(\Omega^{2n+1}S^{2n+1}) \rightarrow \pi_i(P^{2n} \wedge J)$ is surjective. This is true (1.5) outside the stable range except possibly in one Kervaire-invariant dimension. In the stable range ($i < 2n + 1$) the classes not in the image are given in 7.10. The complete calculation of $\pi_*(P^{2n} \wedge J)$ is given; it is quite elementary. To

show that various families of elements are in the image of s_* requires much work, using the preceding results on Y -homotopy and careful consideration of Adams filtration. The Kervaire-invariant question plays a central role; this will be greatly expanded upon in a series of papers by the author, M. Barratt, and J. Jones.

The proofs involve deep methods of unstable homotopy theory such as the Λ -algebra and unstable resolutions. The important ideas involved more than compensate for the occasional lack of clarity. Much of the work is involved in constructing a homomorphism from a spectral sequence of unstable homotopy groups to one of stable groups and in showing that it is, in fact, an isomorphism.

The author's theorem of bo -resolutions [Pacific J. Math. **92** (1981), no. 2, 365–383; [MR0618072](#)] is used in calculating the v_1 -periodic homotopy groups. An addendum to that paper will appear shortly and will show that minor changes required in that theorem do not affect the applications in this paper.

A section of conjectures and problems appears at the end of the paper. Most deal with torsion and v_1 -torsion in homotopy groups. Another discusses the possibility of mimicking some of the results of this paper for v_2 -periodicity. *Donald M. Davis*

© Copyright American Mathematical Society 1983, 2016