

Algebraic Models in Homotopy Theory

Michael A. Mandell

Indiana University

Colloquium
University of Minnesota



Homotopy Theory

Fundamental Question

Are spaces X and Y homotopy equivalent?



Homotopy Theory

Fundamental Question

Are spaces X and Y homotopy equivalent?

Methods



Homotopy Theory

Fundamental Question

Are spaces X and Y homotopy equivalent?

Methods

- Define algebraic invariants



Homotopy Theory

Fundamental Question

Are spaces X and Y homotopy equivalent?

Methods

- Define algebraic invariants
- Compute those invariants



Homotopy Theory

Fundamental Question

Are spaces X and Y homotopy equivalent?

Methods

- Define algebraic invariants
- Compute those invariants

Is it always possible to find an algebraic invariant that distinguishes between non-equivalent spaces?



Homotopy Theory

Fundamental Question

Are spaces X and Y homotopy equivalent?

Methods

- Define algebraic invariants
- Compute those invariants

Is it always possible to find an algebraic invariant that distinguishes between non-equivalent spaces?

For simply connected spaces: **Yes!**

Cochains
EODGA



Outline

- 1 Homotopy, Homology, and Cohomology
- 2 Warm-up Examples
- 3 Rational Homotopy Theory - CDGAs
- 4 Cochains and E_∞ DGAs
- 5 Homotopy Algebras and Homotopy Theory



Outline

- 1 Homotopy, Homology, and Cohomology
- 2 Warm-up Examples
- 3 Rational Homotopy Theory - CDGAs
- 4 Cochains and E_∞ DGAs
- 5 Homotopy Algebras and Homotopy Theory



Outline

- 1 Homotopy, Homology, and Cohomology
- 2 Warm-up Examples
- 3 Rational Homotopy Theory - CDGAs
- 4 Cochains and E_∞ DGAs
- 5 Homotopy Algebras and Homotopy Theory



Outline

- 1 Homotopy, Homology, and Cohomology
- 2 Warm-up Examples
- 3 Rational Homotopy Theory - CDGAs
- 4 Cochains and E_∞ DGAs
- 5 Homotopy Algebras and Homotopy Theory



Outline

- 1 Homotopy, Homology, and Cohomology
- 2 Warm-up Examples
- 3 Rational Homotopy Theory - CDGAs
- 4 Cochains and E_∞ DGAs
- 5 Homotopy Algebras and Homotopy Theory

← commutative
differential
graded
algebra



Outline

- 1 Homotopy, Homology, and Cohomology
- 2 Warm-up Examples
- 3 Rational Homotopy Theory - CDGAs
- 4 Cochains and E_∞ DGAs
- 5 Homotopy Algebras and Homotopy Theory



Outline

- 1 Homotopy, Homology, and Cohomology
- 2 Warm-up Examples
- 3 Rational Homotopy Theory - CDGAs
- 4 Cochains and E_∞ DGAs
- 5 Homotopy Algebras and Homotopy Theory



Algebraic Topology

Fundamental Question

Are spaces X and Y homotopy equivalent?

It is up to you to produce maps in both directions and homotopies between the composite maps.

Whitehead (1949): This simplifies for “nice” spaces.

Nice spaces: Spaces arising for their geometry
CW complexes. Examples:

- Manifolds
- Polytopes, polyhedra
- Simplicial complexes



Algebraic Topology

Fundamental Question

Are spaces X and Y homotopy equivalent?

It is up to you to produce maps in both directions and homotopies between the composite maps.

Whitehead (1949): This simplifies for “nice” spaces.

Nice spaces: Spaces arising from their geometry
CW complexes. Examples:

- Manifolds
- Polytopes, polyhedra
- Simplicial complexes



Algebraic Topology

Fundamental Question

Are spaces X and Y homotopy equivalent?

It is up to you to produce maps in both directions and homotopies between the composite maps.

Whitehead (1949): This simplifies for “nice” spaces.

Nice spaces: Spaces arising from their geometry
CW complexes. Examples:

- Manifolds
- Polytopes, polyhedra
- Simplicial complexes



Algebraic Topology

Fundamental Question

Are spaces X and Y homotopy equivalent?

It is up to you to produce maps in both directions and homotopies between the composite maps.

Whitehead (1949): This simplifies for “nice” spaces.

Nice spaces: Spaces arising from their geometry
CW complexes. Examples:

- Manifolds
- Polytopes, polyhedra
- Simplicial complexes



Algebraic Topology

Fundamental Question

Are spaces X and Y homotopy equivalent?

It is up to you to produce maps in both directions and homotopies between the composite maps.

Whitehead (1949): This simplifies for “nice” spaces.

Nice spaces: Spaces arising for their geometry
CW complexes. Examples:

- Manifolds
- Polytopes, polyhedra
- Simplicial complexes

Now “space” means “nice space”



Algebraic Topology

Fundamental Question

Are spaces X and Y homotopy equivalent?

It is up to you to produce maps in both directions and homotopies between the composite maps.

Whitehead (1949): This simplifies for “nice” spaces.

Nice spaces: Spaces arising from their geometry
CW complexes. Examples:

- Manifolds
- **Polytopes, polyhedra**
- **Simplicial complexes**

Now “space” means “nice space”
plus maybe a finiteness/compactness hypothesis



The Whitehead Theorem

Theorem (The Whitehead Theorem)

A map $X \rightarrow Y$ is a homotopy equivalence if and only if it induces an isomorphism on homotopy groups.

Given a map, you “just” have to check what happens on some algebraic invariants. But can’t usually compute homotopy groups.



The Whitehead Theorem

Theorem (The Whitehead Theorem)

A map $X \rightarrow Y$ is a homotopy equivalence if and only if it induces an isomorphism on homotopy groups.

Given a map, you “just” have to check what happens on some algebraic invariants. But can't usually compute homotopy groups.



The Whitehead Theorem

Theorem (The Whitehead Theorem)

A map $X \rightarrow Y$ is a homotopy equivalence if and only if it induces an isomorphism on homotopy groups.

Given a map, you “just” have to check what happens on some algebraic invariants. But can’t usually compute homotopy groups.



The Whitehead Theorems

Theorem (The Whitehead Theorem)

A map $X \rightarrow Y$ is a homotopy equivalence if and only if it induces an isomorphism on homotopy groups.

Given a map, you “just” have to check what happens on some algebraic invariants. But can’t usually compute homotopy groups.

Theorem (The Whitehead Theorem II)

A map $X \rightarrow Y$ between simply connected spaces is a homotopy equivalence if and only if it induces an isomorphism on or (equivalently) cohomology.

homology

How much does (co)homology say about a simply connected space?



The Whitehead Theorems

Theorem (The Whitehead Theorem)

A map $X \rightarrow Y$ is a homotopy equivalence if and only if it induces an isomorphism on homotopy groups.

Given a map, you “just” have to check what happens on some algebraic invariants. But can’t usually compute homotopy groups.

Theorem (The Whitehead Theorem II)

A map $X \rightarrow Y$ between simply connected spaces is a homotopy equivalence if and only if it induces an isomorphism on or (equivalently) cohomology.

How much does (co)homology say about a simply connected space?



Example: Homology Spheres

Any simply connected space with the homology/cohomology of the sphere S^n ($n > 1$)

dim	0	1	...	$n-1$	n	$n+1$...
H^*	\mathbb{Z}	0	...	0	\mathbb{Z}	0	...

is homotopy equivalent to the sphere.



Example: Homology Spheres

Any simply connected space with the homology/cohomology of the sphere S^n ($n > 1$)

dim	0	1	...	$n-1$	n	$n+1$...
H^*	\mathbb{Z}	0	...	0	\mathbb{Z}	0	...

is homotopy equivalent to the sphere.



Example: Homology Spheres

Any simply connected space with the homology/cohomology of the sphere S^n ($n > 1$)

dim	0	1	...	$n-1$	n	$n+1$...
H^*	\mathbb{Z}	0	...	0	\mathbb{Z}	0	...

is homotopy equivalent to the sphere.

Theorem (Hurewicz Theorem) *for simply conn spaces*

If $H_q X$ is trivial for $1 \leq q < n$, then the Hurewicz map $\pi_n X \rightarrow H_n X$ is an isomorphism.

then $\pi_q = 0$

$$\underbrace{S^n \rightarrow X}$$



Example: $\mathbb{C}P^2$

$H^*(\mathbb{C}P^2)$ looks like: look like this:

dim	0	1	2	3	4	5	6	7	...
H^*	\mathbb{Z}	0	\mathbb{Z}	0	\mathbb{Z}	0	0	0	...

Other spaces also have cohomology like this, e.g., $S^2 \vee S^4$



Example: $\mathbb{C}P^2$

$H^*(\mathbb{C}P^2)$ looks like: look like this:

dim	0	1	2	3	4	5	6	7	...
H^*	\mathbb{Z}	0	\mathbb{Z}	0	\mathbb{Z}	0	0	0	...

Other spaces also have cohomology like this, e.g., $S^2 \vee S^4$



Example: $\mathbb{C}P^2$

$H^*(\mathbb{C}P^2)$ looks like: look like this:

dim	0	1	2	3	4	5	6	7	...
H^*	\mathbb{Z}	0	\mathbb{Z}	0	\mathbb{Z}	0	0	0	...

Other spaces also have cohomology like this, e.g., $S^2 \vee S^4$

Can distinguish these with the cup product



Example: $\mathbb{C}P^2$

$H^*(\mathbb{C}P^2)$ looks like: look like this:

dim	0	1	2	3	4	5	6	7	...
H^*	\mathbb{Z}	0	\mathbb{Z}	0	\mathbb{Z}	0	0	0	...

x

y



Other spaces also have cohomology like this, e.g., $S^2 \vee S^4$

Can distinguish these with the cup product

dim	0	1	2	3	4	5	6	7	...
$\mathbb{C}P^2$	1	0	x	0	$y = x^2$	0	0	0	...
$S^2 \vee S^4$	1	0	x	0	$y, x^2 = 0$	0	0	0	...



Classification

For every n , there is a space X_n with cohomology

dim	0	1	2	3	4	5	6	7	...
X_n	1	0	x	0	<u>$y, x^2 = ny$</u>	0	0	0	...

Every space with cohomology

\mathbb{Z}	0	\mathbb{Z}	0	\mathbb{Z}	0	0	...
--------------	---	--------------	---	--------------	---	---	-----

is homotopy equivalent to one of these.

$$X_m \simeq X_n \text{ if and only if } m = \pm n$$



Classification

For every n , there is a space X_n with cohomology

dim	0	1	2	3	4	5	6	7	...
X_n	1	0	x	0	$y, x^2 = ny$	0	0	0	...

Every space with cohomology

\mathbb{Z}	0	\mathbb{Z}	0	\mathbb{Z}	0	0	...
--------------	---	--------------	---	--------------	---	---	-----

is homotopy equivalent to one of these.

$$X_m \simeq X_n \text{ if and only if } m = \pm n$$



Classification

For every n , there is a space X_n with cohomology

dim	0	1	2	3	4	5	6	7	...
X_n	1	0	x	0	$y, x^2 = ny$	0	0	0	...

Every space with cohomology

\mathbb{Z}	0	\mathbb{Z}	0	\mathbb{Z}	0	0	...
--------------	---	--------------	---	--------------	---	---	-----


is homotopy equivalent to one of these.

$$X_m \simeq X_n \text{ if and only if } m = \pm n$$

$$S^2$$

$$\pi_3 S^2 = \mathbb{Z}$$


$$S^3 \rightarrow S^2$$



[This page intentionally left blank]

Example $\Sigma \mathbb{C}P^2$

Suspension – take $\mathbb{C}P^2 \times [0, 1]$ and collapse each of $\mathbb{C}P^2 \times \{0\}$ and $\mathbb{C}P^2 \times \{1\}$ to a point.



Example: $\Sigma \mathbb{C}P^2$

Suspension – take $\mathbb{C}P^2 \times [0, 1]$ and collapse each of $\mathbb{C}P \times \{0\}$ and $\mathbb{C}P \times \{1\}$ to a point.

This shifts cohomology groups up.

dim	0	1	2	3	4	5	6	7	8	...
H^*	\mathbb{Z}	0	0	\mathbb{Z}	0	\mathbb{Z}	0	0	0	...

It also kills the cup product.

But not the Steenrod operations on $H^*(-; \mathbb{Z}/2)$.



Example: $\Sigma \mathbb{C}P^2$

Suspension – take $\mathbb{C}P^2 \times [0, 1]$ and collapse each of $\mathbb{C}P \times \{0\}$ and $\mathbb{C}P \times \{1\}$ to a point.

This shifts cohomology groups up.

dim	0	1	2	3	4	5	6	7	8	...
H^*	\mathbb{Z}	0	0	\mathbb{Z}	0	\mathbb{Z}	0	0	0	...

It also kills the cup product.

But not the Steenrod operations on $H^*(-; \mathbb{Z}/2)$.



Example: $\Sigma \mathbb{C}P^2$

Suspension – take $\mathbb{C}P^2 \times [0, 1]$ and collapse each of $\mathbb{C}P \times \{0\}$ and $\mathbb{C}P \times \{1\}$ to a point.

This shifts cohomology groups up.

dim	0	1	2	3	4	5	6	7	8	...
H^*	\mathbb{Z}	0	0	\mathbb{Z}	0	\mathbb{Z}	0	0	0	...

It also kills the cup product.

But not the Steenrod operations on $H^*(-; \mathbb{Z}/2)$.

$$Sq^2 : H^n(X; \mathbb{Z}/2) \rightarrow H^{n+2}(X; \mathbb{Z}/2)$$

In this case remembers
cup product from $\mathbb{C}P^2$



Classification

Every space with cohomology

\mathbb{Z}	0	0	\mathbb{Z}	0	\mathbb{Z}	0	0	...
--------------	---	---	--------------	---	--------------	---	---	-----

is homotopy equivalent to exactly one of $\Sigma \mathbb{C}P^2$ or $S^3 \vee S^5$.

$$\begin{array}{ccccccc}
 \pi_2 & 0 & 0 & \pi_4 & 0 & \pi_6 & 0 & 0 & 0 \\
 & & & \searrow & & & & & \\
 & & & Sq^2 & & & & & \\
 & & & 0 & \text{or} & \mathbb{Z} & & & \\
 & & \swarrow & & \searrow & & & & \\
 & & S^3 \vee S^5 & & \Sigma \mathbb{C}P^2 & & & &
 \end{array}$$

$$\pi_4 S^3 = \mathbb{Z}/2$$



The Steenrod / Grothendieck Problem



The Steenrod / Grothendieck Problem

gen rational hty thry

Problem

Find structure on cohomology or cochains that classifies simply connected spaces up to homotopy equivalence.

Solution is E_∞ DGA

Mandell, “Cochains and Homotopy Type”, *Pub. Math. IHÉS*, 2006.

Problem

Given a homotopy invariant (or property or ???), find a structure on cohomology or cochains that determines it. Or vice-versa.



The Steenrod / Grothendieck Problem

Problem

Find structure on cohomology or cochains that classifies simply connected spaces up to homotopy equivalence.

Solution is E_∞ DGA

Mandell, “Cochains and Homotopy Type”, *Pub. Math. IHÉS*, 2006.

Problem

Given a homotopy invariant (or property or ???), find a structure on cohomology or cochains that determines it. Or vice-versa.



The Steenrod / Grothendieck Problem

Problem

Find structure on cohomology or cochains that classifies simply connected spaces up to homotopy equivalence.

Solution is E_∞ DGA

Mandell, “Cochains and Homotopy Type”, *Pub. Math. IHÉS*, 2006.

Problem

Given a homotopy invariant (or property or ???), find a structure on cohomology or cochains that determines it. Or vice-versa.



Rational Homotopy Theory

The Whitehead Theorem: A map $X \rightarrow Y$ of simply connected space is a homotopy equivalence if and only if it induces an isomorphism on integral homology.

Definition (Rational Equivalence)

A *rational equivalence* is a map $X \rightarrow Y$ that induces an isomorphism on rational homology $H_*(X; \mathbb{Q}) \xrightarrow{\cong} H_*(Y; \mathbb{Q})$ or (equivalently) on rational cohomology $H^*(X; \mathbb{Q}) \xrightarrow{\cong} H^*(Y; \mathbb{Q})$

Rational Homotopy Theory: Make rational equivalences into isomorphisms.

Rational Homotopy Category: Category obtained by formally inverting the rational equivalences.



Rational Homotopy Theory

The Whitehead Theorem: A map $X \rightarrow Y$ of simply connected space is a homotopy equivalence if and only if it induces an isomorphism on integral homology.

Definition (Rational Equivalence)

A rational equivalence is a map $X \rightarrow Y$ that induces an isomorphism on rational homology $H_*(X; \mathbb{Q}) \xrightarrow{\cong} H_*(Y; \mathbb{Q})$ or (equivalently) on rational cohomology $H^*(X; \mathbb{Q}) \xrightarrow{\cong} H^*(Y; \mathbb{Q})$

Rational Homotopy Theory: Make rational equivalences into isomorphisms.

Rational Homotopy Category: Category obtained by formally inverting the rational equivalences.



Rational Homotopy Theory

The Whitehead Theorem: A map $X \rightarrow Y$ of simply connected space is a homotopy equivalence if and only if it induces an isomorphism on integral homology.

Definition (Rational Equivalence)

A *rational equivalence* is a map $X \rightarrow Y$ that induces an isomorphism on rational homology $H_*(X; \mathbb{Q}) \xrightarrow{\cong} H_*(Y; \mathbb{Q})$ or (equivalently) on rational cohomology $H^*(X; \mathbb{Q}) \xrightarrow{\cong} H^*(Y; \mathbb{Q})$

Rational Homotopy Theory: Make rational equivalences into isomorphisms.

Rational Homotopy Category: Category obtained by formally inverting the rational equivalences.



Rational Homotopy Theory

The Whitehead Theorem: A map $X \rightarrow Y$ of simply connected space is a homotopy equivalence if and only if it induces an isomorphism on integral homology.

Definition (Rational Equivalence)

A *rational equivalence* is a map $X \rightarrow Y$ that induces an isomorphism on rational homology $H_*(X; \mathbb{Q}) \xrightarrow{\cong} H_*(Y; \mathbb{Q})$ or (equivalently) on rational cohomology $H^*(X; \mathbb{Q}) \xrightarrow{\cong} H^*(Y; \mathbb{Q})$

Rational Homotopy Theory: Make rational equivalences into isomorphisms.

Rational Homotopy Category: Category obtained by formally inverting the rational equivalences.



Rational Invariants

What information about simply connected spaces is left in the rational homotopy category?

(Anything that takes rational equivalences to isomorphisms)

Lots of rational mapping space data, including *rational homotopy groups*.

$$\pi_n X \otimes \mathbb{Q}$$

More or less anything $\otimes \mathbb{Q}$ that can be computed from spectral sequences.

Serre 1950's: Rational invariants are relatively easy to compute.



Rational Invariants

What information about simply connected spaces is left in the rational homotopy category?

(Anything that takes rational equivalences to isomorphisms)

Lots of rational mapping space data, including rational homotopy groups.

$$\pi_n X \otimes \mathbb{Q}$$

More or less anything $\otimes \mathbb{Q}$ that can be computed from spectral sequences.

Serre 1950's: Rational invariants are relatively easy to compute.



Rational Invariants

What information about simply connected spaces is left in the rational homotopy category?

(Anything that takes rational equivalences to isomorphisms)

Lots of rational mapping space data, including *rational homotopy groups*.

$$\pi_n X \otimes \mathbb{Q}$$

More or less anything $\otimes \mathbb{Q}$ that can be computed from spectral sequences.

Serre 1950's: Rational invariants are relatively easy to compute.



Rational Invariants

What information about simply connected spaces is left in the rational homotopy category?

(Anything that takes rational equivalences to isomorphisms)

Lots of rational mapping space data, including *rational homotopy groups*.

$$\pi_n X \otimes \mathbb{Q}$$

More or less anything $\otimes \mathbb{Q}$ that can be computed from spectral sequences.

Serre 1950's: Rational invariants are relatively easy to compute.

Serre's
E-theory



De Rham complex

$H^*(-; \mathbb{Q})$ or $H^*(-; \mathbb{R})$ have a carrier that is a

commutative differential graded algebra (CDGA) A

The De Rham complex of a manifold $\Omega^* M$

$$\omega \wedge \eta = (-1)^{|\omega||\eta|} \eta \wedge \omega$$

$$d\omega$$

$$d(\omega \wedge \eta) = d\omega \wedge \eta + (-1)^{|\omega|} \omega \wedge d\eta$$



De Rham complex

$H^*(-; \mathbb{Q})$ or $H^*(-; \mathbb{R})$ have a carrier that is a

commutative differential graded algebra (CDGA)

The De Rham complex of a manifold $\Omega^* M$

Piecewise smooth version on a triangulation $\Omega_{PS}^* M$

Piecewise polynomial version using polynomials with coefficients in \mathbb{Q}



De Rham complex

$H^*(-; \mathbb{Q})$ or $H^*(-; \mathbb{R})$ have a carrier that is a

commutative differential graded algebra (CDGA)

The De Rham complex of a manifold $\Omega^* M$

Piecewise smooth version on a triangulation $\Omega_{PS}^* M$

Piecewise polynomial version using polynomials with coefficients in \mathbb{Q}

$$\underbrace{x_0 + \dots + x_n}_{=1}$$



De Rham complex

$H^*(-; \mathbb{Q})$ or $H^*(-; \mathbb{R})$ have a carrier that is a
commutative differential graded algebra (CDGA)

The De Rham complex of a manifold $\Omega^* M$

Piecewise smooth version on a triangulation $\Omega_{PS}^* M$

Piecewise polynomial version using polynomials with coefficients in \mathbb{Q}

\Rightarrow Thom–Sullivan De Rham complex $\Omega_{TS}^* M$ $H^* \Omega_{TS} M \simeq H^*(M; \mathbb{Q})$

Makes sense for any simplicial complex / space.

$$H^*(\Omega_{TS}^* X) \cong H^*(X; \mathbb{Q})$$



De Rham complex

$H^*(-; \mathbb{Q})$ or $H^*(-; \mathbb{R})$ have a carrier that is a
commutative differential graded algebra (CDGA)

The De Rham complex of a manifold $\Omega^* M$

Piecewise smooth version on a triangulation $\Omega_{PS}^* M$

Piecewise polynomial version using polynomials with coefficients in \mathbb{Q}

\implies Thom–Sullivan De Rham complex $\Omega_{TS}^* M \quad \sim$

Makes sense for any simplicial complex / space.

$$\underline{H^*(\Omega_{TS}^* X)} \cong H^*(X; \mathbb{Q})$$



Quillen–Sullivan Theorem

Quasi-isomorphism: A map of CDGAs that induces an isomorphism on cohomology.

equiv. rel. gens

$$A \rightarrow B$$



Quillen–Sullivan Theorem

Quasi-isomorphism: A map of CDGAs that induces an isomorphism on cohomology.

Theorem (Quillen / Sullivan)

Simply connected spaces are rationally equivalent if and only if their Thom–Sullivan De Rham complexes are quasi-isomorphic.

The Thom–Sullivan De Rham complex provides an algebraic model for the rational homotopy type

The rational homotopy groups of X are the André–Quillen cohomology groups of $\Omega_{TS}^* X$.



Quillen–Sullivan Theorem

Quasi-isomorphism: A map of CDGAs that induces an isomorphism on cohomology.

Theorem (Quillen / Sullivan)

Simply connected spaces are rationally equivalent if and only if their Thom–Sullivan De Rham complexes are quasi-isomorphic.

The Thom–Sullivan De Rham complex provides an algebraic model for the rational homotopy type

The rational homotopy groups of X are the André–Quillen cohomology groups of $\Omega_{TS}^* X$.



Quillen–Sullivan Theorem

Quasi-isomorphism: A map of CDGAs that induces an isomorphism on cohomology.

Theorem (Quillen / Sullivan)

Simply connected spaces are rationally equivalent if and only if their Thom–Sullivan De Rham complexes are quasi-isomorphic.

The Thom–Sullivan De Rham complex provides an algebraic model for the rational homotopy type

The rational homotopy groups of X are the André–Quillen cohomology groups of $\Omega_{TS}^* X$.



E_∞ Differential Graded Algebras

An E_∞ DGA is a generalization of a commutative DGA.

Instead of requiring the multiplication to be commutative, require it to be *homotopy* commutative up to “all higher homotopies”



E_∞ Differential Graded Algebras

An E_∞ DGA is a generalization of a commutative DGA.

Instead of requiring the multiplication to be commutative, require it to be *homotopy* commutative up to “all higher homotopies”



E_∞ Differential Graded Algebras

An E_∞ DGA is a generalization of a commutative DGA.


Instead of requiring the multiplication to be commutative, require it to be *homotopy* commutative up to “all higher homotopies”


 $xy \bullet$
 $\bullet yx$


E_∞ Differential Graded Algebras

An E_∞ DGA is a generalization of a commutative DGA.


Instead of requiring the multiplication to be commutative, require it to be *homotopy* commutative up to “all higher homotopies”



E_∞ Differential Graded Algebras

An E_∞ DGA is a generalization of a commutative DGA.


Instead of requiring the multiplication to be commutative, require it to be *homotopy* commutative up to “all higher homotopies”



E_∞ Differential Graded Algebras

An E_∞ DGA is a generalization of a commutative DGA.

Instead of requiring the multiplication to be commutative, require it to be *homotopy* commutative up to “all higher homotopies”



Steenrod Operations

E_∞ DGAs admit Steenrod operations


Working over $\mathbb{Z}/2$,

$$dh^n(x, y) = h^{n-1}(x, y) + h^{n-1}(y, x) + h^n(dx, y) + h^n(x, dy)$$

So for $dx \equiv 0 \pmod{2}$,

$$dh^n(x, x) \equiv h^{n-1}(x, x) + h^{n-1}(x, x) + 0 + 0 \equiv 0 \pmod{2}$$

$h^n(x, x)$ is a mod 2 cycle, represents $Sq^{2|x|-n}x$.



Steenrod Operations

E_∞ DGAs admit Steenrod operations


Working over $\mathbb{Z}/2$,

$$\underline{d}h^n(x, y) = \underline{h^{n-1}(x, y)} + \underline{h^{n-1}(y, x)} + \underline{h^n(dx, y)} + \underline{h^n(x, dy)}$$

So for $dx \equiv 0 \pmod 2$,

$$dh^n(x, x) \equiv h^{n-1}(x, x) + h^{n-1}(x, x) + 0 + 0 \equiv 0 \pmod 2$$

$h^n(x, x)$ is a mod 2 cycle, represents $Sq^{2|x|-n}x$.



Steenrod Operations

E_∞ DGAs admit Steenrod operations


Working over $\mathbb{Z}/2$,

$$dh^n(x, y) = \underline{h^{n-1}(x, y)} + \underline{h^{n-1}(y, x)} + \underline{h^n(dx, y)} + \underline{h^n(x, dy)}$$

So for $dx \equiv 0 \pmod{2}$,

$$dh^n(x, x) \equiv h^{n-1}(x, x) + h^{n-1}(x, x) + 0 + 0 \equiv 0 \pmod{2}$$

$h^n(x, x)$ is a mod 2 cycle, represents $Sq^{2|x|-n}x$.



Steenrod Operations

E_∞ DGAs admit Steenrod operations

Working over $\mathbb{Z}/2$,

$$dh^n(x, y) = h^{n-1}(x, y) + h^{n-1}(y, x) + h^n(dx, y) + h^n(x, dy)$$


So for $dx \equiv 0 \pmod{2}$,

$$dh^n(x, x) \equiv h^{n-1}(x, x) + h^{n-1}(x, x) + 0 + 0 \equiv 0 \pmod{2}$$

$h^n(x, x)$ is a mod 2 cycle, represents $Sq^{2|x|-n}x$.

For CDGA $L^n = 0$

so Sq^i is zero



Cochains and E_∞ DGAs

The simplicial (or singular) cochain complex is naturally an E_∞ DGA.

Theorem

Any functor to chain complexes or E_∞ DGAs that satisfies a dimension hypothesis and a weak gluing condition is naturally quasi-isomorphic to the cochain functor with some coefficients.

Example

The Thom–Sullivan De Rham complex $\Omega_{TS}^* X$ is naturally quasi-isomorphic to $C^*(X; \mathbb{Q})$ through maps of E_∞ DGAs.

Consequence

No carrier for integral cohomology can be a CDGA.



Cochains and E_∞ DGAs

The simplicial (or singular) cochain complex is naturally an E_∞ DGA.

Theorem

Any functor to chain complexes or E_∞ DGAs that satisfies a dimension hypothesis and a weak gluing condition is naturally quasi-isomorphic to the cochain functor with some coefficients. *why cond*

Example

The Thom–Sullivan De Rham complex $\Omega_{TS}^* X$ is naturally quasi-isomorphic to $C^*(X; \mathbb{Q})$ through maps of E_∞ DGAs.

Consequence

No carrier for integral cohomology can be a CDGA.



Cochains and E_∞ DGAs

The simplicial (or singular) cochain complex is naturally an E_∞ DGA.

Theorem

Any functor to chain complexes or E_∞ DGAs that satisfies a dimension hypothesis and a weak gluing condition is naturally quasi-isomorphic to the cochain functor with some coefficients.

Example

The Thom–Sullivan De Rham complex $\Omega_{TS}^* X$ is naturally quasi-isomorphic to $C^*(X; \mathbb{Q})$ through maps of E_∞ DGAs.

Consequence

No carrier for integral cohomology can be a CDGA.



Cochains and E_∞ DGAs

The simplicial (or singular) cochain complex is naturally an E_∞ DGA.

Theorem

Any functor to chain complexes or E_∞ DGAs that satisfies a dimension hypothesis and a weak gluing condition is naturally quasi-isomorphic to the cochain functor with some coefficients.

Example

The Thom–Sullivan De Rham complex $\Omega_{TS}^* X$ is naturally quasi-isomorphic to $C^*(X; \mathbb{Q})$ through maps of E_∞ DGAs.

Consequence

No carrier for integral cohomology can be a CDGA.



Algebraic Models for Homotopy Theory

Theorem (2006)

Simply connected spaces are homotopy equivalent if and only if their cochain E_∞ DGAs are quasi-isomorphic.

The cochain complex as an E_∞ DGA provides an algebraic model for homotopy types.

“Can” compute homotopy groups using (e.g.) analogue of the method of Cartan–Serre.

Example

$C^*(S^2)$ easy to describe as an E_∞ DGA. Beyond a certain range, higher homotopy groups are unknown.



Algebraic Models for Homotopy Theory

Theorem

Simply connected spaces are homotopy equivalent if and only if their cochain E_∞ DGAs are quasi-isomorphic.

The cochain complex as an E_∞ DGA provides an algebraic model for homotopy types.

“Can” compute homotopy groups using (e.g.) analogue of the method of Cartan–Serre.

Example

$C^*(S^2)$ easy to describe as an E_∞ DGA. Beyond a certain range, higher homotopy groups are unknown.



Algebraic Models for Homotopy Theory

Theorem

Simply connected spaces are homotopy equivalent if and only if their cochain E_∞ DGAs are quasi-isomorphic.

The cochain complex as an E_∞ DGA provides an algebraic model for homotopy types.

“Can” compute homotopy groups using (e.g.) analogue of the method of Cartan–Serre.

Example

$C^*(S^2)$ easy to describe as an E_∞ DGA. Beyond a certain range, higher homotopy groups are unknown.



Algebraic Models for Homotopy Theory

Theorem

Simply connected spaces are homotopy equivalent if and only if their cochain E_∞ DGAs are quasi-isomorphic.

The cochain complex as an E_∞ DGA provides an algebraic model for homotopy types.

“Can” compute homotopy groups using (e.g.) analogue of the method of Cartan–Serre.

Example

$C^*(S^2)$ easy to describe as an E_∞ DGA. Beyond a certain range, higher homotopy groups are unknown.



Homotopy Algebras and Homotopy Theory

Hierarchy of algebraic structures encoding higher homotopies of commutativity.

$$E_1 \text{ DGAs} \subset E_2 \text{ DGAs} \subset E_3 \text{ DGAs} \subset \cdots \subset E_\infty \text{ DGAs}$$

E_1 DGAs are associative DGAs

E_2 DGAs are homotopy commutative plus a little more

Concise definition in terms of brace operations $x\{y_1, \dots, y_n\}$
 ($x\{y\}$ is the commutativity homotopy)

E_3 and higher are “even more homotopy commutative”



Homotopy Algebras and Homotopy Theory

Hierarchy of algebraic structures encoding higher homotopies of commutativity.

$$E_1 \text{ DGAs} \subset E_2 \text{ DGAs} \subset E_3 \text{ DGAs} \subset \cdots \subset E_\infty \text{ DGAs}$$

E_1 DGAs are associative DGAs

E_2 DGAs are homotopy commutative plus a little more

Concise definition in terms of brace operations $x\{y_1, \dots, y_n\}$
 ($x\{y\}$ is the commutativity homotopy)

E_3 and higher are “even more homotopy commutative”



Homotopy Algebras and Homotopy Theory

Hierarchy of algebraic structures encoding higher homotopies of commutativity.

$$E_1 \text{ DGAs} \subset E_2 \text{ DGAs} \subset E_3 \text{ DGAs} \subset \cdots \subset E_\infty \text{ DGAs}$$

E_1 DGAs are associative DGAs

E_2 DGAs are homotopy commutative plus a little more

Concise definition in terms of brace operations $x\{y_1, \dots, y_n\}$
 ($x\{y\}$ is the commutativity homotopy)

E_3 and higher are “even more homotopy commutative”



Homotopy Algebras and Homotopy Theory

Hierarchy of algebraic structures encoding higher homotopies of commutativity.

$$E_1 \text{ DGAs} \subset E_2 \text{ DGAs} \subset E_3 \text{ DGAs} \subset \cdots \subset E_\infty \text{ DGAs}$$

E_1 DGAs are associative DGAs

E_2 DGAs are homotopy commutative plus a little more

Concise definition in terms of brace operations $x\{y_1, \dots, y_n\}$
 ($x\{y\}$ is the commutativity homotopy)

E_3 and higher are “even more homotopy commutative”



Homotopy Algebras and Homotopy Theory

Hierarchy of algebraic structures encoding higher homotopies of commutativity.

$$E_1 \text{ DGAs} \subset E_2 \text{ DGAs} \subset E_3 \text{ DGAs} \subset \cdots \subset E_\infty \text{ DGAs}$$

E_1 DGAs are associative DGAs

E_2 DGAs are homotopy commutative plus a little more

Concise definition in terms of brace operations $x\{y_1, \dots, y_n\}$
 ($x\{y\}$ is the commutativity homotopy)

E_3 and higher are “even more homotopy commutative”



Invariants of the E_2 Structure

When we regard C^*X as an E_2 DGA, what information about a simply connected space X remains?

- Homology / cohomology of based loop space as a Hopf algebra
- Homology / cohomology of the free loop space as an H^*X -module.
- Homology / cohomology of mapping space X^M where $M = T^2$ or $\Sigma_g^2 \setminus \{p_1, \dots, p_n\}$, $n \geq 1$.



Invariants of the E_2 Structure

When we regard C^*X as an E_2 DGA, what information about a simply connected space X remains?

- Homology / cohomology of based loop space as a Hopf algebra
- Homology / cohomology of the free loop space as an H^*X -module.
- Homology / cohomology of mapping space X^M where $M = T^2$ or $\Sigma_g^2 \setminus \{p_1, \dots, p_n\}$, $n \geq 1$.



Invariants of the E_2 Structure

When we regard C^*X as an E_2 DGA, what information about a simply connected space X remains?

- Homology / cohomology of based loop space as a Hopf algebra
In particular, the rational homotopy groups
- Homology / cohomology of the free loop space as an H^*X -module.
- Homology / cohomology of mapping space X^M where $M = T^2$ or $\Sigma_g^2 \setminus \{p_1, \dots, p_n\}$, $n \geq 1$.



Invariants of the E_2 Structure

When we regard C^*X as an E_2 DGA, what information about a simply connected space X remains?

- Homology / cohomology of based loop space as a Hopf algebra
In particular, the rational homotopy groups
- Homology / cohomology of the free loop space as an H^*X -module.
- Homology / cohomology of mapping space X^M where $M = T^2$ or $\Sigma_g^2 \setminus \{p_1, \dots, p_n\}$, $n \geq 1$.



Invariants of the E_2 Structure


When we regard C^*X as an E_2 DGA, what information about a simply connected space X remains?

- Homology / cohomology of based loop space as a Hopf algebra
In particular, the rational homotopy groups
- Homology / cohomology of the free loop space as an H^*X -module.
- Homology / cohomology of mapping space X^M where $M = T^2$ or $\Sigma_g^2 \setminus \{p_1, \dots, p_n\}$, $n \geq 1$.



Invariants of the E_2 Structure

When we regard C^*X as an E_2 DGA, what information about a simply connected space X remains?

- Homology / cohomology of based loop space as a Hopf algebra
In particular, the rational homotopy groups
- Homology / cohomology of the free loop space as an H^*X -module.
- Homology / cohomology of mapping space X^M where $M = T^2$ or $\Sigma_g^2 \setminus \{p_1, \dots, p_n\}$, $n \geq 1$.
- For X a PD space, the *string topology* BV algebra 



E_2 DGAs, DG Hopf Algebras, and Anick's Theorem

Gerstenhaber–Voronov (IMRN 1995): An E_2 DGA is pretty much the same thing as a DG Hopf algebra.

Bar Construction

E_2 DGA structure on $A \iff$ DG Hopf algebra structure on BA [+/-]

Anick (JAMS 1989) studied BC^*X as a (DG) “Hopf algebra up to homotopy” and proved (for primes p)

Theorem (Anick)

*If X is at least c -connected ($c \geq 1$) and at most pd -dimensional, then after inverting $1, \dots, p-1$ and changing the multiplication up to homotopy, BC^*X is dual to the universal enveloping algebra of a DG Lie algebra.*



E_2 DGAs, DG Hopf Algebras, and Anick's Theorem

Gerstenhaber–Voronov (IMRN 1995): An E_2 DGA is pretty much the same thing as a DG Hopf algebra.

Bar Construction

E_2 DGA structure on $A \iff$ DG Hopf algebra structure on BA [+/-]

Anick (JAMS 1989) studied BC^*X as a (DG) “Hopf algebra up to homotopy” and proved (for primes p)

Theorem (Anick)

*If X is at least c -connected ($c \geq 1$) and at most pd -dimensional, then after inverting $1, \dots, p-1$ and changing the multiplication up to homotopy, BC^*X is dual to the universal enveloping algebra of a DG Lie algebra.*



E_2 DGAs, DG Hopf Algebras, and Anick's Theorem

Gerstenhaber–Voronov (IMRN 1995): An E_2 DGA is pretty much the same thing as a DG Hopf algebra.

Bar Construction

E_2 DGA structure on $A \iff$ DG Hopf algebra structure on BA [+/-]

Anick (JAMS 1989) studied BC^*X as a (DG) “Hopf algebra up to homotopy” and proved (for primes p)

Theorem (Anick)

*If X is at least c -connected ($c \geq 1$) and at most p -dimensional, then after inverting $1, \dots, p-1$ and changing the multiplication up to homotopy, BC^*X is dual to the universal enveloping algebra of a DG Lie algebra.*



An E_2 Version of Anick's Theorem

Conjecture

*If X is at least c -connected ($c \geq 1$) and at most p -dimensional, then after inverting $1, \dots, p-1$, the E_2 DGA C^*X is quasi-isomorphic to a commutative DGA.*

Consequences

- For highly connected / low dimensional spaces, the E_2 information is relatively accessible.
- On a fixed space, for all but finitely primes, the E_2 information is relatively accessible.



An E_2 Version of Anick's Theorem

Conjecture

*If X is at least c -connected ($c \geq 1$) and at most pd -dimensional, then after inverting $1, \dots, p-1$, the E_2 DGA C^*X is quasi-isomorphic to a commutative DGA.*

Consequences

- For highly connected / low dimensional spaces, the E_2 information is relatively accessible.
- On a fixed space, for all but finitely primes, the E_2 information is relatively accessible.



An E_2 Version of Anick's Theorem

Conjecture

*If X is at least c -connected ($c \geq 1$) and at most pd -dimensional, then after inverting $1, \dots, p-1$, the E_2 DGA C^*X is quasi-isomorphic to a commutative DGA.*

Consequences

- For highly connected / low dimensional spaces, the E_2 information is relatively accessible.
- On a fixed space, for all but finitely primes, the E_2 information is relatively accessible.

