

**MR1669268 (2000e:55012)** 55Q40 55N91 55Q10 55Q51

**Arone, Greg (1-CHI); Mahowald, Mark (1-NW)**

**The Goodwillie tower of the identity functor and the unstable periodic homotopy of spheres. (English summary)**

*Invent. Math.* **135** (1999), no. 3, 743–788.

# FEATURED REVIEW.

This is an important and beautiful paper. On the one hand, it confirms the senior author's decades-old vision of how unstable homotopy made "periodic" can be calculated from a small amount of stable information. On the other hand, it illustrates how T. Goodwillie's philosophy of how to resolve homotopy functors is leading to a new generation of combinatorial models in homotopy theory, particularly when applied by the junior author.

To set the stage, Goodwillie's previous work (partially available in [*K-Theory* **4** (1990), no. 1, 1–27; [MR1076523](#); *K-Theory* **5** (1991/92), no. 4, 295–332; [MR1162445](#); "Calculus. III. The Taylor series of a homotopy functor", in preparation]) implies that, given a pointed space  $X$ , there is a unique natural tower of fibrations

$$\cdots \rightarrow P_n(X) \rightarrow P_{n-1}(X) \rightarrow \cdots \rightarrow P_1(X) = QX$$

having the following properties. First, there is a map from  $X$  to the inverse limit, and this map is a homotopy equivalence if  $X$  is connected. Second, there are spectra  $C_n$  with an action of the  $n$ th symmetric group  $\Sigma_n$ , so that  $D_n(X)$ , the fiber of  $P_n(X) \rightarrow P_{n-1}(X)$ , is naturally equivalent to  $\Omega^\infty((C_n \wedge X^{\wedge n})_{h\Sigma_n})$ .

B. Johnson [*Trans. Amer. Math. Soc.* **347** (1995), no. 4, 1295–1321; [MR1297532](#)] had identified  $C_n$  as the equivariant  $S$ -dual of an explicit finite-dimensional  $\Sigma_n$ -space  $\Delta_n$ , so that  $D_n(X) \simeq \Omega^\infty(\text{Map}_S(\Delta_n, X^{\wedge n})_{h\Sigma_n})$ , where  $\text{Map}_S$  denotes function spectrum.

The first thing done in the paper under review is to replace  $\Delta_n$  by a different model better suited for homotopical analysis. Let  $\tilde{K}_n$  denote the category of partitions of a fixed set with  $n$  elements, with the initial and terminal partitions discarded. Let  $K_n$  denote the unreduced suspension of the classifying space of  $\tilde{K}_n$ . Then there is a  $\Sigma_n$ -equivariant map  $SK_n \rightarrow \Delta_n$  which is a nonequivariant equivalence. Thus  $D_n(X) \simeq \Omega^\infty(\text{Map}_S(SK_n, X^{\wedge n})_{h\Sigma_n})$ .

The authors now specialize to the case when  $X$  is an odd-dimensional sphere. By filtering  $K_n$  by skeleta, and computing with homology with rational coefficients and then with mod  $p$  coefficients, they are able to deduce that  $\text{Map}_S(SK_n, S^{n(2s+1)})_{h\Sigma_n}$ , and thus  $D_n(S^{2s+1})$ , is contractible unless  $n$  is a power of a prime  $p$ . Furthermore,  $D_{p^i}(S^{2a+1})$  is  $p$ -local, and, with  $L(i, s)$  denoting  $\text{Map}_S(SK_{p^i}, S^{n(2s+1)})_{h\Sigma_{p^i}}$ ,  $H^*(L(i, s); \mathbf{Z}/p)$  is free over the subalgebra  $A(i-1)$  of the mod  $p$  Steenrod algebra  $A$ .

An immediate consequence of this last fact is that  $D_{p^i}(S^{2a+1})$  has trivial  $v_k$ -periodic homotopy if  $i > k$ . Summarizing, with  $R_{i,s} = P_{p^i}(S^{2s+1})$  and with the  $v_k^{-1}$ -localization functor applied to everything, there is a finite tower  $R_{k,s} \rightarrow R_{k-1,s} \rightarrow \cdots \rightarrow R_{0,s}$  with  $S^{2s+1} \rightarrow R_k$  a  $v_k$ -periodic homotopy equivalence, and with the  $i$ th fiber equal to the 0th space of an explicit spectrum  $L(i, s)$ .

When  $k = 0$  we learn that  $S^{2s+1} \rightarrow QS^{2s+1}$  is a rational equivalence, known since the 1950s. When  $k = 1$  and  $p = 2$ , we learn that  $S^{2s+1}$  is equivalent in  $v_1$ -periodic homotopy to the fiber of the Hopf invariant  $QS^{2s+1} \rightarrow Q(\mathbf{RP}^\infty/\mathbf{RP}^{2s})$ . This was shown previously in major work by Mahowald [*Ann. of Math.* (2) **116** (1982), no. 1, 65–112; [MR0662118](#)] and Mahowald and R. D. Thompson [*Topology* **31** (1992), no. 1, 133–141; [MR1153241](#)].

Making the story even more complete, and giving a second proof of the most technical parts of the paper under review, is subsequent work by Arone and W. Dwyer [“Partition complexes, Tits buildings, and symmetric products”, *Proc. London Math. Soc.*, to appear]. They identify the spectra  $L(i, s)$  as suitable Thom spectra over the “Steinberg module” part of  $B(\mathbf{Z}/p)^i$ . Such a result was presaged by work in the early 1980s by the reviewer, S. Priddy, and S. Mitchell [see, e.g., N. J. Kuhn and S. B. Priddy, *Math. Proc. Cambridge Philos. Soc.* **98** (1985), no. 3, 459–480; [MR0803606](#)]. This older work implies, among other things, that the spectral sequence for computing  $\pi_*(S^1)$  associated to the tower of  $R_{k,0}$ ’s collapses at  $E_2$ . It would be very interesting to know if anything like this carries over to the spectral sequences that calculate  $\pi_*(S^{2s+1})$  for  $s \geq 1$ .

*N. J. Kuhn*

---

### References

1. Anderson, D.W., Davis, D.W.: A vanishing theorem in homological algebra. *Comment. Math. Helv.* **48**, 318–327 (1973) [MR0334207](#)
2. Adem, A., Milgram, J.: *Cohomology of finite groups*. Grundlehren der mathematischen Wissenschaften 309, Springer-Verlag (1995) [MR1317096](#)
3. Arone, G., Kankaanrinta, M.: A functorial model for iterated Snaith splitting with applications to calculus of functors. *Stable and Unstable Homotopy*, Fields Institute Communications, Vol. **19**, 1–30 (1998) [MR1622334](#)
4. Arone, G., Dwyer, W.: Partition posets, Tits buildings and symmetric product spectra (tentative title). in preparation.
5. Bousfield, A.K., Kan, D.M.: *Homotopy limits, completions and localizations*. Springer Lecture Notes, Vol. 304 (1972) [MR0365573](#)
6. Bruner, P.R., May, J.P., McClure, J.E., Steinberg, M.:  $H_\infty$  ring spectra and their applications. Springer Lecture Notes, Vol. 1176 (1986) [MR0836132](#)
7. Cohen, F.R., Lada, T.J., May, J.P.: *The homology of iterated loop spaces*. Springer Lecture Notes, Vol. 533 (1976) [MR0436146](#)
8. Goodwillie, T.G.: Calculus I: the first derivative of pseudoisotopy theory. *K-Theory* **4**, 1–27 (1990) [MR1076523](#)
9. Goodwillie, T.G.: Calculus II: analytic functors. *K-Theory* **5**, 295–332 (1992) [MR1162445](#)
10. Goodwillie, T.G.: Calculus III: the Taylor series of a homotopy functor. in preparation
11. James, I.M.: Reduced product spaces. *Annals of Math.* **62**, 170–197 (1953) [MR0073181](#)
12. Johnson, B.: The derivatives of homotopy theory. *Trans. Amer. Math. Soc.* **347**, Number 4, 1295–1321 (1995) [MR1297532](#)
13. Kahn, D.S., Priddy, S.B.: On the transfer in the homology of symmetric groups. *Math. Proc. Cambridge Philos. Soc.* **83**, 91–101 (1978) [MR0464229](#)
14. Kuhn, N.J.: A Kahn-Priddy sequence and a conjecture of G.W. Whitehead. *Math. Proc. Cambridge Philos. Soc.* **92**, 467–483 (1982) [MR0677471](#)
15. Kuhn, N.J.: Chevalley group theory and the transfer in the homology of symmetric groups. *Topology* **24**, 247–264 (1985) [MR0815479](#)
16. Kuhn, N.J., Priddy, S.B.: The transfer and Whitehead conjecture. *Math. Proc. Cambridge Philos. Soc.* **98**, 459–480 (1985) [MR0803606](#)
17. Mahowald, M.: The image of J in the EHP sequence. *Annals Math* **116**, 65–112 (1982) [MR0662118](#)
18. Mahowald, M., Sadofsky, H.:  $v_n$  telescopes and the Adams spectral sequence. *Duke Math J.* **78**, 101–129 (1995) [MR1328754](#)

19. Mahowald, M., Thompson, R.: The K-theory localization of an unstable sphere. *Topology* **31**, 133–141 (1992) [MR1153241](#)
20. Mahowald, M., Thompson, R.: On the secondary suspension homomorphism. *Amer. J. of Math.* **116**, 179–206 (1994) [MR1262430](#)
21. Margolis, H.R.: Spectra and the Steenrod algebra. North Holland Mathematical Library [MR0738973](#)
22. Miller, H., Wilkerson, C.: Vanishing lines for modules over the Steenrod algebra. *J. Pure Appl. Algebra* **22**, 293–307 (1981) [MR0629336](#)
23. Milnor, J.: On the construction FK. in "Algebraic Topology: A Student's Guide" by J.F. Adams, London Math. Soc. Lecture Note Series 4, Cambridge Univ. Press, Cambridge 119–135 (1972) [MR0445484](#)
24. Nakaoka, M.: Cohomology mod  $p$  of symmetric products of spheres. *J. Inst. Poly., Osaka City Univ.* 9 1–18 (1958) [MR0121788](#)
25. Thompson, R.: The  $v_1$ -periodic homotopy groups of an unstable sphere at odd primes. *Trans. Am. Math. Soc.* **319**, 535–559 (1990) [MR1010890](#)
26. Welcher, P.J.: Symmetric fiber spectra and  $K(n)$ -homology acyclicity. *Indiana Univ. Math. J.* **30**, 801–812 (1981) [MR0632853](#)

*Note: This list reflects references listed in the original paper as accurately as possible with no attempt to correct errors.*

© Copyright American Mathematical Society 2000, 2016