

Longwater watch report

Watch closed · resilient

A one-time snapshot of this Longwater game’s accepted journal. Recorded from Longwater’s game simulation. Net changes include the action, the tide and dawn drift.

This report can be read or printed without the game. It does not resume a watch or update when play continues.

Opening → final readings

Freshwater: 5 → 1. Seed packs: 5 → 1.

Across the completed watch

Cell	Depth (cm)	Salt (ppt)	Oxygen (%)	Life (%)	Canopy (/ 3)
North Bank	34 → 100	34 → 9	64 → 64	44 → 63	0 → 2
Heart Pool	56 → 100	30 → 10	55 → 42	49 → 54	0 → 2
South Reach	67 → 100	40 → 19	43 → 19	38 → 46	0 → 2

Opening readings

Freshwater: 5. Seed packs: 5.

Before the first tide

Cell	Depth (cm)	Salt (ppt)	Oxygen (%)	Life (%)	Canopy (/ 3)
North Bank	34	34	64	44	0
Heart Pool	56	30	55	49	0
South Reach	67	40	43	38	0

Tide 1 · Gate · North Bank

Spring tide

A long lunar push carries brine inland.

Full field notes

- Freshwater gate · North Bank: water 34 → 41 cm, salt 34 → 24 ppt, oxygen 64 → 63%. Freshwater −1 (4 left).
- Spring tide · all three cells: water +14 cm, salt +4 ppt; North Bank oxygen 63 → 60%. A long lunar push carries brine inland.
- Dawn drift · suitable depth/salt/oxygen → biomass +1, dawn oxygen +2; the water-depth/salinity rule shifts salt 0 ppt.
- Net · North Bank: water 34 → 55 cm, salt 34 → 28 ppt, oxygen 64 → 62%, biomass 44 → 45%.

Freshwater: 5 → 4. Seed packs: 5 → 5.

Before → after tide 1

Cell	Depth (cm)	Salt (ppt)	Oxygen (%)	Life (%)	Canopy (/ 3)
North Bank	34 → 55	34 → 28	64 → 62	44 → 45	0 → 0
Heart Pool	56 → 70	30 → 34	55 → 54	49 → 50	0 → 0
South Reach	67 → 81	40 → 43	43 → 40	38 → 38	0 → 0

Tide 2 · Shade · Heart Pool

Neap ebb

A short, quiet ebb leaves pools exposed.

Full field notes

- Reed-bed shade · Heart Pool: canopy 0 → 1, salt 34 → 32 ppt, oxygen 54 → 58%, biomass 50 → 53%. Canopy blunts heat and drying.
- Neap ebb · all three cells: water -5 cm, salt +2 ppt; Heart Pool oxygen 58 → 61%. A short, quiet ebb leaves pools exposed.

- Dawn drift · suitable depth/salt/oxygen → biomass +1, dawn oxygen +2; the water-depth/salinity rule shifts salt 0 ppt.
- Net · Heart Pool: water 70 → 65 cm, salt 34 → 34 ppt, oxygen 54 → 63%, biomass 50 → 54%.

Freshwater: 4 → 4. Seed packs: 5 → 5.

Before → after tide 2

Cell	Depth (cm)	Salt (ppt)	Oxygen (%)	Life (%)	Canopy (/ 3)
North Bank	55 → 50	28 → 30	62 → 67	45 → 46	0 → 0
Heart Pool	70 → 65	34 → 34	54 → 63	50 → 54	0 → 1
South Reach	81 → 76	43 → 45	40 → 43	38 → 38	0 → 0

Tide 3 · Seed · South Reach

Heat pulse

Hot air accelerates evaporation and oxygen loss.

Full field notes

- Reseeding · South Reach: biomass 38 → 50% (+12; deep-water settling rule not triggered), oxygen 43 → 44%. Seed satchels −1 (4 left).
- Heat pulse · all three cells: water -8 cm, salt +7 ppt; South Reach oxygen 44 → 37% (canopy relief +0). Hot air accelerates evaporation and oxygen loss.
- Dawn drift · suitable depth/salt/oxygen → biomass +1, dawn oxygen +2; the water-depth/salinity rule shifts salt 0 ppt.
- Net · South Reach: water 76 → 68 cm, salt 45 → 52 ppt, oxygen 43 → 39%, biomass 38 → 49%.

Freshwater: 4 → 4. Seed packs: 5 → 4.

Before → after tide 3

Cell	Depth (cm)	Salt (ppt)	Oxygen (%)	Life (%)	Canopy (/ 3)
North Bank	50 → 42	30 → 37	67 → 62	46 → 45	0 → 0
Heart Pool	65 → 57	34 → 41	63 → 60	54 → 54	1 → 1
South Reach	76 → 68	45 → 52	43 → 39	38 → 49	0 → 0

Tide 4 · Shade · North Bank

Runoff plume

Fresh upland runoff dilutes the channel edge.

Full field notes

- Reed-bed shade · North Bank: canopy 0 → 1, salt 37 → 35 ppt, oxygen 62 → 66%, biomass 45 → 48%. Canopy blunts heat and drying.
- Runoff plume · all three cells: water +9 cm, salt -11 ppt; North Bank oxygen 66 → 68%. Fresh upland runoff dilutes the channel edge.
- Dawn drift · suitable depth/salt/oxygen → biomass +1, dawn oxygen +2; the water-depth/salinity rule shifts salt 0 ppt.
- Net · North Bank: water 42 → 51 cm, salt 37 → 24 ppt, oxygen 62 → 70%, biomass 45 → 49%.

Freshwater: 4 → 4. Seed packs: 4 → 4.

Before → after tide 4

Cell	Depth (cm)	Salt (ppt)	Oxygen (%)	Life (%)	Canopy (/ 3)
North Bank	42 → 51	37 → 24	62 → 70	45 → 49	0 → 1
Heart Pool	57 → 66	41 → 30	60 → 64	54 → 55	1 → 1
South Reach	68 → 77	52 → 41	39 → 41	49 → 49	0 → 0

Tide 5 · Shade · South Reach

Slackwater

A gentle ebb gives the reed shelves room to breathe.

Full field notes

- Reed-bed shade · South Reach: canopy 0 → 1, salt 41 → 39 ppt, oxygen 41 → 45%, biomass 49 → 52%. Canopy blunts heat and drying.
- Slackwater · all three cells: water -6 cm, salt +1 ppt; South Reach oxygen 45 → 48%. A gentle ebb gives the reed shelves room to breathe.

- Dawn drift · suitable depth/salt/oxygen → biomass +1, dawn oxygen +2; the water-depth/salinity rule shifts salt 0 ppt.
- Net · South Reach: water 77 → 71 cm, salt 41 → 40 ppt, oxygen 41 → 50%, biomass 49 → 53%.

Freshwater: 4 → 4. Seed packs: 4 → 4.

Before → after tide 5

Cell	Depth (cm)	Salt (ppt)	Oxygen (%)	Life (%)	Canopy (/ 3)
North Bank	51 → 45	24 → 25	70 → 75	49 → 50	1 → 1
Heart Pool	66 → 60	30 → 31	64 → 69	55 → 56	1 → 1
South Reach	77 → 71	41 → 40	41 → 50	49 → 53	0 → 1

Tide 6 · Gate · Heart Pool

King tide

An unusually high tide presses across the low marsh.

Full field notes

- Freshwater gate · Heart Pool: water 60 → 67 cm, salt 31 → 21 ppt, oxygen 69 → 68%. Freshwater -1 (3 left).
- King tide · all three cells: water +18 cm, salt +5 ppt; Heart Pool oxygen 68 → 62%. An unusually high tide presses across the low marsh.
- Dawn drift · no growth window → biomass 0, dawn oxygen -2; the water-depth/salinity rule shifts salt -1 ppt.
- Net · Heart Pool: water 60 → 85 cm, salt 31 → 25 ppt, oxygen 69 → 60%, biomass 56 → 56%.

Freshwater: 4 → 3. Seed packs: 4 → 4.

Before → after tide 6

Cell	Depth (cm)	Salt (ppt)	Oxygen (%)	Life (%)	Canopy (/ 3)
North Bank	45 → 63	25 → 30	75 → 71	50 → 51	1 → 1
Heart Pool	60 → 85	31 → 25	69 → 60	56 → 56	1 → 1
South Reach	71 → 89	40 → 44	50 → 42	53 → 51	1 → 1

Tide 7 · Seed · North Bank

Fog veil

Cool fog slows drying and softens the morning light.

Full field notes

- Reseeding · North Bank: biomass 51 → 63% (+12; deep-water settling rule not triggered), oxygen 71 → 72%. Seed satchels -1 (3 left).
- Fog veil · all three cells: water +1 cm, salt -1 ppt; North Bank oxygen 72 → 76%. Cool fog slows drying and softens the morning light.
- Dawn drift · suitable depth/salt/oxygen → biomass +1, dawn oxygen +2; the water-depth/salinity rule shifts salt 0 ppt.
- Net · North Bank: water 63 → 64 cm, salt 30 → 29 ppt, oxygen 71 → 78%, biomass 51 → 64%.

Freshwater: 3 → 3. Seed packs: 4 → 3.

Before → after tide 7

Cell	Depth (cm)	Salt (ppt)	Oxygen (%)	Life (%)	Canopy (/ 3)
North Bank	63 → 64	30 → 29	71 → 78	51 → 64	1 → 1
Heart Pool	85 → 86	25 → 23	60 → 62	56 → 56	1 → 1
South Reach	89 → 90	44 → 42	42 → 44	51 → 49	1 → 1

Tide 8 · Shade · Heart Pool

Dry wind

A warm offshore wind pulls water from the pools.

Full field notes

- Reed-bed shade · Heart Pool: canopy 1 → 2, salt 23 → 21 ppt, oxygen 62 → 66%, biomass 56 → 59%. Canopy blunts heat and drying.
- Dry wind · all three cells: water -7 cm, salt +6 ppt; Heart Pool oxygen 66 → 65% (canopy relief +4). A warm offshore wind pulls water from the pools.

- Dawn drift · no growth window → biomass 0, dawn oxygen 0; the water-depth/salinity rule shifts salt -1 ppt.
- Net · Heart Pool: water 86 → 79 cm, salt 23 → 26 ppt, oxygen 62 → 65%, biomass 56 → 59%.

Freshwater: 3 → 3. Seed packs: 3 → 3.

Before → after tide 8

Cell	Depth (cm)	Salt (ppt)	Oxygen (%)	Life (%)	Canopy (/ 3)
North Bank	64 → 57	29 → 35	78 → 77	64 → 64	1 → 1
Heart Pool	86 → 79	23 → 26	62 → 65	56 → 59	1 → 2
South Reach	90 → 83	42 → 47	44 → 39	49 → 48	1 → 1

Tide 9 · Seed · Heart Pool

Runoff pulse

A second freshwater pulse pushes through the inlet.

Full field notes

- Reseeding · Heart Pool: biomass 59 → 67% (+8; deep-water settling rule applied), oxygen 65 → 66%. Seed satchels -1 (2 left).
- Runoff pulse · all three cells: water +10 cm, salt -10 ppt; Heart Pool oxygen 66 → 67%. A second freshwater pulse pushes through the inlet.
- Dawn drift · stress boundary reached → biomass -2, dawn oxygen -2; the water-depth/salinity rule shifts salt -1 ppt.
- Net · Heart Pool: water 79 → 89 cm, salt 26 → 15 ppt, oxygen 65 → 65%, biomass 59 → 65%.

Freshwater: 3 → 3. Seed packs: 3 → 2.

Before → after tide 9

Cell	Depth (cm)	Salt (ppt)	Oxygen (%)	Life (%)	Canopy (/ 3)
North Bank	57 → 67	35 → 25	77 → 80	64 → 65	1 → 1
Heart Pool	79 → 89	26 → 15	65 → 65	59 → 65	2 → 2
South Reach	83 → 93	47 → 36	39 → 38	48 → 46	1 → 1

Tide 10 · Shade · North Bank

Spring tide

The next spring tide reaches the lower reed line.

Full field notes

- Reed-bed shade · North Bank: canopy 1 → 2, salt 25 → 23 ppt, oxygen 80 → 84%, biomass 65 → 68%. Canopy blunts heat and drying.
- Spring tide · all three cells: water +15 cm, salt +4 ppt; North Bank oxygen 84 → 80%. The next spring tide reaches the lower reed line.
- Dawn drift · no growth window → biomass 0, dawn oxygen 0; the water-depth/salinity rule shifts salt -1 ppt.
- Net · North Bank: water 67 → 82 cm, salt 25 → 26 ppt, oxygen 80 → 80%, biomass 65 → 68%.

Freshwater: 3 → 3. Seed packs: 2 → 2.

Before → after tide 10

Cell	Depth (cm)	Salt (ppt)	Oxygen (%)	Life (%)	Canopy (/ 3)
North Bank	67 → 82	25 → 26	80 → 80	65 → 68	1 → 2
Heart Pool	89 → 100	15 → 18	65 → 59	65 → 63	2 → 2
South Reach	93 → 100	36 → 39	38 → 32	46 → 44	1 → 1

Tide 11 · Gate · South Reach

Silt ribbon

Fine silt clouds the water before settling on the flats.

Full field notes

- Freshwater gate · South Reach: water 100 → 100 cm, salt 39 → 29 ppt, oxygen 32 → 31%. Freshwater -1 (2 left).
- Silt ribbon · all three cells: water +5 cm, salt -5 ppt; South Reach oxygen 31 → 28%. Fine silt clouds the water before settling on the flats.

- Dawn drift · stress boundary reached → biomass -2, dawn oxygen -2; the water-depth/salinity rule shifts salt -1 ppt.
- Net · South Reach: water 100 → 100 cm, salt 39 → 23 ppt, oxygen 32 → 26%, biomass 44 → 41%.

Freshwater: 3 → 2. Seed packs: 2 → 2.

Before → after tide 11

Cell	Depth (cm)	Salt (ppt)	Oxygen (%)	Life (%)	Canopy (/ 3)
North Bank	82 → 87	26 → 20	80 → 75	68 → 67	2 → 2
Heart Pool	100 → 100	18 → 12	59 → 54	63 → 60	2 → 2
South Reach	100 → 100	39 → 23	32 → 26	44 → 41	1 → 1

Tide 12 · Shade · South Reach

Heat pulse

A second hot spell tests every shaded cell.

Full field notes

- Reed-bed shade · South Reach: canopy 1 → 2, salt 23 → 21 ppt, oxygen 26 → 30%, biomass 41 → 44%. Canopy blunts heat and drying.
- Heat pulse · all three cells: water -9 cm, salt +8 ppt; South Reach oxygen 30 → 27% (canopy relief +4). A second hot spell tests every shaded cell.
- Dawn drift · stress boundary reached → biomass -2, dawn oxygen -2; the water-depth/salinity rule shifts salt -1 ppt.
- Net · South Reach: water 100 → 91 cm, salt 23 → 28 ppt, oxygen 26 → 25%, biomass 41 → 42%.

Freshwater: 2 → 2. Seed packs: 2 → 2.

Before → after tide 12

Cell	Depth (cm)	Salt (ppt)	Oxygen (%)	Life (%)	Canopy (/ 3)
North Bank	87 → 78	20 → 28	75 → 72	67 → 67	2 → 2
Heart Pool	100 → 91	12 → 19	54 → 49	60 → 58	2 → 2

Cell	Depth (cm)	Salt (ppt)	Oxygen (%)	Life (%)	Canopy (/ 3)
South Reach	100 → 91	23 → 28	26 → 25	41 → 42	1 → 2

Tide 13 · Seed · South Reach

Freshwater pulse

The last runoff pulse clears a narrow freshwater lane.

Full field notes

- Reseeding · South Reach: biomass 42 → 50% (+8; deep-water settling rule applied), oxygen 25 → 26%. Seed satchels –1 (1 left).
- Freshwater pulse · all three cells: water +12 cm, salt -12 ppt; South Reach oxygen 26 → 28%. The last runoff pulse clears a narrow freshwater lane.
- Dawn drift · stress boundary reached → biomass -2, dawn oxygen -2; the water-depth/salinity rule shifts salt -1 ppt.
- Net · South Reach: water 91 → 100 cm, salt 28 → 15 ppt, oxygen 25 → 26%, biomass 42 → 48%.

Freshwater: 2 → 2. Seed packs: 2 → 1.

Before → after tide 13

Cell	Depth (cm)	Salt (ppt)	Oxygen (%)	Life (%)	Canopy (/ 3)
North Bank	78 → 90	28 → 15	72 → 72	67 → 65	2 → 2
Heart Pool	91 → 100	19 → 6	49 → 49	58 → 56	2 → 2
South Reach	91 → 100	28 → 15	25 → 26	42 → 48	2 → 2

Tide 14 · Gate · North Bank

Moon-pull tide

A final high tide closes the fourteen-day watch.

Full field notes

- Freshwater gate · North Bank: water 90 → 97 cm, salt 15 → 5 ppt, oxygen 72 → 71%. Freshwater −1 (1 left).
- Moon-pull tide · all three cells: water +16 cm, salt +5 ppt; North Bank oxygen 71 → 66%. A final high tide closes the fourteen-day watch.
- Dawn drift · stress boundary reached → biomass -2, dawn oxygen -2; the water-depth/salinity rule shifts salt -1 ppt.
- Net · North Bank: water 90 → 100 cm, salt 15 → 9 ppt, oxygen 72 → 64%, biomass 65 → 63%.

Freshwater: 2 → 1. Seed packs: 1 → 1.

Before → after tide 14

Cell	Depth (cm)	Salt (ppt)	Oxygen (%)	Life (%)	Canopy (/ 3)
North Bank	90 → 100	15 → 9	72 → 64	65 → 63	2 → 2
Heart Pool	100 → 100	6 → 10	49 → 42	56 → 54	2 → 2
South Reach	100 → 100	15 → 19	26 → 19	48 → 46	2 → 2

14 of 14 tides recorded. Only completed tides are included. Print using your browser’s Print command.