

# The semantic account of formal consequence, from Alfred Tarski back to John Buridan

Jacob Archambault

Fordham University

31 Oct 2016 / Logic and Metaphysics workshop, CUNY  
Graduate Center

# Outline

- 1 Introduction
- 2 From classical consequence back to Tarski
  - Classical consequence today
  - Tarskian consequence
    - Tarskian consequences and domain variation
- 3 Buridan's theory of formal consequence
  - Preliminaries
  - Formal and material consequence
- 4 Formal consequence from Tarski back to John Buridan

# Outline

- 1 Introduction
- 2 From classical consequence back to Tarski
  - Classical consequence today
  - Tarskian consequence
    - Tarskian consequences and domain variation
- 3 Buridan's theory of formal consequence
  - Preliminaries
  - Formal and material consequence
- 4 Formal consequence from Tarski back to John Buridan

# Outline

- 1 Introduction
- 2 From classical consequence back to Tarski
  - Classical consequence today
  - Tarskian consequence
    - Tarskian consequences and domain variation
- 3 Buridan's theory of formal consequence
  - Preliminaries
  - Formal and material consequence
- 4 Formal consequence from Tarski back to John Buridan

# Outline

- 1 Introduction
- 2 From classical consequence back to Tarski
  - Classical consequence today
  - Tarskian consequence
    - Tarskian consequences and domain variation
- 3 Buridan's theory of formal consequence
  - Preliminaries
  - Formal and material consequence
- 4 Formal consequence from Tarski back to John Buridan

# Outline

- 1 Introduction
- 2 From classical consequence back to Tarski
  - Classical consequence today
  - Tarskian consequence
    - Tarskian consequences and domain variation
- 3 Buridan's theory of formal consequence
  - Preliminaries
  - Formal and material consequence
- 4 Formal consequence from Tarski back to John Buridan

# Classical consequence today: models

- $M = (D, I)$
- $D$  is a collection of objects: cats and dogs, numbers, members of the Medici family, or whatever else one likes.
- The interpretation function  $I$  then assigns:
  - each name  $a$  to an element in the domain  $D$
  - each  $n$ -ary relation symbol  $R^n$  to a subset of  $D^n$
- Besides interpretations, one also has *valuations*  $\{v, v' \dots\}$  on  $M$ , each of which assigns values in  $D$  to all the variables of the language  $L$

# Classical consequence today: models

- $M = (D, I)$
- $D$  is a collection of objects: cats and dogs, numbers, members of the Medici family, or whatever else one likes.
- The interpretation function  $I$  then assigns:
  - each name  $a$  to an element in the domain  $D$
  - each  $n$ -ary relation symbol  $R^n$  to a subset of  $D^n$
- Besides interpretations, one also has *valuations*  $\{v, v' \dots\}$  on  $M$ , each of which assigns values in  $D$  to all the variables of the language  $L$



# Classical consequence today: models

- $M = (D, I)$
- $D$  is a collection of objects: cats and dogs, numbers, members of the Medici family, or whatever else one likes.
- The interpretation function  $I$  then assigns:
  - each name  $a$  to an element in the domain  $D$
  - each  $n$ -ary relation symbol  $R^n$  to a subset of  $D^n$
- Besides interpretations, one also has *valuations*  $\{v, v' \dots\}$  on  $M$ , each of which assigns values in  $D$  to all the variables of the language  $L$

# Classical consequence today: models

- $M = (D, I)$
- $D$  is a collection of objects: cats and dogs, numbers, members of the Medici family, or whatever else one likes.
- The interpretation function  $I$  then assigns:
  - each name  $a$  to an element in the domain  $D$
  - each  $n$ -ary relation symbol  $R^n$  to a subset of  $D^n$
- Besides interpretations, one also has *valuations*  $\{v, v' \dots\}$  on  $M$ , each of which assigns values in  $D$  to all the variables of the language  $L$

# Classical consequence today: models

- $M = (D, I)$
- $D$  is a collection of objects: cats and dogs, numbers, members of the Medici family, or whatever else one likes.
- The interpretation function  $I$  then assigns:
  - each name  $a$  to an element in the domain  $D$
  - each  $n$ -ary relation symbol  $R^n$  to a subset of  $D^n$
- Besides interpretations, one also has *valuations*  $\{v, v' \dots\}$  on  $M$ , each of which assigns values in  $D$  to all the variables of the language  $L$

# Classical consequence today: models

- $M = (D, I)$
- $D$  is a collection of objects: cats and dogs, numbers, members of the Medici family, or whatever else one likes.
- The interpretation function  $I$  then assigns:
  - each name  $a$  to an element in the domain  $D$
  - each  $n$ -ary relation symbol  $R^n$  to a subset of  $D^n$
- Besides interpretations, one also has *valuations*  $\{v, v' \dots\}$  on  $M$ , each of which assigns values in  $D$  to all the variables of the language  $L$

# Classical Consequence today: semantics

Let  $(v \star I)$  be the operation such that  $v \star I$  agrees with  $v$  on the assignment of variables, and with  $I$  on the assignment of terms, and write  $(v \star I)(t_1, \dots, t_n)$  for  $((v \star I)(t_1), \dots, (v \star I)(t_n))$ . The truth value of different formulae in a model  $M$  - i.e. *truth in a model* - is then recursively determined as follows:

# Classical consequence today: semantics

- 1 For atomic formulae of arity  $n$ ,  $(v \star I)(Rt_1 \dots t_n) = T$  iff  $(v \star I)(t_1, \dots, t_n) \in I(R)$
- 2  $(v \star I)(\neg \phi) = T$  iff  $(v \star I)(\phi) = F$
- 3  $(v \star I)(\phi \wedge \psi) = T$  iff  $(v \star I)(\phi) = (v \star I)(\psi) = T$
- 4  $(v \star I)(\phi \vee \psi) = T$  iff  $(v \star I)(\phi) = T$  or  $(v \star I)(\psi) = T$
- 5  $(v \star I)(\phi \supset \psi) = T$  iff  $(v \star I)(\phi) = F$  or  $(v \star I)(\psi) = T$
- 6  $(v \star I)(\phi \equiv \psi) = T$  iff  $(v \star I)(\phi) = (v \star I)(\psi)$
- 7  $(v \star I)(\forall x \phi) = T$  iff, for every  $x$ -variant  $v'$  of  $v$ ,  $(v' \star I)(\phi) = T$
- 8  $(v \star I)(\exists x \phi) = T$  iff, for some  $x$ -variant  $v'$  of  $v$ ,  $(v' \star I)(\phi) = T$

# Classical consequence today: semantics

- 1 For atomic formulae of arity  $n$ ,  $(v \star I)(Rt_1 \dots t_n) = T$  iff  $(v \star I)(t_1, \dots t_n) \in I(R)$
- 2  $(v \star I)(\neg \phi) = T$  iff  $(v \star I)(\phi) = F$
- 3  $(v \star I)(\phi \wedge \psi) = T$  iff  $(v \star I)(\phi) = (v \star I)(\psi) = T$
- 4  $(v \star I)(\phi \vee \psi) = T$  iff  $(v \star I)(\phi) = T$  or  $(v \star I)(\psi) = T$
- 5  $(v \star I)(\phi \supset \psi) = T$  iff  $(v \star I)(\phi) = F$  or  $(v \star I)(\psi) = T$
- 6  $(v \star I)(\phi \equiv \psi) = T$  iff  $(v \star I)(\phi) = (v \star I)(\psi)$
- 7  $(v \star I)(\forall x \phi) = T$  iff, for every  $x$ -variant  $v'$  of  $v$ ,  $(v' \star I)(\phi) = T$
- 8  $(v \star I)(\exists x \phi) = T$  iff, for some  $x$ -variant  $v'$  of  $v$ ,  $(v' \star I)(\phi) = T$

# Classical consequence today: semantics

- 1 For atomic formulae of arity  $n$ ,  $(v \star I)(Rt_1 \dots t_n) = T$  iff  $(v \star I)(t_1, \dots, t_n) \in I(R)$
- 2  $(v \star I)(\neg \phi) = T$  iff  $(v \star I)(\phi) = F$
- 3  $(v \star I)(\phi \wedge \psi) = T$  iff  $(v \star I)(\phi) = (v \star I)(\psi) = T$
- 4  $(v \star I)(\phi \vee \psi) = T$  iff  $(v \star I)(\phi) = T$  or  $(v \star I)(\psi) = T$
- 5  $(v \star I)(\phi \supset \psi) = T$  iff  $(v \star I)(\phi) = F$  or  $(v \star I)(\psi) = T$
- 6  $(v \star I)(\phi \equiv \psi) = T$  iff  $(v \star I)(\phi) = (v \star I)(\psi)$
- 7  $(v \star I)(\forall x \phi) = T$  iff, for every  $x$ -variant  $v'$  of  $v$ ,  $(v' \star I)(\phi) = T$
- 8  $(v \star I)(\exists x \phi) = T$  iff, for some  $x$ -variant  $v'$  of  $v$ ,  $(v' \star I)(\phi) = T$



# Classical consequence today: semantics

- 1 For atomic formulae of arity  $n$ ,  $(v \star I)(Rt_1 \dots t_n) = T$  iff  $(v \star I)(t_1, \dots t_n) \in I(R)$
- 2  $(v \star I)(\neg \phi) = T$  iff  $(v \star I)(\phi) = F$
- 3  $(v \star I)(\phi \wedge \psi) = T$  iff  $(v \star I)(\phi) = (v \star I)(\psi) = T$
- 4  $(v \star I)(\phi \vee \psi) = T$  iff  $(v \star I)(\phi) = T$  or  $(v \star I)(\psi) = T$
- 5  $(v \star I)(\phi \supset \psi) = T$  iff  $(v \star I)(\phi) = F$  or  $(v \star I)(\psi) = T$
- 6  $(v \star I)(\phi \equiv \psi) = T$  iff  $(v \star I)(\phi) = (v \star I)(\psi)$
- 7  $(v \star I)(\forall x \phi) = T$  iff, for every  $x$ -variant  $v'$  of  $v$ ,  $(v' \star I)(\phi) = T$
- 8  $(v \star I)(\exists x \phi) = T$  iff, for some  $x$ -variant  $v'$  of  $v$ ,  $(v' \star I)(\phi) = T$

# Classical consequence today: semantics

- 1 For atomic formulae of arity  $n$ ,  $(v \star I)(Rt_1 \dots t_n) = T$  iff  $(v \star I)(t_1, \dots, t_n) \in I(R)$
- 2  $(v \star I)(\neg \phi) = T$  iff  $(v \star I)(\phi) = F$
- 3  $(v \star I)(\phi \wedge \psi) = T$  iff  $(v \star I)(\phi) = (v \star I)(\psi) = T$
- 4  $(v \star I)(\phi \vee \psi) = T$  iff  $(v \star I)(\phi) = T$  or  $(v \star I)(\psi) = T$
- 5  $(v \star I)(\phi \supset \psi) = T$  iff  $(v \star I)(\phi) = F$  or  $(v \star I)(\psi) = T$
- 6  $(v \star I)(\phi \equiv \psi) = T$  iff  $(v \star I)(\phi) = (v \star I)(\psi)$
- 7  $(v \star I)(\forall x \phi) = T$  iff, for every  $x$ -variant  $v'$  of  $v$ ,  $(v' \star I)(\phi) = T$
- 8  $(v \star I)(\exists x \phi) = T$  iff, for some  $x$ -variant  $v'$  of  $v$ ,  $(v' \star I)(\phi) = T$

# Classical consequence today: semantics

- 1 For atomic formulae of arity  $n$ ,  $(v \star I)(Rt_1 \dots t_n) = T$  iff  $(v \star I)(t_1, \dots, t_n) \in I(R)$
- 2  $(v \star I)(\neg \phi) = T$  iff  $(v \star I)(\phi) = F$
- 3  $(v \star I)(\phi \wedge \psi) = T$  iff  $(v \star I)(\phi) = (v \star I)(\psi) = T$
- 4  $(v \star I)(\phi \vee \psi) = T$  iff  $(v \star I)(\phi) = T$  or  $(v \star I)(\psi) = T$
- 5  $(v \star I)(\phi \supset \psi) = T$  iff  $(v \star I)(\phi) = F$  or  $(v \star I)(\psi) = T$
- 6  $(v \star I)(\phi \equiv \psi) = T$  iff  $(v \star I)(\phi) = (v \star I)(\psi)$
- 7  $(v \star I)(\forall x \phi) = T$  iff, for every  $x$ -variant  $v'$  of  $v$ ,  $(v' \star I)(\phi) = T$
- 8  $(v \star I)(\exists x \phi) = T$  iff, for some  $x$ -variant  $v'$  of  $v$ ,  $(v' \star I)(\phi) = T$

# Classical consequence today: semantics

- 1 For atomic formulae of arity  $n$ ,  $(v \star I)(Rt_1 \dots t_n) = T$  iff  $(v \star I)(t_1, \dots t_n) \in I(R)$
- 2  $(v \star I)(\neg \phi) = T$  iff  $(v \star I)(\phi) = F$
- 3  $(v \star I)(\phi \wedge \psi) = T$  iff  $(v \star I)(\phi) = (v \star I)(\psi) = T$
- 4  $(v \star I)(\phi \vee \psi) = T$  iff  $(v \star I)(\phi) = T$  or  $(v \star I)(\psi) = T$
- 5  $(v \star I)(\phi \supset \psi) = T$  iff  $(v \star I)(\phi) = F$  or  $(v \star I)(\psi) = T$
- 6  $(v \star I)(\phi \equiv \psi) = T$  iff  $(v \star I)(\phi) = (v \star I)(\psi)$
- 7  $(v \star I)(\forall x \phi) = T$  iff, for every  $x$ -variant  $v'$  of  $v$ ,  $(v' \star I)(\phi) = T$
- 8  $(v \star I)(\exists x \phi) = T$  iff, for some  $x$ -variant  $v'$  of  $v$ ,  $(v' \star I)(\phi) = T$

# Classical consequence today: semantics

- 1 For atomic formulae of arity  $n$ ,  $(v \star I)(Rt_1 \dots t_n) = T$  iff  $(v \star I)(t_1, \dots, t_n) \in I(R)$
- 2  $(v \star I)(\neg \phi) = T$  iff  $(v \star I)(\phi) = F$
- 3  $(v \star I)(\phi \wedge \psi) = T$  iff  $(v \star I)(\phi) = (v \star I)(\psi) = T$
- 4  $(v \star I)(\phi \vee \psi) = T$  iff  $(v \star I)(\phi) = T$  or  $(v \star I)(\psi) = T$
- 5  $(v \star I)(\phi \supset \psi) = T$  iff  $(v \star I)(\phi) = F$  or  $(v \star I)(\psi) = T$
- 6  $(v \star I)(\phi \equiv \psi) = T$  iff  $(v \star I)(\phi) = (v \star I)(\psi)$
- 7  $(v \star I)(\forall x \phi) = T$  iff, for every  $x$ -variant  $v'$  of  $v$ ,  $(v' \star I)(\phi) = T$
- 8  $(v \star I)(\exists x \phi) = T$  iff, for some  $x$ -variant  $v'$  of  $v$ ,  $(v' \star I)(\phi) = T$

# Classical consequence today: formal consequence

- A sentence  $\phi$  is *satisfiable* if it is made true on some model  $M$ . When this happens,  $M$  is said to be a *model* of  $\phi$ .
- $\Gamma$  is satisfiable iff there is some model  $M$  on which every sentence in  $\Gamma$  is satisfiable. Similarly when this occurs,  $M$  is said to be a model of  $\Gamma$ .
- A sentence  $\phi$  is said to be a logical, or formal, consequence of a set  $\Gamma$ , written  $\Gamma \models \phi$ , iff every model of  $\Gamma$  is a model of  $\phi$ .

# Classical consequence today: formal consequence

- A sentence  $\phi$  is *satisfiable* if it is made true on some model  $M$ . When this happens,  $M$  is said to be a *model* of  $\phi$ .
- $\Gamma$  is satisfiable iff there is some model  $M$  on which every sentence in  $\Gamma$  is satisfiable. Similarly when this occurs,  $M$  is said to be a model of  $\Gamma$ .
- A sentence  $\phi$  is said to be a logical, or formal, consequence of a set  $\Gamma$ , written  $\Gamma \models \phi$ , iff every model of  $\Gamma$  is a model of  $\phi$ .

# Classical consequence today: formal consequence

- A sentence  $\phi$  is *satisfiable* if it is made true on some model  $M$ . When this happens,  $M$  is said to be a *model* of  $\phi$ .
- $\Gamma$  is satisfiable iff there is some model  $M$  on which every sentence in  $\Gamma$  is satisfiable. Similarly when this occurs,  $M$  is said to be a model of  $\Gamma$ .
- A sentence  $\phi$  is said to be a logical, or formal, consequence of a set  $\Gamma$ , written  $\Gamma \models \phi$ , iff every model of  $\Gamma$  is a model of  $\phi$ .



# Outline

- 1 Introduction
- 2 From classical consequence back to Tarski
  - Classical consequence today
  - **Tarskian consequence**
    - Tarskian consequences and domain variation
- 3 Buridan's theory of formal consequence
  - Preliminaries
  - Formal and material consequence
- 4 Formal consequence from Tarski back to John Buridan

# Tarskian formal consequence

- Both Tarskian and classical semantic accounts determine formal consequence by way of variation
- Classical languages are left uninterpreted. Tarskian languages are not.
- Where classical consequences varies the interpretations of non-logical constants, Tarski's replaces these with variables.
- A Tarskian model does not include an interpretation. It is simply a sequence of objects

# Tarskian formal consequence

- Both Tarskian and classical semantic accounts determine formal consequence by way of variation
- Classical languages are left uninterpreted. Tarskian languages are not.
- Where classical consequences varies the interpretations of non-logical constants, Tarski's replaces these with variables.
- A Tarskian model does not include an interpretation. It is simply a sequence of objects

# Tarskian formal consequence

- Both Tarskian and classical semantic accounts determine formal consequence by way of variation
- Classical languages are left uninterpreted. Tarskian languages are not.
- Where classical consequences varies the interpretations of non-logical constants, Tarski's replaces these with variables.
- A Tarskian model does not include an interpretation. It is simply a sequence of objects

# Tarskian formal consequence

- Both Tarskian and classical semantic accounts determine formal consequence by way of variation
- Classical languages are left uninterpreted. Tarskian languages are not.
- Where classical consequences varies the interpretations of non-logical constants, Tarski's replaces these with variables.
- A Tarskian model does not include an interpretation. It is simply a sequence of objects

# Tarskian formal consequence

*Let us assume that, in the language which we are considering, to each extra-logical constant correspond certain variable symbols, and this in such a way that, by replacing in an arbitrary sentence a constant by a corresponding variable, we transform this sentence into a sentential function. Let us further consider an arbitrary class of sentences  $L$ , and let us replace all extra-logical constants occurring in the sentences of class  $L$  by corresponding variables (equiform constants by equiform variables, non-equiform by non-equiform); we shall obtain a class of sentential functions  $L'$ .*

# Tarskian formal consequence

*An arbitrary sequence of objects which satisfies each sentential function of the class  $L'$  we shall call a model of the class  $L$  (in just this sense one usually speaks about a model of the system of axioms of a deductive theory); if in particular the class  $L$  consists of only one sentence  $X$ , we will simply speak about a model of the sentence  $X$  (Tarski 2002, pp. 185-186).*

# Tarskian consequences and domain variation

- Tarski's pre-WWII mathematical work makes use of variable domains.
- Where Tarski's use of domain variation is explicit in his post-WWII discussions of consequence, he refers his reader to his earlier work unproblematically.
- Tarski's broadest later discussion of the issues of his 1936 paper also assumes an invariant domain.
- All of this suggests a high level of continuity between Tarski's earlier and later work on the subject.



# Tarskian consequences and domain variation

- Tarski's pre-WWII mathematical work makes use of variable domains.
- Where Tarski's use of domain variation is explicit in his post-WWII discussions of consequence, he refers his reader to his earlier work unproblematically.
- Tarski's broadest later discussion of the issues of his 1936 paper also assumes an invariant domain.
- All of this suggests a high level of continuity between Tarski's earlier and later work on the subject.

# Tarskian consequences and domain variation

- Tarski's pre-WWII mathematical work makes use of variable domains.
- Where Tarski's use of domain variation is explicit in his post-WWII discussions of consequence, he refers his reader to his earlier work unproblematically.
- Tarski's broadest later discussion of the issues of his 1936 paper also assumes an invariant domain.
- All of this suggests a high level of continuity between Tarski's earlier and later work on the subject.

# Tarskian consequences and domain variation

- Tarski's pre-WWII mathematical work makes use of variable domains.
- Where Tarski's use of domain variation is explicit in his post-WWII discussions of consequence, he refers his reader to his earlier work unproblematically.
- Tarski's broadest later discussion of the issues of his 1936 paper also assumes an invariant domain.
- All of this suggests a high level of continuity between Tarski's earlier and later work on the subject.

# Tarskian consequences and domain variation

- Tarski's pre-WWII mathematical work makes use of variable domains.
- Where Tarski's use of domain variation is explicit in his post-WWII discussions of consequence, he refers his reader to his earlier work unproblematically.
- Tarski's broadest later discussion of the issues of his 1936 paper also assumes an invariant domain.
- All of this suggests a high level of continuity between Tarski's earlier and later work on the subject.

# Outline

- 1 Introduction
- 2 From classical consequence back to Tarski
  - Classical consequence today
  - Tarskian consequence
    - Tarskian consequences and domain variation
- 3 Buridan's theory of formal consequence
  - Preliminaries
  - Formal and material consequence
- 4 Formal consequence from Tarski back to John Buridan

# Buridan's Treatise on Consequences

- Book 1: assertoric consequences.
- Book 2: modal consequences
- Book 3: assertoric syllogistic
- Book 4: modal syllogistic

# Buridan's Treatise on Consequences: Book 1

- 1 Truth and falsity of propositions
- 2 Causes of truth
- 3 Definition of consequence
- 4 Division of consequence
- 5 Supposition of terms
- 6 Ampliation of terms
- 7 The matter and form of propositions
- 8 Listing of assertoric consequences

# Buridanian consequence: preliminaries

- Buridan's causes of truth play a role analogous to Tarskian models
- Buridan retains modality and tense
- Propositions remain within the range of semantic referents
- Truthmaker semantics



# Buridanian consequence: preliminaries

- Buridan's causes of truth play a role analogous to Tarskian models
- Buridan retains modality and tense
- Propositions remain within the range of semantic referents
- Truthmaker semantics

# Buridanian consequence: preliminaries

- Buridan's causes of truth play a role analogous to Tarskian models
- Buridan retains modality and tense
- Propositions remain within the range of semantic referents
- Truthmaker semantics

# Buridanian consequence: preliminaries

- Buridan's causes of truth play a role analogous to Tarskian models
- Buridan retains modality and tense
- Propositions remain within the range of semantic referents
- Truthmaker semantics

# Buridanian consequence: preliminaries

*If Colin's horse, which cantered well, is dead, 'Colin's horse cantered well' ... is true because things were in reality as the proposition signifies they were. In the same way 'The Antichrist will preach' is true, not because things are in reality as the proposition signifies, but because things will be in reality as the proposition signifies they will be. Similarly, 'Something that never will be can be' is true, not because things are as the proposition signifies, but because things can be as it signifies them to be.' And so it is clear that it is necessary to assign causes of truth to different types of propositions in different ways [TC I. 1, p. 63].*

# Buridanian consequence: preliminaries

*I say that both terms being undistributed but suppositing determinately, then there are more causes of truth than if one were distributed and the other confused without distribution. This is clear because every cause of truth enough to make 'Every B is A' true is enough to make 'B is A' true, but not vice versa. Therefore a proposition [1] has most causes of truth with each term undistributed [2] and fewer with one term distributed and the other confused without distribution, [3] and fewer still with one distributed and the other used determinately without distribution, [4] and fewest of all with both distributed [TC I. 2, p. 66].*

# Outline

- 1 Introduction
- 2 From classical consequence back to Tarski
  - Classical consequence today
  - Tarskian consequence
    - Tarskian consequences and domain variation
- 3 Buridan's theory of formal consequence
  - Preliminaries
  - Formal and material consequence
- 4 Formal consequence from Tarski back to John Buridan

# Formal and material consequence

*A consequence is a hypothetical proposition composed of an antecedent and consequent, indicating the antecedent to be antecedent and the consequent to be consequent; this designation occurs by the word 'if' or by the word 'therefore' or an equivalent [TC I. 3, p. 67].*

# Formal and material consequence

*A consequence is called 'formal' if it is valid in all terms retaining a similar form. Or, if you want to put it explicitly, a formal consequence is one where every proposition similar in form that might be formed would be a good consequence. ...A material consequence, however, is one where not every proposition similar in form would be a good consequence, or, as it is commonly put, which does not hold in all terms retaining the same form [TC I. 4, p. 68].*



# Formal and material consequence

*A consequence is called 'formal' if it is valid in all terms retaining a similar form. Or, if you want to put it explicitly, a formal consequence is one where every proposition similar in form that might be formed would be a good consequence. ...A material consequence, however, is one where not every proposition similar in form would be a good consequence, or, as it is commonly put, which does not hold in all terms retaining the same form [TC I. 4, p. 68].*

# Formal and material consequence

*It seems to me that no material consequence is evident in inferring except by its reduction to a formal one. Now it is reduced to a formal one by the addition of some necessary proposition or propositions whose addition to the given antecedent produces a formal consequence [TC I. 4, p. 68].*

# Formal and material consequence

*I say that when we speak of matter and form, by the matter of a proposition or consequence we mean the purely categorematic terms, namely the subject and predicate, setting aside the syncategoremes attached to them by which they are [1] conjoined [2] or denied [3] or distributed [4] or given a certain kind of supposition; we say all the rest pertains to form [TC I. 7, p. 74].*

# Formal consequence from Tarski back to John Buridan

- Both modern and Tarskian approaches begin with a partition of all terms of a *language* into logical and non-logical terms; Buridan's partition of terms into categorical and syncategorematic occurs not at the level of a language, but at that of the sentence.
- Tarski's project prioritizes determining the logical terms of a language. Buridan's the categorematic
- Tarskian consequence was designed for recursively defined artificial languages. Buridanian consequence, Buridan's stilted fourteenth-century scholastic Latin.

# Formal consequence from Tarski back to John Buridan

- Both modern and Tarskian approaches begin with a partition of all terms of a *language* into logical and non-logical terms; Buridan's partition of terms into categorical and syncategorematic occurs not at the level of a language, but at that of the sentence.
- Tarski's project prioritizes determining the logical terms of a language. Buridan's the categorematic
- Tarskian consequence was designed for recursively defined artificial languages. Buridanian consequence, Buridan's stilted fourteenth-century scholastic Latin.

# Formal consequence from Tarski back to John Buridan

- Both modern and Tarskian approaches begin with a partition of all terms of a *language* into logical and non-logical terms; Buridan's partition of terms into categorical and syncategorematic occurs not at the level of a language, but at that of the sentence.
- Tarski's project prioritizes determining the logical terms of a language. Buridan's the categorematic
- Tarskian consequence was designed for recursively defined artificial languages. Buridanian consequence, Buridan's stilted fourteenth-century scholastic Latin.

# Formal consequence from Tarski back to John Buridan

- Both modern and Tarskian approaches begin with a partition of all terms of a *language* into logical and non-logical terms; Buridan's partition of terms into categorical and syncategorematic occurs not at the level of a language, but at that of the sentence.
- Tarski's project prioritizes determining the logical terms of a language. Buridan's the categorematic
- Tarskian consequence was designed for recursively defined artificial languages. Buridanian consequence, Buridan's stilted fourteenth-century scholastic Latin.

# Formal consequence from Tarski back to John Buridan

- Buridan counts the copula, negation, modalities, tenses, quantifiers, intentional operators, as well as disjunction, conjunction and negation for terms among the formal parts of a sentence. He does not mention sentential connectives as pertaining to form.
- Tarski distinguishes consequences from hypothetical propositions. Buridan identifies them.
- Where Tarski provides an informative definition of consequence in terms of models, Buridan provides a deflationary one.



# Formal consequence from Tarski back to John Buridan

- Buridan counts the copula, negation, modalities, tenses, quantifiers, intentional operators, as well as disjunction, conjunction and negation for terms among the formal parts of a sentence. He does not mention sentential connectives as pertaining to form.
- Tarski distinguishes consequences from hypothetical propositions. Buridan identifies them.
- Where Tarski provides an informative definition of consequence in terms of models, Buridan provides a deflationary one.

# Formal consequence from Tarski back to John Buridan

- Buridan counts the copula, negation, modalities, tenses, quantifiers, intentional operators, as well as disjunction, conjunction and negation for terms among the formal parts of a sentence. He does not mention sentential connectives as pertaining to form.
- Tarski distinguishes consequences from hypothetical propositions. Buridan identifies them.
- Where Tarski provides an informative definition of consequence in terms of models, Buridan provides a deflationary one.

# Formal consequence from Tarski back to John Buridan

- Buridan counts the copula, negation, modalities, tenses, quantifiers, intentional operators, as well as disjunction, conjunction and negation for terms among the formal parts of a sentence. He does not mention sentential connectives as pertaining to form.
- Tarski distinguishes consequences from hypothetical propositions. Buridan identifies them.
- Where Tarski provides an informative definition of consequence in terms of models, Buridan provides a deflationary one.

# Formal consequence from Tarski back to John Buridan

- On the received classical account, a model of a sentence  $\phi$  in a language  $L$  consists of a domain  $D$  and an interpretation  $I$ , i.e. a mapping of the sentences of  $L$  recursively determined by a mapping of terms to elements in  $D$  and  $n$ -ary predicates to sets of  $n$ -tuples in  $D^n$ .
- On Tarski's account, a model of a set of sentences  $\Gamma$  is a sequence of objects in a fixed domain satisfying the sentential functions obtained by uniformly replacing each non-logical constant in the sentences of  $\Gamma$  with variables of the appropriate order and arity.
- Buridan's account of causes of truth, by contrast, maps hypothetical propositions to states sufficient to make them true on their *intended* interpretation.

# Formal consequence from Tarski back to John Buridan

- On the received classical account, a model of a sentence  $\phi$  in a language  $L$  consists of a domain  $D$  and an interpretation  $I$ , i.e. a mapping of the sentences of  $L$  recursively determined by a mapping of terms to elements in  $D$  and  $n$ -ary predicates to sets of  $n$ -tuples in  $D^n$ .
- On Tarski's account, a model of a set of sentences  $\Gamma$  is a sequence of objects in a fixed domain satisfying the sentential functions obtained by uniformly replacing each non-logical constant in the sentences of  $\Gamma$  with variables of the appropriate order and arity.
- Buridan's account of causes of truth, by contrast, maps hypothetical propositions to states sufficient to make them true on their *intended* interpretation.

# Formal consequence from Tarski back to John Buridan

- On the received classical account, a model of a sentence  $\phi$  in a language  $L$  consists of a domain  $D$  and an interpretation  $I$ , i.e. a mapping of the sentences of  $L$  recursively determined by a mapping of terms to elements in  $D$  and  $n$ -ary predicates to sets of  $n$ -tuples in  $D^n$ .
- On Tarski's account, a model of a set of sentences  $\Gamma$  is a sequence of objects in a fixed domain satisfying the sentential functions obtained by uniformly replacing each non-logical constant in the sentences of  $\Gamma$  with variables of the appropriate order and arity.
- Buridan's account of causes of truth, by contrast, maps hypothetical propositions to states sufficient to make them true on their *intended* interpretation.

# Formal consequence from Tarski back to John Buridan

- On the received classical account, a model of a sentence  $\phi$  in a language  $L$  consists of a domain  $D$  and an interpretation  $I$ , i.e. a mapping of the sentences of  $L$  recursively determined by a mapping of terms to elements in  $D$  and  $n$ -ary predicates to sets of  $n$ -tuples in  $D^n$ .
- On Tarski's account, a model of a set of sentences  $\Gamma$  is a sequence of objects in a fixed domain satisfying the sentential functions obtained by uniformly replacing each non-logical constant in the sentences of  $\Gamma$  with variables of the appropriate order and arity.
- Buridan's account of causes of truth, by contrast, maps hypothetical propositions to states sufficient to make them true on their *intended* interpretation.

# Formal consequence from Tarski back to John Buridan

Those developments that *were* necessary historical conditions for the development of Tarski's account from the Buridanian one are comparatively few, but central.

- the development of the concept of a model
- The concept of a sentential function
- The concept of a function
- The decision to regard antecedents as premise sets closed under entailment, and hence infinite, presupposed the development of recursion and its application to logical entailment.



# Formal consequence from Tarski back to John Buridan

Those developments that *were* necessary historical conditions for the development of Tarski's account from the Buridanian one are comparatively few, but central.

- the development of the concept of a model
- The concept of a sentential function
- The concept of a function
- The decision to regard antecedents as premise sets closed under entailment, and hence infinite, presupposed the development of recursion and its application to logical entailment.

# Formal consequence from Tarski back to John Buridan

Those developments that *were* necessary historical conditions for the development of Tarski's account from the Buridanian one are comparatively few, but central.

- the development of the concept of a model
- The concept of a sentential function
- The concept of a function
- The decision to regard antecedents as premise sets closed under entailment, and hence infinite, presupposed the development of recursion and its application to logical entailment.

# Formal consequence from Tarski back to John Buridan

Those developments that *were* necessary historical conditions for the development of Tarski's account from the Buridanian one are comparatively few, but central.

- the development of the concept of a model
- The concept of a sentential function
- The concept of a function
- The decision to regard antecedents as premise sets closed under entailment, and hence infinite, presupposed the development of recursion and its application to logical entailment.

# Conclusion

If I put something here, does my prior outline screen show up?