


# CSE 311: Foundations of Computing

---

## Lecture 15: Induction



# **More Logic**

## **Induction**

# Mathematical Induction

---

## Method for proving statements about all natural numbers

- A new logical inference rule!
  - It only applies over the natural numbers
  - The idea is to **use** the special structure of the naturals to prove things more easily
- Particularly useful for reasoning about programs!  
**for (int i=0; i < n; n++) { ... }**
  - Show  $P(i)$  holds after  $i$  times through the loop

**Prove**  $\forall a, b, m > 0 \forall k \in \mathbb{N} ((a \equiv_m b) \rightarrow (a^k \equiv_m b^k))$

---

**Let  $a, b, m > 0$  be arbitrary. Let  $k \in \mathbb{N}$  be arbitrary.**

**Suppose that  $a \equiv_m b$ .**

**We know  $((a \equiv_m b) \wedge (a \equiv_m b)) \rightarrow (a^2 \equiv_m b^2)$  by multiplying congruences. So, applying this repeatedly, we have:**

$$\begin{aligned} & ((a \equiv_m b) \wedge (a \equiv_m b)) \rightarrow (a^2 \equiv_m b^2) \\ & ((a^2 \equiv_m b^2) \wedge (a \equiv_m b)) \rightarrow (a^3 \equiv_m b^3) \end{aligned}$$

...

$$((a^{k-1} \equiv_m b^{k-1}) \wedge (a \equiv_m b)) \rightarrow (a^k \equiv_m b^k)$$

The “...”s is a problem! We don’t have a proof rule that allows us to say “do this over and over”.

But there such a property of the natural numbers!

---

Domain: Natural Numbers

$$\frac{P(0) \quad \forall k (P(k) \longrightarrow P(k + 1))}{\therefore \forall n P(n)}$$

# Induction Is A Rule of Inference

---

Domain: Natural Numbers

$$\frac{P(0) \quad \forall k (P(k) \rightarrow P(k + 1))}{\therefore \forall n P(n)}$$


How do the givens prove  $P(3)$ ?

# Induction Is A Rule of Inference

Domain: Natural Numbers

$$\frac{P(0) \quad \forall k (P(k) \rightarrow P(k + 1))}{\therefore \forall n P(n)}$$

How do the givens prove  $P(5)$ ?



First, we have  $P(0)$ .

Since  $P(n) \rightarrow P(n+1)$  for all  $n$ , we have  $P(0) \rightarrow P(1)$ .

Since  $P(0)$  is true and  $P(0) \rightarrow P(1)$ , by Modus Ponens,  $P(1)$  is true.

Since  $P(n) \rightarrow P(n+1)$  for all  $n$ , we have  $P(1) \rightarrow P(2)$ .

Since  $P(1)$  is true and  $P(1) \rightarrow P(2)$ , by Modus Ponens,  $P(2)$  is true.

## Using The Induction Rule In A Formal Proof

---

$$\frac{\begin{array}{c} P(0) \\ \forall k (P(k) \longrightarrow P(k + 1)) \end{array}}{\therefore \forall n P(n)}$$



# Using The Induction Rule In A Formal Proof

---

$$\frac{P(0) \quad \forall k (P(k) \rightarrow P(k + 1))}{\therefore \forall n P(n)}$$

1.  $P(0)$

4.  $\forall k (P(k) \rightarrow P(k+1))$

5.  $\forall n P(n)$

Induction: 1, 4

# Using The Induction Rule In A Formal Proof

---

$$\frac{P(0) \quad \forall k (P(k) \longrightarrow P(k + 1))}{\therefore \forall n P(n)}$$

1.  $P(0)$
2. Let  $k$  be an arbitrary integer  $\geq 0$

3.  $P(k) \longrightarrow P(k+1)$

4.  $\forall k (P(k) \longrightarrow P(k+1))$

5.  $\forall n P(n)$

Intro  $\forall$ : 2, 3

Induction: 1, 4

# Using The Induction Rule In A Formal Proof

---

$$\frac{\begin{array}{c} P(0) \\ \forall k (P(k) \rightarrow P(k + 1)) \end{array}}{\therefore \forall n P(n)}$$

1.  $P(0)$
2. Let  $k$  be an arbitrary integer  $\geq 0$ 
  - 3.1.  $P(k)$  Assumption
  - 3.2. ...
  - 3.3.  $P(k+1)$
3.  $P(k) \rightarrow P(k+1)$  Direct Proof Rule
4.  $\forall k (P(k) \rightarrow P(k+1))$  Intro  $\forall$ : 2, 3
5.  $\forall n P(n)$  Induction: 1, 4

# Translating to an English Proof

---

$$\begin{array}{c} P(0) \\ \forall k (P(k) \rightarrow P(k + 1)) \\ \hline \therefore \forall n P(n) \end{array}$$

1. Prove  $P(0)$

**Base Case**

2. Let  $k$  be an arbitrary integer  $\geq 0$

**Inductive  
Hypothesis**

3.1. Suppose that  $P(k)$  is true

3.2. ...

**Inductive  
Step**

3.3. Prove  $P(k+1)$  is true

3.  $P(k) \rightarrow P(k+1)$

Direct Proof Rule

4.  $\forall k (P(k) \rightarrow P(k+1))$

Intro  $\forall$ : 2, 3

5.  $\forall n P(n)$

Induction: 1, 4

**Conclusion**

# Translating to an English Proof

---

|   |           |                        |
|---|-----------|------------------------|
| 1. Prove $P(0)$                             | Base Case |                        |
| 2. Let $k$ be an arbitrary integer $\geq 0$ |           | Inductive Hypothesis   |
| 3.1. Assume that $P(k)$ is true             |           |                        |
| 3.2. ...                                    |           | Inductive Step         |
| 3.3. Prove $P(k+1)$ is true                 |           |                        |
| 3. $P(k) \rightarrow P(k+1)$                |           | Direct Proof Rule      |
| 4. $\forall k (P(k) \rightarrow P(k+1))$    |           | Intro $\forall$ : 2, 3 |
| 5. $\forall n P(n)$                         |           | Induction: 1, 4        |
| Conclusion                                  |           |                        |

## Induction English Proof Template

*[...Define  $P(n)$ ...]*

We will show that  $P(n)$  is true for every  $n \in \mathbb{N}$  by Induction.

Base Case: *[...proof of  $P(0)$  here...]*

Induction Hypothesis:

Suppose that  $P(k)$  is true for an arbitrary  $k \in \mathbb{N}$ .

Induction Step:

*[...proof of  $P(k + 1)$  here...]*

The proof of  $P(k + 1)$  **must** invoke the IH somewhere.

So, the claim is true by induction.

# Inductive Proofs In 5 Easy Steps

---

## Proof:

1. “Let  $P(n)$  be... . We will show that  $P(n)$  is true for every  $n \geq 0$  by Induction.”

2. “Base Case:” Prove  $P(0)$

3. “Inductive Hypothesis:

Suppose  $P(k)$  is true for an arbitrary integer  $k \geq 0$ ”

4. “Inductive Step:” Prove that  $P(k + 1)$  is true.

*Use the goal to figure out what you need.*

*Make sure you are using I.H. and point out where you are using it. (Don't assume  $P(k + 1)$  !!)*

5. “Conclusion: Result follows by induction”

**What is  $1 + 2 + 4 + \dots + 2^n$  ?**

---

- $1 = 1$
- $1 + 2 = 3$
- $1 + 2 + 4 = 7$
- $1 + 2 + 4 + 8 = 15$
- $1 + 2 + 4 + 8 + 16 = 31$

**It sure looks like this sum is  $2^{n+1} - 1$**

**How can we prove it?**

**We could prove it for  $n = 1, n = 2, n = 3, \dots$  but that would literally take forever.**

**Good that we have induction!**

**Prove  $1 + 2 + 4 + \dots + 2^n = 2^{n+1} - 1$**

---



**Prove  $1 + 2 + 4 + \dots + 2^n = 2^{n+1} - 1$**

---

- 1. Let  $P(n)$  be " $2^0 + 2^1 + \dots + 2^n = 2^{n+1} - 1$ ". We will show  $P(n)$  is true for all natural numbers by induction.**

**Prove  $1 + 2 + 4 + \dots + 2^n = 2^{n+1} - 1$**

---

- 1. Let  $P(n)$  be " $2^0 + 2^1 + \dots + 2^n = 2^{n+1} - 1$ ". We will show  $P(n)$  is true for all natural numbers by induction.**
- 2. Base Case ( $n=0$ ):  $2^0 = 1 = 2 - 1 = 2^{0+1} - 1$  so  $P(0)$  is true.**

**Prove  $1 + 2 + 4 + \dots + 2^n = 2^{n+1} - 1$**

---

- 1. Let  $P(n)$  be " $2^0 + 2^1 + \dots + 2^n = 2^{n+1} - 1$ ". We will show  $P(n)$  is true for all natural numbers by induction.**
- 2. Base Case ( $n=0$ ):  $2^0 = 1 = 2 - 1 = 2^{0+1} - 1$  so  $P(0)$  is true.**
- 3. Induction Hypothesis: Suppose that  $P(k)$  is true for some arbitrary integer  $k \geq 0$ , i.e., that  $2^0 + 2^1 + \dots + 2^k = 2^{k+1} - 1$ .**

**Prove  $1 + 2 + 4 + \dots + 2^n = 2^{n+1} - 1$**

---

- 1. Let  $P(n)$  be “ $2^0 + 2^1 + \dots + 2^n = 2^{n+1} - 1$ ”. We will show  $P(n)$  is true for all natural numbers by induction.**
- 2. Base Case ( $n=0$ ):  $2^0 = 1 = 2 - 1 = 2^{0+1} - 1$  so  $P(0)$  is true.**
- 3. Induction Hypothesis: Suppose that  $P(k)$  is true for some arbitrary integer  $k \geq 0$ , i.e., that  $2^0 + 2^1 + \dots + 2^k = 2^{k+1} - 1$ .**
- 4. Induction Step:**

**Goal: Show  $P(k+1)$ , i.e. show  $2^0 + 2^1 + \dots + 2^k + 2^{k+1} = 2^{k+2} - 1$**

**Prove  $1 + 2 + 4 + \dots + 2^n = 2^{n+1} - 1$**

---

- 1. Let  $P(n)$  be “ $2^0 + 2^1 + \dots + 2^n = 2^{n+1} - 1$ ”. We will show  $P(n)$  is true for all natural numbers by induction.**
- 2. Base Case ( $n=0$ ):  $2^0 = 1 = 2 - 1 = 2^{0+1} - 1$  so  $P(0)$  is true.**
- 3. Induction Hypothesis: Suppose that  $P(k)$  is true for some arbitrary integer  $k \geq 0$ , i.e., that  $2^0 + 2^1 + \dots + 2^k = 2^{k+1} - 1$ .**
- 4. Induction Step:**

$$2^0 + 2^1 + \dots + 2^k = 2^{k+1} - 1 \quad \text{by IH}$$

**Adding  $2^{k+1}$  to both sides, we get:**

$$2^0 + 2^1 + \dots + 2^k + 2^{k+1} = 2^{k+1} + 2^{k+1} - 1$$

**Note that  $2^{k+1} + 2^{k+1} = 2(2^{k+1}) = 2^{k+2}$ .**

**So, we have  $2^0 + 2^1 + \dots + 2^k + 2^{k+1} = 2^{k+2} - 1$ , which is exactly  $P(k+1)$ .**

**Prove  $1 + 2 + 4 + \dots + 2^n = 2^{n+1} - 1$**

---

- 1. Let  $P(n)$  be “ $2^0 + 2^1 + \dots + 2^n = 2^{n+1} - 1$ ”. We will show  $P(n)$  is true for all natural numbers by induction.**
- 2. Base Case ( $n=0$ ):  $2^0 = 1 = 2 - 1 = 2^{0+1} - 1$  so  $P(0)$  is true.**
- 3. Induction Hypothesis: Suppose that  $P(k)$  is true for some arbitrary integer  $k \geq 0$ , i.e., that  $2^0 + 2^1 + \dots + 2^k = 2^{k+1} - 1$ .**
- 4. Induction Step:**

**We can calculate**

$$\begin{aligned} 2^0 + 2^1 + \dots + 2^k + 2^{k+1} &= (2^0 + 2^1 + \dots + 2^k) + 2^{k+1} \\ &= (2^{k+1} - 1) + 2^{k+1} && \text{by the IH} \\ &= 2(2^{k+1}) - 1 \\ &= 2^{k+2} - 1, \end{aligned}$$

**which is exactly  $P(k+1)$ .**

**Alternative way of writing the inductive step**

**Prove  $1 + 2 + 4 + \dots + 2^n = 2^{n+1} - 1$**

---

- 1. Let  $P(n)$  be “ $2^0 + 2^1 + \dots + 2^n = 2^{n+1} - 1$ ”. We will show  $P(n)$  is true for all natural numbers by induction.**
- 2. Base Case ( $n=0$ ):  $2^0 = 1 = 2 - 1 = 2^{0+1} - 1$  so  $P(0)$  is true.**
- 3. Induction Hypothesis: Suppose that  $P(k)$  is true for some arbitrary integer  $k \geq 0$ , i.e., that  $2^0 + 2^1 + \dots + 2^k = 2^{k+1} - 1$ .**
- 4. Induction Step:**

**We can calculate**

$$\begin{aligned} 2^0 + 2^1 + \dots + 2^k + 2^{k+1} &= (2^0 + 2^1 + \dots + 2^k) + 2^{k+1} \\ &= (2^{k+1} - 1) + 2^{k+1} && \text{by the IH} \\ &= 2(2^{k+1}) - 1 \\ &= 2^{k+2} - 1, \end{aligned}$$

**which is exactly  $P(k+1)$ .**

- 5. Thus  $P(n)$  is true for all  $n \in \mathbb{N}$ , by induction.**

**Prove  $1 + 2 + 3 + \dots + n = n(n + 1)/2$**

---



**Prove  $1 + 2 + 3 + \dots + n = n(n + 1)/2$**

---

- 1. Let  $P(n)$  be " $0 + 1 + 2 + \dots + n = n(n+1)/2$ ". We will show  $P(n)$  is true for all natural numbers by induction.**

**Prove  $1 + 2 + 3 + \dots + n = n(n + 1)/2$**

---

- 1. Let  $P(n)$  be “ $0 + 1 + 2 + \dots + n = n(n+1)/2$ ”. We will show  $P(n)$  is true for all natural numbers by induction.**
- 2. Base Case ( $n=0$ ):  $0 = 0(0+1)/2$ . Therefore  $P(0)$  is true.**

**Prove  $1 + 2 + 3 + \dots + n = n(n + 1)/2$**

---

- 1. Let  $P(n)$  be “ $0 + 1 + 2 + \dots + n = n(n+1)/2$ ”. We will show  $P(n)$  is true for all natural numbers by induction.**
- 2. Base Case ( $n=0$ ):  $0 = 0(0+1)/2$ . Therefore  $P(0)$  is true.**
- 3. Induction Hypothesis: Suppose that  $P(k)$  is true for some arbitrary integer  $k \geq 0$ . I.e., suppose  $1 + 2 + \dots + k = k(k+1)/2$**

**Prove  $1 + 2 + 3 + \dots + n = n(n + 1)/2$**

---

- 1. Let  $P(n)$  be “ $0 + 1 + 2 + \dots + n = n(n+1)/2$ ”. We will show  $P(n)$  is true for all natural numbers by induction.**
- 2. Base Case ( $n=0$ ):  $0 = 0(0+1)/2$ . Therefore  $P(0)$  is true.**
- 3. Induction Hypothesis: Suppose that  $P(k)$  is true for some arbitrary integer  $k \geq 0$ . I.e., suppose  $1 + 2 + \dots + k = k(k+1)/2$**
- 4. Induction Step:**

**Goal: Show  $P(k+1)$ , i.e. show  $1 + 2 + \dots + k + (k+1) = (k+1)(k+2)/2$**

**Prove  $1 + 2 + 3 + \dots + n = n(n + 1)/2$**

---

- 1. Let  $P(n)$  be “ $0 + 1 + 2 + \dots + n = n(n+1)/2$ ”. We will show  $P(n)$  is true for all natural numbers by induction.**
- 2. Base Case ( $n=0$ ):  $0 = 0(0+1)/2$ . Therefore  $P(0)$  is true.**
- 3. Induction Hypothesis: Suppose that  $P(k)$  is true for some arbitrary integer  $k \geq 0$ . I.e., suppose  $1 + 2 + \dots + k = k(k+1)/2$**
- 4. Induction Step:**

$$\begin{aligned} 1 + 2 + \dots + k + (k+1) &= (1 + 2 + \dots + k) + (k+1) \\ &= k(k+1)/2 + (k+1) \text{ by IH} \\ &= (k+1)(k/2 + 1) \\ &= (k+1)(k+2)/2 \end{aligned}$$

**So, we have shown  $1 + 2 + \dots + k + (k+1) = (k+1)(k+2)/2$ , which is exactly  $P(k+1)$ .**

- 5. Thus  $P(n)$  is true for all  $n \in \mathbb{N}$ , by induction.**

## Another example of a pattern

---

- $2^0 - 1 = 1 - 1 = 0 = 3 \cdot 0$
- $2^2 - 1 = 4 - 1 = 3 = 3 \cdot 1$
- $2^4 - 1 = 16 - 1 = 15 = 3 \cdot 5$
- $2^6 - 1 = 64 - 1 = 63 = 3 \cdot 21$
- $2^8 - 1 = 256 - 1 = 255 = 3 \cdot 85$
- ...

**Prove:  $3 \mid (2^{2n} - 1)$  for all  $n \geq 0$**

---

**Prove:  $3 \mid (2^{2n} - 1)$  for all  $n \geq 0$**

---

- 1. Let  $P(n)$  be " $3 \mid (2^{2n} - 1)$ ". We will show  $P(n)$  is true for all natural numbers by induction.**



**Prove:  $3 \mid (2^{2n} - 1)$  for all  $n \geq 0$**

---

- 1. Let  $P(n)$  be " $3 \mid (2^{2n} - 1)$ ". We will show  $P(n)$  is true for all natural numbers by induction.**
- 2. Base Case ( $n=0$ ):  $2^{2 \cdot 0} - 1 = 1 - 1 = 0 = 3 \cdot 0$  Therefore  $P(0)$  is true**

**Prove:  $3 \mid (2^{2n} - 1)$  for all  $n \geq 0$**

---

- 1. Let  $P(n)$  be " $3 \mid (2^{2n} - 1)$ ". We will show  $P(n)$  is true for all natural numbers by induction.**
- 2. Base Case ( $n=0$ ):  $2^{2 \cdot 0} - 1 = 1 - 1 = 0 = 3 \cdot 0$  Therefore  $P(0)$  is true.**
- 3. Induction Hypothesis: Suppose that  $P(k)$  is true for some arbitrary integer  $k \geq 0$ . I.e., suppose that  $3 \mid (2^{2k} - 1)$**

**Prove:  $3 \mid (2^{2n} - 1)$  for all  $n \geq 0$**

---

- 1. Let  $P(n)$  be “ $3 \mid (2^{2n} - 1)$ ”. We will show  $P(n)$  is true for all natural numbers by induction.**
- 2. Base Case ( $n=0$ ):  $2^{2 \cdot 0} - 1 = 1 - 1 = 0 = 3 \cdot 0$  Therefore  $P(0)$  is true.**
- 3. Induction Hypothesis: Suppose that  $P(k)$  is true for some arbitrary integer  $k \geq 0$ . I.e., suppose that  $3 \mid (2^{2k} - 1)$**
- 4. Induction Step:**

**Goal: Show  $P(k+1)$ , i.e. show  $3 \mid (2^{2(k+1)} - 1)$**

**Prove:  $3 \mid (2^{2n} - 1)$  for all  $n \geq 0$**

---

- 1. Let  $P(n)$  be “ $3 \mid (2^{2n} - 1)$ ”. We will show  $P(n)$  is true for all natural numbers by induction.**
- 2. Base Case ( $n=0$ ):  $2^{2 \cdot 0} - 1 = 1 - 1 = 0 = 3 \cdot 0$  Therefore  $P(0)$  is true.**
- 3. Induction Hypothesis: Suppose that  $P(k)$  is true for some arbitrary integer  $k \geq 0$ . I.e., suppose that  $3 \mid (2^{2k} - 1)$**
- 4. Induction Step:**

By IH,  $3 \mid (2^{2k} - 1)$  so  $2^{2k} - 1 = 3j$  for some integer  $j$

So  $2^{2(k+1)} - 1 = 2^{2k+2} - 1 = 4(2^{2k}) - 1 = 4(3j+1) - 1$   
 $= 12j+3 = 3(4j+1)$

Therefore  $3 \mid (2^{2(k+1)} - 1)$  which is exactly  $P(k+1)$ .
- 5. Thus  $P(n)$  is true for all  $n \in \mathbb{N}$ , by induction.**