









## □ APPENDIXES

### A Numbers, Inequalities, and Absolute Values


1.  $|5 - 23| = |-18| = 18$
2.  $|5| - |-23| = 5 - 23 = -18$
3.  $|-π| = π$  because  $π > 0$ .
4.  $|\pi - 2| = \pi - 2$  because  $\pi - 2 > 0$ .
5.  $|\sqrt{5} - 5| = -(\sqrt{5} - 5) = 5 - \sqrt{5}$  because  $\sqrt{5} - 5 < 0$ .
6.  $||-2| - |-3|| = |2 - 3| = |-1| = 1$
7. If  $x < 2$ ,  $x - 2 < 0$ , so  $|x - 2| = -(x - 2) = 2 - x$ .
8. If  $x > 2$ ,  $x - 2 > 0$ , so  $|x - 2| = x - 2$ .
9.  $|x + 1| = \begin{cases} x + 1 & \text{if } x + 1 \geq 0 \\ -(x + 1) & \text{if } x + 1 < 0 \end{cases} = \begin{cases} x + 1 & \text{if } x \geq -1 \\ -x - 1 & \text{if } x < -1 \end{cases}$
10.  $|2x - 1| = \begin{cases} 2x - 1 & \text{if } 2x - 1 \geq 0 \\ -(2x - 1) & \text{if } 2x - 1 < 0 \end{cases} = \begin{cases} 2x - 1 & \text{if } x \geq \frac{1}{2} \\ 1 - 2x & \text{if } x < \frac{1}{2} \end{cases}$
11.  $|x^2 + 1| = x^2 + 1$  [since  $x^2 + 1 \geq 0$  for all  $x$ ].
12. Determine when  $1 - 2x^2 < 0 \Leftrightarrow 1 < 2x^2 \Leftrightarrow x^2 > \frac{1}{2} \Leftrightarrow \sqrt{x^2} > \sqrt{\frac{1}{2}} \Leftrightarrow |x| > \sqrt{\frac{1}{2}} \Leftrightarrow$   
 $x < -\frac{1}{\sqrt{2}} \text{ or } x > \frac{1}{\sqrt{2}}$ . Thus,  $|1 - 2x^2| = \begin{cases} 1 - 2x^2 & \text{if } -\frac{1}{\sqrt{2}} \leq x \leq \frac{1}{\sqrt{2}} \\ 2x^2 - 1 & \text{if } x < -\frac{1}{\sqrt{2}} \text{ or } x > \frac{1}{\sqrt{2}} \end{cases}$
13.  $2x + 7 > 3 \Leftrightarrow 2x > -4 \Leftrightarrow x > -2$ , so  $x \in (-2, \infty)$ .
14.  $3x - 11 < 4 \Leftrightarrow 3x < 15 \Leftrightarrow x < 5$ , so  $x \in (-\infty, 5)$ .
15.  $1 - x \leq 2 \Leftrightarrow -x \leq 1 \Leftrightarrow x \geq -1$ , so  $x \in [-1, \infty)$ .
16.  $4 - 3x \geq 6 \Leftrightarrow -3x \geq 2 \Leftrightarrow x \leq -\frac{2}{3}$ , so  $x \in (-\infty, -\frac{2}{3}]$ .
17.  $2x + 1 < 5x - 8 \Leftrightarrow 9 < 3x \Leftrightarrow 3 < x$ , so  $x \in (3, \infty)$ .
18.  $1 + 5x > 5 - 3x \Leftrightarrow 8x > 4 \Leftrightarrow x > \frac{1}{2}$ , so  $x \in (\frac{1}{2}, \infty)$ .
19.  $-1 < 2x - 5 < 7 \Leftrightarrow 4 < 2x < 12 \Leftrightarrow 2 < x < 6$ , so  $x \in (2, 6)$ .




20.  $1 < 3x + 4 \leq 16 \Leftrightarrow -3 < 3x \leq 12 \Leftrightarrow -1 < x \leq 4$ , so  $x \in (-1, 4]$ .




21.  $0 \leq 1 - x < 1 \Leftrightarrow -1 \leq -x < 0 \Leftrightarrow 1 \geq x > 0$ , so  $x \in (0, 1]$ .




22.  $-5 \leq 3 - 2x \leq 9 \Leftrightarrow -8 \leq -2x \leq 6 \Leftrightarrow 4 \geq x \geq -3$ , so  $x \in [-3, 4]$ .



23.  $4x < 2x + 1 \leq 3x + 2$ . So  $4x < 2x + 1 \Leftrightarrow 2x < 1 \Leftrightarrow x < \frac{1}{2}$ , and  
 $2x + 1 \leq 3x + 2 \Leftrightarrow -1 \leq x$ . Thus,  $x \in [-1, \frac{1}{2})$ .



24.  $2x - 3 < x + 4 < 3x - 2$ . So  $2x - 3 < x + 4 \Leftrightarrow x < 7$ , and  
 $x + 4 < 3x - 2 \Leftrightarrow 6 < 2x \Leftrightarrow 3 < x$ , so  $x \in (3, 7)$ .




25.  $(x - 1)(x - 2) > 0$ .

Case 1: (both factors are positive, so their product is positive)  $x - 1 > 0 \Leftrightarrow x > 1$ ,  
 and  $x - 2 > 0 \Leftrightarrow x > 2$ , so  $x \in (2, \infty)$ .

Case 2: (both factors are negative, so their product is positive)  $x - 1 < 0 \Leftrightarrow x < 1$ ,  
 and  $x - 2 < 0 \Leftrightarrow x < 2$ , so  $x \in (-\infty, 1)$ .

Thus, the solution set is  $(-\infty, 1) \cup (2, \infty)$ .




26.  $(2x + 3)(x - 1) \geq 0$ .

Case 1:  $2x + 3 \geq 0 \Leftrightarrow x \geq -\frac{3}{2}$ , and  $x - 1 \geq 0 \Leftrightarrow x \geq 1$ , so  $x \in [1, \infty)$ .

Case 2:  $2x + 3 \leq 0 \Leftrightarrow x \leq -\frac{3}{2}$ , and  $x - 1 \leq 0 \Leftrightarrow x \leq 1$ , so  $x \in (-\infty, -\frac{3}{2}]$ .

Thus, the solution set is  $(-\infty, -\frac{3}{2}] \cup [1, \infty)$ .




27.  $2x^2 + x \leq 1 \Leftrightarrow 2x^2 + x - 1 \leq 0 \Leftrightarrow (2x - 1)(x + 1) \leq 0$ .

Case 1:  $2x - 1 \geq 0 \Leftrightarrow x \geq \frac{1}{2}$ , and  $x + 1 \leq 0 \Leftrightarrow x \leq -1$ ,  
 which is an impossible combination.

Case 2:  $2x - 1 \leq 0 \Leftrightarrow x \leq \frac{1}{2}$ , and  $x + 1 \geq 0 \Leftrightarrow x \geq -1$ , so  $x \in [-1, \frac{1}{2}]$ .

Thus, the solution set is  $[-1, \frac{1}{2}]$ .




28.  $x^2 < 2x + 8 \Leftrightarrow x^2 - 2x - 8 < 0 \Leftrightarrow (x - 4)(x + 2) < 0$ .

Case 1:  $x > 4$  and  $x < -2$ , which is impossible.

Case 2:  $x < 4$  and  $x > -2$ .

Thus, the solution set is  $(-2, 4)$ .



29.  $x^2 + x + 1 > 0 \Leftrightarrow x^2 + x + \frac{1}{4} + \frac{3}{4} > 0 \Leftrightarrow (x + \frac{1}{2})^2 + \frac{3}{4} > 0$ . But since

$(x + \frac{1}{2})^2 \geq 0$  for every real  $x$ , the original inequality will be true for all real  $x$  as well.

Thus, the solution set is  $(-\infty, \infty)$ .




30.  $x^2 + x > 1 \Leftrightarrow x^2 + x - 1 > 0$ . Using the quadratic formula, we obtain

$$x^2 + x - 1 = \left(x - \frac{-1-\sqrt{5}}{2}\right) \left(x - \frac{-1+\sqrt{5}}{2}\right) > 0.$$

Case 1:  $x - \frac{-1-\sqrt{5}}{2} > 0$  and  $x - \frac{-1+\sqrt{5}}{2} > 0$ , so that  $x > \frac{-1+\sqrt{5}}{2}$ .

Case 2:  $x - \frac{-1-\sqrt{5}}{2} < 0$  and  $x - \frac{-1+\sqrt{5}}{2} < 0$ , so that  $x < \frac{-1-\sqrt{5}}{2}$ .

Thus, the solution set is  $\left(-\infty, \frac{-1-\sqrt{5}}{2}\right) \cup \left(\frac{-1+\sqrt{5}}{2}, \infty\right)$ .




31.  $x^2 < 3 \Leftrightarrow x^2 - 3 < 0 \Leftrightarrow (x - \sqrt{3})(x + \sqrt{3}) < 0$ .

Case 1:  $x > \sqrt{3}$  and  $x < -\sqrt{3}$ , which is impossible.

Case 2:  $x < \sqrt{3}$  and  $x > -\sqrt{3}$ .

Thus, the solution set is  $(-\sqrt{3}, \sqrt{3})$ .




Another method:  $x^2 < 3 \Leftrightarrow |x| < \sqrt{3} \Leftrightarrow -\sqrt{3} < x < \sqrt{3}$ .

32.  $x^2 \geq 5 \Leftrightarrow x^2 - 5 \geq 0 \Leftrightarrow (x - \sqrt{5})(x + \sqrt{5}) \geq 0$ .

Case 1:  $x \geq \sqrt{5}$  and  $x \geq -\sqrt{5}$ , so  $x \in [\sqrt{5}, \infty)$ .

Case 2:  $x \leq \sqrt{5}$  and  $x \leq -\sqrt{5}$ , so  $x \in (-\infty, -\sqrt{5}]$ .


Thus, the solution set is  $(-\infty, -\sqrt{5}] \cup [\sqrt{5}, \infty)$ .



Another method:  $x^2 \geq 5 \Leftrightarrow |x| \geq \sqrt{5} \Leftrightarrow x \geq \sqrt{5}$  or  $x \leq -\sqrt{5}$ .

33.  $x^3 - x^2 \leq 0 \Leftrightarrow x^2(x - 1) \leq 0$ . Since  $x^2 \geq 0$  for all  $x$ , the inequality is satisfied when  $x - 1 \leq 0 \Leftrightarrow x \leq 1$ .

Thus, the solution set is  $(-\infty, 1]$ .




34.  $(x + 1)(x - 2)(x + 3) = 0 \Leftrightarrow x = -1, 2$ , or  $-3$ . Construct a chart:

Interval	$x + 1$	$x - 2$	$x + 3$	$(x + 1)(x - 2)(x + 3)$
$x < -3$	-	-	-	-
$-3 < x < -1$	-	-	+	+
$-1 < x < 2$	+	-	+	-
$x > 2$	+	+	+	+

Thus,  $(x + 1)(x - 2)(x + 3) \geq 0$  on  $[-3, -1]$  and  $[2, \infty)$ , and the solution set


is  $[-3, -1] \cup [2, \infty)$ .



35.  $x^3 > x \Leftrightarrow x^3 - x > 0 \Leftrightarrow x(x^2 - 1) > 0 \Leftrightarrow x(x - 1)(x + 1) > 0$ . Construct a chart:

Interval	$x$	$x - 1$	$x + 1$	$x(x - 1)(x + 1)$
$x < -1$	-	-	-	-
$-1 < x < 0$	-	-	+	+
$0 < x < 1$	+	-	+	-
$x > 1$	+	+	+	+


Since  $x^3 > x$  when the last column is positive, the solution set is  $(-1, 0) \cup (1, \infty)$ .




36.  $x^3 + 3x < 4x^2 \Leftrightarrow x^3 - 4x^2 + 3x < 0 \Leftrightarrow x(x^2 - 4x + 3) < 0 \Leftrightarrow x(x-1)(x-3) < 0$ .

Interval	$x$	$x-1$	$x-3$	$x(x-1)(x-3)$
$x < 0$	-	-	-	-
$0 < x < 1$	+	-	-	+
$1 < x < 3$	+	+	-	-
$x > 3$	+	+	+	+

Thus, the solution set is  $(-\infty, 0) \cup (1, 3)$ .




37.  $1/x < 4$ . This is clearly true for  $x < 0$ . So suppose  $x > 0$ . then  $1/x < 4 \Leftrightarrow 1 < 4x \Leftrightarrow \frac{1}{4} < x$ . Thus, the solution set is  $(-\infty, 0) \cup (\frac{1}{4}, \infty)$ .



38.  $-3 < 1/x \leq 1$ . We solve the two inequalities separately and take the intersection of the solution sets. First,  $-3 < 1/x$  is clearly true for  $x > 0$ . So suppose  $x < 0$ . Then  $-3 < 1/x \Leftrightarrow -3x > 1 \Leftrightarrow x < -\frac{1}{3}$ , so for this inequality, the solution set is  $(-\infty, -\frac{1}{3}) \cup (0, \infty)$ . Now  $1/x \leq 1$  is clearly true if  $x < 0$ . So suppose  $x > 0$ . Then  $1/x \leq 1 \Leftrightarrow 1 \leq x$ , and the solution set here is  $(-\infty, 0) \cup [1, \infty)$ .

Taking the intersection of the two solution sets gives the final solution set:

$(-\infty, -\frac{1}{3}) \cup [1, \infty)$ .



39.  $C = \frac{5}{9}(F - 32) \Rightarrow F = \frac{9}{5}C + 32$ . So  $50 \leq F \leq 95 \Rightarrow 50 \leq \frac{9}{5}C + 32 \leq 95 \Rightarrow 18 \leq \frac{9}{5}C \leq 63 \Rightarrow 10 \leq C \leq 35$ . So the interval is  $[10, 35]$ .

40. Since  $20 \leq C \leq 30$  and  $C = \frac{5}{9}(F - 32)$ , we have  $20 \leq \frac{5}{9}(F - 32) \leq 30 \Rightarrow 36 \leq F - 32 \leq 54 \Rightarrow 68 \leq F \leq 86$ . So the interval is  $[68, 86]$ .

41. (a) Let  $T$  represent the temperature in degrees Celsius and  $h$  the height in km.  $T = 20$  when  $h = 0$  and  $T$  decreases by  $10^\circ\text{C}$  for every km ( $1^\circ\text{C}$  for each 100-m rise). Thus,  $T = 20 - 10h$  when  $0 \leq h \leq 12$ .

(b) From part (a),  $T = 20 - 10h \Rightarrow 10h = 20 - T \Rightarrow h = 2 - T/10$ . So  $0 \leq h \leq 5 \Rightarrow 0 \leq 2 - T/10 \leq 5 \Rightarrow -2 \leq -T/10 \leq 3 \Rightarrow -20 \leq -T \leq 30 \Rightarrow 20 \geq T \geq -30 \Rightarrow -30 \leq T \leq 20$ . Thus, the range of temperatures (in  $^\circ\text{C}$ ) to be expected is  $[-30, 20]$ .

42. The ball will be at least 32 ft above the ground if  $h \geq 32 \Leftrightarrow 128 + 16t - 16t^2 \geq 32 \Leftrightarrow 16t^2 - 16t - 96 \leq 0 \Leftrightarrow 16(t-3)(t+2) \leq 0$ .  $t = 3$  and  $t = -2$  are endpoints of the interval we're looking for, and constructing a table gives  $-2 \leq t \leq 3$ . But  $t \geq 0$ , so the ball will be at least 32 ft above the ground in the time interval  $[0, 3]$ .

43.  $|2x| = 3 \Leftrightarrow$  either  $2x = 3$  or  $2x = -3 \Leftrightarrow x = \frac{3}{2}$  or  $x = -\frac{3}{2}$ .

44.  $|3x + 5| = 1 \Leftrightarrow$  either  $3x + 5 = 1$  or  $-1$ . In the first case,  $3x = -4 \Leftrightarrow x = -\frac{4}{3}$ , and in the second case,  $3x = -6 \Leftrightarrow x = -2$ . So the solutions are  $-2$  and  $-\frac{4}{3}$ .

45.  $|x + 3| = |2x + 1| \Leftrightarrow$  either  $x + 3 = 2x + 1$  or  $x + 3 = -(2x + 1)$ . In the first case,  $x = 2$ , and in the second case,  $x + 3 = -2x - 1 \Leftrightarrow 3x = -4 \Leftrightarrow x = -\frac{4}{3}$ . So the solutions are  $-\frac{4}{3}$  and 2.
46.  $\left|\frac{2x-1}{x+1}\right| = 3 \Leftrightarrow$  either  $\frac{2x-1}{x+1} = 3$  or  $\frac{2x-1}{x+1} = -3$ . In the first case,  $2x - 1 = 3x + 3 \Leftrightarrow x = -4$ , and in the second case,  $2x - 1 = -3x - 3 \Leftrightarrow x = -\frac{2}{5}$ .
47. By Property 5 of absolute values,  $|x| < 3 \Leftrightarrow -3 < x < 3$ , so  $x \in (-3, 3)$ .
48. By Properties 4 and 6 of absolute values,  $|x| \geq 3 \Leftrightarrow x \leq -3$  or  $x \geq 3$ , so  $x \in (-\infty, -3] \cup [3, \infty)$ .
49.  $|x - 4| < 1 \Leftrightarrow -1 < x - 4 < 1 \Leftrightarrow 3 < x < 5$ , so  $x \in (3, 5)$ .
50.  $|x - 6| < 0.1 \Leftrightarrow -0.1 < x - 6 < 0.1 \Leftrightarrow 5.9 < x < 6.1$ , so  $x \in (5.9, 6.1)$ .
51.  $|x + 5| \geq 2 \Leftrightarrow x + 5 \geq 2$  or  $x + 5 \leq -2 \Leftrightarrow x \geq -3$  or  $x \leq -7$ , so  $x \in (-\infty, -7] \cup [-3, \infty)$ .
52.  $|x + 1| \geq 3 \Leftrightarrow x + 1 \geq 3$  or  $x + 1 \leq -3 \Leftrightarrow x \geq 2$  or  $x \leq -4$ , so  $x \in (-\infty, -4] \cup [2, \infty)$ .
53.  $|2x - 3| \leq 0.4 \Leftrightarrow -0.4 \leq 2x - 3 \leq 0.4 \Leftrightarrow 2.6 \leq 2x \leq 3.4 \Leftrightarrow 1.3 \leq x \leq 1.7$ , so  $x \in [1.3, 1.7]$ .
54.  $|5x - 2| < 6 \Leftrightarrow -6 < 5x - 2 < 6 \Leftrightarrow -4 < 5x < 8 \Leftrightarrow -\frac{4}{5} < x < \frac{8}{5}$ , so  $x \in (-\frac{4}{5}, \frac{8}{5})$ .
55.  $1 \leq |x| \leq 4$ . So either  $1 \leq x \leq 4$  or  $1 \leq -x \leq 4 \Leftrightarrow -1 \geq x \geq -4$ . Thus,  $x \in [-4, -1] \cup [1, 4]$ .
56.  $0 < |x - 5| < \frac{1}{2}$ . Clearly  $0 < |x - 5|$  for  $x \neq 5$ . Now  $|x - 5| < \frac{1}{2} \Leftrightarrow -\frac{1}{2} < x - 5 < \frac{1}{2} \Leftrightarrow 4.5 < x < 5.5$ . So the solution set is  $(4.5, 5) \cup (5, 5.5)$ .
57.  $a(bx - c) \geq bc \Leftrightarrow bx - c \geq \frac{bc}{a} \Leftrightarrow bx \geq \frac{bc}{a} + c = \frac{bc + ac}{a} \Leftrightarrow x \geq \frac{bc + ac}{ab}$
58.  $a \leq bx + c < 2a \Leftrightarrow a - c \leq bx < 2a - c \Leftrightarrow \frac{a - c}{b} \leq x < \frac{2a - c}{b}$  (since  $b > 0$ )
59.  $ax + b < c \Leftrightarrow ax < c - b \Leftrightarrow x > \frac{c - b}{a}$  [since  $a < 0$ ]
60.  $\frac{ax + b}{c} \leq b \Leftrightarrow ax + b \geq bc$  [since  $c < 0$ ]  $\Leftrightarrow ax \geq bc - b \Leftrightarrow x \leq \frac{b(c - 1)}{a}$  [since  $a < 0$ ]
61.  $|(x + y) - 5| = |(x - 2) + (y - 3)| \leq |x - 2| + |y - 3| < 0.01 + 0.04 = 0.05$
62. Use the Triangle Inequality:  $|x + 3| < \frac{1}{2} \Rightarrow$   
 $|4x + 13| = |4(x + 3) + 1| \leq |4(x + 3)| + |1| = 4|x + 3| + 1 < 4\left(\frac{1}{2}\right) + 1 = 3$   
*Another method:*  $|x + 3| < \frac{1}{2} \Rightarrow -\frac{1}{2} < x + 3 < \frac{1}{2} \Rightarrow -2 < 4x + 12 < 2 \Rightarrow -1 < 4x + 13 < 3 \Rightarrow$   
 $|4x + 13| < 3$
63. If  $a < b$  then  $a + a < a + b$  and  $a + b < b + b$ . So  $2a < a + b < 2b$ . Dividing by 2, we get  $a < \frac{1}{2}(a + b) < b$ .

64. If  $0 < a < b$ , then  $\frac{1}{ab} > 0$ . So  $a < b \Rightarrow \frac{1}{ab} \cdot a < \frac{1}{ab} \cdot b \Leftrightarrow \frac{1}{b} < \frac{1}{a}$ .

65.  $|ab| = \sqrt{(ab)^2} = \sqrt{a^2b^2} = \sqrt{a^2} \sqrt{b^2} = |a| |b|$

66.  $\left|\frac{a}{b}\right| |b| = \left|\frac{a}{b} \cdot b\right| = |a|$  [using the result of Exercise 65]. Dividing the equation through by  $|b|$  gives  $\left|\frac{a}{b}\right| = \frac{|a|}{|b|}$ .

67. If  $0 < a < b$ , then  $a \cdot a < a \cdot b$  and  $a \cdot b < b \cdot b$  [using Rule 3 of Inequalities]. So  $a^2 < ab < b^2$  and hence  $a^2 < b^2$ .

68. Following the hint, the Triangle Inequality becomes  $|(x - y) + y| \leq |x - y| + |y| \Leftrightarrow |x| \leq |x - y| + |y| \Leftrightarrow |x - y| \geq |x| - |y|$ .

69. Observe that the sum, difference and product of two integers is always an integer. Let the rational numbers be represented

by  $r = m/n$  and  $s = p/q$  (where  $m, n, p$  and  $q$  are integers with  $n \neq 0, q \neq 0$ ). Now  $r + s = \frac{m}{n} + \frac{p}{q} = \frac{mq + pn}{nq}$ ,

but  $mq + pn$  and  $nq$  are both integers, so  $\frac{mq + pn}{nq} = r + s$  is a rational number by definition. Similarly,

$r - s = \frac{m}{n} - \frac{p}{q} = \frac{mq - pn}{nq}$  is a rational number. Finally,  $r \cdot s = \frac{m}{n} \cdot \frac{p}{q} = \frac{mp}{nq}$  but  $mp$  and  $nq$  are both integers, so

$\frac{mp}{nq} = r \cdot s$  is a rational number by definition.

70. (a) No. Consider the case of  $\sqrt{2}$  and  $-\sqrt{2}$ . Both are irrational numbers, yet  $\sqrt{2} + (-\sqrt{2}) = 0$  and 0, being an integer, is not irrational.

(b) No. Consider the case of  $\sqrt{2}$  and  $\sqrt{2}$ . Both are irrational numbers, yet  $\sqrt{2} \cdot \sqrt{2} = 2$  is not irrational.

## B Coordinate Geometry and Lines

1. Use the distance formula with  $P_1(x_1, y_1) = (1, 1)$  and  $P_2(x_2, y_2) = (4, 5)$  to get

$$|P_1P_2| = \sqrt{(4-1)^2 + (5-1)^2} = \sqrt{3^2 + 4^2} = \sqrt{25} = 5$$

2. The distance from  $(1, -3)$  to  $(5, 7)$  is  $\sqrt{(5-1)^2 + [7-(-3)]^2} = \sqrt{4^2 + 10^2} = \sqrt{116} = 2\sqrt{29}$ .

3. The distance from  $(6, -2)$  to  $(-1, 3)$  is  $\sqrt{(-1-6)^2 + [3-(-2)]^2} = \sqrt{(-7)^2 + 5^2} = \sqrt{74}$ .

4. The distance from  $(1, -6)$  to  $(-1, -3)$  is  $\sqrt{(-1-1)^2 + [-3-(-6)]^2} = \sqrt{(-2)^2 + 3^2} = \sqrt{13}$ .

5. The distance from  $(2, 5)$  to  $(4, -7)$  is  $\sqrt{(4-2)^2 + (-7-5)^2} = \sqrt{2^2 + (-12)^2} = \sqrt{148} = 2\sqrt{37}$ .

6. The distance from  $(a, b)$  to  $(b, a)$  is  $\sqrt{(b-a)^2 + (a-b)^2} = \sqrt{(a-b)^2 + (a-b)^2} = \sqrt{2(a-b)^2} = \sqrt{2}|a-b|$ .

7. The slope  $m$  of the line through  $P(1, 5)$  and  $Q(4, 11)$  is  $m = \frac{11-5}{4-1} = \frac{6}{3} = 2$ .

8. The slope  $m$  of the line through  $P(-1, 6)$  and  $Q(4, -3)$  is  $m = \frac{-3-6}{4-(-1)} = -\frac{9}{5}$ .

9. The slope  $m$  of the line through  $P(-3, 3)$  and  $Q(-1, -6)$  is  $m = \frac{-6 - 3}{-1 - (-3)} = -\frac{9}{2}$ .
10. The slope  $m$  of the line through  $P(-1, -4)$  and  $Q(6, 0)$  is  $m = \frac{0 - (-4)}{6 - (-1)} = \frac{4}{7}$ .
11. Using  $A(0, 2)$ ,  $B(-3, -1)$ , and  $C(-4, 3)$ , we have  $|AC| = \sqrt{(-4 - 0)^2 + (3 - 2)^2} = \sqrt{(-4)^2 + 1^2} = \sqrt{17}$  and  $|BC| = \sqrt{[-4 - (-3)]^2 + [3 - (-1)]^2} = \sqrt{(-1)^2 + 4^2} = \sqrt{17}$ , so the triangle has two sides of equal length, and is isosceles.
12. (a) Using  $A(6, -7)$ ,  $B(11, -3)$ , and  $C(2, -2)$ , we have  
 $|AB| = \sqrt{(11 - 6)^2 + [-3 - (-7)]^2} = \sqrt{5^2 + 4^2} = \sqrt{41}$ ,  
 $|AC| = \sqrt{(2 - 6)^2 + [-2 - (-7)]^2} = \sqrt{(-4)^2 + 5^2} = \sqrt{41}$ , and  
 $|BC| = \sqrt{(2 - 11)^2 + [-2 - (-3)]^2} = \sqrt{(-9)^2 + 1^2} = \sqrt{82}$ .  
 Thus,  $|AB|^2 + |AC|^2 = 41 + 41 = 82 = |BC|^2$  and so  $\triangle ABC$  is a right triangle.
- (b)  $m_{AB} = \frac{-3 - (-7)}{11 - 6} = \frac{4}{5}$  and  $m_{AC} = \frac{-2 - (-7)}{2 - 6} = -\frac{5}{4}$ . Thus  $m_{AB} \cdot m_{AC} = -1$  and so  $AB$  is perpendicular to  $AC$  and  $\triangle ABC$  must be a right triangle.
- (c) Taking lengths from part (a), the base is  $\sqrt{41}$  and the height is  $\sqrt{41}$ . Thus the area is  $\frac{1}{2}bh = \frac{1}{2}\sqrt{41}\sqrt{41} = \frac{41}{2}$ .
13. Using  $A(-2, 9)$ ,  $B(4, 6)$ ,  $C(1, 0)$ , and  $D(-5, 3)$ , we have  
 $|AB| = \sqrt{[4 - (-2)]^2 + (6 - 9)^2} = \sqrt{6^2 + (-3)^2} = \sqrt{45} = \sqrt{9} \sqrt{5} = 3\sqrt{5}$ ,  
 $|BC| = \sqrt{(1 - 4)^2 + (0 - 6)^2} = \sqrt{(-3)^2 + (-6)^2} = \sqrt{45} = \sqrt{9} \sqrt{5} = 3\sqrt{5}$ ,  
 $|CD| = \sqrt{(-5 - 1)^2 + (3 - 0)^2} = \sqrt{(-6)^2 + 3^2} = \sqrt{45} = \sqrt{9} \sqrt{5} = 3\sqrt{5}$ , and  
 $|DA| = \sqrt{[-2 - (-5)]^2 + (9 - 3)^2} = \sqrt{3^2 + 6^2} = \sqrt{45} = \sqrt{9} \sqrt{5} = 3\sqrt{5}$ . So all sides are of equal length and we have a rhombus. Moreover,  $m_{AB} = \frac{6 - 9}{4 - (-2)} = -\frac{1}{2}$ ,  $m_{BC} = \frac{0 - 6}{1 - 4} = 2$ ,  $m_{CD} = \frac{3 - 0}{-5 - 1} = -\frac{1}{2}$ , and  $m_{DA} = \frac{9 - 3}{-2 - (-5)} = 2$ , so the sides are perpendicular. Thus,  $A$ ,  $B$ ,  $C$ , and  $D$  are vertices of a square.
14. (a) Using  $A(-1, 3)$ ,  $B(3, 11)$ , and  $C(5, 15)$ , we have  
 $|AB| = \sqrt{[3 - (-1)]^2 + (11 - 3)^2} = \sqrt{4^2 + 8^2} = \sqrt{80} = 4\sqrt{5}$ ,  
 $|BC| = \sqrt{(5 - 3)^2 + (15 - 11)^2} = \sqrt{2^2 + 4^2} = \sqrt{20} = 2\sqrt{5}$ , and  
 $|AC| = \sqrt{[5 - (-1)]^2 + (15 - 3)^2} = \sqrt{6^2 + 12^2} = \sqrt{180} = 6\sqrt{5}$ . Thus,  $|AC| = |AB| + |BC|$ .
- (b)  $m_{AB} = \frac{11 - 3}{3 - (-1)} = \frac{8}{4} = 2$  and  $m_{AC} = \frac{15 - 3}{5 - (-1)} = \frac{12}{6} = 2$ . Since the segments  $AB$  and  $AC$  have the same slope,  $A$ ,  $B$  and  $C$  must be collinear.

15. For the vertices  $A(1, 1)$ ,  $B(7, 4)$ ,  $C(5, 10)$ , and  $D(-1, 7)$ , the slope of the line segment  $AB$  is  $\frac{4-1}{7-1} = \frac{1}{2}$ , the slope of  $CD$  is  $\frac{7-10}{-1-5} = \frac{1}{2}$ , the slope of  $BC$  is  $\frac{10-4}{5-7} = -3$ , and the slope of  $DA$  is  $\frac{1-7}{1-(-1)} = -3$ . So  $AB$  is parallel to  $CD$  and


$BC$  is parallel to  $DA$ . Hence  $ABCD$  is a parallelogram.

16. For the vertices  $A(1, 1)$ ,  $B(11, 3)$ ,  $C(10, 8)$ , and  $D(0, 6)$ , the slopes of the four sides are  $m_{AB} = \frac{3-1}{11-1} = \frac{1}{5}$ ,


$m_{BC} = \frac{8-3}{10-11} = -5$ ,  $m_{CD} = \frac{6-8}{0-10} = \frac{1}{5}$ , and  $m_{DA} = \frac{1-6}{1-0} = -5$ . Hence  $AB \parallel CD$ ,  $BC \parallel DA$ ,  $AB \perp BC$ ,

$BC \perp CD$ ,  $CD \perp DA$ , and  $DA \perp AB$ , and so  $ABCD$  is a rectangle.


17. The graph of the equation  $x = 3$  is a vertical line with  $x$ -intercept 3. The line does not have a slope.




18. The graph of the equation  $y = -2$  is a horizontal line with  $y$ -intercept  $-2$ . The line has slope 0.



19.  $xy = 0 \Leftrightarrow x = 0$  or  $y = 0$ . The graph consists of the coordinate axes.



20.  $|y| = 1 \Leftrightarrow y = 1$  or  $y = -1$



21. By the point-slope form of the equation of a line, an equation of the line through  $(2, -3)$  with slope 6 is

$$y - (-3) = 6(x - 2) \text{ or } y = 6x - 15.$$

22.  $y - 4 = -3[x - (-1)]$  or  $y = -3x + 1$

23.  $y - 7 = \frac{2}{3}(x - 1)$  or  $y = \frac{2}{3}x + \frac{19}{3}$

24.  $y - (-5) = -\frac{7}{2}[x - (-3)]$  or  $y = -\frac{7}{2}x - \frac{31}{2}$

25. The slope of the line through  $(2, 1)$  and  $(1, 6)$  is  $m = \frac{6-1}{1-2} = -5$ , so an equation of the line is

$$y - 1 = -5(x - 2) \text{ or } y = -5x + 11.$$



26. For  $(-1, -2)$  and  $(4, 3)$ ,  $m = \frac{3 - (-2)}{4 - (-1)} = 1$ . An equation of the line is  $y - 3 = 1(x - 4)$  or  $y = x - 1$ .

27. By the slope-intercept form of the equation of a line, an equation of the line is  $y = 3x - 2$ .

28. By the slope-intercept form of the equation of a line, an equation of the line is  $y = \frac{2}{5}x + 4$ .

29. Since the line passes through  $(1, 0)$  and  $(0, -3)$ , its slope is  $m = \frac{-3 - 0}{0 - 1} = 3$ , so an equation is  $y = 3x - 3$ .

*Another method:* From Exercise 61,  $\frac{x}{1} + \frac{y}{-3} = 1 \Rightarrow -3x + y = -3 \Rightarrow y = 3x - 3$ .

30. For  $(-8, 0)$  and  $(0, 6)$ ,  $m = \frac{6 - 0}{0 - (-8)} = \frac{3}{4}$ . So an equation is  $y = \frac{3}{4}x + 6$ .

*Another method:* From Exercise 61,  $\frac{x}{-8} + \frac{y}{6} = 1 \Rightarrow -3x + 4y = 24 \Rightarrow y = \frac{3}{4}x + 6$ .

31. The line is parallel to the  $x$ -axis, so it is horizontal and must have the form  $y = k$ . Since it goes through the point  $(x, y) = (4, 5)$ , the equation is  $y = 5$ .

32. The line is parallel to the  $y$ -axis, so it is vertical and must have the form  $x = k$ . Since it goes through the point  $(x, y) = (4, 5)$ , the equation is  $x = 4$ .


33. Putting the line  $x + 2y = 6$  into its slope-intercept form gives us  $y = -\frac{1}{2}x + 3$ , so we see that this line has slope  $-\frac{1}{2}$ . Thus, we want the line of slope  $-\frac{1}{2}$  that passes through the point  $(1, -6)$ :  $y - (-6) = -\frac{1}{2}(x - 1) \Leftrightarrow y = -\frac{1}{2}x - \frac{11}{2}$ .

34.  $2x + 3y + 4 = 0 \Leftrightarrow y = -\frac{2}{3}x - \frac{4}{3}$ , so  $m = -\frac{2}{3}$  and the required line is  $y = -\frac{2}{3}x + 6$ .


35.  $2x + 5y + 8 = 0 \Leftrightarrow y = -\frac{2}{5}x - \frac{8}{5}$ . Since this line has slope  $-\frac{2}{5}$ , a line perpendicular to it would have slope  $\frac{5}{2}$ , so the required line is  $y - (-2) = \frac{5}{2}[x - (-1)] \Leftrightarrow y = \frac{5}{2}x + \frac{1}{2}$ .

36.  $4x - 8y = 1 \Leftrightarrow y = \frac{1}{2}x - \frac{1}{8}$ . Since this line has slope  $\frac{1}{2}$ , a line perpendicular to it would have slope  $-2$ , so the required line is  $y - (-\frac{2}{3}) = -2(x - \frac{1}{2}) \Leftrightarrow y = -2x + \frac{1}{3}$ .


37.  $x + 3y = 0 \Leftrightarrow y = -\frac{1}{3}x$ , so the slope is  $-\frac{1}{3}$  and the  $y$ -intercept is 0.



38.  $2x - 5y = 0 \Leftrightarrow y = \frac{2}{5}x$ , so the slope is  $\frac{2}{5}$  and the  $y$ -intercept is 0.




39.  $y = -2$  is a horizontal line with slope 0 and  $y$ -intercept  $-2$ .




40.  $2x - 3y + 6 = 0 \Leftrightarrow$

$y = \frac{2}{3}x + 2$ , so the slope is  $\frac{2}{3}$   
and the  $y$ -intercept is 2.




41.  $3x - 4y = 12 \Leftrightarrow$

$y = \frac{3}{4}x - 3$ , so the slope is  $\frac{3}{4}$   
and the  $y$ -intercept is  $-3$ .




42.  $4x + 5y = 10 \Leftrightarrow$


$y = -\frac{4}{5}x + 2$ , so the slope is  $-\frac{4}{5}$   
and the  $y$ -intercept is 2.



43.  $\{(x, y) \mid x < 0\}$




44.  $\{(x, y) \mid y > 0\}$




45.  $\{(x, y) \mid xy < 0\} =$


$\{(x, y) \mid x < 0 \text{ and } y > 0\}$   
 $\cup \{(x, y) \mid x > 0 \text{ and } y < 0\}$




46.  $\{(x, y) \mid x \geq 1 \text{ and } y < 3\}$




47.  $\{(x, y) \mid |x| \leq 2\} =$   
 $\{(x, y) \mid -2 \leq x \leq 2\}$




48.  $\{(x, y) \mid |x| < 3 \text{ and } |y| < 2\}$




49.  $\{(x, y) \mid 0 \leq y \leq 4, x \leq 2\}$




50.  $\{(x, y) \mid y > 2x - 1\}$



51.  $\{(x, y) \mid 1 + x \leq y \leq 1 - 2x\}$



52.  $\{(x, y) \mid -x \leq y < \frac{1}{2}(x+3)\}$



53. Let  $P(0, y)$  be a point on the  $y$ -axis. The distance from  $P$  to  $(5, -5)$  is

$$\begin{aligned} \sqrt{(5-0)^2 + (-5-y)^2} &= \sqrt{5^2 + (y+5)^2}. \text{ The distance from } P \text{ to } (1, 1) \text{ is} \\ \sqrt{(1-0)^2 + (1-y)^2} &= \sqrt{1^2 + (y-1)^2}. \text{ We want these distances to be equal:} \\ \sqrt{5^2 + (y+5)^2} &= \sqrt{1^2 + (y-1)^2} \Leftrightarrow 5^2 + (y+5)^2 = 1^2 + (y-1)^2 \Leftrightarrow \\ 25 + (y^2 + 10y + 25) &= 1 + (y^2 - 2y + 1) \Leftrightarrow 12y = -48 \Leftrightarrow y = -4. \\ \text{So the desired point is } (0, -4). \end{aligned}$$

54. Let  $M$  be the point  $\left(\frac{x_1+x_2}{2}, \frac{y_1+y_2}{2}\right)$ . Then

$$\begin{aligned} |MP_1|^2 &= \left(x_1 - \frac{x_1+x_2}{2}\right)^2 + \left(y_1 - \frac{y_1+y_2}{2}\right)^2 = \left(\frac{x_1-x_2}{2}\right)^2 + \left(\frac{y_1-y_2}{2}\right)^2 \\ |MP_2|^2 &= \left(x_2 - \frac{x_1+x_2}{2}\right)^2 + \left(y_2 - \frac{y_1+y_2}{2}\right)^2 = \left(\frac{x_2-x_1}{2}\right)^2 + \left(\frac{y_2-y_1}{2}\right)^2 \end{aligned}$$

Hence,  $|MP_1| = |MP_2|$ ; that is,  $M$  is equidistant from  $P_1$  and  $P_2$ .

55. (a) Using the midpoint formula from Exercise 54 with  $(1, 3)$  and  $(7, 15)$ , we get  $\left(\frac{1+7}{2}, \frac{3+15}{2}\right) = (4, 9)$ .

(b) Using the midpoint formula from Exercise 54 with  $(-1, 6)$  and  $(8, -12)$ , we get  $\left(\frac{-1+8}{2}, \frac{6+(-12)}{2}\right) = \left(\frac{7}{2}, -3\right)$ .

56. With  $A(1, 0)$ ,  $B(3, 6)$ , and  $C(8, 2)$ , the midpoint  $M_1$  of  $AB$  is  $\left(\frac{1+3}{2}, \frac{0+6}{2}\right) = (2, 3)$ , the midpoint  $M_2$  of  $BC$  is  $\left(\frac{3+8}{2}, \frac{6+2}{2}\right) = \left(\frac{11}{2}, 4\right)$ , and the midpoint  $M_3$  of  $CA$  is  $\left(\frac{8+1}{2}, \frac{2+0}{2}\right) = \left(\frac{9}{2}, 1\right)$ . The lengths of the medians are

$$\begin{aligned} |AM_2| &= \sqrt{\left(\frac{11}{2} - 1\right)^2 + (4 - 0)^2} = \sqrt{\left(\frac{9}{2}\right)^2 + 4^2} = \sqrt{\frac{145}{4}} = \frac{\sqrt{145}}{2} \\ |BM_3| &= \sqrt{\left(\frac{9}{2} - 3\right)^2 + (1 - 6)^2} = \sqrt{\left(\frac{3}{2}\right)^2 + (-5)^2} = \sqrt{\frac{109}{4}} = \frac{\sqrt{109}}{2} \\ |CM_1| &= \sqrt{(2 - 8)^2 + (3 - 2)^2} = \sqrt{(-6)^2 + 1^2} = \sqrt{37} \end{aligned}$$

57.  $2x - y = 4 \Leftrightarrow y = 2x - 4 \Rightarrow m_1 = 2$  and  $6x - 2y = 10 \Leftrightarrow 2y = 6x - 10 \Leftrightarrow y = 3x - 5 \Rightarrow m_2 = 3$ . Since  $m_1 \neq m_2$ , the two lines are not parallel. To find the point of intersection:  $2x - 4 = 3x - 5 \Leftrightarrow x = 1 \Rightarrow y = -2$ . Thus, the point of intersection is  $(1, -2)$ .

58.  $3x - 5y + 19 = 0 \Leftrightarrow 5y = 3x + 19 \Leftrightarrow y = \frac{3}{5}x + \frac{19}{5} \Rightarrow m_1 = \frac{3}{5}$  and  $10x + 6y - 50 = 0 \Leftrightarrow 6y = -10x + 50 \Leftrightarrow y = -\frac{5}{3}x + \frac{25}{3} \Rightarrow m_2 = -\frac{5}{3}$ . Since  $m_1 m_2 = \frac{3}{5} \left(-\frac{5}{3}\right) = -1$ , the two lines are perpendicular. To find the point of intersection:  $\frac{3}{5}x + \frac{19}{5} = -\frac{5}{3}x + \frac{25}{3} \Leftrightarrow 9x + 57 = -25x + 125 \Leftrightarrow 34x = 68 \Leftrightarrow x = 2 \Rightarrow y = \frac{3}{5} \cdot 2 + \frac{19}{5} = \frac{25}{5} = 5$ . Thus, the point of intersection is  $(2, 5)$ .

59. With  $A(1, 4)$  and  $B(7, -2)$ , the slope of segment  $AB$  is  $\frac{-2-4}{7-1} = -1$ , so its perpendicular bisector has slope 1. The midpoint of  $AB$  is  $\left(\frac{1+7}{2}, \frac{4+(-2)}{2}\right) = (4, 1)$ , so an equation of the perpendicular bisector is  $y - 1 = 1(x - 4)$  or  $y = x - 3$ .

60. (a) Side  $PQ$  has slope  $\frac{4-0}{3-1} = 2$ , so its equation is  $y - 0 = 2(x - 1) \Leftrightarrow y = 2x - 2$ . Side  $QR$  has slope  $\frac{6-4}{-1-3} = -\frac{1}{2}$ , so its equation is  $y - 4 = -\frac{1}{2}(x - 3) \Leftrightarrow y = -\frac{1}{2}x + \frac{11}{2}$ . Side  $RP$  has slope  $\frac{0-6}{1-(-1)} = -3$ , so its equation is  $y - 0 = -3(x - 1) \Leftrightarrow y = -3x + 3$ .


(b)  $M_1$  (the midpoint of  $PQ$ ) has coordinates  $(\frac{1+3}{2}, \frac{0+4}{2}) = (2, 2)$ .  $M_2$  (the midpoint of  $QR$ ) has coordinates  $(\frac{3-1}{2}, \frac{4+6}{2}) = (1, 5)$ .  $M_3$  (the midpoint of  $RP$ ) has coordinates  $(\frac{1-1}{2}, \frac{0+6}{2}) = (0, 3)$ .  $RM_1$  has slope  $\frac{2-6}{2-(-1)} = -\frac{4}{3}$  and hence equation  $y - 2 = -\frac{4}{3}(x - 2) \Leftrightarrow y = -\frac{4}{3}x + \frac{14}{3}$ .  $PM_2$  is a vertical line with equation  $x = 1$ .  $QM_3$  has slope  $\frac{3-4}{0-3} = \frac{1}{3}$  and hence equation  $y - 3 = \frac{1}{3}(x - 0) \Leftrightarrow y = \frac{1}{3}x + 3$ .  $PM_2$  and  $RM_1$  intersect where  $x = 1$  and  $y = -\frac{4}{3}(1) + \frac{14}{3} = \frac{10}{3}$ , or at  $(1, \frac{10}{3})$ .  $PM_2$  and  $QM_3$  intersect where  $x = 1$  and  $y = \frac{1}{3}(1) + 3 = \frac{10}{3}$ , or at  $(1, \frac{10}{3})$ , so this is the point where all three medians intersect.

61. (a) Since the  $x$ -intercept is  $a$ , the point  $(a, 0)$  is on the line, and similarly since the  $y$ -intercept is  $b$ ,  $(0, b)$  is on the line. Hence, the slope of the line is  $m = \frac{b-0}{0-a} = -\frac{b}{a}$ . Substituting into  $y = mx + b$  gives  $y = -\frac{b}{a}x + b \Leftrightarrow \frac{b}{a}x + y = b \Leftrightarrow \frac{x}{a} + \frac{y}{b} = 1$ .

(b) Letting  $a = 6$  and  $b = -8$  gives  $\frac{x}{6} + \frac{y}{-8} = 1 \Leftrightarrow -8x + 6y = -48$  [multiply by  $-48$ ]  $\Leftrightarrow 6y = 8x - 48 \Leftrightarrow 3y = 4x - 24 \Leftrightarrow y = \frac{4}{3}x - 8$ .

62. (a) Let  $d$  = distance traveled (in miles) and  $t$  = time elapsed (in hours). At  $t = 0$ ,  $d = 0$  and at  $t = 50$  minutes  $= 50 \cdot \frac{1}{60} = \frac{5}{6}$  h,  $d = 40$ . Thus, we have two points:  $(0, 0)$  and  $(\frac{5}{6}, 40)$ , so  $m = \frac{40-0}{5/6-0} = 48$  and  $d = 48t$ .

(c) The slope is 48 and represents the car's speed in mi/h.



## C Graphs of Second-Degree Equations

1. An equation of the circle with center  $(3, -1)$  and radius 5 is  $(x - 3)^2 + (y + 1)^2 = 5^2 = 25$ .

2. An equation of the circle with center  $(-2, -8)$  and radius 10 is  $(x + 2)^2 + (y + 8)^2 = 10^2 = 100$ .

3. The equation has the form  $x^2 + y^2 = r^2$ . Since  $(4, 7)$  lies on the circle, we have  $4^2 + 7^2 = r^2 \Rightarrow r^2 = 65$ . So the required equation is  $x^2 + y^2 = 65$ .

4. The equation has the form  $(x + 1)^2 + (y - 5)^2 = r^2$ . Since  $(-4, -6)$  lies on the circle, we have  $r^2 = (-4 + 1)^2 + (-6 - 5)^2 = 130$ . So an equation is  $(x + 1)^2 + (y - 5)^2 = 130$ .

5.  $x^2 + y^2 - 4x + 10y + 13 = 0 \Leftrightarrow x^2 - 4x + y^2 + 10y = -13 \Leftrightarrow (x^2 - 4x + 4) + (y^2 + 10y + 25) = -13 + 4 + 25 = 16 \Leftrightarrow (x - 2)^2 + (y + 5)^2 = 4^2$ . Thus, we have a circle with center  $(2, -5)$  and radius 4.

6.  $x^2 + y^2 + 6y + 2 = 0 \Leftrightarrow x^2 + (y^2 + 6y + 9) = -2 + 9 \Leftrightarrow x^2 + (y + 3)^2 = 7$ . Thus, we have a circle with center  $(0, -3)$  and radius  $\sqrt{7}$ .


7.  $x^2 + y^2 + x = 0 \Leftrightarrow (x^2 + x + \frac{1}{4}) + y^2 = \frac{1}{4} \Leftrightarrow (x + \frac{1}{2})^2 + y^2 = (\frac{1}{2})^2$ . Thus, we have a circle with center  $(-\frac{1}{2}, 0)$  and radius  $\frac{1}{2}$ .

8.  $16x^2 + 16y^2 + 8x + 32y + 1 = 0 \Leftrightarrow 16(x^2 + \frac{1}{2}x + \frac{1}{16}) + 16(y^2 + 2y + 1) = -1 + 1 + 16 \Leftrightarrow 16(x + \frac{1}{4})^2 + 16(y + 1)^2 = 16 \Leftrightarrow (x + \frac{1}{4})^2 + (y + 1)^2 = 1$ . Thus, we have a circle with center  $(-\frac{1}{4}, -1)$  and radius 1.


9.  $2x^2 + 2y^2 - x + y = 1 \Leftrightarrow 2(x^2 - \frac{1}{2}x + \frac{1}{16}) + 2(y^2 + \frac{1}{2}y + \frac{1}{16}) = 1 + \frac{1}{8} + \frac{1}{8} \Leftrightarrow 2(x - \frac{1}{4})^2 + 2(y + \frac{1}{4})^2 = \frac{5}{4} \Leftrightarrow (x - \frac{1}{4})^2 + (y + \frac{1}{4})^2 = \frac{5}{8}$ . Thus, we have a circle with center  $(\frac{1}{4}, -\frac{1}{4})$  and radius  $\frac{\sqrt{5}}{2\sqrt{2}} = \frac{\sqrt{10}}{4}$ .

10.  $x^2 + y^2 + ax + by + c = 0 \Leftrightarrow (x^2 + ax + \frac{1}{4}a^2) + (y^2 + by + \frac{1}{4}b^2) = -c + \frac{1}{4}a^2 + \frac{1}{4}b^2 \Leftrightarrow (x + \frac{1}{2}a)^2 + (y + \frac{1}{2}b)^2 = \frac{1}{4}(a^2 + b^2 - 4c)$ . For this to represent a nondegenerate circle,  $\frac{1}{4}(a^2 + b^2 - 4c) > 0$  or  $a^2 + b^2 > 4c$ . If this condition is satisfied, the circle has center  $(-\frac{1}{2}a, -\frac{1}{2}b)$  and radius  $\frac{1}{2}\sqrt{a^2 + b^2 - 4c}$ .


11.  $y = -x^2$ . Parabola




12.  $y^2 - x^2 = 1$ . Hyperbola




13.  $x^2 + 4y^2 = 16 \Leftrightarrow \frac{x^2}{16} + \frac{y^2}{4} = 1$ . Ellipse




14.  $x = -2y^2$ . Parabola




15.  $16x^2 - 25y^2 = 400 \Leftrightarrow \frac{x^2}{25} - \frac{y^2}{16} = 1$ . Hyperbola




16.  $25x^2 + 4y^2 = 100 \Leftrightarrow \frac{x^2}{4} + \frac{y^2}{25} = 1$ . Ellipse




17.  $4x^2 + y^2 = 1 \Leftrightarrow \frac{x^2}{1/4} + y^2 = 1$ . Ellipse




18.  $y = x^2 + 2$ . Parabola with vertex at (0, 2)




19.  $x = y^2 - 1$ . Parabola with vertex at (-1, 0)




20.  $9x^2 - 25y^2 = 225 \Leftrightarrow \frac{x^2}{25} - \frac{y^2}{9} = 1$ . Hyperbola




21.  $9y^2 - x^2 = 9 \Leftrightarrow y^2 - \frac{x^2}{9} = 1$ . Hyperbola



22.  $2x^2 + 5y^2 = 10 \Leftrightarrow \frac{x^2}{5} + \frac{y^2}{2} = 1$ . Ellipse




23.  $xy = 4$ . Hyperbola




24.  $y = x^2 + 2x = (x^2 + 2x + 1) - 1 = (x + 1)^2 - 1$ .

Parabola with vertex at (-1, -1)



25.  $9(x - 1)^2 + 4(y - 2)^2 = 36 \Leftrightarrow$


$\frac{(x - 1)^2}{4} + \frac{(y - 2)^2}{9} = 1$ . Ellipse centered at (1, 2)



26.  $16x^2 + 9y^2 - 36y = 108 \Leftrightarrow$


$16x^2 + 9(y^2 - 4y + 4) = 108 + 36 = 144 \Leftrightarrow$

$\frac{x^2}{9} + \frac{(y - 2)^2}{16} = 1$ . Ellipse centered at (0, 2)



27.  $y = x^2 - 6x + 13 = (x^2 - 6x + 9) + 4 = (x - 3)^2 + 4$ .


Parabola with vertex at (3, 4)




28.  $x^2 - y^2 - 4x + 3 = 0 \Leftrightarrow$

$(x^2 - 4x + 4) - y^2 = -3 + 4 = 1 \Leftrightarrow$

$(x - 2)^2 - y^2 = 1$ . Hyperbola centered at (2, 0)




29.  $x = 4 - y^2 = -y^2 + 4$ . Parabola with vertex at (4, 0)



30.  $y^2 - 2x + 6y + 5 = 0 \Leftrightarrow y^2 + 6y + 9 = 2x + 4 \Leftrightarrow$


$(y + 3)^2 = 2(x + 2)$ . Parabola with vertex (-2, -3)



31.  $x^2 + 4y^2 - 6x + 5 = 0 \Leftrightarrow$

$(x^2 - 6x + 9) + 4y^2 = -5 + 9 = 4 \Leftrightarrow$


$\frac{(x - 3)^2}{4} + y^2 = 1$ . Ellipse centered at (3, 0)



32.  $4x^2 + 9y^2 - 16x + 54y + 61 = 0 \Leftrightarrow$


$4(x^2 - 4x + 4) + 9(y^2 + 6y + 9) = -61 + 16 + 81 = 36$

$\Leftrightarrow \frac{(x - 2)^2}{9} + \frac{(y + 3)^2}{4} = 1$ . Ellipse centered at (2, -3)



33.  $y = 3x$  and  $y = x^2$  intersect where  $3x = x^2 \Leftrightarrow$

$0 = x^2 - 3x = x(x - 3)$ , that is, at (0, 0) and (3, 9).




34.  $y = 4 - x^2$ ,  $x - 2y = 2$ . Substitute  $y$  from the first

equation into the second:  $x - 2(4 - x^2) = 2 \Leftrightarrow$

$2x^2 + x - 10 = 0 \Leftrightarrow (2x + 5)(x - 2) = 0 \Leftrightarrow$

$x = -\frac{5}{2}$  or 2. So the points of intersection are  $(-\frac{5}{2}, -\frac{9}{4})$  and (2, 0).



35. The parabola must have an equation of the form  $y = a(x - 1)^2 - 1$ . Substituting  $x = 3$  and  $y = 3$  into the equation gives  $3 = a(3 - 1)^2 - 1$ , so  $a = 1$ , and the equation is  $y = (x - 1)^2 - 1 = x^2 - 2x$ . Note that using the other point  $(-1, 3)$  would have given the same value for  $a$ , and hence the same equation.

36. The ellipse has an equation of the form  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ . Substituting  $x = 1$  and  $y = -\frac{10\sqrt{2}}{3}$  gives


$$\frac{1^2}{a^2} + \frac{(-10\sqrt{2}/3)^2}{b^2} = \frac{1}{a^2} + \frac{200}{9b^2} = 1. \text{ Substituting } x = -2 \text{ and } y = \frac{5\sqrt{5}}{3} \text{ gives } \frac{(-2)^2}{a^2} + \frac{(5\sqrt{5}/3)^2}{b^2} = \frac{4}{a^2} + \frac{125}{9b^2} = 1.$$

From the first equation,  $\frac{1}{a^2} = 1 - \frac{200}{9b^2}$ . Putting this into the second equation gives  $4\left(1 - \frac{200}{9b^2}\right) + \frac{125}{9b^2} = 1 \Leftrightarrow$


$$3 = \frac{675}{9b^2} \Leftrightarrow b^2 = \frac{675}{27} = 25, \text{ so } b = 5. \text{ Hence } \frac{1}{a^2} = 1 - \frac{200}{9(5)^2} = \frac{1}{9} \text{ and so } a = 3. \text{ The equation of the ellipse}$$

$$\text{is } \frac{x^2}{9} + \frac{y^2}{25} = 1.$$


37.  $\{(x, y) \mid x^2 + y^2 \leq 1\}$




38.  $\{(x, y) \mid x^2 + y^2 > 4\}$



39.  $\{(x, y) \mid y \geq x^2 - 1\}$



40.  $\{(x, y) \mid x^2 + 4y^2 \leq 4\}$



## D Trigonometry

1.  $210^\circ = 210^\circ \left(\frac{\pi}{180^\circ}\right) = \frac{7\pi}{6} \text{ rad}$

2.  $300^\circ = 300^\circ \left(\frac{\pi}{180^\circ}\right) = \frac{5\pi}{3} \text{ rad}$

3.  $9^\circ = 9^\circ \left(\frac{\pi}{180^\circ}\right) = \frac{\pi}{20} \text{ rad}$

4.  $-315^\circ = -315^\circ \left(\frac{\pi}{180^\circ}\right) = -\frac{7\pi}{4} \text{ rad}$

5.  $900^\circ = 900^\circ \left(\frac{\pi}{180^\circ}\right) = 5\pi \text{ rad}$

6.  $36^\circ = 36^\circ \left(\frac{\pi}{180^\circ}\right) = \frac{\pi}{5} \text{ rad}$

7.  $4\pi \text{ rad} = 4\pi \left(\frac{180^\circ}{\pi}\right) = 720^\circ$

8.  $-\frac{7\pi}{2} \text{ rad} = -\frac{7\pi}{2} \left(\frac{180^\circ}{\pi}\right) = -630^\circ$

9.  $\frac{5\pi}{12} \text{ rad} = \frac{5\pi}{12} \left(\frac{180^\circ}{\pi}\right) = 75^\circ$

10.  $\frac{8\pi}{3} \text{ rad} = \frac{8\pi}{3} \left(\frac{180^\circ}{\pi}\right) = 480^\circ$



11.  $-\frac{3\pi}{8} \text{ rad} = -\frac{3\pi}{8} \left( \frac{180^\circ}{\pi} \right) = -67.5^\circ$








12.  $5 \text{ rad} = 5 \left( \frac{180^\circ}{\pi} \right) = \left( \frac{900}{\pi} \right)^\circ$

13. Using Formula 3,  $a = r\theta = 36 \cdot \frac{\pi}{12} = 3\pi \text{ cm}$ .

14. Using Formula 3,  $a = r\theta = 10 \cdot 72^\circ \left( \frac{\pi}{180^\circ} \right) = 4\pi \text{ cm}$ .

15. Using Formula 3,  $\theta = a/r = \frac{1}{1.5} = \frac{2}{3} \text{ rad} = \frac{2}{3} \left( \frac{180^\circ}{\pi} \right) = \left( \frac{120}{\pi} \right)^\circ \approx 38.2^\circ$ .


16.  $a = r\theta \Rightarrow r = \frac{a}{\theta} = \frac{6}{3\pi/4} = \frac{8}{\pi} \text{ cm}$



From the diagram we see that a point on the terminal side is  $P(-1, 1)$ .

Therefore, taking  $x = -1$ ,  $y = 1$ ,  $r = \sqrt{2}$  in the definitions of the trigonometric ratios, we have  $\sin \frac{3\pi}{4} = \frac{1}{\sqrt{2}}$ ,  $\cos \frac{3\pi}{4} = -\frac{1}{\sqrt{2}}$ ,

$\tan \frac{3\pi}{4} = -1$ ,  $\csc \frac{3\pi}{4} = \sqrt{2}$ ,  $\sec \frac{3\pi}{4} = -\sqrt{2}$ , and  $\cot \frac{3\pi}{4} = -1$ .




From the diagram and Figure 8, we see that a point on the terminal side is

$P(-1, -\sqrt{3})$ . Therefore, taking  $x = -1$ ,  $y = -\sqrt{3}$ ,  $r = 2$  in the

definitions of the trigonometric ratios, we have  $\sin \frac{4\pi}{3} = -\frac{\sqrt{3}}{2}$ ,

$\cos \frac{4\pi}{3} = -\frac{1}{2}$ ,  $\tan \frac{4\pi}{3} = \sqrt{3}$ ,  $\csc \frac{4\pi}{3} = -\frac{2}{\sqrt{3}}$ ,  $\sec \frac{4\pi}{3} = -2$ , and

$\cot \frac{4\pi}{3} = \frac{1}{\sqrt{3}}$ .



From the diagram we see that a point on the terminal side is  $P(0, 1)$ .


Therefore taking  $x = 0$ ,  $y = 1$ ,  $r = 1$  in the definitions of the

trigonometric ratios, we have  $\sin \frac{9\pi}{2} = 1$ ,  $\cos \frac{9\pi}{2} = 0$ ,  $\tan \frac{9\pi}{2} = y/x$  is

undefined since  $x = 0$ ,  $\csc \frac{9\pi}{2} = 1$ ,  $\sec \frac{9\pi}{2} = r/x$  is undefined since


$x = 0$ , and  $\cot \frac{9\pi}{2} = 0$ .

26.




From the diagram, we see that a point on the terminal side is  $P(-1, 0)$ . Therefore taking  $x = -1$ ,  $y = 0$ ,  $r = 1$  in the definitions of the trigonometric ratios we have  $\sin(-5\pi) = 0$ ,  $\cos(-5\pi) = -1$ ,  $\tan(-5\pi) = 0$ ,  $\csc(-5\pi)$  is undefined,  $\sec(-5\pi) = -1$ , and  $\cot(-5\pi)$  is undefined.

27.



Using Figure 8 we see that a point on the terminal side is  $P(-\sqrt{3}, 1)$ . Therefore taking  $x = -\sqrt{3}$ ,  $y = 1$ ,  $r = 2$  in the definitions of the trigonometric ratios, we have  $\sin \frac{5\pi}{6} = \frac{1}{2}$ ,  $\cos \frac{5\pi}{6} = -\frac{\sqrt{3}}{2}$ ,  $\tan \frac{5\pi}{6} = -\frac{1}{\sqrt{3}}$ ,  $\csc \frac{5\pi}{6} = 2$ ,  $\sec \frac{5\pi}{6} = -\frac{2}{\sqrt{3}}$ , and  $\cot \frac{5\pi}{6} = -\sqrt{3}$ .

28.



From the diagram, we see that a point on the terminal side is  $P(-1, 1)$ . Therefore taking  $x = -1$ ,  $y = 1$ ,  $r = \sqrt{2}$  in the definitions of the trigonometric ratios we have  $\sin \frac{11\pi}{4} = \frac{1}{\sqrt{2}}$ ,  $\cos \frac{11\pi}{4} = -\frac{1}{\sqrt{2}}$ ,  $\tan \frac{11\pi}{4} = -1$ ,  $\csc \frac{11\pi}{4} = \sqrt{2}$ ,  $\sec \frac{11\pi}{4} = -\sqrt{2}$ , and  $\cot \frac{11\pi}{4} = -1$ .

29.  $\sin \theta = y/r = \frac{3}{5} \Rightarrow y = 3$ ,  $r = 5$ , and  $x = \sqrt{r^2 - y^2} = 4$  (since  $0 < \theta < \frac{\pi}{2}$ ). Therefore taking  $x = 4$ ,  $y = 3$ ,  $r = 5$  in the definitions of the trigonometric ratios, we have  $\cos \theta = \frac{4}{5}$ ,  $\tan \theta = \frac{3}{4}$ ,  $\csc \theta = \frac{5}{3}$ ,  $\sec \theta = \frac{5}{4}$ , and  $\cot \theta = \frac{4}{3}$ .

30. Since  $0 < \alpha < \frac{\pi}{2}$ ,  $\alpha$  is in the first quadrant where  $x$  and  $y$  are both positive. Therefore,  $\tan \alpha = y/x = \frac{2}{1} \Rightarrow y = 2$ ,  $x = 1$ , and  $r = \sqrt{x^2 + y^2} = \sqrt{5}$ . Taking  $x = 1$ ,  $y = 2$ ,  $r = \sqrt{5}$  in the definitions of the trigonometric ratios, we have  $\sin \alpha = \frac{2}{\sqrt{5}}$ ,  $\cos \alpha = \frac{1}{\sqrt{5}}$ ,  $\csc \alpha = \frac{\sqrt{5}}{2}$ ,  $\sec \alpha = \sqrt{5}$ , and  $\cot \alpha = \frac{1}{2}$ .

31.  $\frac{\pi}{2} < \phi < \pi \Rightarrow \phi$  is in the second quadrant, where  $x$  is negative and  $y$  is positive. Therefore  $\sec \phi = r/x = -1.5 = -\frac{3}{2} \Rightarrow r = 3$ ,  $x = -2$ , and  $y = \sqrt{r^2 - x^2} = \sqrt{5}$ . Taking  $x = -2$ ,  $y = \sqrt{5}$ , and  $r = 3$  in the definitions of the trigonometric ratios, we have  $\sin \phi = \frac{\sqrt{5}}{3}$ ,  $\cos \phi = -\frac{2}{3}$ ,  $\tan \phi = -\frac{\sqrt{5}}{2}$ ,  $\csc \phi = \frac{3}{\sqrt{5}}$ , and  $\cot \theta = -\frac{2}{\sqrt{5}}$ .

32. Since  $\pi < x < \frac{3\pi}{2}$ ,  $x$  is in the third quadrant where  $x$  and  $y$  are both negative. Therefore  $\cos x = x/r = -\frac{1}{3} \Rightarrow x = -1$ ,  $r = 3$ , and  $y = -\sqrt{r^2 - x^2} = -\sqrt{8} = -2\sqrt{2}$ . Taking  $x = -1$ ,  $r = 3$ ,  $y = -2\sqrt{2}$  in the definitions of the trigonometric ratios, we have  $\sin x = -\frac{2\sqrt{2}}{3}$ ,  $\tan x = 2\sqrt{2}$ ,  $\csc x = -\frac{3}{2\sqrt{2}}$ ,  $\sec x = -3$ , and  $\cot x = \frac{1}{2\sqrt{2}}$ .

33.  $\pi < \beta < 2\pi$  means that  $\beta$  is in the third or fourth quadrant where  $y$  is negative. Also since  $\cot \beta = x/y = 3$  which is positive,  $x$  must also be negative. Therefore  $\cot \beta = x/y = \frac{3}{1} \Rightarrow x = -3$ ,  $y = -1$ , and  $r = \sqrt{x^2 + y^2} = \sqrt{10}$ . Taking  $x = -3$ ,  $y = -1$  and  $r = \sqrt{10}$  in the definitions of the trigonometric ratios, we have  $\sin \beta = -\frac{1}{\sqrt{10}}$ ,  $\cos \beta = -\frac{3}{\sqrt{10}}$ ,  $\tan \beta = \frac{1}{3}$ ,  $\csc \beta = -\sqrt{10}$ , and  $\sec \beta = -\frac{\sqrt{10}}{3}$ .


34. Since  $\frac{3\pi}{2} < \theta < 2\pi$ ,  $\theta$  is in the fourth quadrant where  $x$  is positive and  $y$  is negative. Therefore  $\csc \theta = r/y = -\frac{4}{3} \Rightarrow r = 4$ ,  $y = -3$ , and  $x = \sqrt{r^2 - y^2} = \sqrt{7}$ . Taking  $x = \sqrt{7}$ ,  $y = -3$ , and  $r = 4$  in the definitions of the trigonometric ratios, we have  $\sin \theta = -\frac{3}{4}$ ,  $\cos \theta = \frac{\sqrt{7}}{4}$ ,  $\tan \theta = -\frac{3}{\sqrt{7}}$ ,  $\sec \theta = \frac{4}{\sqrt{7}}$ , and  $\cot \theta = -\frac{\sqrt{7}}{3}$ .

35.  $\sin 35^\circ = \frac{x}{10} \Rightarrow x = 10 \sin 35^\circ \approx 5.73576 \text{ cm}$

36.  $\cos 40^\circ = \frac{x}{25} \Rightarrow x = 25 \cos 40^\circ \approx 19.15111 \text{ cm}$

37.  $\tan \frac{2\pi}{5} = \frac{x}{8} \Rightarrow x = 8 \tan \frac{2\pi}{5} \approx 24.62147 \text{ cm}$

38.  $\cos \frac{3\pi}{8} = \frac{22}{x} \Rightarrow x = \frac{22}{\cos \frac{3\pi}{8}} \approx 57.48877 \text{ cm}$



(a) From the diagram we see that  $\sin \theta = \frac{y}{r} = \frac{a}{c}$ , and  $\sin(-\theta) = \frac{-a}{c} = -\frac{a}{c} = -\sin \theta$ .

(b) Again from the diagram we see that  $\cos \theta = \frac{x}{r} = \frac{b}{c} = \cos(-\theta)$ .

40. (a) Using (12a) and (12b), we have

$$\tan(x+y) = \frac{\sin(x+y)}{\cos(x+y)} = \frac{\sin x \cos y + \cos x \sin y}{\cos x \cos y - \sin x \sin y} = \frac{\frac{\sin x \cos y}{\cos x \cos y} + \frac{\cos x \sin y}{\cos x \cos y}}{\frac{\cos x \cos y}{\cos x \cos y} - \frac{\sin x \sin y}{\cos x \cos y}} = \frac{\tan x + \tan y}{1 - \tan x \tan y}$$

- (b) From (10a) and (10b), we have  $\tan(-\theta) = -\tan \theta$ , so (14a) implies that

$$\tan(x-y) = \tan(x+(-y)) = \frac{\tan x + \tan(-y)}{1 - \tan x \tan(-y)} = \frac{\tan x - \tan y}{1 + \tan x \tan y}$$

41. (a) Using (12a) and (13a), we have

$$\frac{1}{2}[\sin(x+y) + \sin(x-y)] = \frac{1}{2}[\sin x \cos y + \cos x \sin y + \sin x \cos y - \cos x \sin y] = \frac{1}{2}(2 \sin x \cos y) = \sin x \cos y.$$

- (b) This time, using (12b) and (13b), we have

$$\frac{1}{2}[\cos(x+y) + \cos(x-y)] = \frac{1}{2}[\cos x \cos y - \sin x \sin y + \cos x \cos y + \sin x \sin y] = \frac{1}{2}(2 \cos x \cos y) = \cos x \cos y.$$

- (c) Again using (12b) and (13b), we have

$$\begin{aligned} \frac{1}{2}[\cos(x-y) - \cos(x+y)] &= \frac{1}{2}[\cos x \cos y + \sin x \sin y - \cos x \cos y + \sin x \sin y] \\ &= \frac{1}{2}(2 \sin x \sin y) = \sin x \sin y \end{aligned}$$

42. Using (13b),  $\cos\left(\frac{\pi}{2} - x\right) = \cos \frac{\pi}{2} \cos x + \sin \frac{\pi}{2} \sin x = 0 \cdot \cos x + 1 \cdot \sin x = \sin x$ .

43. Using (12a), we have  $\sin\left(\frac{\pi}{2} + x\right) = \sin \frac{\pi}{2} \cos x + \cos \frac{\pi}{2} \sin x = 1 \cdot \cos x + 0 \cdot \sin x = \cos x$ .

44. Using (13a), we have  $\sin(\pi - x) = \sin \pi \cos x - \cos \pi \sin x = 0 \cdot \cos x - (-1) \sin x = \sin x$ .

45. Using (6), we have  $\sin \theta \cot \theta = \sin \theta \cdot \frac{\cos \theta}{\sin \theta} = \cos \theta$ .

46.  $(\sin x + \cos x)^2 = \sin^2 x + 2 \sin x \cos x + \cos^2 x = (\sin^2 x + \cos^2 x) + \sin 2x$  [by (15a)]  $= 1 + \sin 2x$  [by (7)]

$$47. \sec y - \cos y = \frac{1}{\cos y} - \cos y \text{ [by (6)]} = \frac{1 - \cos^2 y}{\cos y} = \frac{\sin^2 y}{\cos y} \text{ [by (7)]} = \frac{\sin y}{\cos y} \sin y = \tan y \sin y \text{ [by (6)]}$$

$$48. \tan^2 \alpha - \sin^2 \alpha = \frac{\sin^2 \alpha}{\cos^2 \alpha} - \sin^2 \alpha = \frac{\sin^2 \alpha - \sin^2 \alpha \cos^2 \alpha}{\cos^2 \alpha} = \frac{\sin^2 \alpha (1 - \cos^2 \alpha)}{\cos^2 \alpha} = \tan^2 \alpha \sin^2 \alpha \text{ [by (6), (7)]}$$

$$\begin{aligned} 49. \cot^2 \theta + \sec^2 \theta &= \frac{\cos^2 \theta}{\sin^2 \theta} + \frac{1}{\cos^2 \theta} \text{ [by (6)]} = \frac{\cos^2 \theta \cos^2 \theta + \sin^2 \theta}{\sin^2 \theta \cos^2 \theta} \\ &= \frac{(1 - \sin^2 \theta)(1 - \sin^2 \theta) + \sin^2 \theta}{\sin^2 \theta \cos^2 \theta} \text{ [by (7)]} = \frac{1 - \sin^2 \theta + \sin^4 \theta}{\sin^2 \theta \cos^2 \theta} \\ &= \frac{\cos^2 \theta + \sin^4 \theta}{\sin^2 \theta \cos^2 \theta} \text{ [by (7)]} = \frac{1}{\sin^2 \theta} + \frac{\sin^2 \theta}{\cos^2 \theta} = \csc^2 \theta + \tan^2 \theta \text{ [by (6)]} \end{aligned}$$

$$50. 2 \csc 2t = \frac{2}{\sin 2t} = \frac{2}{2 \sin t \cos t} \text{ [by (15a)]} = \frac{1}{\sin t \cos t} = \sec t \csc t$$

$$51. \text{ Using (14a), we have } \tan 2\theta = \tan(\theta + \theta) = \frac{\tan \theta + \tan \theta}{1 - \tan \theta \tan \theta} = \frac{2 \tan \theta}{1 - \tan^2 \theta}.$$

$$52. \frac{1}{1 - \sin \theta} + \frac{1}{1 + \sin \theta} = \frac{1 + \sin \theta + 1 - \sin \theta}{(1 - \sin \theta)(1 + \sin \theta)} = \frac{2}{1 - \sin^2 \theta} = \frac{2}{\cos^2 \theta} \text{ [by (7)]} = 2 \sec^2 \theta$$

53. Using (15a) and (16a),

$$\begin{aligned} \sin x \sin 2x + \cos x \cos 2x &= \sin x (2 \sin x \cos x) + \cos x (2 \cos^2 x - 1) = 2 \sin^2 x \cos x + 2 \cos^3 x - \cos x \\ &= 2(1 - \cos^2 x) \cos x + 2 \cos^3 x - \cos x \text{ [by (7)]} \\ &= 2 \cos x - 2 \cos^3 x + 2 \cos^3 x - \cos x = \cos x \end{aligned}$$

$$\text{Or: } \sin x \sin 2x + \cos x \cos 2x = \cos(2x - x) \text{ [by 13(b)]} = \cos x$$

54. We start with the right side using equations (12a) and (13a):

$$\begin{aligned} \sin(x + y) \sin(x - y) &= (\sin x \cos y + \cos x \sin y)(\sin x \cos y - \cos x \sin y) \\ &= \sin^2 x \cos^2 y - \sin x \cos y \cos x \sin y + \cos x \sin y \sin x \cos y - \cos^2 x \sin^2 y \\ &= \sin^2 x (1 - \sin^2 y) - (1 - \sin^2 x) \sin^2 y \text{ [by (7)]} \\ &= \sin^2 x - \sin^2 x \sin^2 y - \sin^2 y + \sin^2 x \sin^2 y = \sin^2 x - \sin^2 y \end{aligned}$$

$$\begin{aligned} 55. \frac{\sin \phi}{1 - \cos \phi} &= \frac{\sin \phi}{1 - \cos \phi} \cdot \frac{1 + \cos \phi}{1 + \cos \phi} = \frac{\sin \phi (1 + \cos \phi)}{1 - \cos^2 \phi} = \frac{\sin \phi (1 + \cos \phi)}{\sin^2 \phi} \text{ [by (7)]} \\ &= \frac{1 + \cos \phi}{\sin \phi} = \frac{1}{\sin \phi} + \frac{\cos \phi}{\sin \phi} = \csc \phi + \cot \phi \text{ [by (6)]} \end{aligned}$$

$$56. \tan x + \tan y = \frac{\sin x}{\cos x} + \frac{\sin y}{\cos y} = \frac{\sin x \cos y + \cos x \sin y}{\cos x \cos y} = \frac{\sin(x + y)}{\cos x \cos y} \text{ [by (12a)]}$$

57. Using (12a),

$$\begin{aligned} \sin 3\theta + \sin \theta &= \sin(2\theta + \theta) + \sin \theta = \sin 2\theta \cos \theta + \cos 2\theta \sin \theta + \sin \theta \\ &= \sin 2\theta \cos \theta + (2 \cos^2 \theta - 1) \sin \theta + \sin \theta \text{ [by (16a)]} \\ &= \sin 2\theta \cos \theta + 2 \cos^2 \theta \sin \theta - \sin \theta + \sin \theta = \sin 2\theta \cos \theta + \sin 2\theta \cos \theta \text{ [by (15a)]} \\ &= 2 \sin 2\theta \cos \theta \end{aligned}$$


58. We use (12b) with  $x = 2\theta$ ,  $y = \theta$  to get

$$\begin{aligned}\cos 3\theta &= \cos(2\theta + \theta) = \cos 2\theta \cos \theta - \sin 2\theta \sin \theta \\ &= (2\cos^2 \theta - 1)\cos \theta - 2\sin^2 \theta \cos \theta \quad [\text{by (16a) and (15a)}] \\ &= (2\cos^2 \theta - 1)\cos \theta - 2(1 - \cos^2 \theta)\cos \theta \quad [\text{by (7)}] \\ &= 2\cos^3 \theta - \cos \theta - 2\cos \theta + 2\cos^3 \theta = 4\cos^3 \theta - 3\cos \theta\end{aligned}$$

59. Since  $\sin x = \frac{1}{3}$  we can label the opposite side as having length 1, the hypotenuse as having length 3, and use the Pythagorean Theorem to get that the adjacent side has length  $\sqrt{8}$ . Then, from the diagram,

$\cos x = \frac{\sqrt{8}}{3}$ . Similarly we have that  $\sin y = \frac{3}{5}$ . Now use (12a):

$$\sin(x + y) = \sin x \cos y + \cos x \sin y = \frac{1}{3} \cdot \frac{4}{5} + \frac{\sqrt{8}}{3} \cdot \frac{3}{5} = \frac{4}{15} + \frac{3\sqrt{8}}{15} = \frac{4+6\sqrt{2}}{15}.$$



60. Use (12b) and the values for  $\sin y$  and  $\cos x$  obtained in Exercise 59 to get

$$\cos(x + y) = \cos x \cos y - \sin x \sin y = \frac{\sqrt{8}}{3} \cdot \frac{4}{5} - \frac{1}{3} \cdot \frac{3}{5} = \frac{8\sqrt{2}-3}{15}$$

61. Using (13b) and the values for  $\cos x$  and  $\sin y$  obtained in Exercise 59, we have

$$\cos(x - y) = \cos x \cos y + \sin x \sin y = \frac{\sqrt{8}}{3} \cdot \frac{4}{5} + \frac{1}{3} \cdot \frac{3}{5} = \frac{8\sqrt{2}+3}{15}$$

62. Using (13a) and the values for  $\sin y$  and  $\cos x$  obtained in Exercise 59, we get

$$\sin(x - y) = \sin x \cos y - \cos x \sin y = \frac{1}{3} \cdot \frac{4}{5} - \frac{\sqrt{8}}{3} \cdot \frac{3}{5} = \frac{4-6\sqrt{2}}{15}$$

63. Using (15a) and the values for  $\sin y$  and  $\cos y$  obtained in Exercise 59, we have  $\sin 2y = 2 \sin y \cos y = 2 \cdot \frac{3}{5} \cdot \frac{4}{5} = \frac{24}{25}$ .

64. Using (16a) with  $\cos y = \frac{4}{5}$ , we have  $\cos 2y = 2 \cos^2 y - 1 = 2\left(\frac{4}{5}\right)^2 - 1 = \frac{32}{25} - 1 = \frac{7}{25}$ .

65.  $2 \cos x - 1 = 0 \Leftrightarrow \cos x = \frac{1}{2} \Rightarrow x = \frac{\pi}{3}, \frac{5\pi}{3}$  for  $x \in [0, 2\pi]$ .

66.  $3 \cot^2 x = 1 \Leftrightarrow 3 = 1/\cot^2 x \Leftrightarrow \tan^2 x = 3 \Leftrightarrow \tan x = \pm\sqrt{3} \Rightarrow x = \frac{\pi}{3}, \frac{2\pi}{3}, \frac{4\pi}{3}, \text{ and } \frac{5\pi}{3}$ .

67.  $2 \sin^2 x = 1 \Leftrightarrow \sin^2 x = \frac{1}{2} \Leftrightarrow \sin x = \pm\frac{1}{\sqrt{2}} \Rightarrow x = \frac{\pi}{4}, \frac{3\pi}{4}, \frac{5\pi}{4}, \frac{7\pi}{4}$ .

68.  $|\tan x| = 1 \Leftrightarrow \tan x = -1 \text{ or } \tan x = 1 \Leftrightarrow x = \frac{3\pi}{4}, \frac{7\pi}{4} \text{ or } x = \frac{\pi}{4}, \frac{5\pi}{4}$ .

69. Using (15a), we have  $\sin 2x = \cos x \Leftrightarrow 2 \sin x \cos x - \cos x = 0 \Leftrightarrow \cos x(2 \sin x - 1) = 0 \Leftrightarrow \cos x = 0$  or  $2 \sin x - 1 = 0 \Rightarrow x = \frac{\pi}{2}, \frac{3\pi}{2}$  or  $\sin x = \frac{1}{2} \Rightarrow x = \frac{\pi}{6}$  or  $\frac{5\pi}{6}$ . Therefore, the solutions are  $x = \frac{\pi}{6}, \frac{\pi}{2}, \frac{5\pi}{6}, \frac{3\pi}{2}$ .

70. By (15a),  $2 \cos x + \sin 2x = 0 \Leftrightarrow 2 \cos x + 2 \sin x \cos x = 0 \Leftrightarrow 2 \cos x(1 + \sin x) = 0 \Leftrightarrow \cos x = 0$  or  $1 + \sin x = 0 \Leftrightarrow x = \frac{\pi}{2}, \frac{3\pi}{2}$  or  $\sin x = -1 \Rightarrow x = \frac{3\pi}{2}$ . So the solutions are  $x = \frac{\pi}{2}, \frac{3\pi}{2}$ .

71.  $\sin x = \tan x \Leftrightarrow \sin x - \tan x = 0 \Leftrightarrow \sin x - \frac{\sin x}{\cos x} = 0 \Leftrightarrow \sin x \left(1 - \frac{1}{\cos x}\right) = 0 \Leftrightarrow \sin x = 0$  or

$1 - \frac{1}{\cos x} = 0 \Rightarrow x = 0, \pi, 2\pi$  or  $1 = \frac{1}{\cos x} \Rightarrow \cos x = 1 \Rightarrow x = 0, 2\pi$ . Therefore the solutions are  $x = 0, \pi, 2\pi$ .

72. By (16a),  $2 + \cos 2x = 3 \cos x \Leftrightarrow 2 + 2 \cos^2 x - 1 = 3 \cos x \Leftrightarrow 2 \cos^2 x - 3 \cos x + 1 = 0 \Leftrightarrow$

$$(2 \cos x - 1)(\cos x - 1) = 0 \Leftrightarrow \cos x = 1 \text{ or } \cos x = \frac{1}{2} \Rightarrow x = 0, 2\pi \text{ or } x = \frac{\pi}{3}, \frac{5\pi}{3}.$$

73. We know that  $\sin x = \frac{1}{2}$  when  $x = \frac{\pi}{6}$  or  $\frac{5\pi}{6}$ , and from Figure 14(a), we see that  $\sin x \leq \frac{1}{2} \Rightarrow 0 \leq x \leq \frac{\pi}{6}$  or

$$\frac{5\pi}{6} \leq x \leq 2\pi \text{ for } x \in [0, 2\pi].$$


74.  $2 \cos x + 1 > 0 \Rightarrow 2 \cos x > -1 \Rightarrow \cos x > -\frac{1}{2}$ .  $\cos x = -\frac{1}{2}$  when  $x = \frac{2\pi}{3}, \frac{4\pi}{3}$  and from Figure 14(b), we see that

$$\cos x > -\frac{1}{2} \text{ when } 0 \leq x < \frac{2\pi}{3}, \frac{4\pi}{3} < x \leq 2\pi.$$


75.  $\tan x = -1$  when  $x = \frac{3\pi}{4}, \frac{7\pi}{4}$ , and  $\tan x = 1$  when  $x = \frac{\pi}{4}$  or  $\frac{5\pi}{4}$ . From Figure 15(a) we see that  $-1 < \tan x < 1 \Rightarrow$

$$0 \leq x < \frac{\pi}{4}, \frac{3\pi}{4} < x < \frac{5\pi}{4}, \text{ and } \frac{7\pi}{4} < x \leq 2\pi.$$


76. We know that  $\sin x = \cos x$  when  $x = \frac{\pi}{4}, \frac{5\pi}{4}$ , and from the diagram we see that  $\sin x > \cos x$  when  $\frac{\pi}{4} < x < \frac{5\pi}{4}$ .




77.  $y = \cos(x - \frac{\pi}{3})$ . We start with the graph of  $y = \cos x$  and shift it  $\frac{\pi}{3}$  units to the right.




78.  $y = \tan 2x$ . Start with the graph of  $y = \tan x$  with period  $\pi$  and compress it to a period of  $\frac{\pi}{2}$ .




79.  $y = \frac{1}{3} \tan(x - \frac{\pi}{2})$ . We start with the graph of  $y = \tan x$ , shift it  $\frac{\pi}{2}$  units to the right and compress it to  $\frac{1}{3}$  of its original vertical size.




80.  $y = 1 + \sec x$ . Start with the graph of  $y = \sec x$  and raise it by one unit.



81.  $y = |\sin x|$ . We start with the graph of  $y = \sin x$  and reflect the parts below the  $x$ -axis about the  $x$ -axis.



82.  $y = 2 + \sin(x + \frac{\pi}{4})$ . Start with the graph of  $y = \sin x$ , and shift it  $\frac{\pi}{4}$  units to the left and 2 units up.



83. From the figure in the text, we see that  $x = b \cos \theta$ ,  $y = b \sin \theta$ , and from the distance formula we have that the distance  $c$  from  $(x, y)$  to  $(a, 0)$  is  $c = \sqrt{(x - a)^2 + (y - 0)^2} \Rightarrow$

$$\begin{aligned} c^2 &= (b \cos \theta - a)^2 + (b \sin \theta)^2 = b^2 \cos^2 \theta - 2ab \cos \theta + a^2 + b^2 \sin^2 \theta \\ &= a^2 + b^2(\cos^2 \theta + \sin^2 \theta) - 2ab \cos \theta = a^2 + b^2 - 2ab \cos \theta \quad [\text{by (7)}] \end{aligned}$$


84.  $|AB|^2 = |AC|^2 + |BC|^2 - 2|AC||BC| \cos \angle C = (820)^2 + (910)^2 - 2(820)(910) \cos 103^\circ \approx 1,836,217 \Rightarrow$   
 $|AB| \approx 1355 \text{ m}$

85. Using the Law of Cosines, we have  $c^2 = 1^2 + 1^2 - 2(1)(1) \cos(\alpha - \beta) = 2[1 - \cos(\alpha - \beta)]$ . Now, using the distance formula,  $c^2 = |AB|^2 = (\cos \alpha - \cos \beta)^2 + (\sin \alpha - \sin \beta)^2$ . Equating these two expressions for  $c^2$ , we get  
 $2[1 - \cos(\alpha - \beta)] = \cos^2 \alpha + \sin^2 \alpha + \cos^2 \beta + \sin^2 \beta - 2 \cos \alpha \cos \beta - 2 \sin \alpha \sin \beta \Rightarrow$   
 $1 - \cos(\alpha - \beta) = 1 - \cos \alpha \cos \beta - \sin \alpha \sin \beta \Rightarrow \cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta.$

86.  $\cos(x + y) = \cos(x - (-y)) = \cos x \cos(-y) + \sin x \sin(-y)$   
 $= \cos x \cos y - \sin x \sin y \quad [\text{using Equations (10a) and (10b)}]$

87. In Exercise 86 we used the subtraction formula for cosine to prove the addition formula for cosine. Using that formula with  $x = \frac{\pi}{2} - \alpha$ ,  $y = \beta$ , we get  $\cos[(\frac{\pi}{2} - \alpha) + \beta] = \cos(\frac{\pi}{2} - \alpha) \cos \beta - \sin(\frac{\pi}{2} - \alpha) \sin \beta \Rightarrow$   
 $\cos[\frac{\pi}{2} - (\alpha - \beta)] = \cos(\frac{\pi}{2} - \alpha) \cos \beta - \sin(\frac{\pi}{2} - \alpha) \sin \beta$ . Now we use the identities given in the problem,  
 $\cos(\frac{\pi}{2} - \theta) = \sin \theta$  and  $\sin(\frac{\pi}{2} - \theta) = \cos \theta$ , to get  $\sin(\alpha - \beta) = \sin \alpha \cos \beta - \cos \alpha \sin \beta$ .

88. If  $0 < \theta < \frac{\pi}{2}$ , we have the case depicted in the first diagram. In this case, we see that the height of the triangle is  $h = a \sin \theta$ . If  $\frac{\pi}{2} \leq \theta < \pi$ , we have the case depicted in the second diagram. In this case, the height of the triangle is  $h = a \sin(\pi - \theta) = a \sin \theta$  (by the identity proved in Exercise 44). So in either case, the area of the triangle is  
 $\frac{1}{2}bh = \frac{1}{2}ab \sin \theta$ .



89. Using the formula from Exercise 88, the area of the triangle is  $\frac{1}{2}(10)(3) \sin 107^\circ \approx 14.34457 \text{ cm}^2$ .

## E Sigma Notation

1.  $\sum_{i=1}^5 \sqrt{i} = \sqrt{1} + \sqrt{2} + \sqrt{3} + \sqrt{4} + \sqrt{5}$
2.  $\sum_{i=1}^6 \frac{1}{i+1} = \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \frac{1}{5} + \frac{1}{6} + \frac{1}{7}$
3.  $\sum_{i=4}^6 3^i = 3^4 + 3^5 + 3^6$
4.  $\sum_{i=4}^6 i^3 = 4^3 + 5^3 + 6^3$
5.  $\sum_{k=0}^4 \frac{2k-1}{2k+1} = -1 + \frac{1}{3} + \frac{3}{5} + \frac{5}{7} + \frac{7}{9}$
6.  $\sum_{k=5}^8 x^k = x^5 + x^6 + x^7 + x^8$
7.  $\sum_{i=1}^n i^{10} = 1^{10} + 2^{10} + 3^{10} + \cdots + n^{10}$
8.  $\sum_{j=n}^{n+3} j^2 = n^2 + (n+1)^2 + (n+2)^2 + (n+3)^2$
9.  $\sum_{j=0}^{n-1} (-1)^j = 1 - 1 + 1 - 1 + \cdots + (-1)^{n-1}$
10.  $\sum_{i=1}^n f(x_i) \Delta x_i = f(x_1) \Delta x_1 + f(x_2) \Delta x_2 + f(x_3) \Delta x_3 + \cdots + f(x_n) \Delta x_n$
11.  $1 + 2 + 3 + 4 + \cdots + 10 = \sum_{i=1}^{10} i$
12.  $\sqrt{3} + \sqrt{4} + \sqrt{5} + \sqrt{6} + \sqrt{7} = \sum_{i=3}^7 \sqrt{i}$
13.  $\frac{1}{2} + \frac{2}{3} + \frac{3}{4} + \frac{4}{5} + \cdots + \frac{19}{20} = \sum_{i=1}^{19} \frac{i}{i+1}$
14.  $\frac{3}{7} + \frac{4}{8} + \frac{5}{9} + \frac{6}{10} + \cdots + \frac{23}{27} = \sum_{i=3}^{23} \frac{i}{i+4}$
15.  $2 + 4 + 6 + 8 + \cdots + 2n = \sum_{i=1}^n 2i$
16.  $1 + 3 + 5 + 7 + \cdots + (2n-1) = \sum_{i=1}^n (2i-1)$
17.  $1 + 2 + 4 + 8 + 16 + 32 = \sum_{i=0}^5 2^i$
18.  $\frac{1}{1} + \frac{1}{4} + \frac{1}{9} + \frac{1}{16} + \frac{1}{25} + \frac{1}{36} = \sum_{i=1}^6 \frac{1}{i^2}$
19.  $x + x^2 + x^3 + \cdots + x^n = \sum_{i=1}^n x^i$
20.  $1 - x + x^2 - x^3 + \cdots + (-1)^n x^n = \sum_{i=0}^n (-1)^i x^i$
21.  $\sum_{i=4}^8 (3i-2) = [3(4)-2] + [3(5)-2] + [3(6)-2] + [3(7)-2] + [3(8)-2] = 10 + 13 + 16 + 19 + 22 = 80$
22.  $\sum_{i=3}^6 i(i+2) = 3 \cdot 5 + 4 \cdot 6 + 5 \cdot 7 + 6 \cdot 8 = 15 + 24 + 35 + 48 = 122$
23.  $\sum_{j=1}^6 3^{j+1} = 3^2 + 3^3 + 3^4 + 3^5 + 3^6 + 3^7 = 9 + 27 + 81 + 243 + 729 + 2187 = 3276$   
(For a more general method, see Exercise 47.)
24.  $\sum_{k=0}^8 \cos k\pi = \cos 0 + \cos \pi + \cos 2\pi + \cos 3\pi + \cos 4\pi + \cos 5\pi + \cos 6\pi + \cos 7\pi + \cos 8\pi$   
 $= 1 - 1 + 1 - 1 + 1 - 1 + 1 - 1 + 1 = 1$
25.  $\sum_{n=1}^{20} (-1)^n = -1 + 1 - 1 + 1 - 1 + 1 - 1 + 1 - 1 + 1 - 1 + 1 - 1 + 1 - 1 + 1 - 1 + 1 - 1 + 1 = 0$
26.  $\sum_{i=1}^{100} 4 = \underbrace{4 + 4 + 4 + \cdots + 4}_{(100 \text{ summands})} = 100 \cdot 4 = 400$



$$27. \sum_{i=0}^4 (2^i + i^2) = (1 + 0) + (2 + 1) + (4 + 4) + (8 + 9) + (16 + 16) = 61$$

$$28. \sum_{i=-2}^4 2^{3-i} = 2^5 + 2^4 + 2^3 + 2^2 + 2^1 + 2^0 + 2^{-1} = 63.5$$

$$29. \sum_{i=1}^n 2i = 2 \sum_{i=1}^n i = 2 \cdot \frac{n(n+1)}{2} \quad [\text{by Theorem 3(c)}] = n(n+1)$$

$$30. \sum_{i=1}^n (2 - 5i) = \sum_{i=1}^n 2 - \sum_{i=1}^n 5i = 2n - 5 \sum_{i=1}^n i = 2n - \frac{5n(n+1)}{2} = \frac{4n}{2} - \frac{5n^2 + 5n}{2} = -\frac{n(5n+1)}{2}$$

$$\begin{aligned} 31. \sum_{i=1}^n (i^2 + 3i + 4) &= \sum_{i=1}^n i^2 + 3 \sum_{i=1}^n i + \sum_{i=1}^n 4 = \frac{n(n+1)(2n+1)}{6} + \frac{3n(n+1)}{2} + 4n \\ &= \frac{1}{6}[(2n^3 + 3n^2 + n) + (9n^2 + 9n) + 24n] = \frac{1}{6}(2n^3 + 12n^2 + 34n) = \frac{1}{3}n(n^2 + 6n + 17) \end{aligned}$$

$$\begin{aligned} 32. \sum_{i=1}^n (3 + 2i)^2 &= \sum_{i=1}^n (9 + 12i + 4i^2) = \sum_{i=1}^n 9 + 12 \sum_{i=1}^n i + 4 \sum_{i=1}^n i^2 = 9n + 6n(n+1) + \frac{2n(n+1)(2n+1)}{3} \\ &= \frac{27n + 18n^2 + 18n + 4n^3 + 6n^2 + 2n}{3} = \frac{1}{3}(4n^3 + 24n^2 + 47n) = \frac{1}{3}n(4n^2 + 24n + 47) \end{aligned}$$

$$\begin{aligned} 33. \sum_{i=1}^n (i+1)(i+2) &= \sum_{i=1}^n (i^2 + 3i + 2) = \sum_{i=1}^n i^2 + 3 \sum_{i=1}^n i + \sum_{i=1}^n 2 = \frac{n(n+1)(2n+1)}{6} + \frac{3n(n+1)}{2} + 2n \\ &= \frac{n(n+1)}{6} [(2n+1) + 9] + 2n = \frac{n(n+1)}{3} (n+5) + 2n \\ &= \frac{n}{3} [(n+1)(n+5) + 6] = \frac{n}{3} (n^2 + 6n + 11) \end{aligned}$$

$$\begin{aligned} 34. \sum_{i=1}^n i(i+1)(i+2) &= \sum_{i=1}^n (i^3 + 3i^2 + 2i) = \sum_{i=1}^n i^3 + 3 \sum_{i=1}^n i^2 + 2 \sum_{i=1}^n i \\ &= \left[ \frac{n(n+1)}{2} \right]^2 + \frac{3n(n+1)(2n+1)}{6} + \frac{2n(n+1)}{2} \\ &= n(n+1) \left[ \frac{n(n+1)}{4} + \frac{2n+1}{2} + 1 \right] = \frac{n(n+1)}{4} (n^2 + n + 4n + 2 + 4) \\ &= \frac{n(n+1)}{4} (n^2 + 5n + 6) = \frac{n(n+1)(n+2)(n+3)}{4} \end{aligned}$$

$$\begin{aligned} 35. \sum_{i=1}^n (i^3 - i - 2) &= \sum_{i=1}^n i^3 - \sum_{i=1}^n i - \sum_{i=1}^n 2 = \left[ \frac{n(n+1)}{2} \right]^2 - \frac{n(n+1)}{2} - 2n \\ &= \frac{1}{4}n(n+1)[n(n+1) - 2] - 2n = \frac{1}{4}n(n+1)(n+2)(n-1) - 2n \\ &= \frac{1}{4}n[(n+1)(n-1)(n+2) - 8] = \frac{1}{4}n[(n^2 - 1)(n+2) - 8] = \frac{1}{4}n(n^3 + 2n^2 - n - 10) \end{aligned}$$

$$\begin{aligned} 36. \text{ By Theorem 3(c) we have that } \sum_{i=1}^n i &= \frac{n(n+1)}{2} = 78 \Leftrightarrow n(n+1) = 156 \Leftrightarrow n^2 + n - 156 = 0 \Leftrightarrow \\ &(n+13)(n-12) = 0 \Leftrightarrow n = 12 \text{ or } -13. \text{ But } n = -13 \text{ produces a negative answer for the sum, so } n = 12. \end{aligned}$$

$$37. \text{ By Theorem 2(a) and Example 3, } \sum_{i=1}^n c = c \sum_{i=1}^n 1 = cn.$$

38. Let  $S_n$  be the statement that  $\sum_{i=1}^n i^3 = \left[ \frac{n(n+1)}{2} \right]^2$ .

1.  $S_1$  is true because  $1^3 = \left( \frac{1 \cdot 2}{2} \right)^2$ .

2. Assume  $S_k$  is true. Then  $\sum_{i=1}^k i^3 = \left[ \frac{k(k+1)}{2} \right]^2$ , so

$$\sum_{i=1}^{k+1} i^3 = \left[ \frac{k(k+1)}{2} \right]^2 + (k+1)^3 = \frac{(k+1)^2}{4} [k^2 + 4(k+1)] = \frac{(k+1)^2}{4} (k+2)^2 = \left( \frac{(k+1)[(k+1)+1]}{2} \right)^2$$

showing that  $S_{k+1}$  is true.

Therefore,  $S_n$  is true for all  $n$  by mathematical induction.

39. 
$$\sum_{i=1}^n [(i+1)^4 - i^4] = (2^4 - 1^4) + (3^4 - 2^4) + (4^4 - 3^4) + \cdots + [(n+1)^4 - n^4]$$
$$= (n+1)^4 - 1^4 = n^4 + 4n^3 + 6n^2 + 4n$$

On the other hand,

$$\begin{aligned} \sum_{i=1}^n [(i+1)^4 - i^4] &= \sum_{i=1}^n (4i^3 + 6i^2 + 4i + 1) = 4 \sum_{i=1}^n i^3 + 6 \sum_{i=1}^n i^2 + 4 \sum_{i=1}^n i + \sum_{i=1}^n 1 \\ &= 4S + n(n+1)(2n+1) + 2n(n+1) + n \quad \left[ \text{where } S = \sum_{i=1}^n i^3 \right] \\ &= 4S + 2n^3 + 3n^2 + n + 2n^2 + 2n + n = 4S + 2n^3 + 5n^2 + 4n \end{aligned}$$

Thus,  $n^4 + 4n^3 + 6n^2 + 4n = 4S + 2n^3 + 5n^2 + 4n$ , from which it follows that

$$4S = n^4 + 2n^3 + n^2 = n^2(n^2 + 2n + 1) = n^2(n+1)^2 \text{ and } S = \left[ \frac{n(n+1)}{2} \right]^2.$$

40. The area of  $G_i$  is

$$\begin{aligned} \left( \sum_{k=1}^i k \right)^2 - \left( \sum_{k=1}^{i-1} k \right)^2 &= \left[ \frac{i(i+1)}{2} \right]^2 - \left[ \frac{(i-1)i}{2} \right]^2 = \frac{i^2}{4} [(i+1)^2 - (i-1)^2] \\ &= \frac{i^2}{4} [(i^2 + 2i + 1) - (i^2 - 2i + 1)] = \frac{i^2}{4} (4i) = i^3 \end{aligned}$$

Thus, the area of  $ABCD$  is  $\sum_{i=1}^n i^3 = \left[ \frac{n(n+1)}{2} \right]^2$ .

41. (a)  $\sum_{i=1}^n [i^4 - (i-1)^4] = (1^4 - 0^4) + (2^4 - 1^4) + (3^4 - 2^4) + \cdots + [n^4 - (n-1)^4] = n^4 - 0 = n^4$

(b)  $\sum_{i=1}^{100} (5^i - 5^{i-1}) = (5^1 - 5^0) + (5^2 - 5^1) + (5^3 - 5^2) + \cdots + (5^{100} - 5^{99}) = 5^{100} - 5^0 = 5^{100} - 1$

(c)  $\sum_{i=3}^{99} \left( \frac{1}{i} - \frac{1}{i+1} \right) = \left( \frac{1}{3} - \frac{1}{4} \right) + \left( \frac{1}{4} - \frac{1}{5} \right) + \left( \frac{1}{5} - \frac{1}{6} \right) + \cdots + \left( \frac{1}{99} - \frac{1}{100} \right) = \frac{1}{3} - \frac{1}{100} = \frac{97}{300}$

(d)  $\sum_{i=1}^n (a_i - a_{i-1}) = (a_1 - a_0) + (a_2 - a_1) + (a_3 - a_2) + \cdots + (a_n - a_{n-1}) = a_n - a_0$

42. Summing the inequalities  $-|a_i| \leq a_i \leq |a_i|$  for  $i = 1, 2, \dots, n$ , we get  $-\sum_{i=1}^n |a_i| \leq \sum_{i=1}^n a_i \leq \sum_{i=1}^n |a_i|$ . Since  $|x| \leq c \Leftrightarrow$

$-c \leq x \leq c$ , we have  $\left| \sum_{i=1}^n a_i \right| \leq \sum_{i=1}^n |a_i|$ . Another method: Use mathematical induction.

$$43. \lim_{n \rightarrow \infty} \sum_{i=1}^n \frac{1}{n} \left( \frac{i}{n} \right)^2 = \lim_{n \rightarrow \infty} \frac{1}{n^3} \sum_{i=1}^n i^2 = \lim_{n \rightarrow \infty} \frac{1}{n^3} \frac{n(n+1)(2n+1)}{6} = \lim_{n \rightarrow \infty} \frac{1}{6} \left( 1 + \frac{1}{n} \right) \left( 2 + \frac{1}{n} \right) = \frac{1}{6}(1)(2) = \frac{1}{3}$$

$$44. \lim_{n \rightarrow \infty} \sum_{i=1}^n \frac{1}{n} \left[ \left( \frac{i}{n} \right)^3 + 1 \right] = \lim_{n \rightarrow \infty} \sum_{i=1}^n \left[ \frac{i^3}{n^4} + \frac{1}{n} \right] = \lim_{n \rightarrow \infty} \left[ \frac{1}{n^4} \sum_{i=1}^n i^3 + \frac{1}{n} \sum_{i=1}^n 1 \right] = \lim_{n \rightarrow \infty} \left[ \frac{1}{n^4} \left( \frac{n(n+1)}{2} \right)^2 + \frac{1}{n}(n) \right]$$

$$= \lim_{n \rightarrow \infty} \frac{1}{4} \left( 1 + \frac{1}{n} \right)^2 + 1 = \frac{1}{4} + 1 = \frac{5}{4}$$

$$45. \lim_{n \rightarrow \infty} \sum_{i=1}^n \frac{2}{n} \left[ \left( \frac{2i}{n} \right)^3 + 5 \left( \frac{2i}{n} \right) \right] = \lim_{n \rightarrow \infty} \sum_{i=1}^n \left[ \frac{16}{n^4} i^3 + \frac{20}{n^2} i \right] = \lim_{n \rightarrow \infty} \left[ \frac{16}{n^4} \sum_{i=1}^n i^3 + \frac{20}{n^2} \sum_{i=1}^n i \right]$$

$$= \lim_{n \rightarrow \infty} \left[ \frac{16}{n^4} \frac{n^2(n+1)^2}{4} + \frac{20}{n^2} \frac{n(n+1)}{2} \right] = \lim_{n \rightarrow \infty} \left[ \frac{4(n+1)^2}{n^2} + \frac{10n(n+1)}{n^2} \right]$$

$$= \lim_{n \rightarrow \infty} \left[ 4 \left( 1 + \frac{1}{n} \right)^2 + 10 \left( 1 + \frac{1}{n} \right) \right] = 4 \cdot 1 + 10 \cdot 1 = 14$$

$$46. \lim_{n \rightarrow \infty} \sum_{i=1}^n \frac{3}{n} \left[ \left( 1 + \frac{3i}{n} \right)^3 - 2 \left( 1 + \frac{3i}{n} \right) \right] = \lim_{n \rightarrow \infty} \sum_{i=1}^n \frac{3}{n} \left[ 1 + \frac{9i}{n} + \frac{27i^2}{n^2} + \frac{27i^3}{n^3} - 2 - \frac{6i}{n} \right]$$

$$= \lim_{n \rightarrow \infty} \sum_{i=1}^n \left[ \frac{81}{n^4} i^3 + \frac{81}{n^3} i^2 + \frac{9}{n^2} i - \frac{3}{n} \right]$$

$$= \lim_{n \rightarrow \infty} \left[ \frac{81}{n^4} \frac{n^2(n+1)^2}{4} + \frac{81}{n^3} \frac{n(n+1)(2n+1)}{6} + \frac{9}{n^2} \frac{n(n+1)}{2} - \frac{3}{n} n \right]$$

$$= \lim_{n \rightarrow \infty} \left[ \frac{81}{4} \left( 1 + \frac{1}{n} \right)^2 + \frac{27}{2} \left( 1 + \frac{1}{n} \right) \left( 2 + \frac{1}{n} \right) + \frac{9}{2} \left( 1 + \frac{1}{n} \right) - 3 \right]$$

$$= \frac{81}{4} + \frac{54}{2} + \frac{9}{2} - 3 = \frac{195}{4}$$

$$47. \text{ Let } S = \sum_{i=1}^n ar^{i-1} = a + ar + ar^2 + \cdots + ar^{n-1}. \text{ Multiplying both sides by } r \text{ gives us}$$

$$rS = ar + ar^2 + \cdots + ar^{n-1} + ar^n. \text{ Subtracting the first equation from the second, we find}$$

$$(r-1)S = ar^n - a = a(r^n - 1), \text{ so } S = \frac{a(r^n - 1)}{r - 1} \quad [\text{since } r \neq 1].$$

$$48. \sum_{i=1}^n \frac{3}{2^{i-1}} = 3 \sum_{i=1}^n \left( \frac{1}{2} \right)^{i-1} = \frac{3 \left[ \left( \frac{1}{2} \right)^n - 1 \right]}{\frac{1}{2} - 1} \quad [\text{using Exercise 47 with } a = 3 \text{ and } r = \frac{1}{2}] = 6 \left[ 1 - \left( \frac{1}{2} \right)^n \right]$$

$$49. \sum_{i=1}^n (2i + 2^i) = 2 \sum_{i=1}^n i + \sum_{i=1}^n 2 \cdot 2^{i-1} = 2 \frac{n(n+1)}{2} + \frac{2(2^n - 1)}{2 - 1} = 2^{n+1} + n^2 + n - 2.$$

For the first sum we have used Theorems 2(a) and 3(c), and for the second, Exercise 47 with  $a = r = 2$ .

$$50. \sum_{i=1}^m \left[ \sum_{j=1}^n (i+j) \right] = \sum_{i=1}^m \left[ \sum_{j=1}^n i + \sum_{j=1}^n j \right] \quad [\text{Theorem 2(b)}] = \sum_{i=1}^m \left[ ni + \frac{n(n+1)}{2} \right] \quad [\text{Theorem 3(b) and 3(c)}]$$

$$= \sum_{i=1}^m ni + \sum_{i=1}^m \frac{n(n+1)}{2} = \frac{nm(m+1)}{2} + \frac{nm(n+1)}{2} = \frac{nm}{2} (m+n+2)$$

**G Complex Numbers**

1.  $(5 - 6i) + (3 + 2i) = (5 + 3) + (-6 + 2)i = 8 + (-4)i = 8 - 4i$
2.  $(4 - \frac{1}{2}i) - (9 + \frac{5}{2}i) = (4 - 9) + (-\frac{1}{2} - \frac{5}{2})i = -5 + (-3)i = -5 - 3i$
3.  $(2 + 5i)(4 - i) = 2(4) + 2(-i) + (5i)(4) + (5i)(-i) = 8 - 2i + 20i - 5i^2 = 8 + 18i - 5(-1)$   
 $= 8 + 18i + 5 = 13 + 18i$
4.  $(1 - 2i)(8 - 3i) = 8 - 3i - 16i + 6(-1) = 2 - 19i$
5.  $\overline{12 + 7i} = 12 - 7i$
6.  $2i(\frac{1}{2} - i) = i - 2(-1) = 2 + i \Rightarrow \overline{2i(\frac{1}{2} - i)} = \overline{2 + i} = 2 - i$
7.  $\frac{1 + 4i}{3 + 2i} = \frac{1 + 4i}{3 + 2i} \cdot \frac{3 - 2i}{3 - 2i} = \frac{3 - 2i + 12i - 8(-1)}{3^2 + 2^2} = \frac{11 + 10i}{13} = \frac{11}{13} + \frac{10}{13}i$
8.  $\frac{3 + 2i}{1 - 4i} = \frac{3 + 2i}{1 - 4i} \cdot \frac{1 + 4i}{1 + 4i} = \frac{3 + 12i + 2i + 8(-1)}{1^2 + 4^2} = \frac{-5 + 14i}{17} = -\frac{5}{17} + \frac{14}{17}i$
9.  $\frac{1}{1 + i} = \frac{1}{1 + i} \cdot \frac{1 - i}{1 - i} = \frac{1 - i}{1 - (-1)} = \frac{1 - i}{2} = \frac{1}{2} - \frac{1}{2}i$
10.  $\frac{3}{4 - 3i} = \frac{3}{4 - 3i} \cdot \frac{4 + 3i}{4 + 3i} = \frac{12 + 9i}{16 - 9(-1)} = \frac{12}{25} + \frac{9}{25}i$
11.  $i^3 = i^2 \cdot i = (-1)i = -i$
12.  $i^{100} = (i^2)^{50} = (-1)^{50} = 1$
13.  $\sqrt{-25} = \sqrt{25}i = 5i$
14.  $\sqrt{-3}\sqrt{-12} = \sqrt{3}i\sqrt{12}i = \sqrt{3 \cdot 12}i^2 = \sqrt{36}(-1) = -6$
15.  $\overline{12 - 5i} = 12 + 5i$  and  $|12 - 5i| = \sqrt{12^2 + (-5)^2} = \sqrt{144 + 25} = \sqrt{169} = 13$
16.  $\overline{-1 + 2\sqrt{2}i} = -1 - 2\sqrt{2}i$  and  $|-1 + 2\sqrt{2}i| = \sqrt{(-1)^2 + (2\sqrt{2})^2} = \sqrt{1 + 8} = \sqrt{9} = 3$
17.  $\overline{-4i} = 0 - 4i = 0 + 4i = 4i$  and  $|-4i| = \sqrt{0^2 + (-4)^2} = \sqrt{16} = 4$
18. Let  $z = a + bi$  and  $w = c + di$ .
  - (a)  $\overline{z + w} = \overline{(a + bi) + (c + di)} = \overline{(a + c) + (b + d)i} = (a + c) - (b + d)i = (a - bi) + (c - di) = \overline{z} + \overline{w}$
  - (b)  $\overline{zw} = \overline{(a + bi)(c + di)} = \overline{(ac - bd) + (ad + bc)i} = (ac - bd) - (ad + bc)i$ .  
 On the other hand,  $\overline{z} \overline{w} = (a - bi)(c - di) = (ac - bd) - (ad + bc)i = \overline{zw}$ .

(c) Use mathematical induction and part (b): Let  $S_n$  be the statement that  $\overline{z^n} = \overline{z}^n$ .  $S_1$  is true because  $\overline{z^1} = \overline{z} = \overline{z}^1$ .

Assume  $S_k$  is true, that is  $\overline{z^k} = \overline{z}^k$ . Then  $\overline{z^{k+1}} = \overline{z^{1+k}} = \overline{z z^k} = \overline{z} \overline{z^k}$  [part (b) with  $w = z^k$ ]  $= \overline{z} \overline{z}^k = \overline{z}^{1+k} = \overline{z}^{k+1}$ , which shows that  $S_{k+1}$  is true. Therefore, by mathematical induction,  $\overline{z^n} = \overline{z}^n$  for every positive integer  $n$ .

*Another proof:* Use part (b) with  $w = z$ , and mathematical induction.

$$19. 4x^2 + 9 = 0 \Leftrightarrow 4x^2 = -9 \Leftrightarrow x^2 = -\frac{9}{4} \Leftrightarrow x = \pm\sqrt{-\frac{9}{4}} = \pm\sqrt{\frac{9}{4}}i = \pm\frac{3}{2}i.$$

$$20. x^4 = 1 \Leftrightarrow x^4 - 1 = 0 \Leftrightarrow (x^2 - 1)(x^2 + 1) = 0 \Leftrightarrow x^2 - 1 = 0 \text{ or } x^2 + 1 = 0 \Leftrightarrow x = \pm 1 \text{ or } x = \pm i.$$

$$21. \text{ By the quadratic formula, } x^2 + 2x + 5 = 0 \Leftrightarrow x = \frac{-2 \pm \sqrt{2^2 - 4(1)(5)}}{2(1)} = \frac{-2 \pm \sqrt{-16}}{2} = \frac{-2 \pm 4i}{2} = -1 \pm 2i.$$

$$22. 2x^2 - 2x + 1 = 0 \Leftrightarrow x = \frac{-(-2) \pm \sqrt{(-2)^2 - 4(2)(1)}}{2(2)} = \frac{2 \pm \sqrt{-4}}{4} = \frac{2 \pm 2i}{4} = \frac{1}{2} \pm \frac{1}{2}i$$

$$23. \text{ By the quadratic formula, } z^2 + z + 2 = 0 \Leftrightarrow z = \frac{-1 \pm \sqrt{1^2 - 4(1)(2)}}{2(1)} = \frac{-1 \pm \sqrt{-7}}{2} = -\frac{1}{2} \pm \frac{\sqrt{7}}{2}i.$$

$$24. z^2 + \frac{1}{2}z + \frac{1}{4} = 0 \Leftrightarrow 4z^2 + 2z + 1 = 0 \Leftrightarrow$$

$$z = \frac{-2 \pm \sqrt{2^2 - 4(4)(1)}}{2(4)} = \frac{-2 \pm \sqrt{-12}}{8} = \frac{-2 \pm 2\sqrt{3}i}{8} = -\frac{1}{4} \pm \frac{\sqrt{3}}{4}i$$

$$25. \text{ For } z = -3 + 3i, r = \sqrt{(-3)^2 + 3^2} = 3\sqrt{2} \text{ and } \tan \theta = \frac{3}{-3} = -1 \Rightarrow \theta = \frac{3\pi}{4} \text{ (since } z \text{ lies in the second quadrant).}$$

$$\text{Therefore, } -3 + 3i = 3\sqrt{2}(\cos \frac{3\pi}{4} + i \sin \frac{3\pi}{4}).$$

$$26. \text{ For } z = 1 - \sqrt{3}i, r = \sqrt{1^2 + (-\sqrt{3})^2} = 2 \text{ and } \tan \theta = \frac{-\sqrt{3}}{1} = -\sqrt{3} \Rightarrow \theta = \frac{5\pi}{3} \text{ (since } z \text{ lies in the fourth quadrant).}$$

$$\text{Therefore, } 1 - \sqrt{3}i = 2(\cos \frac{5\pi}{3} + i \sin \frac{5\pi}{3}).$$

$$27. \text{ For } z = 3 + 4i, r = \sqrt{3^2 + 4^2} = 5 \text{ and } \tan \theta = \frac{4}{3} \Rightarrow \theta = \tan^{-1}(\frac{4}{3}) \text{ (since } z \text{ lies in the first quadrant). Therefore,}$$

$$3 + 4i = 5[\cos(\tan^{-1} \frac{4}{3}) + i \sin(\tan^{-1} \frac{4}{3})].$$

$$28. \text{ For } z = 8i, r = \sqrt{0^2 + 8^2} = 8 \text{ and } \tan \theta = \frac{8}{0} \text{ is undefined, so } \theta = \frac{\pi}{2} \text{ (since } z \text{ lies on the positive imaginary axis). Therefore,}$$

$$8i = 8(\cos \frac{\pi}{2} + i \sin \frac{\pi}{2}).$$

$$29. \text{ For } z = \sqrt{3} + i, r = \sqrt{(\sqrt{3})^2 + 1^2} = 2 \text{ and } \tan \theta = \frac{1}{\sqrt{3}} \Rightarrow \theta = \frac{\pi}{6} \Rightarrow z = 2(\cos \frac{\pi}{6} + i \sin \frac{\pi}{6}).$$

$$\text{For } w = 1 + \sqrt{3}i, r = 2 \text{ and } \tan \theta = \sqrt{3} \Rightarrow \theta = \frac{\pi}{3} \Rightarrow w = 2(\cos \frac{\pi}{3} + i \sin \frac{\pi}{3}).$$

$$\text{Therefore, } zw = 2 \cdot 2[\cos(\frac{\pi}{6} + \frac{\pi}{3}) + i \sin(\frac{\pi}{6} + \frac{\pi}{3})] = 4(\cos \frac{\pi}{2} + i \sin \frac{\pi}{2}),$$

$$z/w = \frac{2}{2}[\cos(\frac{\pi}{6} - \frac{\pi}{3}) + i \sin(\frac{\pi}{6} - \frac{\pi}{3})] = \cos(-\frac{\pi}{6}) + i \sin(-\frac{\pi}{6}), \text{ and } 1 = 1 + 0i = 1(\cos 0 + i \sin 0) \Rightarrow$$

$1/z = \frac{1}{2} [\cos(0 - \frac{\pi}{6}) + i \sin(0 - \frac{\pi}{6})] = \frac{1}{2} [\cos(-\frac{\pi}{6}) + i \sin(-\frac{\pi}{6})]$ . For  $1/z$ , we could also use the formula that precedes Example 5 to obtain  $1/z = \frac{1}{2} (\cos \frac{\pi}{6} - i \sin \frac{\pi}{6})$ .

30. For  $z = 4\sqrt{3} - 4i$ ,  $r = \sqrt{(4\sqrt{3})^2 + (-4)^2} = \sqrt{64} = 8$  and  $\tan \theta = \frac{-4}{4\sqrt{3}} = -\frac{1}{\sqrt{3}} \Rightarrow \theta = \frac{11\pi}{6} \Rightarrow$   
 $z = 8(\cos \frac{11\pi}{6} + i \sin \frac{11\pi}{6})$ . For  $w = 8i$ ,  $r = \sqrt{0^2 + 8^2} = 8$  and  $\tan \theta = \frac{8}{0}$  is undefined, so  $\theta = \frac{\pi}{2} \Rightarrow$   
 $w = 8(\cos \frac{\pi}{2} + i \sin \frac{\pi}{2})$ . Therefore,  $zw = 8 \cdot 8 [\cos(\frac{11\pi}{6} + \frac{\pi}{2}) + i \sin(\frac{11\pi}{6} + \frac{\pi}{2})] = 64(\cos \frac{\pi}{3} + i \sin \frac{\pi}{3})$ ,  
 $z/w = \frac{8}{8} [\cos(\frac{11\pi}{6} - \frac{\pi}{2}) + i \sin(\frac{11\pi}{6} - \frac{\pi}{2})] = \cos \frac{4\pi}{3} + i \sin \frac{4\pi}{3}$ , and  
 $1 = 1 + 0i = 1(\cos 0 + i \sin 0) \Rightarrow 1/z = \frac{1}{8} [\cos(0 - \frac{11\pi}{6}) + i \sin(0 - \frac{11\pi}{6})] = \frac{1}{8} [\cos(\frac{\pi}{6}) + i \sin(\frac{\pi}{6})]$ .  
 For  $1/z$ , we could also use the formula that precedes Example 5 to obtain  $1/z = \frac{1}{8} (\cos \frac{11\pi}{6} - i \sin \frac{11\pi}{6})$ .

31. For  $z = 2\sqrt{3} - 2i$ ,  $r = \sqrt{(2\sqrt{3})^2 + (-2)^2} = 4$  and  $\tan \theta = \frac{-2}{2\sqrt{3}} = -\frac{1}{\sqrt{3}} \Rightarrow \theta = -\frac{\pi}{6} \Rightarrow$   
 $z = 4[\cos(-\frac{\pi}{6}) + i \sin(-\frac{\pi}{6})]$ . For  $w = -1 + i$ ,  $r = \sqrt{2}$ ,  $\tan \theta = \frac{1}{-1} = -1 \Rightarrow \theta = \frac{3\pi}{4} \Rightarrow$   
 $w = \sqrt{2}(\cos \frac{3\pi}{4} + i \sin \frac{3\pi}{4})$ . Therefore,  $zw = 4\sqrt{2} [\cos(-\frac{\pi}{6} + \frac{3\pi}{4}) + i \sin(-\frac{\pi}{6} + \frac{3\pi}{4})] = 4\sqrt{2} (\cos \frac{7\pi}{12} + i \sin \frac{7\pi}{12})$ ,  
 $z/w = \frac{4}{\sqrt{2}} [\cos(-\frac{\pi}{6} - \frac{3\pi}{4}) + i \sin(-\frac{\pi}{6} - \frac{3\pi}{4})] = \frac{4}{\sqrt{2}} [\cos(-\frac{11\pi}{12}) + i \sin(-\frac{11\pi}{12})] = 2\sqrt{2} (\cos \frac{13\pi}{12} + i \sin \frac{13\pi}{12})$ , and  
 $1/z = \frac{1}{4} [\cos(-\frac{\pi}{6}) - i \sin(-\frac{\pi}{6})] = \frac{1}{4} (\cos \frac{\pi}{6} + i \sin \frac{\pi}{6})$ .

32. For  $z = 4(\sqrt{3} + i) = 4\sqrt{3} + 4i$ ,  $r = \sqrt{(4\sqrt{3})^2 + 4^2} = \sqrt{64} = 8$  and  $\tan \theta = \frac{4}{4\sqrt{3}} = \frac{1}{\sqrt{3}} \Rightarrow \theta = \frac{\pi}{6} \Rightarrow$   
 $z = 8(\cos \frac{\pi}{6} + i \sin \frac{\pi}{6})$ . For  $w = -3 - 3i$ ,  $r = \sqrt{(-3)^2 + (-3)^2} = \sqrt{18} = 3\sqrt{2}$  and  $\tan \theta = \frac{-3}{-3} = 1 \Rightarrow \theta = \frac{5\pi}{4} \Rightarrow$   
 $w = 3\sqrt{2} (\cos \frac{5\pi}{4} + i \sin \frac{5\pi}{4})$ . Therefore,  $zw = 8 \cdot 3\sqrt{2} [\cos(\frac{\pi}{6} + \frac{5\pi}{4}) + i \sin(\frac{\pi}{6} + \frac{5\pi}{4})] = 24\sqrt{2} (\cos \frac{17\pi}{12} + i \sin \frac{17\pi}{12})$ ,  
 $z/w = \frac{8}{3\sqrt{2}} [\cos(\frac{\pi}{6} - \frac{5\pi}{4}) + i \sin(\frac{\pi}{6} - \frac{5\pi}{4})] = \frac{4\sqrt{2}}{3} [\cos(-\frac{13\pi}{12}) + i \sin(-\frac{13\pi}{12})]$ , and  $1/z = \frac{1}{8} (\cos \frac{\pi}{6} - i \sin \frac{\pi}{6})$ .

33. For  $z = 1 + i$ ,  $r = \sqrt{2}$  and  $\tan \theta = \frac{1}{1} = 1 \Rightarrow \theta = \frac{\pi}{4} \Rightarrow z = \sqrt{2} (\cos \frac{\pi}{4} + i \sin \frac{\pi}{4})$ . So by De Moivre's Theorem,

$$\begin{aligned} (1 + i)^{20} &= [\sqrt{2} (\cos \frac{\pi}{4} + i \sin \frac{\pi}{4})]^{20} = (2^{1/2})^{20} (\cos \frac{20 \cdot \pi}{4} + i \sin \frac{20 \cdot \pi}{4}) = 2^{10} (\cos 5\pi + i \sin 5\pi) \\ &= 2^{10} [-1 + i(0)] = -2^{10} = -1024 \end{aligned}$$

34. For  $z = 1 - \sqrt{3}i$ ,  $r = \sqrt{1^2 + (-\sqrt{3})^2} = 2$  and  $\tan \theta = \frac{-\sqrt{3}}{1} = -\sqrt{3} \Rightarrow \theta = \frac{5\pi}{3} \Rightarrow z = 2 (\cos \frac{5\pi}{3} + i \sin \frac{5\pi}{3})$ .

So by De Moivre's Theorem,

$$\begin{aligned} (1 - \sqrt{3}i)^5 &= [2(\cos \frac{5\pi}{3} + i \sin \frac{5\pi}{3})]^5 = 2^5 (\cos \frac{5 \cdot 5\pi}{3} + i \sin \frac{5 \cdot 5\pi}{3}) = 2^5 (\cos \frac{\pi}{3} + i \sin \frac{\pi}{3}) \\ &= 32 (\frac{1}{2} + \frac{\sqrt{3}}{2}i) = 16 + 16\sqrt{3}i \end{aligned}$$

35. For  $z = 2\sqrt{3} + 2i$ ,  $r = \sqrt{(2\sqrt{3})^2 + 2^2} = \sqrt{16} = 4$  and  $\tan \theta = \frac{2}{2\sqrt{3}} = \frac{1}{\sqrt{3}} \Rightarrow \theta = \frac{\pi}{6} \Rightarrow z = 4(\cos \frac{\pi}{6} + i \sin \frac{\pi}{6})$ .

So by De Moivre's Theorem,

$$(2\sqrt{3} + 2i)^5 = [4(\cos \frac{\pi}{6} + i \sin \frac{\pi}{6})]^5 = 4^5 (\cos \frac{5\pi}{6} + i \sin \frac{5\pi}{6}) = 1024 [-\frac{\sqrt{3}}{2} + \frac{1}{2}i] = -512\sqrt{3} + 512i.$$

36. For  $z = 1 - i$ ,  $r = \sqrt{2}$  and  $\tan \theta = \frac{-1}{1} = -1 \Rightarrow \theta = \frac{7\pi}{4} \Rightarrow z = \sqrt{2} \left( \cos \frac{7\pi}{4} + i \sin \frac{7\pi}{4} \right) \Rightarrow$

$$(1 - i)^8 = \left[ \sqrt{2} \left( \cos \frac{7\pi}{4} + i \sin \frac{7\pi}{4} \right) \right]^8 = 2^4 \left( \cos \frac{8 \cdot 7\pi}{4} + i \sin \frac{8 \cdot 7\pi}{4} \right) = 16(\cos 14\pi + i \sin 14\pi) = 16(1 + 0i) = 16.$$

37.  $1 = 1 + 0i = 1(\cos 0 + i \sin 0)$ . Using Equation 3 with  $r = 1$ ,  $n = 8$ , and  $\theta = 0$ , we have


$$w_k = 1^{1/8} \left[ \cos \left( \frac{0 + 2k\pi}{8} \right) + i \sin \left( \frac{0 + 2k\pi}{8} \right) \right] = \cos \frac{k\pi}{4} + i \sin \frac{k\pi}{4}, \text{ where } k = 0, 1, 2, \dots, 7.$$

$$w_0 = 1(\cos 0 + i \sin 0) = 1, w_1 = 1 \left( \cos \frac{\pi}{4} + i \sin \frac{\pi}{4} \right) = \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}}i,$$

$$w_2 = 1 \left( \cos \frac{\pi}{2} + i \sin \frac{\pi}{2} \right) = i, w_3 = 1 \left( \cos \frac{3\pi}{4} + i \sin \frac{3\pi}{4} \right) = -\frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}}i,$$

$$w_4 = 1(\cos \pi + i \sin \pi) = -1, w_5 = 1 \left( \cos \frac{5\pi}{4} + i \sin \frac{5\pi}{4} \right) = -\frac{1}{\sqrt{2}} - \frac{1}{\sqrt{2}}i,$$

$$w_6 = 1 \left( \cos \frac{3\pi}{2} + i \sin \frac{3\pi}{2} \right) = -i, w_7 = 1 \left( \cos \frac{7\pi}{4} + i \sin \frac{7\pi}{4} \right) = \frac{1}{\sqrt{2}} - \frac{1}{\sqrt{2}}i$$



38.  $32 = 32 + 0i = 32(\cos 0 + i \sin 0)$ . Using Equation 3 with  $r = 32$ ,  $n = 5$ , and  $\theta = 0$ , we have

$$w_k = 32^{1/5} \left[ \cos \left( \frac{0 + 2k\pi}{5} \right) + i \sin \left( \frac{0 + 2k\pi}{5} \right) \right] = 2 \left( \cos \frac{2\pi k}{5} + i \sin \frac{2\pi k}{5} \right), \text{ where } k = 0, 1, 2, 3, 4.$$


$$w_0 = 2(\cos 0 + i \sin 0) = 2$$

$$w_1 = 2 \left( \cos \frac{2\pi}{5} + i \sin \frac{2\pi}{5} \right)$$

$$w_2 = 2 \left( \cos \frac{4\pi}{5} + i \sin \frac{4\pi}{5} \right)$$

$$w_3 = 2 \left( \cos \frac{6\pi}{5} + i \sin \frac{6\pi}{5} \right)$$

$$w_4 = 2 \left( \cos \frac{8\pi}{5} + i \sin \frac{8\pi}{5} \right)$$




39.  $i = 0 + i = 1 \left( \cos \frac{\pi}{2} + i \sin \frac{\pi}{2} \right)$ . Using Equation 3 with  $r = 1$ ,  $n = 3$ , and  $\theta = \frac{\pi}{2}$ , we have

$$w_k = 1^{1/3} \left[ \cos \left( \frac{\frac{\pi}{2} + 2k\pi}{3} \right) + i \sin \left( \frac{\frac{\pi}{2} + 2k\pi}{3} \right) \right], \text{ where } k = 0, 1, 2.$$

$$w_0 = \left( \cos \frac{\pi}{6} + i \sin \frac{\pi}{6} \right) = \frac{\sqrt{3}}{2} + \frac{1}{2}i$$

$$w_1 = \left( \cos \frac{5\pi}{6} + i \sin \frac{5\pi}{6} \right) = -\frac{\sqrt{3}}{2} + \frac{1}{2}i$$

$$w_2 = \left( \cos \frac{9\pi}{6} + i \sin \frac{9\pi}{6} \right) = -i$$




40.  $1 + i = \sqrt{2} \left( \cos \frac{\pi}{4} + i \sin \frac{\pi}{4} \right)$ . Using Equation 3 with  $r = \sqrt{2}$ ,  $n = 3$ , and  $\theta = \frac{\pi}{4}$ , we have

$$w_k = (\sqrt{2})^{1/3} \left[ \cos \left( \frac{\frac{\pi}{4} + 2k\pi}{3} \right) + i \sin \left( \frac{\frac{\pi}{4} + 2k\pi}{3} \right) \right], \text{ where } k = 0, 1, 2.$$

$$w_0 = 2^{1/6} \left( \cos \frac{\pi}{12} + i \sin \frac{\pi}{12} \right)$$

$$w_1 = 2^{1/6} \left( \cos \frac{3\pi}{4} + i \sin \frac{3\pi}{4} \right) = 2^{1/6} \left( -\frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}}i \right) = -2^{-1/3} + 2^{-1/3}i$$

$$w_2 = 2^{1/6} \left( \cos \frac{17\pi}{12} + i \sin \frac{17\pi}{12} \right)$$



41. Using Euler's formula (6) with  $y = \frac{\pi}{2}$ , we have  $e^{i\pi/2} = \cos \frac{\pi}{2} + i \sin \frac{\pi}{2} = 0 + 1i = i$ .

42. Using Euler's formula (6) with  $y = 2\pi$ , we have  $e^{2\pi i} = \cos 2\pi + i \sin 2\pi = 1$ .

43. Using Euler's formula (6) with  $y = \frac{\pi}{3}$ , we have  $e^{i\pi/3} = \cos \frac{\pi}{3} + i \sin \frac{\pi}{3} = \frac{1}{2} + \frac{\sqrt{3}}{2}i$ .

44. Using Euler's formula (6) with  $y = -\pi$ , we have  $e^{-i\pi} = \cos(-\pi) + i \sin(-\pi) = -1$ .

45. Using Equation 7 with  $x = 2$  and  $y = \pi$ , we have  $e^{2+i\pi} = e^2 e^{i\pi} = e^2(\cos \pi + i \sin \pi) = e^2(-1 + 0) = -e^2$ .

46. Using Equation 7 with  $x = \pi$  and  $y = 1$ , we have  $e^{\pi+i} = e^\pi \cdot e^{1i} = e^\pi(\cos 1 + i \sin 1) = e^\pi \cos 1 + (e^\pi \sin 1)i$ .

47. Take  $r = 1$  and  $n = 3$  in De Moivre's Theorem to get

$$\begin{aligned} [1(\cos \theta + i \sin \theta)]^3 &= 1^3(\cos 3\theta + i \sin 3\theta) \\ (\cos \theta + i \sin \theta)^3 &= \cos 3\theta + i \sin 3\theta \\ \cos^3 \theta + 3(\cos^2 \theta)(i \sin \theta) + 3(\cos \theta)(i \sin \theta)^2 + (i \sin \theta)^3 &= \cos 3\theta + i \sin 3\theta \\ \cos^3 \theta + (3 \cos^2 \theta \sin \theta)i - 3 \cos \theta \sin^2 \theta - (\sin^3 \theta)i &= \cos 3\theta + i \sin 3\theta \\ (\cos^3 \theta - 3 \sin^2 \theta \cos \theta) + (3 \sin \theta \cos^2 \theta - \sin^3 \theta)i &= \cos 3\theta + i \sin 3\theta \end{aligned}$$

Equating real and imaginary parts gives  $\cos 3\theta = \cos^3 \theta - 3 \sin^2 \theta \cos \theta$  and  $\sin 3\theta = 3 \sin \theta \cos^2 \theta - \sin^3 \theta$ .

48. Using Formula 6,

$$e^{ix} + e^{-ix} = (\cos x + i \sin x) + [\cos(-x) + i \sin(-x)] = \cos x + i \sin x + \cos x - i \sin x = 2 \cos x$$

Thus,  $\cos x = \frac{e^{ix} + e^{-ix}}{2}$ . Similarly,

$$e^{ix} - e^{-ix} = (\cos x + i \sin x) - [\cos(-x) + i \sin(-x)] = \cos x + i \sin x - \cos x - (-i \sin x) = 2i \sin x$$

Therefore,  $\sin x = \frac{e^{ix} - e^{-ix}}{2i}$ .

49.  $F(x) = e^{rx} = e^{(a+bi)x} = e^{ax+bx i} = e^{ax}(\cos bx + i \sin bx) = e^{ax} \cos bx + i(e^{ax} \sin bx) \Rightarrow$

$$\begin{aligned} F'(x) &= (e^{ax} \cos bx)' + i(e^{ax} \sin bx)' \\ &= (ae^{ax} \cos bx - be^{ax} \sin bx) + i(ae^{ax} \sin bx + be^{ax} \cos bx) \\ &= a[e^{ax}(\cos bx + i \sin bx)] + b[e^{ax}(-\sin bx + i \cos bx)] \\ &= ae^{rx} + b[e^{ax}(i^2 \sin bx + i \cos bx)] \\ &= ae^{rx} + bi[e^{ax}(\cos bx + i \sin bx)] = ae^{rx} + bie^{rx} = (a + bi)e^{rx} = re^{rx} \end{aligned}$$

50. (a) From Exercise 49,  $F(x) = e^{(1+i)x} \Rightarrow F'(x) = (1+i)e^{(1+i)x}$ . So

$$\int e^{(1+i)x} dx = \frac{1}{1+i} \int F'(x) dx = \frac{1}{1+i} F(x) + C = \frac{1-i}{2} F(x) + C = \frac{1-i}{2} e^{(1+i)x} + C$$



$$(b) \int e^{(1+i)x} dx = \int e^x e^{ix} dx = \int e^x (\cos x + i \sin x) dx = \int e^x \cos x dx + i \int e^x \sin x dx \quad (1).$$

Also,

$$\begin{aligned} \frac{1-i}{2} e^{(1+i)x} &= \frac{1}{2} e^{(1+i)x} - \frac{1}{2} i e^{(1+i)x} = \frac{1}{2} e^{x+ix} - \frac{1}{2} i e^{x+ix} \\ &= \frac{1}{2} e^x (\cos x + i \sin x) - \frac{1}{2} i e^x (\cos x + i \sin x) \\ &= \frac{1}{2} e^x \cos x + \frac{1}{2} e^x \sin x + \frac{1}{2} i e^x \sin x - \frac{1}{2} i e^x \cos x \\ &= \frac{1}{2} e^x (\cos x + \sin x) + i \left[ \frac{1}{2} e^x (\sin x - \cos x) \right] \quad (2) \end{aligned}$$

Equating the real and imaginary parts in (1) and (2), we see that  $\int e^x \cos x dx = \frac{1}{2} e^x (\cos x + \sin x) + C$  and

$$\int e^x \sin x dx = \frac{1}{2} e^x (\sin x - \cos x) + C.$$

NOT FOR SALE

INSTRUCTOR USE ONLY

© Cengage Learning. All Rights Reserved.