

ECE 311 Lab 4

Jacob Hutter

March 26, 2017

Report Item 1

```
1 function [ ] = filters(N,wc,w0 )
    d = zeros(1,N*2 + 1);
3    d(N+1) = 1; % delta function
    n = N*2+1;
5    w = fftshift((0:n-1)/n*2*pi);
    w(1:n/2) = w(1:n/2) - 2*pi;
7    N = linspace(-N,N,(N*2)+1); % create -N to N array
    lpi = (wc/pi).*sinc(wc.*N./pi);
9    lpm = fftshift(fft(lpi));
    hpi = d-lpi;
11    hpm = fftshift(fft(hpi));
    bpi = cos(w0.*N).*lpi;
13    bpm = fftshift(fft(bpi));





15    figure;
    subplot(211);
17    stem(N,lpi);
    title('Low Pass Filter impulse Response');
19    ylabel('h[n]');
    xlabel('n');
21    subplot(212);
    plot(w,abs(lpm));
23    title('Low Pass Filter Magnitude Response');
    ylabel('|Hd(w)|');
25    xlabel('w');





27    figure;
    subplot(211);
29    stem(N,hpi);
    title('High Pass Filter impulse Response');
31    ylabel('h[n]');
    xlabel('n');
33    subplot(212);
    plot(w,abs(hpm));
35    title('High Pass Filter Magnitude Response');
    ylabel('|Hd(w)|');
37    xlabel('w');





39    figure;
    subplot(211);
41    stem(N,bpi);
    title('Band Pass Filter impulse Response');
```

```
43     ylabel('h[n]');  
44     xlabel('n');  
45     subplot(212);  
46     plot(w,abs(bpm));  
47     title('Band Pass Filter Magnitude Response');  
48     ylabel('|Hd(w)|');  
49     xlabel('w');  
end
```

filters.m








```

1 load impulseresponse.mat
  % variable name is h
3 figure;
  subplot
5 subplot(311);
  stem(h);
7 n = 74;
  w = fftshift((0:n-1)/n*2*pi);
9 w(1:n/2) = w(1:n/2) - 2*pi;
  title('Impulse Response of h');
11 xlabel('n');
  ylabel('h[n]');
13 subplot(312);
  h_m = abs(fftshift(fft(h)));
15 h_m = mag2db(h_m);
  plot(w,h_m);
17 title('Magnitude Response of h');
  xlabel('w');
19 ylabel('db');
  subplot(313);
21 h_p = angle(fftshift(fft(h)));
  plot(w,h_p);
23 title('Phase Response of h');
  xlabel('w');
25 ylabel('angle of Hd(w)');

27 %find pass band ripple
  top = max(h_m);
29 bottom_range = h_m(28:48);
  bottom = min(bottom_range);
31 passband_ripple = top - bottom;
  % result is 8.0126
33 %passband edge is approximately .75 rad to 1.25 rad so .5 rad

```

impresp.m




```

N = 25;
M = (N-1)/2;
w = fftshift((0:N-1)/N*2*pi); % 1. define omega as you would for
    FFT
w(1:N/2) = w(1:N/2) - 2*pi; %
i=sqrt(-1);
for j=1:N
    if(abs(w(j)) < pi/3), % 2.
        g_w(j) = 1 * exp(-i*M*w(j));
    else
        g_w(j) = 0;
    end
end

g_n = ifft(fftshift(g_w)); % 3. find g[n], should be shifted
w_n = hamming(N)'; % window (transposed)
h_n = g_n .* w_n; % h_n is impulse response
figure;
subplot(311);
plot(abs(h_n));
title('Magnitude of h[n]');
xlabel('n');
ylabel('abs(h[n])');
subplot(312);
plot(w, mag2db(abs(fftshift(fft(h_n)))));
title('Magnitude of Hd(w)');
xlabel('w');
ylabel('db');
subplot(313);
plot(w, angle(fftshift(fft(h_n))));
title('Phase of Hd(w)');
xlabel('w');
ylabel('radians');

```

FIR_FILTER.m




```

1  clc , clear all , close all
   f = [54,64];
3  a = [1,0];
   rp = [2];
5  rs = 50;
   fs = 300;
7  dev = [(10^(rp/20)-1)/(10^(rp/20)+1)  10^(-rs/20)];
   [n,fo,mo,w] = firpmord(f, a, dev, fs);
9  b = firpm(n,fo,mo,w);
   freqz(b,1);
11 figure
    impz(b,1);

```

report4.m






```

function [a, b, c] = mySTDFT(x, M, D, P, f_s )
2 a_dim1 = ceil(P/2);
a_dim2 = floor(((length(x) - M)/D) + 1);
4 dft_mat = zeros(a_dim1,a_dim2);
time_shifts = zeros(1,a_dim2);
6 for i=1:a_dim2,
    time_shifts(i) = (i*D)/f_s; % vector of shifts
8     time_slice = x((1+D*(i-1)):((i-1)*D)+M)); % get time slice
    time_slice = fft(time_slice,P); % dft zero padded to P
10    time_slice = time_slice(1,1:a_dim1); % only take half of P
    dft_mat(:,i) = time_slice; %assign column in dft_mat
12 end
a = dft_mat; % output 1
14 b = linspace(0,f_s,a_dim1); % output 2
c = time_shifts; % output 3
16 imagesc(c,b,abs(a));
ylabel('Frequency');
18 xlabel('time shifts');
str = sprintf('Spectrogram at M = %f',M);
20 title(str);
end

```

mySTDFT.m



```

2  [y fs] = audioread('sound1.wav');

4  N = length(y);
   w = fftshift((0:N-1)/N*2*pi); % define omega as you would for FFT
6  w(1:N/2) = w(1:N/2) - 2*pi; %
   y_w = fftshift(fft(y));
8  figure;
   plot(w,abs(y_w));
10 ylabel('magnitude');
   xlabel('radians');
12 title('magnitude spectrum of sound1.wav before filter');



14
16 f= [0 .4 .5 1];
   a = [1 1 0 0];
   b = firpm(50,f,a);
18 b_w = fftshift(fft(b,length(y)))'; % get filter

20 y_w = b_w .* y_w; % apply filter
   figure;
22 plot(w,abs(y_w));
   ylabel('magnitude');
24 xlabel('radians');
   title('magnitude spectrum of sound1.wav after filter');

26
   y = ifft(ifftshift(y_w));
28 soundsc(y);
   filename = 'filtered1.wav';
30 audiowrite(filename,y,fs);

```

report6.m



```



[y fs] = audioread('sound2.wav');
2 %soundsc(y);
N = length(y);
4 w = fftshift((0:N-1)/N*2*pi); % define omega as you would for FFT
w(1:N/2) = w(1:N/2) - 2*pi; %
6
yw = fftshift(fft(y));
8 ys = spectrogram(y);
figure;
10 plot(w,abs(yw));
title('sound2.wav magnitude response prior to filtering');
12 xlabel('radians');
ylabel('magnitude');
14 figure;
imagesc(abs(ys));
16
f = [0 .1 .2 1];
18 a = [1 0 0 0];
b = firpm(50,f,a);
20 b_w = fftshift(fft(b,length(y)))'; % get filter

22 yw = b_w .* yw; % apply filter
figure;
24 plot(w,abs(yw));
ylabel('magnitude');
26 xlabel('radians');
title('magnitude spectrum of sound2.wav after filter');
28

30 y = ifft(ifftshift(yw));
ys = spectrogram(y);
32 figure;
imagesc(abs(ys));
34 soundsc(y);
filename = 'filtered2.wav';
36 audiowrite(filename,y,fs);

```

report7.m



```

2 y = load('speechsig.mat');
  x = y.x;
4 xnoise = y.xnoise;

6 N = length(xnoise);
  w = fftshift((0:N-1)/N*2*pi); % define omega as you would for FFT
8 w(1:N/2) = w(1:N/2) - 2*pi; %

10 xnoisew = fftshift(fft(xnoise));

12 h = hamming(N);
  xnoisewh = xnoisew .* h;
14
  xnoise = ifft(ifftshift(xnoisewh),N);
16 %soundsc(real(xnoise));

18 r = zeros(N,1);
  r(2000:4999,1) = rectwin(3000);
20
  xnoisewr = xnoisew .* r;
22 plot(abs(xnoisew));
  xnoise = ifft(ifftshift(xnoisewr));
24
26 %soundsc(real(xnoise));

```

report8.m