

Algebra

Applied Mathematics

Calculus and Analysis

Discrete Mathematics

Foundations of Mathematics

Geometry

History and Terminology

Number Theory

Probability and Statistics

Recreational Mathematics

Topology

Alphabetical Index

Interactive Entries

Random Entry

New in MathWorld

MathWorld Classroom

About MathWorld

Contribute to MathWorld

Send a Message to the Team

MathWorld Book

Wolfram Web Resources »

13,651 entries
Last updated: Fri Jun 29 2018

Created, developed, and
nurtured by Eric Weisstein
at Wolfram Research

Discrete Mathematics > Combinatorics > Permutations >
Interactive Entries > Interactive Demonstrations >

Permutation

DOWNLOAD

Wolfram Notebook

EXPLORE THIS TOPIC IN

The MathWorld Classroom

A permutation, also called an "arrangement number" or "order," is a rearrangement of the elements of an ordered list S into a [one-to-one correspondence](#) with S itself. The number of permutations on a set of n elements is given by $n!$ (n factorial; Uspensky 1937, p. 18). For example, there are $2! = 2 \cdot 1 = 2$ permutations of $\{1, 2\}$, namely $\{1, 2\}$ and $\{2, 1\}$, and $3! = 3 \cdot 2 \cdot 1 = 6$ permutations of $\{1, 2, 3\}$, namely $\{1, 2, 3\}$, $\{1, 3, 2\}$, $\{2, 1, 3\}$, $\{2, 3, 1\}$, $\{3, 1, 2\}$, and $\{3, 2, 1\}$. The permutations of a list can be found in the [Wolfram Language](#) using the command `Permutations[list]`. A list of length n can be tested to see if it is a permutation of $1, \dots, n$ in the [Wolfram Language](#) using the command `PermutationListQ[list]`.

Sedgewick (1977) summarizes a number of algorithms for generating permutations, and identifies the minimum change permutation algorithm of Heap (1963) to be generally the fastest (Skiena 1990, p. 10). Another method of enumerating permutations was given by Johnson (1963; Séroul 2000, pp. 213-218).

The number of ways of obtaining an *ordered* subset of k elements from a set of n elements is given by

$${}_n P_k \equiv \frac{n!}{(n-k)!}$$

(1)

(Uspensky 1937, p. 18), where $n!$ is a [factorial](#). For example, there are $4!/2! = 12$ 2-subsets of $\{1, 2, 3, 4\}$, namely $\{1, 2\}$, $\{1, 3\}$, $\{1, 4\}$, $\{2, 1\}$, $\{2, 3\}$, $\{2, 4\}$, $\{3, 1\}$, $\{3, 2\}$, $\{3, 4\}$, $\{4, 1\}$, $\{4, 2\}$, and $\{4, 3\}$. The *unordered* subsets containing k elements are known as the *k-subsets* of a given set.

A representation of a permutation as a product of [permutation cycles](#) is unique (up to the ordering of the cycles). An example of a cyclic decomposition is the permutation $\{4, 2, 1, 3\}$ of $\{1, 2, 3, 4\}$. This is denoted $(2)(143)$, corresponding to the disjoint permutation cycles (2) and (143) . There is a great deal of freedom in picking the representation of a cyclic decomposition since (1) the cycles are disjoint and can therefore be specified in any order, and (2) any rotation of a given cycle specifies the same cycle (Skiena 1990, p. 20). Therefore, $(431)(2)$, $(314)(2)$, $(143)(2)$, $(2)(431)$, $(2)(314)$, and $(2)(143)$ all describe the same permutation.

Another notation that explicitly identifies the positions occupied by elements before and after application of a permutation on n elements uses a $2 \times n$ matrix, where the first row is $(123 \dots n)$ and the second row is the new arrangement. For example, the permutation which switches elements 1 and 2 and fixes 3 would be written as

$$\begin{bmatrix} 1 & 2 & 3 \\ 2 & 1 & 3 \end{bmatrix}.$$

(2)

Any permutation is also a product of [transpositions](#). Permutations are commonly denoted in [lexicographic](#) or [transposition order](#). There is a correspondence between a permutation and a pair of [Young tableaux](#) known as the [Schensted correspondence](#).

The number of wrong permutations of n objects is $[n!/e]$ where $[x]$ is the [nearest integer function](#). A permutation of n ordered objects in which no object is in its natural place is called a [derangement](#) (or sometimes, a complete permutation) and the number of such permutations is given by the [subfactorial](#) $!n$.

Using

$$(x+y)^n = \sum_{r=0}^n \binom{n}{r} x^{n-r} y^r$$

(3)

with $x = y = 1$ gives

$$2^n = \sum_{r=0}^n \binom{n}{r},$$

(4)

so the number of ways of choosing 0, 1, ..., or n at a time is 2^n .

The set of all permutations of a set of elements $1, \dots, n$ can be obtained using the following recursive procedure

$$\begin{array}{c} 1 \ 2 \\ / \\ 2 \ 1 \end{array} \qquad \begin{array}{c} 1 \ 2 \ 3 \\ / \quad / \\ 1 \ 3 \ 2 \end{array} \qquad \begin{array}{c} 1 \ 2 \ 3 \\ / \quad / \quad / \\ 3 \ 1 \ 2 \end{array}$$

(5)

$$\begin{array}{c} 1 \ 2 \ 3 \\ / \quad / \quad / \\ 3 \ 1 \ 2 \\ | \\ 3 \ 2 \ 1 \\ \backslash \quad \backslash \\ 2 \ 3 \ 1 \\ \quad \backslash \\ 2 \quad 1 \ 3 \end{array}$$

(6)

Consider permutations in which no pair of consecutive elements (i.e., rising or falling successions) occur. For $n = 1, 2, \dots$ elements, the numbers of such permutations are 1, 0, 0, 2, 14, 90, 646, 5242, 47622, ... (OEIS [A002464](#)).

Let the set of integers $1, 2, \dots, N$ be permuted and the resulting sequence be divided into increasing [runs](#). Denote the average length of the n th run as N approaches infinity, L_n . The first few values are summarized in the following table, where e is the base of the [natural logarithm](#) (Le Lionnais 1983, pp. 41-42; Knuth 1998).

n	L_n	Sloane	approximate
1	$e - 1$	A091131	1.7182818...
2	$e^2 - 2e$	A091132	1.9524...
3		A091133	1.9957...

binomial theorem

THINGS TO TRY:

- = binomial theorem
- = symmetric group
- = permutation (1 3 5)(2 4)(6

Interactive knowledge apps from
Wolfram Demonstrations Project

Generating Permutations by Sequence of Transpositions
George Beck



Random Permutation of a Given Length
Neil Chriss

Permutations, Derangements, Other Forbidden Position Problems Using Non-Attacking Rooks
Marc Brodie

Permutation Tree
Michael Schreiber

1 > 0

∞ > 1

 > 

Mathematica Student Edition

$$e^3 - 3e^2 + \frac{3}{2}e$$

SEE ALSO:
[Alternating Permutation](#), [Ball Picking](#), [Binomial Coefficient](#), [Choose](#), [Circular Permutation](#), [Combination](#), [Derangement](#), [Eulerian Number](#), [Even Permutation](#), [k-Subset](#), [Linear Extension](#), [Married Couples Problem](#), [Multichoose](#), [Multinomial Coefficient](#), [Multiset](#), [Odd Permutation](#), [Permutation Ascent](#), [Permutation Cycle](#), [Permutation Inversion](#), [Permutation Matrix](#), [Permutation Pattern](#), [Permutation Run](#), [Permutation Symbol](#), [Random Permutation](#), [String](#), [Subfactorial](#), [Transposition](#)

REFERENCES:
Bogomolny, A. "Graphs." http://www.cut-the-knot.org/do_you_know/permutation.shtml.
Conway, J. H. and Guy, R. K. "Arrangement Numbers." In *The Book of Numbers*. New York: Springer-Verlag, p. 66, 1996.
Dickau, R. M. "Permutation Diagrams." <http://mathforum.org/advanced/robertd/permutations.html>.
Heap, B. R. "Permutations by Interchanges." *Computer J.* **6**, 293-294, 1963.
Johnson, S. M. "Generation of Permutations by Adjacent Transpositions." *Math. Comput.* **17**, 282-285, 1963.
Knuth, D. E. *The Art of Computer Programming, Vol. 3: Sorting and Searching, 2nd ed.* Reading, MA: Addison-Wesley, pp. 38-43, 1998.
Kraitchik, M. "The Linear Permutations of n Different Things." §10.1 in *Mathematical Recreations*. New York: W. W. Norton, pp. 239-240, 1942.
Le Lionnais, F. *Les nombres remarquables*. Paris: Hermann, 1983.
Ruskey, F. "Information on Permutations." <http://www.theory.csc.uvic.ca/~cos/inf/per/PermlInfo.html>.
Sedgewick, R. "Permutation Generation Methods." *Comput. Surveys* **9**, 137-164, 1977.
Séroul, R. "Permutations: Johnson's [sic] Algorithm." §8.15 in *Programming for Mathematicians*. Berlin: Springer-Verlag, pp. 213-218, 2000.
Skiena, S. "Permutations." §1.1 in *Implementing Discrete Mathematics: Combinatorics and Graph Theory with Mathematica*. Reading, MA: Addison-Wesley, pp. 3-16, 1990.
Sloane, N. J. A. Sequences [A000142/M1675](#), [A002464/M2070](#), [A091131](#), [A091132](#), and [A091133](#) in "The On-Line Encyclopedia of Integer Sequences."
Trotter, H. F. "Perm (Algorithm 115)." *Comm. ACM* **5**, 434-435, 1962.
Uspensky, J. V. *Introduction to Mathematical Probability*. New York: McGraw-Hill, p. 18, 1937.

Referenced on Wolfram|Alpha: [Permutation](#)

CITE THIS AS:
Weisstein, Eric W. "Permutation." From *MathWorld*--A Wolfram Web Resource. <http://mathworld.wolfram.com/Permutation.html>

Wolfram Web Resources

Mathematica » The #1 tool for creating Demonstrations and anything technical.	Wolfram Alpha » Explore anything with the first computational knowledge engine.	Wolfram Demonstrations Project » Explore thousands of free applications across science, mathematics, engineering, technology, business, art, finance, social sciences, and more.
Computerbasedmath.org » Join the initiative for modernizing math education.	Online Integral Calculator » Solve integrals with Wolfram Alpha.	Step-by-step Solutions » Walk through homework problems step-by-step from beginning to end. Hints help you try the next step on your own.
Wolfram Problem Generator » Unlimited random practice problems and answers with built-in Step-by-step solutions. Practice online or make a printable study sheet.	Wolfram Education Portal » Collection of teaching and learning tools built by Wolfram education experts: dynamic textbook, lesson plans, widgets, interactive Demonstrations, and more.	Wolfram Language » Knowledge-based programming for everyone.