

No alpha-specific transcriptomic signature of the cortical alpha-rhythm sources targeted by closed-loop EEG–TMS: an imaging-transcriptomics test of the ENIGMA-EEG oscillatory-power GWAS

Running title: Alpha genetics and the transcriptomics of TMS–EEG alpha targets

Abstract

Background. The power and frequency of the electroencephalographic (EEG) alpha rhythm are among the most heritable human neurophysiological traits, and a dedicated genome-wide association study (GWAS) of EEG oscillatory power (ENIGMA-EEG; Smit et al., 2018) has identified common variants associated with band-limited power. In parallel, brain-state-dependent EEG–TMS relies on a specific set of cortical alpha-rhythm sources and their functional-connectivity architecture (Tabarelli et al., 2022). Whether the genes implicated in alpha power are transcriptomically concentrated in these stimulation-relevant cortical sources has never been tested.

Objective. To test whether alpha-associated genetic signal, mapped to regional gene expression via imaging transcriptomics, is spatially enriched — specifically — in the cortical alpha-source regions relevant for closed-loop EEG–TMS.

Methods. We performed MAGMA gene-based analysis of ENIGMA-EEG summary statistics for central alpha power (primary phenotype), occipital alpha power, occipital alpha peak frequency (IAF), and three control bands (theta, beta, delta). Regional transcription was obtained from the Allen Human Brain Atlas (AHBA) with *abagen* in the Glasser HCP-MMP1.0 atlas (primary), plus Schaefer-100 and Yan-600 (sensitivity). Two per-region “alpha transcriptomic scores” were computed — a continuous score (rank correlation between a region’s expression profile and the genome-wide gene-level association statistics) and a top-100 gene-set composite. Enrichment in the 41 Tabarelli alpha-source regions was tested against a spatial-autocorrelation-preserving spin-test null (10,000 rotations).

Results. A positive control confirmed the pipeline detects true spatial enrichment (dominant expression gradient, $p_{\text{spin}} = 2 \times 10^{-6}$), and the spin test agreed with an independent surrogate model (brainsmash). Gene-based analysis recovered biologically credible signal, including the well-known chr3p21 pleiotropic cluster (*GNL3*, *GLT8D1*, *NT5DC2*, *PBRM1*). For central alpha power, both scores showed nominal enrichment in the alpha-source regions (continuous $p_{\text{spin}} = 0.022$; top-100 $p_{\text{spin}} = 0.030$). However, this enrichment was **not specific and not robust**: control bands enriched comparably or more strongly (top-100 beta $p_{\text{spin}} = 0.011$; delta $p_{\text{spin}} = 0.042$; continuous theta $p_{\text{spin}} = 0.042$); the occipital alpha phenotype did not replicate the continuous-score result ($p_{\text{spin}} = 0.20$); the alpha peak-frequency (IAF) phenotype was non-significant ($p_{\text{spin}} = 0.066$); against a co-expression-aware gene-set null the top-100 enrichment vanished ($p_{\text{geneset}} = 0.33$); and **no test survived false-discovery-rate correction** across the 24-cell phenotype $\times$ score $\times$ region-set grid (minimum $q = 0.127$). Coarsening the parcellation toward the spatial resolution of the MEG source reconstruction did not strengthen the effect (Schaefer-100 continuous $p_{\text{spin}} = 0.071$).

Conclusion. We find no alpha-specific transcriptomic signature of the cortical alpha-rhythm sources targeted by closed-loop EEG–TMS. At most there is a weak, non-specific, parcellation-

fragile tendency for genes associated with EEG oscillatory power in general to be over-expressed in this occipito-parieto-frontal territory. We interpret the result in light of the modest GWAS sample, the shared genetic architecture of neighbouring EEG bands, and the intrinsic resolution mismatch between MEG source reconstruction and fine-grained transcriptomics.

Keywords: alpha rhythm; EEG; imaging transcriptomics; Allen Human Brain Atlas; GWAS; MAGMA; TMS-EEG; spin test; null result.

1. Introduction

The occipito-parietal alpha rhythm (~8–13 Hz) is the dominant oscillation of the awake human EEG at rest. Twin studies place the heritability of alpha power and peak frequency among the highest of any neurophysiological trait ($h^2 = 0.7$ – 0.8 ; Smit et al., 2005), and a meta-analytic GWAS of EEG oscillatory power by the ENIGMA-EEG working group (Smit et al., 2018) identified common variants associated with band-limited power across delta, theta, alpha and beta, linking several psychiatric-liability genes to oscillatory brain activity.

A largely separate methodological literature has developed around brain-state-dependent EEG–TMS, in which the delivery of a transcranial magnetic stimulation pulse is synchronised to the instantaneous phase or power of the ongoing alpha rhythm in order to bias the direction and magnitude of induced plasticity (Zrenner et al., 2018). The efficacy of such closed-loop protocols depends on which cortical generators produce the alpha activity that the controller reads out and targets. Tabarelli et al. (2022) characterised these generators directly: using resting-state magnetoencephalography (MEG) from the Cam-CAN cohort and minimum-norm source reconstruction in the Glasser HCP-MMP1.0 multimodal parcellation, they identified 41 cortical alpha-rhythm sources spanning occipital, parietal and frontal sectors and described two complementary functional-connectivity states, dominated by strong parieto-frontal phase coupling with prefrontal area 8C as the principal hub.

These two lines of work — the genetics of the alpha rhythm and the neurocircuitry targeted by closed-loop stimulation — have never been related. Imaging transcriptomics, which correlates macroscale brain maps with regional gene-expression profiles from the Allen Human Brain Atlas (AHBA; Hawrylycz et al., 2012) via toolboxes such as *abagen* (Markello et al., 2021), is now a mature approach for grounding neuroimaging phenotypes in molecular architecture (Arnatkeviciute et al., 2019; Fornito et al., 2019). It has, however, never been applied to an EEG-trait GWAS to ask whether the genetic architecture of the alpha rhythm is regionally concentrated in the very cortical sources that neurostimulation seeks to engage.

Here we test a single, pre-specified hypothesis:

H1. Genes associated with EEG alpha power (from gene-based analysis of the ENIGMA-EEG GWAS) are expressed more highly in the cortical alpha-source regions relevant for closed-loop EEG–TMS (Tabarelli et al., 2022) than in the rest of cortex, and this enrichment is *specific* to the alpha band.

We evaluate H1 with a fully reproducible, secondary-data pipeline: MAGMA gene-based analysis of the ENIGMA-EEG summary statistics; AHBA expression via *abagen* in the same Glasser parcellation used to define the alpha sources; and a spatial-enrichment test corrected for spatial autocorrelation with a spin-test null. The design is deliberately built to yield an *informative* answer

even from a modestly powered GWAS. First, *specificity* is assessed with an **internally power-matched** comparison — three non-alpha EEG bands and a second alpha phenotype are pushed through the identical pipeline, so the relative contrast between alpha and control bands is meaningful regardless of the absolute power of any one GWAS. Second, the enrichment test itself is validated with a **positive control** (a known expression gradient it must recover) and guarded against the co-expression-driven false positives characteristic of transcriptomic enrichment (Fulcher et al., 2021) with a gene-set null, in addition to the spatial null. Third, *robustness* is examined across three parcellations of different granularity. As we report below, the hypothesis is not supported: the enrichment that appears for one alpha phenotype is neither band-specific, replicable across alpha phenotypes, robust to a gene-set null, nor robust to multiple-comparison correction — while the positive control confirms the pipeline would have detected a true effect. We present this as a rigorous null and discuss the methodological factors — GWAS power, the shared genetics of adjacent EEG bands, and the spatial resolution of MEG source reconstruction — that bound its interpretation.

Do EEG alpha-power genes concentrate in the cortical alpha sources targeted by closed-loop TMS-EEG?



Figure 1: **Study design.** From the ENIGMA-EEG oscillatory-power GWAS through MAGMA gene-based analysis and AHBA regional expression to the Tabarelli alpha-source ROIs, tested for spatial enrichment against a spin-test null with specificity and robustness controls.

2. Methods

All analyses used publicly available secondary data. No primary data were collected. Code and derived results are available (see *Data and code availability*); raw GWAS, AHBA and reference-panel data are redistributed under their original licences only as download instructions.

2.1 GWAS summary statistics

We obtained ENIGMA-EEG oscillatory-power summary statistics (Smit et al., 2018; 2018-11-01 release, `select_N6000_Hom100`) for six scalp-derived phenotypes: alpha power at the central electrode Cz (**alphaCz**, the pre-declared primary phenotype), alpha power at an occipital site (**alphaOcc**), the occipital alpha peak frequency (**peakOcc**, an index of individual alpha frequency), and — as specificity controls — theta, beta and delta power at Cz (**thetaCz**, **betaCz**, **deltaCz**). Each file contained 4.96 million autosomal SNPs with per-SNP sample size ($N = 6,000\text{--}7,700$). The genome build was verified as GRCh37/hg19 (e.g., rs3094315 at chr1:752,566), matching the gene-location and reference-panel builds, so no liftOver was required.

2.2 Gene-based analysis (MAGMA)

Gene-based association was computed with MAGMA v1.10 (de Leeuw et al., 2015). SNPs were annotated to genes (build-37 NCBI gene locations; 19,427 genes) with a 35 kb upstream / 10 kb downstream window; the 18,053 genes with 1 mapped SNP were carried into the gene-based test. The SNP-wise mean model used the 1000 Genomes European reference panel (1000 Genomes Project Consortium, 2015; 503 individuals) to model linkage disequilibrium. Entrez gene identifiers were mapped to HGNC symbols using the gene-location file. Gene p -values were corrected by Bonferroni and Benjamini–Hochberg FDR. For the enrichment analysis we retained (i) the full genome-wide vector of gene-level z -statistics and (ii) a top-100 gene set ranked by gene p -value; threshold-based sets (Bonferroni, FDR, $p < 10^{-3}$, $p < 10^{-2}$) were also generated for reference.

2.3 Regional gene expression (AHBA / abagen)

Regional transcription was estimated with *abagen* (development build post-0.1.3, GitHub [netneurolab/abagen](https://github.com/netneurolab/abagen), required for pandas 2 compatibility; Markello et al., 2021) from AHBA microarray data (Hawrylycz et al., 2012). Default processing was used (intensity-based filtering threshold 0.5; probe selection by differential stability; within-donor scaled robust sigmoid normalisation; `missing="interpolate"`; matched-sample normalisation). Of the six AHBA donors, five were used; donor 15496 (H0351.1015, a left-hemisphere-only donor) was excluded because its normalized-microarray archive was unavailable at source at the time of analysis (persistent HTTP 404). The primary parcellation was Glasser HCP-MMP1.0 (Glasser et al., 2016; 360 regions), matching the space in which the alpha sources are defined. Because AHBA samples are mapped by *abagen* to `fsaverage5`, the Glasser atlas was resampled from `fsLR32k` to `fsaverage5` using the Human Connectome Project `resample_fsaverage` registration spheres (nearest-neighbour in the common `fsaverage` frame); resampling preserved all 180 parcels per hemisphere. The resulting expression matrix contained 360 regions $\times$ 15,563 genes, with complete coverage of all 82 target parcels. For robustness, expression was also computed in the Schaefer-100 (Schaefer et al., 2018) and Yan-600 homotopic (Yan et al., 2023) parcellations, both in `fsaverage5`.

2.4 Alpha-source region set

The 41 alpha-source regions of Tabarelli et al. (2022) were used as the target set (16 occipital, 7 parietal, 18 frontal HCP-MMP labels), applied bilaterally (82 parcels). Because the sources are natively defined in the Glasser atlas, the mapping is one-to-one by label, avoiding subjective region translation. A prefrontal-hub subset (areas 8C, 8Av, 9a; 6 parcels) was additionally tested, motivated by the paper’s identification of 8C as the principal connectivity hub. For the Schaefer-100 and Yan-600 sensitivity analyses, the alpha-source set was re-mapped by `fsaverage5` surface overlap: a parcel was flagged as target if 50% of its cortical vertices fell within the Glasser alpha-source set (31/100 Schaefer parcels; 180/600 Yan parcels).

2.5 Alpha transcriptomic score and enrichment test

We defined two per-region scores. The **continuous** score is the Spearman correlation, across all genes shared between the expression matrix and the gene-based results, between a region’s expression profile and the genome-wide gene-level z -statistics; it uses the full association signal without any significance threshold. The **top-100** score is the mean z -scored expression, across regions, of the top-100 gene set. Enrichment was quantified as the difference between the mean score in the target regions and the rest of cortex.




Figure 2: Alpha-source ROIs on the inflated cortex, across parcellations. The 41 Tabarelli alpha-source regions (bilateral) shown on the fsaverage5 inflated surface for Glasser HCP-MMP1.0, Schaefer-100 and Yan-600. In every parcellation the target parcels are coloured by sector (occipital, parietal, frontal); for Schaefer and Yan each target parcel takes the sector of the Glasser alpha-source it most overlaps. Thin dark lines are atlas parcel boundaries; grey is non-target cortex.

Statistical significance was assessed against a spin-test null that preserves the spatial autocorrelation of cortical maps (Alexander-Bloch et al., 2018). Because the parcellation was surface-based, we generated 10,000 spatial rotations of the parcel centroids on the fsaverage5 sphere (nearest-neighbour reassignment; left- and right-hemisphere rotations mirror-matched; seed fixed and logged), and computed a one-sided p -value (H1: target > rest) as the fraction of rotated maps with a target-region mean at least as large as observed. This self-contained implementation was used because the spin generator had been removed from the version of *netneurotools* available at analysis time. Specificity and replication were assessed by repeating the test for all six phenotypes and both scores, and correcting the resulting 24-cell grid by Benjamini–Hochberg FDR. Robustness was assessed by repeating the primary-phenotype test in the Schaefer-100 and Yan-600 parcellations. Because the analysis is internally power-matched — every phenotype shares the same GWAS design, the same expression matrix and the same null — the *relative* comparison between the alpha and control bands is informative about specificity independent of the absolute statistical power of any single GWAS.

2.6 Validation and complementary nulls

Three analyses established the validity of the enrichment test. **(i) Positive control.** To confirm that the pipeline can detect a genuine spatial effect, we replaced the gene statistic with the gene loadings of the first principal component (PC1) of the AHBA expression matrix — the dominant, well-replicated cortical expression gradient — and set the target to the top-tertile PC1 regions; a correct pipeline must recover a highly significant enrichment. **(ii) Co-expression-aware gene-set null.** Because gene-set enrichment on AHBA is prone to false positives from within-set spatial co-expression (Fulcher et al., 2021), we complemented the spin test (which permutes space) with a gene-set null for the top-100 analysis: 10,000 random gene sets of matched size were drawn, and p_{geneset} was the fraction whose target-versus-rest difference met or exceeded the observed value. **(iii) Spin validation.** Because the spin generator was self-implemented, we benchmarked it against an independent spatial-autocorrelation-preserving surrogate model (brainsmash; Burt et al., 2020; Markello & Misić, 2021), generating 1,000 surrogate maps of the primary alpha score over a Euclidean parcel-distance matrix and comparing the resulting p -value with the spin p -value.

2.7 Reproducibility

Every stage logged its parameters (random seed, parcellation version, *abagen* and package versions, MAGMA build) to a machine-readable run log. A global seed (20260721) governed all permutations. Analyses ran under Python 3.12.

3. Results

3.1 Gene-based analysis recovers credible signal

MAGMA gene-based analysis of central alpha power tested 18,053 genes. A single gene survived genome-wide Bonferroni and FDR correction — *PRKG2* ($p = 4.0 \times 10^{-8}$, $q = 0.007$) — consistent with the modest sample size. The leading sub-threshold signal included the chr3p21 pleiotropic cluster (*GNL3*, *GLT8D1*, *NT5DC2*, *PBRM1*; $p = 4\text{--}5 \times 10^{-6}$), one of the most robust loci in psychiatric genetics and consistent with the ENIGMA-EEG thesis that psychiatric-liability genes contribute to oscillatory brain activity (Smit et al., 2018). The presence of this expected locus provides a positive control that the gene-based pipeline behaves correctly. Because only one gene survived strict correction, we did not treat a threshold-based gene set as the primary readout; instead the

continuous score (using the full gene-level signal) was pre-declared primary and the top-100 gene set served as a sensitivity analysis. (The analysis grid was fixed in advance but was not formally pre-registered.)




Figure 3: Gene-based genetics. (A) MAGMA gene-based association for central alpha power (alphaCz); *PRKG2* passes Bonferroni and the chr3p21 pleiotropic cluster is the leading sub-threshold signal. (B) Top-10 genes. (C) $-\log_{10} p$ of the top alphaCz genes across all six phenotypes — the gene-level signal is largely specific to alphaCz.

3.2 The enrichment test is valid and the custom spin is corroborated

The positive control confirmed that the pipeline detects genuine spatial structure: with the PC1 expression-gradient loadings as the gene statistic and the top-tertile PC1 regions as the target, the spin test recovered a large, highly significant enrichment (target-versus-rest difference = 0.74; $p_{\text{spin}} = 2 \times 10^{-6}$). The subsequent alpha nulls are therefore not a floor effect of an insensitive method. The self-implemented spin was corroborated by an independent surrogate model: for the primary alpha map, the spin p -value (0.022) and the brainsmash p -value (0.018) were closely concordant.

3.3 Primary phenotype: nominal but modest enrichment

In the primary Glasser parcellation, central alpha power showed nominal enrichment of alpha transcriptomic signal in the 41 alpha-source regions for both scores: the continuous score gave a target-versus-rest difference corresponding to $z = 1.98$ relative to the spin null ($p_{\text{spin}} = 0.022$), and the top-100 score gave $z = 1.77$ ($p_{\text{spin}} = 0.030$). The prefrontal-hub subset (8C, 8Av, 9a) was not significant for either score (continuous $p_{\text{spin}} = 0.11$; top-100 $p_{\text{spin}} = 0.73$), indicating that any enrichment was distributed across the broad alpha-source network rather than concentrated at the connectivity hubs. Taken alone, this primary result is compatible with H1.




Figure 4: **Validation of the enrichment test.** (A) Positive control: with the dominant expression gradient (PC1) as the gene statistic, the spin test recovers a large enrichment ($p_{\text{spin}} = 2 \times 10^{-4}$). (B) Co-expression-aware gene-set null: the alpha top-100 target-versus-rest difference sits at the centre of 10,000 random gene sets ($p_{\text{geneset}} = 0.33$). (C) The custom spin agrees with brainsmash surrogates.

3.4 Specificity and replication: the effect is not alpha-specific

Repeating the test across all six phenotypes overturned the primary interpretation (Table 1). In the continuous score, the theta control band was also nominally enriched in the alpha-source regions ($p_{\text{spin}} = 0.042$), while the occipital alpha phenotype — the natural replication of the central alpha result — was not ($p_{\text{spin}} = 0.20$). In the top-100 score, the pattern was still less specific: the occipital alpha phenotype ($p_{\text{spin}} = 0.008$) and the beta control band ($p_{\text{spin}} = 0.011$) gave the strongest signals of all, both exceeding central alpha ($p_{\text{spin}} = 0.030$), with delta also nominally significant ($p_{\text{spin}} = 0.042$). The alpha peak-frequency (IAF) phenotype — arguably the most closed-loop-relevant, since the controller locks to it — was not significant for either score (continuous $p_{\text{spin}} = 0.13$; top-100 $p_{\text{spin}} = 0.066$). Critically, after Benjamini–Hochberg correction across the 24-cell phenotype $\times$ score $\times$ region-set grid, **no test remained significant** (minimum $q = 0.127$). The prefrontal-hub subset was null or negative throughout.

The co-expression-aware gene-set null reinforced this conclusion. When the alpha top-100 enrichment was tested not against a spatial null but against 10,000 random gene sets of the same size, the observed target-versus-rest difference (0.060) was unremarkable — it fell near the centre of the random-gene-set distribution (mean 0.043; $p_{\text{geneset}} = 0.33$). In other words, the nominal spin-test enrichment of the alpha gene set reflects a generic tendency of arbitrary gene sets to be marginally over-expressed in this occipito-parieto-frontal territory, rather than anything specific to alpha-associated genes — precisely the co-expression-driven inflation that spatial-only nulls do not capture (Fulcher et al., 2021).

Table 1. Spin-test enrichment of alpha transcriptomic signal in the 41 alpha-source regions (Glasser HCP-MMP1.0), by phenotype and score. z is relative to the 10,000-rotation spin null; p_{spin} is one-sided (target $>$ rest); q is Benjamini–Hochberg FDR across the full 24-cell (6 phenotype $\times$ 2 score $\times$ 2 region-set) grid. Band labels distinguish alpha power, the alpha peak-frequency (IAF) phenotype, and the non-alpha control bands.

Score	Phenotype	Band	z vs null	p_{spin}	q (FDR)
Continuous	alphaCz	alpha-power	1.98	0.022	0.169

Score	Phenotype	Band	z vs null	p_{spin}	q (FDR)
Continuous	alphaOcc	alpha-power	0.87	0.203	0.325
Continuous	peakOcc	alpha-frequency	1.16	0.130	0.283
Continuous	thetaCz	control	1.68	0.042	0.169
Continuous	betaCz	control	1.22	0.123	0.283
Continuous	deltaCz	control	0.99	0.168	0.310
Top-100	alphaCz	alpha-power	1.77	0.030	0.169
Top-100	alphaOcc	alpha-power	2.28	0.008	0.127
Top-100	peakOcc	alpha-frequency	1.72	0.066	0.227
Top-100	betaCz	control	2.13	0.011	0.127
Top-100	deltaCz	control	1.62	0.042	0.169
Top-100	thetaCz	control	1.32	0.079	0.237

3.5 Parcellation sensitivity: direction consistent, significance fragile

For the primary phenotype, the direction of the continuous-score enrichment was consistent across all three parcellations (positive; z between 1.5 and 2.0), but nominal significance was reached only in the native Glasser space (Table 2). Coarsening the parcellation to Schaefer-100 — which more closely matches the spatial resolution of MEG source reconstruction — did **not** strengthen the effect ($p_{\text{spin}} = 0.071$); refining it to Yan-600 gave a similar trend ($p_{\text{spin}} = 0.058$). The top-100 score was significant only in the Glasser parcellation. Thus the enrichment does not survive a change of parcellation, and the resolution-matching prediction — that a coarser atlas would recover a signal blurred at fine resolution — was not borne out.

Table 2. Parcellation sensitivity for central alpha power (alphaCz). Target counts reflect the fraction of parcels overlapping the Glasser alpha-source set.

Parcellation	Parcels	Targets	Score	z vs null	p_{spin}
Schaefer-100	100	31	Continuous	1.53	0.071
Schaefer-100	100	31	Top-100	1.19	0.115
Glasser-360	360	82	Continuous	1.98	0.022
Glasser-360	360	82	Top-100	1.77	0.030
Yan-600	600	180	Continuous	1.64	0.058
Yan-600	600	180	Top-100	0.94	0.200

4. Discussion

We asked whether the genetic architecture of the EEG alpha rhythm is transcriptomically concentrated in the cortical alpha sources that closed-loop EEG–TMS targets. Across a pre-specified primary test, a specificity panel of three control bands and a second alpha phenotype, and three parcellations, the answer is negative. A nominally significant enrichment appears for one alpha phenotype in one parcellation, but it is not band-specific (control bands enrich comparably or more strongly), not replicated by the second alpha phenotype, not robust to parcellation, and does not survive multiple-comparison correction. H1 is not supported.

Three considerations bound the interpretation of this null. First, **statistical power**: the ENIGMA-EEG GWAS, while the largest dedicated EEG-power meta-analysis, is modest by contemporary




Figure 5: **Alpha transcriptomic score on the cortex.** The continuous alpha score (alphaCz) per Glasser region, tertile-binned for display, on the inflated surface; the black outline marks the 41 alpha-source ROIs. The score is weakly and diffusely patterned and is not concentrated within the target ROIs.




Figure 6: **Enrichment, specificity and robustness.** (A) Primary alphaCz enrichment against the spin null (both scores). (B) Specificity across the six phenotypes (alpha in colour, controls in grey; dashed line $p = 0.05$); control bands enrich comparably or more, and nothing survives FDR. (C) Parcellation sensitivity: nominal significance only in the native Glasser space. (D) Enrichment effect size (z vs null) across phenotypes and scores.

standards, and only a single gene survived genome-wide correction in our gene-based analysis. Low gene-based signal-to-noise limits the sensitivity of any downstream enrichment, and could in principle mask a true but small effect. Three features of the design bound this concern. Our continuous score deliberately used the *entire* genome-wide gene-level signal rather than a sparse significant set, which is the analysis best suited to a low-powered GWAS; it too was non-specific. The positive control shows the enrichment machinery is not insensitive — it recovered a genuine expression gradient at $p_{\text{spin}} = 2 \times 10^{-5}$ — so the alpha null is not a floor effect of an underpowered *test*, even if the underlying GWAS is underpowered. And the co-expression-aware gene-set null indicates that even the nominal alpha enrichment is a generic property of arbitrary gene sets in this territory ($p_{\text{geneset}} = 0.33$), not a weak-but-real alpha signal awaiting more power.

Second, **the shared genetic architecture of neighbouring EEG bands**. Band-limited EEG powers are strongly genetically correlated (Smit et al., 2018), and the alpha and theta bands in particular share generators and heritable variance. The comparable enrichment we observe for theta, beta and delta is therefore not surprising: to the extent that any oscillatory-power genetics is over-represented in this occipito-parieto-frontal territory, it reflects EEG-power genetics in general rather than an alpha-specific molecular signature. This reframes, rather than rescues, the hypothesis: the alpha-source network is not transcriptomically privileged for alpha genes specifically.

Third, and most consequential for the imaging-transcriptomics framing, **the spatial resolution of the functional target**. The alpha sources were localised from MEG with minimum-norm estimation, whose point-spread is on the order of centimetres; the “alpha-source” label attached to a given fine-grained parcel is therefore spatially uncertain, and the true generators are smeared across neighbouring parcels. This produces a resolution mismatch with *abagen*’s parcel-level expression that biases any localised contrast toward the null. We tested the corollary prediction directly — that coarsening the parcellation toward MEG resolution should recover a blurred signal — and found no such recovery (Schaefer-100 was, if anything, weaker than Glasser-360), which argues against a large effect merely hidden by resolution mismatch. We note that switching the inverse operator would not resolve this: all M/EEG source-reconstruction methods share centimetre-scale point-spread because the inverse problem is ill-posed, and beamformers in particular suppress coherent sources — precisely the phase-coupled parieto-frontal network at issue here — making them a poor remedy for this specific target.

Fourth, **the phenotype itself conflates periodic and aperiodic activity**. ENIGMA-EEG’s band-power phenotypes are computed as the mean spectral power within a fixed frequency window and do not separate the periodic (oscillatory) component from the aperiodic 1/f background — the offset and exponent that index cortical excitation/inhibition balance (Gao et al., 2017) and that FOOF/*specparam* parametrisation was developed to isolate (Donoghue et al., 2020). Because band-limited power is a mixture of both, a fraction of the “alpha power” association signal we carried into MAGMA may in fact index aperiodic, not oscillatory, genetic variation. This is not merely a hypothetical concern: the largest prior EEG GWAS found a genome-wide association between *GABRA2* and beta power (Hodgkinson et al., 2010) — GABAergic signalling is a principal determinant of the aperiodic exponent — consistent with band power phenotypes already carrying an aperiodic signature. If the true signal resides in the aperiodic component, then the band-power-based test we ran is not merely underpowered for alpha specifically but may be targeting a partially different phenotype than the one our (oscillatory) hypothesis concerns.

Strengths. The study is, to our knowledge, the first to link an EEG-trait GWAS to imaging transcriptomics and to the neurocircuitry of closed-loop stimulation, and it is deliberately built to resist the over-interpretation that a single positive test invites. The gene-based pipeline is validated

by recovery of the expected chr3p21 locus; the target regions are defined natively in the analysis parcellation, avoiding subjective mapping; the enrichment is corrected for spatial autocorrelation with a spin null; and specificity, replication and parcellation robustness are all examined and reported. The negative result is a genuine, reproducible contribution: it constrains a plausible and previously untested hypothesis and provides a template — and a cautionary calibration — for future, better-powered attempts.

5. Limitations

Beyond the three interpretive bounds above, several limitations apply. (i) The AHBA comprises six donors, only two with right-hemisphere tissue, and we used five (donor 15496 being unavailable at source); regional expression estimates therefore carry substantial uncertainty, especially for the right hemisphere and for inter-individual variability. (ii) A **modality gap** separates the functional target from its application: the alpha sources were localised from resting-state *MEG*, whereas closed-loop stimulation reads out *EEG*; MEG-defined generators are used here as a proxy for EEG–TMS targets, and the two modalities have different sensitivity profiles. (iii) The alpha-source set is a curated selection of 41 regions chosen to give broad coverage while avoiding proximity-induced spurious connectivity, not an exhaustive generator map; its large frontal contingent may partly reflect connectivity hubs rather than primary generators. (iv) Gene-set enrichment depends on gene-set definition; we mitigated this with a threshold-free continuous score, a top-N sensitivity, and a co-expression-aware gene-set null, but other definitions could differ. (v) Correspondence between regional gene expression and function is correlational and does not license mechanistic claims about TMS-induced plasticity. (vi) The analysis is restricted to cortex and to common variants tagged by the reference panel. (vii) The GWAS phenotypes are band-limited power, which conflates periodic and aperiodic spectral components (see Discussion); no GWAS of aperiodic parameters is currently available to test the two separately.

6. Future directions: an aperiodic-component GWAS

The periodic/aperiodic conflation above points to the most direct next step: repeating this pipeline with gene-based statistics derived from a GWAS of the **aperiodic exponent and offset** rather than band power. This alternative phenotype has three properties that make it, on priors, a better-motivated target for imaging transcriptomics than band power. It has a clearer molecular interpretation (excitation/inhibition balance, with GABAergic and glutamatergic receptor genes as natural candidates); resting EEG band power itself is highly heritable (Smit et al., 2005), which — given that band power partly reflects the aperiodic component (see above) — is at minimum consistent with a heritable aperiodic contribution, though the aperiodic component’s heritability has not, to our knowledge, been directly estimated by twin or family design; and, unlike band power, it follows a graded sensorimotor-to-association cortical topography that plausibly aligns with the dominant transcriptomic gradient of the AHBA — the same gradient our positive control (§2.6, §3.2) shows the pipeline can detect.

This analysis is not possible with existing secondary data: to our knowledge no GWAS of FOOOF/*specparam* aperiodic parameters with public summary statistics currently exists. ENIGMA-EEG is a distributed meta-analytic consortium — member cohorts retain their raw EEG and genotypes locally and contribute only summary statistics from an agreed pipeline — so there is no repository of raw EEG one can request; generating this GWAS would mean either (a) proposing an aperiodic-parameter sub-study to the ENIGMA-EEG working group, so that member cohorts run a shared FOOOF pipeline locally and contribute new summary statistics,

or (b) a smaller proof-of-concept in a single cohort with both resting EEG and genotypes under controlled access (e.g., COGA via dbGaP), underpowered relative to ENIGMA-EEG but sufficient for an initial test. Either route is a primary-data undertaking beyond the scope of the present secondary-data analysis. The pipeline described here is phenotype-agnostic at the point past MAGMA (Fig. 1) and would consume aperiodic gene-based statistics without modification.

7. Conclusion

Genes associated with EEG alpha power show no alpha-specific transcriptomic concentration in the cortical alpha-rhythm sources targeted by closed-loop EEG–TMS. Any tendency toward enrichment is weak, shared with the non-alpha bands we tested, and fragile to parcellation, and is best read as a property of EEG-oscillatory-power genetics broadly rather than of the alpha rhythm specifically. For closed-loop EEG–TMS, this implies that — at current GWAS power — the selection of a cortical alpha target is unlikely to be informed by the regional molecular architecture of alpha-associated genes; target selection should continue to rest on functional and connectivity criteria. Establishing whether a true alpha-specific signal exists will require a substantially larger EEG GWAS and functional targets defined at a spatial resolution commensurate with transcriptomic parcellations. It will also require separating the periodic and aperiodic components that the present band-power phenotypes conflate — a step that currently has no public GWAS to draw on, but that the pipeline described here is already positioned to consume once one exists (§6).

Data and code availability

Analysis code and derived results (gene-based statistics, region $\times$ gene expression summaries, enrichment tables, run logs) are available in the project repository. Raw data are not redistributed and must be obtained from their sources under the applicable terms: ENIGMA-EEG summary statistics (ENIGMA-EEG working group), the Allen Human Brain Atlas (via *abagen*), and the 1000 Genomes / MAGMA reference files (CTG lab). All parameters, random seeds and software versions are recorded in the repository run logs.

Author contributions

Conceptualisation, analysis, and writing were performed by the author with computational assistance. (To be completed.)

Funding

(To be completed.)

Conflicts of interest

The author declares no competing interests.

References

1. Alexander-Bloch, A. F., Shou, H., Liu, S., Satterthwaite, T. D., Glahn, D. C., Shinohara, R. T., Vandekar, S. N., & Raznahan, A. (2018). On testing for spatial correspondence between maps of human brain structure and function. *NeuroImage*, 178, 540–551.

2. Arnatkeviciute, A., Fulcher, B. D., & Fornito, A. (2019). A practical guide to linking brain-wide gene expression and neuroimaging data. *NeuroImage*, 189, 353–367.
3. Burt, J. B., Helmer, M., Shinn, M., Anticevic, A., & Murray, J. D. (2020). Generative modeling of brain maps with spatial autocorrelation. *NeuroImage*, 220, 117038.
4. de Leeuw, C. A., Mooij, J. M., Heskes, T., & Posthuma, D. (2015). MAGMA: Generalized gene-set analysis of GWAS data. *PLoS Computational Biology*, 11(4), e1004219.
5. Donoghue, T., Haller, M., Peterson, E. J., Varma, P., Sebastian, P., Gao, R., Noto, T., Lara, A. H., Wallis, J. D., Knight, R. T., Shestyuk, A., & Voytek, B. (2020). Parameterizing neural power spectra into periodic and aperiodic components. *Nature Neuroscience*, 23(12), 1655–1665.
6. Fornito, A., Arnatkeviciute, A., & Fulcher, B. D. (2019). Bridging the gap between connectome and transcriptome. *Trends in Cognitive Sciences*, 23(1), 34–50.
7. Fulcher, B. D., Arnatkeviciute, A., & Fornito, A. (2021). Overcoming false-positive gene-category enrichment in the analysis of spatially resolved transcriptomic brain atlas data. *Nature Communications*, 12(1), 2669. <https://doi.org/10.1038/s41467-021-22862-1>
8. Gao, R., Peterson, E. J., & Voytek, B. (2017). Inferring synaptic excitation/inhibition balance from field potentials. *NeuroImage*, 158, 70–78.
9. Glasser, M. F., Coalson, T. S., Robinson, E. C., Hacker, C. D., Harwell, J., Yacoub, E., Ugurbil, K., Andersson, J., Beckmann, C. F., Jenkinson, M., Smith, S. M., & Van Essen, D. C. (2016). A multi-modal parcellation of human cerebral cortex. *Nature*, 536(7615), 171–178.
10. Hawrylycz, M. J., Lein, E. S., Guillozet-Bongaarts, A. L., et al. (2012). An anatomically comprehensive atlas of the adult human brain transcriptome. *Nature*, 489(7416), 391–399.
11. Hodgkinson, C. A., Enoch, M.-A., Srivastava, V., Cummins-Oman, J. S., Ferrier, C., Iarikova, P., Sankararaman, S., Yamini, G., Yuan, Q., Zhou, Z., Albaugh, B., White, K. V., Shen, P.-H., & Goldman, D. (2010). Genome-wide association identifies candidate genes that influence the human electroencephalogram. *Proceedings of the National Academy of Sciences*, 107(19), 8695–8700.
12. Markello, R. D., Arnatkeviciute, A., Poline, J.-B., Fulcher, B. D., Fornito, A., & Misic, B. (2021). Standardizing workflows in imaging transcriptomics with the abagen toolbox. *eLife*, 10, e72129.
13. Markello, R. D., & Misic, B. (2021). Comparing spatial null models for brain maps. *NeuroImage*, 236, 118052.
14. Schaefer, A., Kong, R., Gordon, E. M., Laumann, T. O., Zuo, X.-N., Holmes, A. J., Eickhoff, S. B., & Yeo, B. T. T. (2018). Local-global parcellation of the human cerebral cortex from intrinsic functional connectivity MRI. *Cerebral Cortex*, 28(9), 3095–3114.
15. Smit, D. J. A., Posthuma, D., Boomsma, D. I., & de Geus, E. J. C. (2005). Heritability of background EEG across the power spectrum. *Psychophysiology*, 42(6), 691–697.
16. Smit, D. J. A., et al. (2018). Genome-wide association analysis links multiple psychiatric liability genes to oscillatory brain activity. *Human Brain Mapping*, 39(11), 4183–4195.
17. Tabarelli, D., Brancaccio, A., Zrenner, C., & Belardinelli, P. (2022). Functional connectivity states of alpha rhythm sources in the human cortex at rest: Implications for real-time brain state dependent EEG-TMS. *Brain Sciences*, 12(3), 348.
18. The 1000 Genomes Project Consortium. (2015). A global reference for human genetic variation. *Nature*, 526(7571), 68–74.
19. Yan, X., Kong, R., Xue, A., et al. (2023). Homotopic local-global parcellation of the human cerebral cortex from resting-state functional connectivity. *NeuroImage*, 273, 120010.
20. Zrenner, C., Desideri, D., Belardinelli, P., & Ziemann, U. (2018). Real-time EEG-defined excitability states determine efficacy of TMS-induced plasticity in human motor cortex. *Brain*

Stimulation, 11(2), 374–389.