

Towards Specifying Symbolic Computation^{*}

Jacques Carette and William M. Farmer

Computing and Software, McMaster University, Canada

<http://www.cas.mcmaster.ca/~carette>

<http://imps.mcmaster.ca/wmfarmer>

Abstract. ??

1 Introduction

2 Background

Let e be a mathematical expression and D be a domain of mathematical values. We say e is *defined in* D if e denotes an element in D . When e is defined in D , the *value of e in D* , written $\text{val}_D(e)$, is the element in D that e denotes. When e is undefined in D , the value of e in D and $\text{val}_D(e)$ are undefined. Two expressions e and e' are *equal in* D , written $e =_D e'$, if e and e' are defined in D and $\text{val}_D(e) = \text{val}_D(e')$ and are *quasi-equal in* D , written $e \simeq_D e'$, if either $e =_D e'$ or e and e' are both undefined in D .

3 Rational Expressions, Rational Functions

3.1 Rational Expressions

Let e be an expression in the language \mathcal{L} of the field $\mathbb{Q}(x)$, that is, a well-formed expression built from the symbols $x, 0, 1, +, *, -, ^{-1}$, elements of \mathbb{Q} and parentheses (as necessary). For greater readability, we will take the liberty of using fractional notation for $^{-1}$ and the exponential notation x^n for $x * \dots * x$ (n times). e can be something simple like $\frac{x^4-1}{x^2-1}$ or something more complicated like

$$\frac{\frac{1-x}{3/2x^{18}+x+17}}{\frac{1}{9834*x^{19393874}-1/5}} + 3 * x - \frac{12}{x}.$$

We assume that $\mathbb{Q} \subseteq \mathbb{Q}[x] \subseteq \mathbb{Q}(x)$ so that the field of rational numbers and the ring of polynomials in x are included in $\mathbb{Q}(x)$. The expressions in \mathcal{L} are intended to denote elements in $\mathbb{Q}(x)$. Of course, expressions like $x/0$ are undefined in $\mathbb{Q}(x)$. We will call members of \mathcal{L} *rational expressions (over \mathbb{Q})*.

We are taught that, like for members of \mathbb{Q} (such as $5/15$), there is a *normal form* for rational expressions. This is typically defined to be a rational expression

^{*} This research is supported by NSERC.

p/q for two polynomials $p, q \in \mathbb{Q}[x]$ such that p and q are themselves in polynomial normal form and $\gcd(p, q) = 1$. The motivation for the latter property is that we usually want to write $\frac{x^4-1}{x^2-1}$ as $x^2 + 1$ just as we usually want to write $5/15$ as $1/3$. Thus, the normal forms of $\frac{x^4-1}{x^2-1}$ and $\frac{x}{x}$ are $x^2 + 1$ and 1 , respectively. This definition of normal form is based on the characteristic that the elements of the *field of fractions* of a ring R can be written as quotients r/s of elements of R where $r_0/s_0 = r_1/s_1$ if and only if $r_0 * s_1 = r_1 * s_0$ in R .

Every computer algebra system implements a function that *normalizes* expressions that denote elements of $\mathbb{Q}(x)$ (including elements of \mathbb{Q} and $\mathbb{Q}[x]$). Let `normRatExpr` be the name of the algorithm that implements this normalization function on \mathcal{L} . Thus the signature of `normRatExpr` is $\mathcal{L} \rightarrow \mathcal{L}$ and the specification of `normRatExpr` is that, for all $e \in \mathcal{L}$, (A) `normRatExpr`(e) is a normal form and (B) $e \simeq_{\mathbb{Q}(x)} \text{normRatExpr}(e)$. `normRatExpr` is an example of an SBMA. (A) is the syntactic component of its specification, and (B) is the semantic component.

3.2 Rational Functions

Let \mathcal{L}' be the set of expressions of the form $\lambda x : \mathbb{Q} . e$ where $e \in \mathcal{L}$. We will call members of \mathcal{L}' *rational functions (over \mathbb{Q})*. That is, a rational function is a lambda expression whose body is a rational expression.

If $f_i = \lambda x : \mathbb{Q} . e_i$ are rational functions for $i = 1, 2$, one might think that $f_1 =_{\mathbb{Q} \rightarrow \mathbb{Q}} f_2$ if $e_1 =_{\mathbb{Q}(x)} e_2$. But this is not the case. For example, the rational functions $\lambda x : \mathbb{Q} . x/x$ and $\lambda x : \mathbb{Q} . 1$ are not equal since $\lambda x : \mathbb{Q} . x/x$ is undefined at 0 while $\lambda x : \mathbb{Q} . 1$ is defined everywhere. But $x/x =_{\mathbb{Q}(x)} 1$! Similarly, $\lambda x : \mathbb{Q} . (1/x - 1/x) \neq_{\mathbb{Q} \rightarrow \mathbb{Q}} \lambda x : \mathbb{Q} . 0$ and $(1/x - 1/x) =_{\mathbb{Q}(x)} 0$. Note that, in some contexts, we might want to say that $\lambda x : \mathbb{Q} . x/x$ and $\lambda x : \mathbb{Q} . 1$ do indeed denote the same function by invoking the concept of *removable singularities*.

I don't see why this reasoning is less clear as a justification that $\lambda x : \mathbb{Q} . (1/x - 1/x)$ and $\lambda x : \mathbb{Q} . 0$ are equal.

As we have just seen, we cannot normalize a rational function by normalizing its body, but we can normalize rational functions if we are careful not to remove points of undefinedness. Let a *quasinormal form* be a rational expression p/q for two polynomials $p, q \in \mathbb{Q}[x]$ such that p and q are themselves in polynomial normal form and there is no irreducible polynomial $r \in \mathbb{Q}[x]$ of degree ≥ 2 that divides both p and q . We can then normalize a rational function by quasinormalizing its body. Let `normRatFun` be the name of the algorithm that implements this normalization function on \mathcal{L}' . Thus the signature of `normRatFun` is $\mathcal{L}' \rightarrow \mathcal{L}'$ and the specification of `normRatFun` is that, for all $\lambda x : \mathbb{Q} . e \in \mathcal{L}'$, (A) `normRatFun`($\lambda x : \mathbb{Q} . e$) = $\lambda x : \mathbb{Q} . e'$ where e' is a quasinormal form and (B) $\lambda x : \mathbb{Q} . e \simeq_{\mathbb{Q} \rightarrow \mathbb{Q}} \text{normRatFun}(\lambda x : \mathbb{Q} . e)$. `normRatFun` is another example of an SBMA. (A) is the syntactic component of its specification, and (B) is the semantic component.

3.3 The Problem Here

So why are we concerned about rational expressions and rational functions? The reason is that computer algebra systems make little distinction between the

two: a rational expression can be interpreted sometimes as a rational expression and sometimes as a rational function. For example, one can always *evaluate* an expression by assigning values to its free variables or even convert it to a function. In Maple¹, these are done respectively via `eval(e, x = 0)` and `unapply(e, x)`. We can exhibit the problematic behaviour as follows: In fact, there is an even more pervasive, one could even say *obnoxious*, way of doing this: as the underlying language is *imperative*, it is possible to do:

insert some Maple code
with output here

```
e := (x^4-1)/(x^2-1);
# many, many more lines of 'code'
x := 1;
try to use 'e'
```

Hence, if an expression e is interpreted as a function, then it is not valid to simplify the function by applying `normRatExpr` to e , but computer algebra systems let the user do exactly this because usually there is no distinction made between e as a rational expression and e as representing a rational function, as we have already mentioned.

To avoid unsound applications of `normRatExpr`, `normRatFun`, and other SBMAs in mathematical systems, we need to carefully, if not formally, specify what these algorithms are intended to do. This is not a straightforward task to do in a traditional logic since SBMAs involve an interplay of syntax and semantics and algorithms like `normRatExpr` and `normRatFun` are very sensitive to definedness considerations. In the next subsection we will show how these two algorithms can be specified in a version of formal logic with undefinedness, quotation, and evaluation.

I don't know why we need
to say this: "Of course,
given some symbol y , $f(y)$
is in \mathcal{L} ."

3.4 The Formal Specification of `normRatExpr` and `normRatFun`

4 Related Work

5 Conclusion

¹ Mathematica has similar commands.

Todo list

■ I don't see why this reasoning is less clear as a justification that $\lambda x : \mathbb{Q} . (1/x - 1/x)$ and $\lambda x : \mathbb{Q} . 0$ are equal.	2
■ insert some Maple code with output here	3
■ I don't know why we need to say this: “Of course, given some symbol y , $f(y)$ is in \mathcal{L} .”	3