

Digital Calculus Frameworks and Comparative Evaluation of their Laplace-Beltrami operators

C. Weill–Duflos, D. Coeurjolly and J-O Lachaud

DGMM 2024

April 18, 2023

The Laplacian

In 1D :

$$\Delta f = \frac{d^2 f}{dx^2}$$

In 2D :

$$\Delta f = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2}$$

Etc...

Laplace Beltrami operator

More general definitions :

$$\Delta f = \operatorname{div}(\operatorname{grad}(f))$$

$$\Delta f = \delta df$$

Example : Sphere Laplacian

$$\Delta = \frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial}{\partial r} \right) + \frac{1}{r^2 \sin(\theta)} \frac{\partial}{\partial \theta} \left(\sin(\theta) \frac{\partial}{\partial \theta} \right) + \frac{1}{r^2 \sin^2(\theta)} \frac{\partial^2}{\partial \phi^2}$$

A Few Properties (why do we care about the Laplacian?)

Integration by parts

$$\forall \Phi, f, \int_{\Omega} \nabla f \cdot \nabla \Phi = - \int_{\Omega} \Delta f \Phi + \int_{\partial\Omega} \Phi \langle \nabla f, \boldsymbol{n} \rangle$$

A Few Properties (why do we care about the Laplacian?)

Integration by parts

$$\forall \Phi, f, \int_{\Omega} \nabla f \cdot \nabla \Phi = - \int_{\Omega} \Delta f \Phi + \int_{\partial\Omega} \Phi \langle \nabla f, \mathbf{n} \rangle$$

Eigen Vectors


In one dimension, real eigenvectors have the form : $\cos(\omega x + \varphi)$

In 2D : $\cos(\omega_x x + \varphi_x) \cos(\omega_y y + \varphi_y)$


Fourier basis!

Geometry Processing usages

- Filtering [Sorkine, O.: Laplacian mesh processing. (2005)]



- Geodesics [Crane, K., Weischedel, C., Wardetzky, M.: The heat method for distance computation. (2017)]



- Shape matching... [Lombaert, H., Grady, L., Polimeni, J.R, Cheriet, F.: FOCUSR: Feature Oriented Correspondence using Spectral Regularization–A Method for Precise Surface Matching(2013)]

Laplacian evaluations

Bunge, A., Botsch, M.: A survey on discrete Laplacians for general polygonal meshes. (2023)

Many different methods for polygonal mesh



Experimental ways to evaluate them : compare expected solutions of known problems, spherical eigenvectors, spherical eigenvalues...

- We adapt an existing method (the Finite Element Method) on surfels to build a Laplace-Beltrami operator
- We also provide a method using interpolated normals (detailed in the paper)
- We compare previous methods to ours, with experiments suggesting convergence
 - Heat Kernel [Caissard, T., Coeurjolly, D., Lachaud, J.O., Roussillon, T.: Laplace-Beltrami operator on digital surfaces. (2019)]
 - Corrected PolyDEC [Coeurjolly, D., Lachaud, J.O.: A simple discrete calculus for digital surfaces. (October 2022)]

Corrected FEM Laplacian

Digital Surface

Surfaces of voxels



Where is our data defined?

Our mesh is defined by a set of Vertices V and a set of faces F

It depends ! Usually :

- Scalar fields on vertices
- Vector fields on faces

Reason : vertices are where the values are “sampled”

On a triangle, using linear interpolation on the values at vertices the gradient is constant

Data representation

We have a vector holding the values at all the vertices

We like to have our operators be matrices, such that : This allows us to invert these operations (to solve a Poisson problem for example)

I want to have L a Laplace Beltrami operator such that, for any vector t holding some value at vertices (for example a temperature) Lt is its laplacian.

Data representation

$$L \cdot \begin{pmatrix} t(0) \\ t(1) \\ \vdots \\ t(|V| - 1) \end{pmatrix} = \begin{pmatrix} \Delta t(0) \\ \Delta t(1) \\ \vdots \\ \Delta t(|V| - 1) \end{pmatrix}$$

$$L \in R^{|V| \times |V|}$$

Finite Element Method


PDE approach, weak Poisson problem formulation :

$$\forall \Phi, \int_{\Omega} \nabla f \cdot \nabla \Phi = - \int_{\Omega} \Delta f \Phi + \int_{\partial\Omega} \Phi \langle \nabla f, \mathbf{n} \rangle$$



We want to solve $\Delta f = g$ for a given g (we assume no boundary) :

$$\forall \Phi, \int_{\Omega} \nabla f \cdot \nabla \Phi = - \int_{\Omega} g \Phi$$

Why corrected ?





How do we correct ?



$$\ell = \int_{\Omega} \langle \mathbf{u}_{\text{naive}} \mid \mathbf{u}_{\text{corrected}} \rangle \ell_{\text{cell,naive}} \rightarrow 2\pi$$

How do we correct ?




$$\ell = \int_{\Omega} \langle u_{\text{naive}} \mid u_{\text{corrected}} \rangle \ell_{\text{cell,naive}} \rightarrow 2\pi$$

We have convergent estimators of normals and curvature! [Lachaud, J.O., Coeurjolly, D., Levallois, J.: Robust and convergent curvature and normal estimators with digital integral invariants. (2017)]

How do we correct :

Metric G :

$$G = \begin{pmatrix} 1 - (u^x)^2 & -u^x u^y \\ -u^x u^y & 1 - (u^y)^2 \end{pmatrix}$$



Used in : $\langle v, w \rangle_G = v^t G w$

Penalizes dot product in a direction by how much the normal is aligned to that direction

How do we correct :

We assume an input vector field \mathbf{u} , constant over each face


$$\nabla f = \frac{1}{(\mathbf{u}^z)^2} \begin{pmatrix} \left(1 - (\mathbf{u}^y)^2\right) \frac{\partial f}{\partial s} + \mathbf{u}^x \mathbf{u}^y \frac{\partial f}{\partial t} \\ \left(1 - (\mathbf{u}^x)^2\right) \frac{\partial f}{\partial t} + \mathbf{u}^x \mathbf{u}^y \frac{\partial f}{\partial s} \end{pmatrix}$$

$$\Delta f = \frac{1}{(\mathbf{u}^z)^2} \left(\left(1 - (\mathbf{u}^y)^2\right) \frac{\partial^2 f}{\partial s^2} + 2\mathbf{u}^x \mathbf{u}^y \frac{\partial^2 f}{\partial s \partial t} + \left(1 - (\mathbf{u}^x)^2\right) \frac{\partial^2 f}{\partial t^2} \right)$$

$$\int_{\square} f = \int_{[0,1] \times [0,1]} f(s, t) \mathbf{u}^z \, ds \, dt$$


We work on each element separately first

Our element here is :



We work on each element separately first


Our element here is : **Basis functions**



- $\Phi_0 = (1 - s)(1 - t)$
- $\Phi_1 = s(1 - t)$
- $\Phi_2 = st$
- $\Phi_3 = (1 - s)t$

We work on each element separately first

Our element here is : **Basis functions**




- $\Phi_0 = (1 - s)(1 - t)$
- $\Phi_1 = s(1 - t)$
- $\Phi_2 = st$
- $\Phi_3 = (1 - s)t$

$$f = [f_0, f_1, f_2, f_2]$$

We work on each element separately first

Our element here is : **Basis functions**



- $\Phi_0 = (1 - s)(1 - t)$

- $\Phi_1 = s(1 - t)$

- $\Phi_2 = st$

- $\Phi_3 = (1 - s)t$

$$\mathbf{f} = [f_0, f_1, f_2, f_3]$$

$$f = f_0\Phi_0 + f_1\Phi_1 + f_2\Phi_2 + f_3\Phi_3$$

Find L_M and M_M such that $\forall f, p$ bilinear, $\int_{\square} \langle \nabla f, \nabla p \rangle = \mathbf{f}^T L_M \mathbf{p}$
and $\int_{\square} fp = \mathbf{f}^T M_M \mathbf{p}$

$$L_{Mi,j} = \int_{\square} \langle \nabla \Phi_i, \nabla \Phi_j \rangle$$

$$M_{Mi,j} = \int_{\square} \Phi_i \Phi_j$$

Without metric

$$L_M = \frac{1}{6} \begin{pmatrix} 4 & -1 & -2 & -1 \\ -1 & 4 & -1 & -2 \\ -2 & -1 & 4 & -1 \\ -1 & -2 & -1 & 4 \end{pmatrix}$$

$$M_M = \frac{1}{36} \begin{pmatrix} 4 & 2 & 1 & 2 \\ 2 & 4 & 2 & 1 \\ 1 & 2 & 4 & 2 \\ 2 & 1 & 2 & 4 \end{pmatrix}$$

With metric

$$L_M = \frac{1}{6u^z} \begin{pmatrix} 3u^x u^y + 2 + 2(u^z)^2 & 2(u^y)^2 - 1 - (u^x)^2 & -1 - 3u^x u^y - (u^z)^2 & 2(u^x)^2 - 1 - (u^y)^2 \\ 2(u^y)^2 - 1 - (u^x)^2 & -3u^x u^y + 2 + 2(u^z)^2 & 2(u^x)^2 - 1 - (u^y)^2 & -1 + 3u^x u^y - (u^z)^2 \\ -1 - 3u^x u^y - (u^z)^2 & 2(u^x)^2 - 1 - (u^y)^2 & 3u^x u^y + 2 + 2(u^z)^2 & 2(u^y)^2 - 1 - (u^x)^2 \\ 2(u^x)^2 - 1 - (u^y)^2 & -1 + 3u^x u^y - (u^z)^2 & 2(u^y)^2 - 1 - (u^x)^2 & -3u^x u^y + 2 + 2(u^z)^2 \end{pmatrix}$$

$$M_M = \frac{u^z}{36} \begin{pmatrix} 4 & 2 & 1 & 2 \\ 2 & 4 & 2 & 1 \\ 1 & 2 & 4 & 2 \\ 2 & 1 & 2 & 4 \end{pmatrix}$$

Back to global

We sum our local L_M and M_M into a big L and M and consider Φ and f global now

We define Φ_i as $[0, \dots, 0, 1, 0, \dots, 0]$, bilinear inside each face

Our problem

$$\forall \Phi, \int_{\Omega} \nabla f \cdot \nabla \Phi = - \int_{\Omega} g \Phi$$

Becomes : $\forall \Phi, \Phi^T L f = \Phi^T M g$

Back to global

We sum our local L_M and M_M into a big L and M and consider Φ and f global now

We define Φ_i as $[0, \dots, 0, 1, 0, \dots, 0]$, bilinear inside each face

Our problem

$$\forall \Phi, \int_{\Omega} \nabla f \cdot \nabla \Phi = - \int_{\Omega} g \Phi$$

Becomes : $\forall \Phi, \Phi^T L f = \Phi^T M g \Leftrightarrow L f = M g$

Poisson problem

$$\mathbf{f} = \mathbf{L}^{-1} \mathbf{M} \mathbf{g}$$

- \mathbf{L} is SPD, which is nice to invert
- Has a kernel of dimension 1, made of constant functions
- Problem can be fixed to a unique solution using boundary conditions

Forward evaluation

Why not use it the other way? Knowing f , find $g = \Delta f$

$$M^{-1} L f = g$$

Results

The laplacian has a known expression on the sphere as well as known eigenvalues

We can compare our results to the expected one

We average results over random discretization of the sphere
(random centers)

The laplacian has a known expression on the sphere as well as known eigenvalues


We can compare our results to the expected one

We average results over random discretization of the sphere (random centers)

As we decrease the gridstep h , error should decrease too (and eventually converge to 0)

Naive Forward evaluation

We try computing $g = M^{-1} L f$




$$f(x) = e^x$$


Fixed Forward evaluation

Apply some diffusion:


$g = M^{-1}L\mathbf{f}$ becomes $g = (M - dtL)^{-1}L\mathbf{f}$ with $dt = 0.035h$



$$f(x) = x^2$$




$$f(x) = e^x$$




$$f : \text{SH}(6, 6)$$


Backward evaluation



$$f(x) = x^2$$







$$f(x) = e^x$$



$$f : \text{SH}(6, 6)$$

Eigenvalues



Perspectives

- Proof of convergence
- Play with curvature : better metric? recover position from normal and mean curvature $\Delta p = -2Hn$
- Explore use for other geometries (triangle mesh with normal maps)