

Voronoi-based Geometry Estimator for 3D Digital Surfaces

Louis Cuel, Jacques-Olivier Lachaud, Boris Thibert

Université de Grenoble
Laboratoire Jean Kuntzman

Université de Savoie
LAMA


September 12th 2014

Differential quantities estimation

Normal




Curvature



Curvature Direction



Sharp Feature




Differential quantities estimation

Normal



Curvature




We want our method to be :

- Accurate
- Stable under noise
- Efficient
- Generic

Curvature Direction





Sharp Feature



Inference for different datas

Unorganized Point Clouds


Digital sets




Contributions

Our Approach :

- A method from computational geometry (VCM) :
 - Voronoi
 - Covariance
- Feed it with digital points
- Show multigrid convergence for geometric quantities




$$\int_{\Omega} (x - p)(x - p)^{\mathbf{t}} \, dx.$$




Contributions

Our Approach :

- A method from computational geometry (VCM) :
 - Voronoi
 - Covariance
- Feed it with digital points
- Show multigrid convergence for geometric quantities



$$\int_{\Omega} (x - p)(x - p)^{\mathbf{t}} \, dx.$$



Difficulties :

- Digitization of the VCM \Rightarrow Digitization error
- Achieve local (even ponctual) stability \Rightarrow Localization error
- Fixing parameters to achieve convergence


Plan

- 1) **Previous Voronoi-based Methods**
- 2) Voronoi Covariance Measure on Point cloud
- 3) Multigrid Convergence of the vcm estimators
- 4) Experiments


Voronoi-Based Methods

Voronoi-based algorithms :

- Amenta et Bern, 1999, Surface reconstruction by Voronoi filtering
- Cohen-Steiner et al., 2007, Voronoi-based Variational Reconstruction
- G. M. O., 2009, Robust Voronoi-based Curvature and Feature Estimation



1st previous Voronoi-based method




$$P = \{p_1, \dots, p_N\} \subseteq \mathbb{R}^d$$

Definition :

Voronoi cell: $\text{Vor}^P(q) = \{ \text{points whose closest point in } P \text{ is } q \}$

1st previous Voronoi-based method

Poles method



$$P = \{p_1, \dots, p_N\} \subseteq \mathbb{R}^d$$

Definition :


Voronoi cell: $\text{Vor}^P(q) = \{ \text{points whose closest point in } P \text{ is } q \}$

$\text{pole}_P(p) := \text{farthest point of } p \text{ in } \text{Vor}_P(p)$

Amenta, Bern, *Discrete and Computational Geometry* **22** (1999)

1st previous Voronoi-based method

Poles method



$$P = \{p_1, \dots, p_N\} \subseteq \mathbb{R}^d$$

Definition :


Voronoi cell: $\text{Vor}^P(q) = \{ \text{points whose closest point in } P \text{ is } q \}$

$\text{pole}_P(p) := \text{farthest point of } p \text{ in } \text{Vor}_P(p)$

Amenta, Bern, *Discrete and Computational Geometry* **22** (1999)

1st previous Voronoi-based method

Poles method



$$P = \{p_1, \dots, p_N\} \subseteq \mathbb{R}^d$$

Definition :

Voronoi cell: $\text{Vor}^P(q) = \{ \text{points whose closest point in } P \text{ is } q \}$

$\text{pole}_P(p) := \text{farthest point of } p \text{ in } \text{Vor}_P(p)$

Amenta, Bern, *Discrete and Computational Geometry* **22** (1999)


Idea : integrate to obtain stability.

2nd previous method : Voronoi Covariance

Alliez, Cohen-Steiner, Tong, Desbruns, Proc. Symposium Geometry Processing 2007

Covariance Matrix: $\text{cov}_p(\Omega) := \int_{\Omega} (x - p)(x - p)^{\mathbf{t}} \, \mathrm{d}x$.

Eigenvectors of $\text{cov}_p(\Omega)$ are the **principal axes** of Ω (viewed from p).




2nd previous method : Voronoi Covariance

Alliez, Cohen-Steiner, Tong, Desbruns, Proc. Symposium Geometry Processing 2007

Covariance Matrix: $\text{cov}_p(\Omega) := \int_{\Omega} (x - p)(x - p)^{\mathbf{t}} \, dx$.

Eigenvectors of $\text{cov}_p(\Omega)$ are the **principal axes** of Ω (viewed from p).

Algorithm:




- They take : $\Omega = \text{Vor}_P(p_i) \cap E$
- The normal is estimated by the eigenvector corresponding to the largest eigenvalue (in red).
- Stability to noise : Sum matrices over a neighborhood.

Plan


- 1) Previous Voronoi-based Methods
- 2) **Voronoi Covariance Measure**
- 3) Multigrid Convergence of the vcm estimators
- 4) Experiments

Voronoi covariance measure



Definition: Offset of P of radius R :
 $P^R = \cup_{p \in P} B(p, R)$.

Voronoi covariance measure




Definition: Offset of P of radius R :
 $P^R = \cup_{p \in P} B(p, R)$.

$$A(p) := \text{cov}_p(\text{Vor}_P(p_i) \cap P^R)$$

Definition: The *Voronoi covariance measure* of P of offset radius R is :

$$\mathcal{V}(P, R) := \sum_{i=1}^N A(p) \delta_p$$

Voronoi covariance measure



The VCM is defined for all compacts.

Definition: Offset of P of radius R :
 $P^R = \cup_{p \in P} B(p, R)$.


$$A(p) := \text{cov}_p(\text{Vor}_P(p_i) \cap P^R)$$

Definition: The *Voronoi covariance measure* of P of offset radius R is :

$$\mathcal{V}(P, R) := \sum_{i=1}^N A(p) \delta_p$$

$$\mathcal{V}(P, R) * \chi_r(p) := \sum_{p_i \in B(p, r)} A(p_i)$$

Voronoi covariance measure



The VCM is defined for all compacts.

Definition: Offset of P of radius R :
 $P^R = \cup_{p \in P} B(p, R)$.


$$A(p) := \text{cov}_p(\text{Vor}_P(p_i) \cap P^R)$$

Definition: The *Voronoi covariance measure* of P of offset radius R is :

$$\mathcal{V}(P, R) := \sum_{i=1}^N A(p) \delta_p$$

$$\mathcal{V}(P, R) * \chi_r(p) := \sum_{p_i \in B(p, r)} A(p_i)$$

Voronoi covariance measure



Definition: Offset of P of radius R :
 $P^R = \cup_{p \in P} B(p, R)$.

$$A(p) := \text{cov}_p(\text{Vor}_P(p_i) \cap P^R)$$

Definition: The *Voronoi covariance measure* of P of offset radius R is :

$$\mathcal{V}(P, R) := \sum_{i=1}^N A(p) \delta_p$$


$$\mathcal{V}(P, R) * \chi_r(p) := \sum_{p_i \in B(p, r)} A(p_i)$$

The VCM is defined for all compacts.

Theorem: Let P, K be two compacts and $p \in \mathbb{R}^d$.


$$\|\mathcal{V}(P, R) * \chi_r(p) - \mathcal{V}(K, R) * \chi_r(p)\| = O(d_H(K, P)^{\frac{1}{2}})$$

Digital Voronoi covariance measure




- X : an object
- $Dig_h(X)$: his Gauss digitization
- $Z_h = \partial_h X \cap h(\mathbb{Z} + \frac{1}{2})^3$

Digital Voronoi covariance measure



$\partial_h X$  Z_h 



- X : an object
- $Dig_h(X)$: his Gauss digitization
- $Z_h = \partial_h X \cap h(\mathbb{Z} + \frac{1}{2})^3$

We approach the VCM of the point cloud Z_h :

$$\Omega_h^R = \{x \in Z_h^R \cap h(\mathbb{Z} + \frac{1}{2})^3, vox(x) \subset Z_h^R\}$$


Definition: The VCM digital estimator

$$\hat{V}_{Z_h, R}(\chi) := \sum_{x \in \Omega_h^R} h^3 (x - p_{Z_h}(x))(x - p_{Z_h}(x))^t \chi(p_{Z_h}(x)),$$

Plan

- 1) Previous Voronoi-based Methods
- 2) Voronoi Covariance Measure
- 3) **Multigrid Convergence**
- 4) Experiments

Multigrid Convergence



Multigrid Convergence : $\|\hat{E}(h) - E\| \rightarrow 0$ when $h \rightarrow 0$

VCM and Normal Multigrid Convergence


Theorem: If ∂X is C^2 , $R < \frac{\rho}{2}$, $\text{diam}(\text{supp}(\chi)) < r$, $h \leq \min\left(R, \frac{r^2}{32\rho}\right)$

$$\begin{aligned} \|\mathcal{V}_{\partial X, R}(\chi) - \hat{\mathcal{V}}_{Z_h, R}(\chi)\|_{op} &= O\left(\text{Lip}(\chi) \times [(r^3 R^{\frac{5}{2}} + r^2 R^3 + r R^{\frac{9}{2}}) h^{\frac{1}{2}}] \right. \\ &\quad \left. + \|\chi\|_{\infty} \times [(r^3 R^{\frac{3}{2}} + r^2 R^2 + r R^{\frac{7}{2}}) h^{\frac{1}{2}} + r^2 R h] \right) \end{aligned}$$

Quantitative
and
localized
bound



Parameters
choice



VCM and Normal Multigrid Convergence

Theorem: If ∂X is C^2 , $R < \frac{\rho}{2}$, $\text{diam}(\text{supp}(\chi)) < r$, $h \leq \min\left(R, \frac{r^2}{32\rho}\right)$

$$\begin{aligned} \|\mathcal{V}_{\partial X, R}(\chi) - \hat{\mathcal{V}}_{Z_h, R}(\chi)\|_{op} &= O\left(\text{Lip}(\chi) \times [(r^3 R^{\frac{5}{2}} + r^2 R^3 + r R^{\frac{9}{2}}) h^{\frac{1}{2}}] \right. \\ &\quad \left. + \|\chi\|_{\infty} \times [(r^3 R^{\frac{3}{2}} + r^2 R^2 + r R^{\frac{7}{2}}) h^{\frac{1}{2}} + r^2 R h] \right) \end{aligned}$$

p_0 a point of ∂X

χ the hat function centered in p_0


with radius r

$\hat{n}_{r, R}(\hat{p}_0) = \text{first eigenvector of } \hat{\mathcal{V}}_{Z_h, R}(\chi)$

Corollary: normal estimator

If $r = R = ah^{\frac{1}{3}}$. Then

$$\langle \hat{n}_{r, R}(\hat{p}_0), n(p_0) \rangle = O\left(h^{\frac{1}{6}}\right).$$



Theorem, sketch of proof

$$\begin{aligned} & \|\mathcal{V}_{\partial X,R}(\chi) - \hat{\mathcal{V}}_{Z_h,R}(\chi)\|_{op} \leq \\ & \|\mathcal{V}_{\partial X,R}(\chi) - \mathcal{V}_{Z_h,R}(\chi)\|_{op} + \|\mathcal{V}_{Z_h,R}(\chi) - \hat{\mathcal{V}}_{Z_h,R}(\chi)\|_{op}. \end{aligned}$$

Localization error


Digitization error

Theorem, sketch of proof

$$\|\mathcal{V}_{\partial X, R}(\chi) - \hat{\mathcal{V}}_{Z_h, R}(\chi)\|_{op} \leq \|\mathcal{V}_{\partial X, R}(\chi) - \mathcal{V}_{Z_h, R}(\chi)\|_{op} + \|\mathcal{V}_{Z_h, R}(\chi) - \hat{\mathcal{V}}_{Z_h, R}(\chi)\|_{op}.$$

Localization error

Digitization error




Theorem, sketch of proof

$$\|\mathcal{V}_{\partial X, R}(\chi) - \hat{\mathcal{V}}_{Z_h, R}(\chi)\|_{op} \leq \|\mathcal{V}_{\partial X, R}(\chi) - \mathcal{V}_{Z_h, R}(\chi)\|_{op} + \|\mathcal{V}_{Z_h, R}(\chi) - \hat{\mathcal{V}}_{Z_h, R}(\chi)\|_{op}.$$


Localization error

Digitization error




Localization error
[Gui, Mé, Ovs, 09]

Digitization error



$\hat{\mathcal{V}}_{Z_h, R}(\chi)$







Plan


- 1) Previous Voronoi-based Methods
- 2) Voronoi Covariance Measure
- 3) Multigrid Convergence
- 4) **Experiments**

Experiments, Convergence Validation




Implemented on DGtal library :




Experiments, Normal Visualization



$$r = R = 3h^{0.5}$$




Experiments, Sharp Feature Estimation





$$r = R = 3h^{0.5}$$

Experiments, Sharp Feature Estimation



$$r = R = 3h^{0.5}$$



Conclusion

Summary:

We have several geometric estimators

- For normals, curvatures and sharp features :
- Theoretically stable
- Efficient
- Convergent and accurate in practice



Conclusion


Summary:

We have several geometric estimators

- For normals, curvatures and sharp features :
- Theoretically stable
- Efficient
- Convergent and accurate in practice

Perspectives:

- Distance to a measure + Voronoi Covariance
= δ -VCM robust to outliers



Even with outliers

