

# Math120 Spring 2016

## Final Review

Solution by Yajie Zhang

Study Notes:

| $p$ | $q$ | $\neg p$ | $p \wedge q$ | $p \vee q$ | $p \rightarrow q$ | $p \leftrightarrow q$ |
|-----|-----|----------|--------------|------------|-------------------|-----------------------|
| T   | T   | F        | T            | T          | T                 | T                     |
| F   | T   | T        | F            | T          | T                 | F                     |
| T   | F   |          | F            | T          | F                 | F                     |
| F   | F   |          | F            | F          | T                 | T                     |

1. Determine the truth values for each of the following statements.

(a) The sum of 4 and 6 is prime if and only if either 4 or 6 is prime.

T, since  $F \leftrightarrow (F \vee F) \equiv F \leftrightarrow F \equiv T$

(b) If 3 divides 20, then 6 divides 20.

T, since  $F \rightarrow F \equiv T$

2. Show that the following statements are tautology using truth tables.

(a)  $p \wedge (p \rightarrow q) \rightarrow q$

| $p$ | $q$ | $p \rightarrow q$ | $p \wedge (p \rightarrow q)$ | $p \wedge (p \rightarrow q) \rightarrow q$ |
|-----|-----|-------------------|------------------------------|--|
| T   | T   | T                 | T                            | T  |
| T   | F   | F                 | F                            | T  |
| F   | T   | T                 | F                            | T  |
| F   | F   | T                 | F                            | T  |

(b)  $((p \vee q) \wedge (\neg p \vee r)) \rightarrow (q \vee r)$

| $p$ | $q$ | $r$ | $\neg p$ | $p \vee q$ | $\neg p \vee r$ | $(p \vee q) \wedge (\neg p \vee r)$ | $q \vee r$ | $((p \vee q) \wedge (\neg p \vee r)) \rightarrow (q \vee r)$ |
|-----|-----|-----|----------|------------|-----------------|-------------------------------------|------------|--|
| T   | T   | T   | F        | T          | T               | T                                   | T          | T  |
| T   | T   | F   | F        | T          | F               | F                                   | T          | T  |
| T   | F   | T   | F        | T          | T               | T                                   | T          | T  |
| F   | T   | T   | T        | T          | T               | T                                   | T          | T  |
| T   | F   | F   | F        | T          | F               | F                                   | F          | T  |
| F   | T   | F   | T        | T          | T               | T                                   | T          | T  |
| F   | F   | T   | T        | F          | T               | F                                   | T          | T  |
| F   | F   | F   | T        | F          | F               | F                                   | F          | T  |

3. Show that the following statements are logically equivalent using truth tables.

(a)  $p \vee (p \wedge q) \equiv p$

| $p$ | $q$ | $p \wedge q$ | $p \vee (p \wedge q)$ |
|-----|-----|--------------|-----------------------|
| T   | T   | T            | T                     |
| T   | F   | F            | T                     |
| F   | T   | F            | F                     |
| F   | F   | F            | F                     |

(b)  $q \rightarrow (p \vee r) \equiv \neg p \rightarrow (q \rightarrow r)$

| $p$ | $q$ | $r$ | $p \vee r$ | $q \rightarrow (p \vee r)$ | $\neg p$ | $q \rightarrow r$ | $\neg p \rightarrow (q \rightarrow r)$ |
|-----|-----|-----|------------|----------------------------|----------|-------------------|--|
| T   | T   | T   | T          | T                          | F        | T                 | T                                      |
| T   | T   | F   | T          | T                          | F        | F                 | T                                      |
| T   | F   | T   | T          | T                          | F        | T                 | T                                      |
| F   | T   | T   | T          | T                          | T        | T                 | T                                      |
| T   | F   | F   | T          | T                          | F        | T                 | T                                      |
| F   | T   | F   | F          | F                          | T        | F                 | F                                      |
| F   | F   | T   | T          | T                          | T        | T                 | T                                      |
| F   | F   | F   | F          | T                          | T        | T                 | T                                      |

---

Study Notes:

$$p \wedge T \equiv p \vee F \equiv p \wedge p \equiv p \vee p \equiv \neg(\neg p) \equiv p$$

$$p \vee T \equiv p \vee \neg p \equiv T \quad \text{Commutative Law} \quad \begin{cases} p \wedge q \equiv q \wedge p \\ p \vee q \equiv q \vee p \end{cases}$$

$$p \wedge F \equiv p \wedge \neg p \equiv F \quad \text{Associative Law} \quad \begin{cases} (p \wedge q) \wedge r \equiv q \wedge (p \wedge r) \\ (p \vee q) \vee r \equiv q \vee (p \vee r) \end{cases}$$

$$p \rightarrow q \equiv \neg p \vee q \quad \text{Distributive Law} \quad \begin{cases} p \vee (q \wedge r) \equiv (p \vee q) \wedge (p \vee r) \\ p \wedge (q \vee r) \equiv (p \wedge q) \vee (p \wedge r) \end{cases}$$

$$p \leftrightarrow q \equiv (p \rightarrow q) \wedge (q \rightarrow p) \quad \text{De Morgan's Law} \quad \begin{cases} \neg(p \wedge q) \equiv \neg p \vee \neg q \\ \neg(p \vee q) \equiv \neg p \wedge \neg q \end{cases}$$

4. Re-do #2 and #3 using equivalent formulas.

(a)  $(p \wedge (p \rightarrow q)) \rightarrow q$

$$\equiv \neg(p \wedge (p \rightarrow q)) \vee q$$

$$\equiv \neg p \vee \neg(p \rightarrow q) \vee q$$

$$\equiv (\neg p \vee q) \vee \neg(p \rightarrow q)$$

$$\equiv (p \rightarrow q) \vee \neg(p \rightarrow q)$$

$$\equiv T$$

$$(b) ((p \vee q) \wedge (\neg p \vee r)) \rightarrow (q \vee r)$$

$$\begin{aligned} &\equiv \neg((p \vee q) \wedge (\neg p \vee r)) \vee (q \vee r) \\ &\equiv \neg(p \vee q) \vee \neg(\neg p \vee r) \vee q \vee r \\ &\equiv ((\neg p \wedge \neg q) \vee q) \vee ((p \wedge \neg r) \vee r) \\ &\equiv ((\neg p \vee q) \wedge (\neg q \vee q)) \vee ((p \vee r) \wedge (\neg r \vee r)) \\ &\equiv ((\neg p \vee q) \wedge T) \vee ((p \vee r) \wedge T) \\ &\equiv \neg p \vee q \vee p \vee r \\ &\equiv (\neg p \vee p) \vee (q \vee r) \\ &\equiv T \vee (q \vee r) \\ &\equiv T \end{aligned}$$

$$(c) p \vee (p \wedge q) \equiv p$$

$$(p \wedge T) \vee (p \wedge q) \equiv p \wedge (T \vee q) \equiv p \wedge T \equiv p$$

$$(d) q \rightarrow (p \vee r) \equiv \neg p \rightarrow (q \rightarrow r)$$

$$\begin{aligned} LHS &\equiv \neg q \vee p \vee r \\ RHS &\equiv \neg(\neg p) \vee (\neg q \vee r) \\ &\equiv p \vee \neg q \vee r \\ &\equiv \neg q \vee p \vee r \end{aligned}$$

5. For each of the following statements, determine its truth value and write a proof to support your conclusion. Assume the domain is the set of all real numbers.

$$(a) \forall x \exists y (\sqrt{x+y} = x) \quad \text{F}$$

Proof by negation:  $\exists x \forall y (\sqrt{x+y} \neq x)$

Let  $x = -1$ , then  $\forall y \sqrt{-1+y} \geq 0 > -1 \neq -1$

$$(b) \exists x \forall y (x^2 + 2y^2 \geq 10) \quad \text{T}$$

Proof:  $\exists x = \sqrt{10}, \forall y (y^2 \geq 0) \Rightarrow 2y^2 \geq 0 \Rightarrow (\sqrt{10})^2 + 2y^2 \geq 10$

$$(c) \exists x \exists y ((x < y) \rightarrow (x^3 > y^3)) \quad \text{T}$$

Proof: Let  $x = 1, y = 1$ , then  $F \rightarrow F \equiv T$

$$(d) \forall x \forall y ((x^y = y^x) \rightarrow (x = y)) \quad \text{F}$$

Proof by negation:  $\exists x \exists y (x^y = y^x \wedge x \neq y)$ , let  $x = 4, y = 2$ , then  $4^2 = 2^4 \wedge 4 \neq 2$

$$(e) \exists x \forall y \exists z \left( \frac{1}{x+y+1} = \frac{1}{y+z+2} \right) \quad \text{F}$$

Proof by negation:  $\forall x \exists y \forall z \left( \frac{1}{x+y+1} \neq \frac{1}{y+z+2} \right)$


Let  $y = -x - 1, \forall x \forall z : \frac{1}{x-x-1+1} = \frac{1}{0} = \text{undefined} \neq \frac{1}{y+z+2}$

$$(f) \quad \forall x \exists y \exists z \left( \sqrt{x-y} + x = \frac{1}{z} \right) \quad \text{T}$$


$$\text{Proof: } \forall x \begin{cases} x \neq 0, \text{ let } y = x, z = \frac{1}{x} \Rightarrow \sqrt{x-x} + x = \frac{1}{\frac{1}{x}} = x \\ x = 0, \text{ let } y = -1, z = 1 \Rightarrow \sqrt{0-(-1)} + 0 = \frac{1}{1} = 1 \end{cases}$$

6. Let  $A$ ,  $B$  and  $C$  be sets. For each of the following sets, draw a Venn diagram to represent the set.

(a)  $\overline{A} \cap (\overline{B} - C)$



(b)  $\overline{A - B} \cap (B \cup C)$



7. Suppose the universal set  $U = [0, 9]$ . Let  $A = (0, 4)$ ,  $B = [3, 9]$ , and  $C = \{1, 6\}$ . Find each of the following sets.

(a)  $\overline{A} \cup B = \{0\} \cup [3, 9]$   
 $\overline{A} = \{0\} \cup [4, 9]$

(b)  $A \cap (\overline{B} - C) = (0, 1) \cup (1, 3)$   
 $\overline{B} = [0, 3] \Rightarrow \overline{B} - C = [0, 1) \cup (1, 3)$

(c)  $(A \cup B) - C = (0, 1) \cup (1, 6) \cup (6, 9]$   
 $A \cup B = (0, 9]$

8. (a) Let function  $f : [-4, \infty) \rightarrow R$  be defined as  $f(x) = \sqrt{x+4}$ . Find each of the following sets.

i.  $f(\{0, 5\}) = \{2, 3\}$

ii.  $f((0, 5)) = (2, 3)$

iii.  $f([-4, 0]) = [0, 2]$

iv.  $f^{-1}([3, 5]) = [5, 21]$

$$\sqrt{x+4} = 3 \Rightarrow x+4 = 9 \Rightarrow x = 5$$

$$\sqrt{x+4} = 5 \Rightarrow x+4 = 25 \Rightarrow x = 21$$

$$\begin{aligned} \text{v. } f^{-1}((-4, 4)) &= [-4, 12] \\ \sqrt{x+4} = 0 &\Rightarrow x+4 = 0 \Rightarrow x = -4 \\ \sqrt{x+4} = 4 &\Rightarrow x+4 = 16 \Rightarrow x = 12 \\ \text{vi. } f^{-1}(\{-2, -1, 0, 1, 2\}) &= \{-4, -3, 0\} \\ \sqrt{x+4} = 0 &\Rightarrow x+4 = 0 \Rightarrow x = -4 \\ \sqrt{x+4} = 1 &\Rightarrow x+4 = 1 \Rightarrow x = -3 \\ \sqrt{x+4} = 2 &\Rightarrow x+4 = 4 \Rightarrow x = 0 \end{aligned}$$

(b) Let function  $f : \mathbb{R} \rightarrow \mathbb{R}$  be defined as  $f(x) = \left\lfloor \frac{3x+1}{2} \right\rfloor$ . Find each of the following sets.

$$\begin{aligned} \text{i. } f((2, 9)) &= \{3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13\} \\ \text{ii. } f([2, 9]) &= \{3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14\} \\ \text{iii. } f(\{0, 2, 4, 6, 8\}) &= \{0, 3, 6, 9, 12\} \\ \text{iv. } f^{-1}([3, 5]) &= \left[ \frac{5}{3}, \frac{11}{3} \right) \\ 3 \leq \frac{3x+1}{2} < 6 &\Rightarrow 6 \leq 3x+1 < 12 \Rightarrow \frac{5}{3} \leq x < \frac{11}{3} \\ \text{v. } f^{-1}((2, 3)) &= \emptyset \\ \text{vi. } f^{-1}((1.5, 3.2)) &= \left[ 1, \frac{7}{3} \right) \\ 2 \leq \frac{3x+1}{2} < 4 &\Rightarrow 4 \leq 3x+1 < 8 \Rightarrow 1 \leq x < \frac{7}{3} \end{aligned}$$

---

Study Notes:

$$f : A \rightarrow B$$

$$\text{1-1: } \forall a_1, a_2 \in A \quad (f(a_1) = f(a_2) \rightarrow a_1 = a_2)$$

$$\text{Not 1-1: } \exists a_1, a_2 \in A \quad (f(a_1) = f(a_2) \text{ but } a_1 \neq a_2)$$

$$\text{Onto: } \forall b \in B, \exists a \in A \quad (f(a) = b)$$

$$\text{Not Onto: } \exists b \in B, \forall a \in A \quad (f(a) \neq b)$$

9. (a) For each of the following function  $f : D \rightarrow \mathbb{R}$ , where  $D = \{x \in \mathbb{R} \mid f(x) \text{ is defined}\}$ , determine whether it's one-to-one and write a proof to support your conclusion.

$$\begin{aligned} \text{i. } f(x) &= \sqrt{1-2x} \quad \text{1-1} \\ \text{Proof: } \forall a_1, a_2 \in D, \text{ assume } f(a_1) &= f(a_2) \\ \Rightarrow \sqrt{1-2a_1} &= \sqrt{1-2a_2} \\ \Rightarrow 1-2a_1 &= 1-2a_2 \\ \Rightarrow -2a_1 &= -2a_2 \\ \Rightarrow a_1 &= a_2 \\ \text{Therefore, } f(x) &= \sqrt{1-2x} \text{ is 1-1} \end{aligned}$$

ii.  $f(x) = x^3 - x + 1$  Not 1-1

Proof: Let  $a_1 = 0, a_2 = 1$

Then  $\begin{cases} f(a_1) = 0^3 - 0 + 1 = 1 \\ f(a_2) = 1^3 - 1 + 1 = 1 \end{cases}$

Hence,  $f(a_1) = f(a_2)$ , but  $a_1 \neq a_2$

Therefore,  $f(x) = x^3 - x + 1$  is not 1-1

iii.  $f(x) = \frac{x+1}{x+2}$  1-1

Proof:  $\forall a_1, a_2 \in D$ , assume  $f(a_1) = f(a_2)$

$$\Rightarrow \frac{a_1 + 1}{a_1 + 2} = \frac{a_2 + 1}{a_2 + 2}$$

$$\Rightarrow (a_1 + 1)(a_2 + 2) = (a_2 + 1)(a_1 + 2)$$

$$\Rightarrow \cancel{a_1 a_2} + a_2 + 2a_1 + \cancel{2} = \cancel{a_1 a_2} + a_1 + 2a_2 + \cancel{2}$$

$$\Rightarrow 2a_1 - a_1 = 2a_2 - a_2$$

$$\Rightarrow a_1 = a_2$$

Therefore,  $f(x) = \frac{x+1}{x+2}$  is 1-1

(b) For each of the following function  $f : D \rightarrow \mathbb{R}$ , where  $D = \{x \in \mathbb{R} \mid f(x) \text{ is defined}\}$ , determine whether it's onto and write a proof to support your conclusion.

i.  $f(x) = \sqrt[3]{4x+3}$  Onto

Proof:  $\forall b \in \mathbb{R}$ , let  $a = \frac{1}{4}(b^3 - 3)$

$$\begin{aligned} \text{then } f(a) &= \sqrt[3]{4a+3} \\ &= \sqrt[3]{4 \cdot \frac{1}{4}(b^3 - 3) + 3} \\ &= \sqrt[3]{b^3 - 3 + 3} \\ &= b \end{aligned}$$

ii.  $f(x) = 4 - 3^x$  Not Onto

Proof: Let  $b = 5 \in \mathbb{R}$ , then  $\forall a \in D$

$$3^a \geq 0$$

$$\Rightarrow -3^a \leq 0$$

$$\Rightarrow 4 - 3^a \leq 4$$

$$\Rightarrow f(a) \leq 4$$

Since  $b = 5$ , then  $f(a) \neq b$

iii.  $f(x) = \frac{2x-3}{x}$  Not Onto

Proof: Let  $b = 2 \in \mathbb{R}$ , then  $\forall a \in D$

$$f(a) = \frac{2a-3}{a} = \frac{2a}{a} - \frac{3}{a} = 2 - \frac{3}{a}$$

Since  $\frac{3}{a} \neq 0$ , then  $2 - \frac{3}{a} \neq 2$

Hence,  $f(a) \neq b$

---

Study Notes:

$$\sum_{i=1}^n c = cn \quad \sum_{i=1}^n i = \frac{n(n+1)}{2} \quad \sum_{i=1}^n i^2 = \frac{n(n+1)(2n+1)}{6} \quad \sum_{i=1}^n i^3 = \left[ \frac{n(n+1)}{2} \right]^2$$

10. Evaluate each of the following sums, and write the answers without sigma notation.  
(No need to simplify)

(a)  $\sum_{n=1}^{60} (2n^3 + 3n^2 - 4n)$

$$\begin{aligned} &= 2 \sum_{n=1}^{60} n^3 + 3 \sum_{n=1}^{60} n^2 - 4 \sum_{n=1}^{60} n \\ &= 2 \cdot \left[ \frac{60 \cdot 61}{2} \right]^2 + 3 \cdot \frac{60 \cdot 61 \cdot (2 \cdot 60 + 1)}{6} - 4 \cdot \frac{60 \cdot 61}{2} \\ &= 6697800 + 221430 - 7320 \\ &= 6911910 \end{aligned}$$

(b)  $\sum_{k=100}^{150} (5k^3 - 1)$

$$\begin{aligned} &= \sum_{k=1}^{150} (5k^3 - 1) - \sum_{k=1}^{99} (5k^3 - 1) \\ &= 5 \sum_{k=1}^{150} k^3 - 150 - \left( 5 \sum_{k=1}^{99} k^3 - 99 \right) \\ &= 5 \cdot \left[ \frac{150 \cdot 151}{2} \right]^2 - 150 - 5 \cdot \left[ \frac{99 \cdot 100}{2} \right]^2 + 99 \\ &= 641278125 - 150 - 122512500 + 99 \\ &= 518765565 \end{aligned}$$

$$\begin{aligned}
(c) \quad & \sum_{i=1}^{2n} \sum_{j=1}^{n+1} (8ij - 1) \\
&= \sum_{i=1}^{2n} \left( 8i \sum_{j=1}^{n+1} j - (n+1) \right) \\
&= \sum_{i=1}^{2n} \left( 8i \cdot \frac{(n+1)(n+2)}{2} - (n+1) \right) \\
&= \sum_{i=1}^{2n} (4i \cdot (n+1)(n+2) - (n+1)) \\
&= 4(n+1)(n+2) \sum_{i=1}^{2n} i - 2n(n+1) \\
&= 4(n+1)(n+2) \frac{2n(2n+1)}{2} - 2n(n+1) \\
&= 4n(n+1)(n+2)(2n+1) - 2n(n+1)
\end{aligned}$$

$$\begin{aligned}
(d) \quad & \sum_{i=1}^n \sum_{j=1}^i (24j^2 + 12i) \\
&= \sum_{i=1}^n \left( 24 \sum_{j=1}^i j^2 + 12i^2 \right) \\
&= \sum_{i=1}^n \left( 24 \frac{i(i+1)(2i+1)}{6} + 12i^2 \right) \\
&= \sum_{i=1}^n (4i(i+1)(2i+1) + 12i^2) \\
&= \sum_{i=1}^n (4i(2i^2 + 2i + i + 1) + 12i^2) \\
&= \sum_{i=1}^n (8i^3 + 12i^2 + 4i + 12i^2) \\
&= \sum_{i=1}^n (8i^3 + 24i^2 + 4i) \\
&= 8 \sum_{i=1}^n i^3 + 24 \sum_{i=1}^n i^2 + 4 \sum_{i=1}^n i \\
&= 8 \left[ \frac{n(n+1)}{2} \right]^2 + 24 \frac{n(n+1)(2n+1)}{6} + 4 \frac{n(n+1)}{2} \\
&= 2n^2(n+1)^2 + 4n(n+1)(2n+1) + 2n(n+1)
\end{aligned}$$



11. Let  $A, B, C$  and  $D$  be zero-one matrices, where

$$A = \begin{bmatrix} 0 & 1 \\ 1 & 1 \\ 0 & 0 \\ 1 & 0 \end{bmatrix}, \quad B = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \end{bmatrix}, \quad C = \begin{bmatrix} 1 & 1 & 0 & 1 \\ 0 & 1 & 1 & 0 \\ 1 & 0 & 1 & 0 \end{bmatrix}, \quad D = \begin{bmatrix} 1 & 1 \\ 0 & 0 \\ 1 & 0 \\ 0 & 1 \end{bmatrix}$$

Find each of the following matrices.

$$(a) (A \vee D) \odot B = \begin{bmatrix} 1 & 1 \\ 1 & 1 \\ 1 & 0 \\ 1 & 1 \end{bmatrix} \odot B = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 0 \\ 1 & 1 & 1 \end{bmatrix}$$

$$(b) C \odot (A \wedge D) = C \odot \begin{bmatrix} 0 & 1 \\ 0 & 0 \\ 0 & 0 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 0 & 0 \\ 0 & 0 \\ 0 & 1 \end{bmatrix}$$

$$(c) B \odot C \odot D = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 0 \end{bmatrix} \odot D = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$$

$$(d) (A \odot B \odot C)^{[2]} = \left( \begin{bmatrix} 0 & 1 & 1 \\ 1 & 1 & 1 \\ 0 & 0 & 0 \\ 1 & 1 & 0 \end{bmatrix} \odot C \right)^{[2]} = \begin{bmatrix} 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 1 \end{bmatrix} \odot \begin{bmatrix} 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 1 \end{bmatrix}$$

12. Conversion.

(a) Convert  $(3AC.8)_{14}$  to base 10.

$$\begin{aligned} (3AC.8)_{14} &= 3 \times 14^2 + 10 \times 14^1 + 12 \times 14^0 + 8 \times 14^{-1} \\ &= 588 + 140 + 12 + 0.\overline{571428} \\ &= (740.\overline{571428})_{10} \end{aligned}$$

(b) Convert  $(782.75)_{10}$  to base 2.

$$\begin{array}{r} \begin{array}{r} \overline{391} \\ 2 \overline{)782} \\ \underline{600} \\ 182 \\ \underline{180} \\ 2 \\ \underline{2} \\ 0 \end{array} \quad \begin{array}{r} \overline{195} \\ 2 \overline{)391} \\ \underline{200} \\ 191 \\ \underline{180} \\ 11 \\ \underline{10} \\ 1 \end{array} \quad \begin{array}{r} \overline{97} \\ 2 \overline{)195} \\ \underline{180} \\ 15 \\ \underline{14} \\ 1 \end{array} \quad \begin{array}{r} \overline{48} \\ 2 \overline{)97} \\ \underline{80} \\ 17 \\ \underline{16} \\ 1 \end{array} \quad \begin{array}{r} \overline{24} \\ 2 \overline{)48} \\ \underline{40} \\ 8 \\ \underline{8} \\ 0 \end{array} \quad \begin{array}{r} \overline{12} \\ 2 \overline{)24} \\ \underline{20} \\ 4 \\ \underline{4} \\ 0 \end{array} \quad \begin{array}{r} \overline{6} \\ 2 \overline{)12} \\ \underline{12} \\ 0 \end{array} \quad \begin{array}{r} \overline{3} \\ 2 \overline{)6} \\ \underline{6} \\ 0 \end{array} \end{array}$$

$$\begin{array}{r} \times 0.75 \\ 2 \\ \hline 1.50 \end{array} \quad \begin{array}{r} \times 0.5 \\ 2 \\ \hline 1.0 \end{array}$$

$$(782.75)_{10} = (11\ 0000\ 1110.11)_2$$

(c) Convert  $(B93.A2)_{12}$  to base 6.

$$\begin{aligned} (B93.A2)_{12} &= 11 \times 12^2 + 9 \times 12^1 + 3 \times 12^0 + 10 \times 12^{-1} + 2 \times 12^{-2} \\ &= 1584 + 108 + 3 + 10/12 + 2/12^2 \\ &= 1695 + \frac{10 \cdot 12 + 2}{12^2} \\ &= 1695 + 0.847\bar{2} \\ &= (1695.847\bar{2})_{10} \end{aligned}$$

$$\begin{array}{r} 282 \\ 6 \overline{)1695} \\ \underline{1200} \\ 495 \\ \underline{480} \\ 15 \\ \underline{12} \\ 3 \end{array} \quad \begin{array}{r} 47 \\ 6 \overline{)282} \\ \underline{240} \\ 42 \\ \underline{42} \\ 0 \end{array} \quad \begin{array}{r} 7 \\ 6 \overline{)47} \\ \underline{42} \\ 5 \end{array} \quad \begin{array}{r} 1 \\ 6 \overline{)7} \\ \underline{6} \\ 1 \end{array} \quad \begin{array}{r} 0 \\ 6 \overline{)1} \end{array}$$

$$0.847\bar{2} \times 6 = 5.08\bar{3} \quad 0.08\bar{3} \times 6 = 0.5 \quad \begin{array}{r} \times 0.5 \\ 6 \\ \hline 3.0 \end{array}$$

$$(B93.A2)_{12} = (11503.503)_6$$

(d) Convert  $(11\ 1011\ 0101.1101)_2$  to base 8.

$$(11\ 1011\ 0101.1101)_2 = (1665.64)_8$$

(e) Convert  $(5CF.B4)_{16}$  to base 2.

$$(5CF.B4)_{16} = (101\ 1100\ 1111.1011\ 01)_2$$

(f) Convert  $(3673.52)_{16}$  to base 8.

$$(3673.52)_{16} = (11\ 0110\ 0111\ 0011.0101\ 001)_2 = (33163.244)_8$$

13. Perform indicated operations.

$$(a) (892.7B)_{13} + (7A.9)_{13} = (940.3B)_{13}$$

$$\begin{array}{r} \phantom{0}^{\frac{1}{8}} \phantom{0}^{\frac{1}{9}} \phantom{0}^{\frac{1}{2}} \phantom{0}^{\frac{7}{9}} \phantom{0}^{\frac{B}{9}} \\ + \phantom{0}^{\frac{7}{9}} \phantom{0}^{\frac{A}{9}} \phantom{0}^{\frac{9}{9}} \phantom{0}^{\frac{13}{9}} \\ \hline 9 \phantom{0}^{\frac{4}{9}} \phantom{0}^{\frac{0}{9}} \phantom{0}^{\frac{3}{9}} \phantom{0}^{\frac{B}{9}} \end{array} \quad \begin{array}{r} \phantom{0}^{\frac{1}{13}} \\ 13 \overline{)16} \\ \underline{13} \\ 3 \end{array} \quad \begin{array}{r} \phantom{0}^{\frac{1}{13}} \\ 13 \overline{)13} \\ \underline{13} \\ 0 \end{array} \quad \begin{array}{r} \phantom{0}^{\frac{1}{13}} \\ 13 \overline{)17} \\ \underline{13} \\ 4 \end{array}$$



$$\begin{aligned}
1 &= 10 - 9 \\
&= 10 - 19 + 10 \\
&= 2(10) - 19 \\
&= 2(48 - 2(19)) - 19 \\
&= 2(48) - 5(19) \\
&= 2(48) - 5(595 - 12(48)) \\
&= 62(48) - 5(595)
\end{aligned}$$

$$GCD(595, 48) = -5 \cdot 595 + 62 \cdot 48$$

(b)  $a = 50209, b = 1426$

|       |                    |    |   |   |   |  |
|-------|--------------------|----|---|---|---|--|
| 50209 |                    |    |   |   |   |  |
| 1426  |                    |    |   |   |   |  |
| 299   | = 50209 - 35(1426) |    |   |   |   |  |
| 230   | = 1426 - 4(299)    | 35 |   |   |   |  |
| 69    | = 299 - 230        | 4  | 1 | 3 |   |  |
| 23    | = 230 - 3(69)      | 1  | 4 | 3 | 3 |  |

|       |      |     |      |     |      |     |
|-------|------|-----|------|-----|------|-----|
| 50209 | 1426 | 299 | 1426 | 299 | 1426 | 299 |
| 42780 | 1196 | 230 | 230  | 69  | 207  | 23  |
| 7429  | 230  |     |      |     |      |     |
| 7130  |      |     |      |     |      |     |
| 299   |      |     |      |     |      |     |

$$\begin{aligned}
23 &= 230 - 3(69) \\
&= 230 - 3(299 - 230) \\
&= 4(230) - 3(299) \\
&= 4(1426 - 4(299)) - 3(299) \\
&= 4(1426) - 19(299) \\
&= 4(1426) - 19(50209 - 35(1426)) \\
&= 669(1426) - 19(50209)
\end{aligned}$$

$$GCD(50209, 1426) = -19 \cdot 50209 + 669 \cdot 1426$$

15. Find the least non-negative residue.

(a)  $x \equiv 22^{4999} \pmod{108}$

$$\begin{aligned}
22^1 &\equiv 22 \pmod{108} \\
22^2 &\equiv 484 \equiv 52 \pmod{108} \\
22^4 &\equiv 52^2 \equiv 4 \pmod{108} \\
22^8 &\equiv 4^2 \equiv 16 \pmod{108} \\
22^{16} &\equiv 16^2 \equiv 40 \pmod{108} \\
22^{32} &\equiv 40^2 \equiv 88 \pmod{108} \\
22^{64} &\equiv 88^2 \equiv 76 \pmod{108} \\
22^{128} &\equiv 76^2 \equiv 52 \pmod{108} \\
22^{256} &\equiv 4 \pmod{108}
\end{aligned}$$

$$22^{512} \equiv 16 \pmod{108}$$

$$22^{1024} \equiv 40 \pmod{108}$$

$$22^{2048} \equiv 88 \pmod{108}$$

$$22^{4096} \equiv 76 \pmod{108}$$

$$\begin{aligned} 4999 - 4096 &= 903 - 512 = 391 - 256 = 135 - 128 \\ &= 7 - 4 = 3 - 2 = 1 - 1 = 0 \end{aligned}$$

$$\begin{aligned} x &\equiv 22^{4999} \equiv 22^{4096} \cdot 22^{512} \cdot 22^{256} \cdot 22^{128} \cdot 22^4 \cdot 22^2 \cdot 22 \\ &\equiv 76 \cdot 16 \cdot 4 \cdot 52 \cdot 4 \cdot 52 \cdot 22 \equiv 1157398528 \equiv 4 \pmod{108} \end{aligned}$$

$$LNR = 4$$

$$(b) \ x \equiv 21^{2345} \pmod{57}$$

$$x \equiv (21^5)^{469} \equiv 51^{469} \pmod{57}$$

$$51^1 \equiv 51 \pmod{57}$$

$$51^2 \equiv 2601 \equiv 36 \pmod{57}$$

$$51^4 \equiv 36^2 \equiv 42 \pmod{57}$$

$$51^8 \equiv 42^2 \equiv 54 \pmod{57}$$

$$51^{16} \equiv 54^2 \equiv 9 \pmod{57}$$

$$51^{32} \equiv 9^2 \equiv 24 \pmod{57}$$

$$51^{64} \equiv 24^2 \equiv 6 \pmod{57}$$

$$51^{128} \equiv 6^2 \equiv 36 \pmod{57}$$

$$51^{256} \equiv 42 \pmod{57}$$

$$469 - 256 = 213 - 128 = 85 - 64 = 21 - 16 = 5 - 4 = 1 - 1 = 0$$

$$\begin{aligned} x &\equiv 51^{469} \equiv 51^{256} \cdot 51^{128} \cdot 51^{64} \cdot 51^{16} \cdot 51^4 \cdot 51 \equiv 42 \cdot 36 \cdot 6 \cdot 9 \cdot 42 \cdot 51 \\ &\equiv 174890016 \equiv 51 \pmod{57} \end{aligned}$$

$$LNR = 51$$

16. Solve for  $x$ .

$$(a) \ \begin{cases} x \equiv 1 \pmod{17} \\ x \equiv 5 \pmod{13} \\ x \equiv 2 \pmod{5} \end{cases}$$

$$x = 1 + 17t, t \in \mathbb{Z}$$

$$1 + 17t \equiv 5 \pmod{13} \Rightarrow 17t \equiv 5 - 1 \equiv 4 \pmod{13} \Rightarrow 4t \equiv 4 \pmod{13}$$

$$1 = 13 - 3(4)$$

$$4^{-1} \equiv -3 + 13 \equiv 10 \pmod{13}$$

$$t \equiv 40 \equiv 1 \pmod{13}$$

$$t = 1 + 13u, u \in \mathbb{Z}$$

$$x = 1 + 17(1 + 13u) = 1 + 17 + 221u = 18 + 221u$$

$$18 + 221u \equiv 2 \pmod{5} \Rightarrow 221u \equiv 2 - 18 \equiv -16 + 4(5) \equiv 4 \pmod{5}$$

$$\Rightarrow u \equiv 4 \pmod{5}$$

$$u = 4 + 5v, v \in \mathbb{Z}$$

$$x = 18 + 221(4 + 5v) = 18 + 884 + 1105v = 902 + 1105v, v \in \mathbb{Z}$$

$$(b) \begin{cases} x \equiv 3 \pmod{22} \\ x \equiv 7 \pmod{36} \\ x \equiv 1 \pmod{51} \end{cases}$$

$$x = 1 + 51t, t \in \mathbb{Z}$$

$$1 + 51t \equiv 7 \pmod{36} \Rightarrow 51t \equiv 6 \pmod{36} \Rightarrow 15t \equiv 6 \pmod{36}$$

$$5t \equiv 2 \pmod{12}$$

$$1 = 5(5) - 2(12)$$

$$5^{-1} \equiv 5 \pmod{12}$$

$$t \equiv 10 \pmod{12}$$

$$t = 10 + 12u, u \in \mathbb{Z}$$

$$x = 1 + 51(10 + 12u) = 511 + 612u$$

$$511 + 612u \equiv 3 \pmod{22} \Rightarrow 612u \equiv -508 + 24(22) \equiv 20 \pmod{22}$$

$$\Rightarrow 18u \equiv 20 \pmod{22}$$

$$9u \equiv 10 \pmod{11}$$

$$1 = 5(9) - 4(11)$$

$$9^{-1} \equiv 5 \pmod{11}$$

$$u \equiv 50 \equiv 6 \pmod{11}$$

$$u = 6 + 11v, v \in \mathbb{Z}$$

$$x = 511 + 612(6 + 11v) = 511 + 3672 + 6732v = 4183 + 6732v, v \in \mathbb{Z}$$

17. Prove each of the following statements.

---

Study Notes:

$$\text{Addition} \begin{cases} \text{Even} + \text{Even} = \text{Even} \\ \text{Odd} + \text{Odd} = \text{Even} \\ \text{Even} + \text{Odd} = \text{Odd} \end{cases}$$

$$\text{Subtraction} \begin{cases} \text{Even} - \text{Even} = \text{Even} \\ \text{Odd} - \text{Odd} = \text{Even} \\ \text{Even} - \text{Odd} = \text{Odd} \end{cases}$$

$$\text{Multiplication} \begin{cases} \text{Even} \times \text{Odd} = \text{Even} \\ \text{Even} \times \text{Even} = \text{Even} \\ \text{Odd} \times \text{Odd} = \text{Odd} \end{cases}$$

$$x^2 \text{ rule} \begin{cases} x^2 = \text{Even} \leftrightarrow x = \text{Even} \\ x^2 = \text{Odd} \leftrightarrow x = \text{Odd} \end{cases}$$

(a) Let  $x$  and  $y$  be integers. Show that  $3x + y$  is odd if and only if  $x^2 + 5y - 7$  is even.

Proof:  $3x + y \rightarrow x^2 + 5y - 7$

Assume  $3x + y$  is odd,

then exactly one of  $3x$  and  $y$  is odd, and the other one is even.

Case 1:  $3x$  is odd, and  $y$  is even

$3x$  is odd  $\Rightarrow$  both 3 and  $x$  are odd

$x$  is odd  $\Rightarrow x^2$  is odd

$y$  is even  $\Rightarrow 5y$  is even

Hence  $x^2 + 5y$  is odd

Since 7 is odd, then  $x^2 + 5y - 7$  is even

Case 2:  $3x$  is even, and  $y$  is odd

$3x$  is even  $\Rightarrow$  at least one of 3 and  $x$  is even

Since 3 is odd, then  $x$  must be even

$x$  is even  $\Rightarrow x^2$  is even

$y$  is odd  $\Rightarrow 5y$  is odd

Hence  $x^2 + 5y$  is odd

Since 7 is odd, then  $x^2 + 5y - 7$  is even

Proof:  $x^2 + 5y - 7 \rightarrow 3x + y$

Assume  $x^2 + 5y - 7$  is even,

then  $x^2$  and  $5y - 7$  are either both odd or both even.

Case 1:  $x^2$  and  $5y - 7$  are both odd

$x^2$  is odd  $\Rightarrow x$  is odd  $\Rightarrow 3x$  is odd

$5y - 7$  is odd, then exactly one of  $5y$  and 7 is odd, and the other is even

Since 7 is odd, then  $5y$  must be even

then at least one of 5 and  $y$  must be even

since 5 is odd,  $y$  must be even

since  $3x$  is odd,  $y$  is even, then  $3x + y$  must be odd

Case 2:  $x^2$  and  $5y - 7$  are both even

$x^2$  is even  $\Rightarrow x$  is even  $\Rightarrow 3x$  is even

$5y - 7$  is even, then  $5y$  and 7 are both odd or both even

Since 7 is odd, then  $5y$  must be odd

then both 5 and  $y$  must be odd

since  $3x$  is even,  $y$  is odd, then  $3x + y$  must be odd

(b) Let  $A, B$  and  $C$  be non-empty sets. Show that if  $A \times B = B \times C$ , then  $A = B = C$ .

Proof:  $A \times B = B \times C \rightarrow A = B = C$  ( $A \subseteq B \subseteq C$  and  $C \subseteq B \subseteq A$ )

Let  $x \in A$

Since  $B$  is non-empty, then  $\exists b \in B$

Hence,  $(x, b) \in A \times B$

Since  $A \times B = B \times C$ , then  $(x, b) \in B \times C$

Hence,  $x \in B$

Therefore,  $A \subseteq B$

Let  $x \in B$

Since  $A$  is non-empty, then  $\exists a \in A$

Hence,  $(a, x) \in A \times B$

Since  $A \times B = B \times C$ , then  $(a, x) \in B \times C$

Hence,  $x \in C$

Therefore,  $B \subseteq C$

Let  $x \in C$   
 Since  $B$  is non-empty, then  $\exists b \in B$   
 Hence,  $(b, x) \in B \times C$   
 Since  $A \times B = B \times C$ , then  $(b, x) \in A \times B$   
 Hence,  $x \in B$   
 Therefore,  $C \subseteq B$

Let  $x \in B$   
 Since  $C$  is non-empty, then  $\exists c \in C$   
 Hence,  $(x, c) \in B \times C$   
 Since  $A \times B = B \times C$ , then  $(x, c) \in A \times B$   
 Hence,  $x \in A$   
 Therefore,  $B \subseteq A$

- (c) Let  $f$  be a one-to-one function from  $A$  to  $B$  and let  $S$  and  $T$  be subsets of  $A$ . Show that if  $S$  and  $T$  are disjoint, then their images  $f(S)$  and  $f(T)$  must be disjoint.

Proof by contradiction:

Assume  $f(S)$  and  $f(T)$  are not disjoint  
 Then  $\exists b \in f(S) \cap f(T)$   
 Hence,  $b \in f(S)$  and  $b \in f(T)$   
 $b \in f(S) \Rightarrow \exists a_1 \in S$ , such that  $f(a_1) = b$   
 $b \in f(T) \Rightarrow \exists a_2 \in T$ , such that  $f(a_2) = b$   
 Hence,  $f(a_1) = f(a_2)$   
 Since  $f$  is 1-1, then  $a_1 = a_2$   
 Since  $a_1 \in S, a_2 \in T$  and  $a_1 = a_2$ , then  $a_1 = a_2 \in S \cap T$   
 Hence,  $S$  and  $T$  are not disjoint. (contradiction)

- (d) Show that if  $x$  is an odd integer, then  $8 \mid (x^2 - 1)$

Proof: Assume  $x$  is odd, then  $x = 2n + 1, n \in \mathbb{Z}$

$$\begin{aligned} x^2 - 1 &= (2n + 1)^2 - 1 \\ &= 4n^2 + 4n + 1 - 1 \\ &= 4n(n + 1) \end{aligned}$$

|  |  |
|--|--|
| Case 1: If $n$ is even,<br>then $n = 2k, k \in \mathbb{Z}$<br>Hence, $4n(n + 1)$<br>$= 4(2k)(n + 1)$<br>$= 8k(n + 1)$<br>Hence, $8 \mid x^2 - 1$ | Case 2: If $n$ is odd,<br>then $n + 1$ is even<br>Hence, $n + 1 = 2m, m \in \mathbb{Z}$<br>Hence, $4n(n + 1)$<br>$= 4n2m = 8nm$<br>Hence, $8 \mid x^2 - 1$ |
|--|--|

18. Prove or disprove each of the following statements.

- (a) If both  $1/x$  and  $1/(x + y)$  are rational, then  $y$  is also rational. T

Proof:  $\frac{1}{x}$  and  $\frac{1}{x + y}$  are rational



$$\Rightarrow \left\{ \begin{array}{l} \frac{1}{x} = \frac{a}{b} \\ \frac{1}{x+y} = \frac{c}{d} \end{array} \right., \text{ where } a, b, c, d \in \mathbb{Z}, \text{ and } b, d \neq 0$$

$$\frac{1}{x} = \frac{a}{b} \Rightarrow b = ax, \text{ since } b \neq 0, \text{ then } a \neq 0$$

$$\frac{1}{x+y} = \frac{c}{d} \Rightarrow d = c(x+y), \text{ since } d \neq 0, \text{ then } c \neq 0$$

$$\left. \begin{array}{l} b = ax \Rightarrow x = \frac{b}{a} \\ d = c(x+y) \Rightarrow x+y = \frac{d}{c} \end{array} \right\} \Rightarrow y = \frac{d}{c} - \frac{b}{a} = \frac{ad-bc}{ac}$$

Since  $a, b, c, d \in \mathbb{Z}$ , then  $(ad-bc), (ac) \in \mathbb{Z}$   
Since  $a \neq 0, c \neq 0$ , then  $ac \neq 0$   
Hence,  $y$  is rational.

- (b) Let  $A, B$  and  $C$  be sets. If  $A - C \subseteq B - C$ , then  $A \subseteq B$  F

Counter example:  $A = \{1\}, B = \{2\}, C = \{1, 2\}$

$$A - C = \emptyset, \text{ and } B - C = \emptyset$$

Hence,  $A - C \subseteq B - C$ , but  $A \not\subseteq B$

- (c) Let  $A$  and  $B$  be sets. Then  $\mathcal{P}(A) \cap \mathcal{P}(B) = \mathcal{P}(A \cap B)$  T

Proof:  $S \in \mathcal{P}(A) \cap \mathcal{P}(B)$

$$\Leftrightarrow S \in \mathcal{P}(A) \text{ and } S \in \mathcal{P}(B)$$

$$\Leftrightarrow S \subseteq A \text{ and } S \subseteq B$$

$$\Leftrightarrow S \subseteq A \cap B$$

$$\Leftrightarrow S \in \mathcal{P}(A \cap B)$$

Hence,  $\mathcal{P}(A) \cap \mathcal{P}(B) \subseteq \mathcal{P}(A \cap B)$  and  $\mathcal{P}(A \cap B) \subseteq \mathcal{P}(A) \cap \mathcal{P}(B)$

Therefore,  $\mathcal{P}(A) \cap \mathcal{P}(B) = \mathcal{P}(A \cap B)$

- (d) Let  $x$  and  $m$  be integers with  $m \geq 1$ . If  $x^2 \equiv 1 \pmod{m}$ , then either  $x \equiv 1 \pmod{m}$  or  $x \equiv -1 \pmod{m}$  F

Let  $x = 10, m = 99$

Then  $x^2 \equiv 100 \equiv 1 \pmod{99}$

But  $10 \not\equiv 1 \pmod{99}$  and  $10 \not\equiv -1 \pmod{99}$

19. Prove each of the following statements using mathematical inductions.

- (a) Show that  $\frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \dots + \frac{1}{2^n} = 1 - \frac{1}{2^n}$  for all integers  $n \geq 1$ .

Basic Step:  $n = 1$

$$LHS = \frac{1}{2} = 1 - \frac{1}{2} = \frac{1}{2} = RHS$$

Inductive Step: Assume  $n = k$

$$\frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \dots + \frac{1}{2^k} = 1 - \frac{1}{2^k}$$

Show  $n = k + 1$

$$\frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \dots + \frac{1}{2^{k+1}} = 1 - \frac{1}{2^{k+1}}$$

$$\begin{aligned}
LHS &= \frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \dots + \frac{1}{2^k} + \frac{1}{2^{k+1}} \\
&= 1 - \frac{1}{2^k} + \frac{1}{2^{k+1}} \\
&= 1 - \frac{2}{2^k \cdot 2} + \frac{1}{2^{k+1}} \\
&= 1 - \frac{1}{2^{k+1}} \\
&= RHS
\end{aligned}$$

(b) Show that  $89 \mid (5^{3n} - 6^{2n})$  for all integer  $n \geq 0$ .

Basic Step:  $n = 0$

$$89 \mid 5^0 - 6^0 = 0$$

Inductive Step: Assume  $n = k$

$$89 \mid 5^{3k} - 6^{2k}$$

Show  $n = k + 1$

$$89 \mid 5^{3(k+1)} - 6^{2(k+1)}$$

$$\begin{aligned}
5^{3(k+1)} - 6^{2(k+1)} &= 5^{3k+3} - 6^{2k+2} \\
&= 5^3 \cdot 5^{3k} - 6^2 \cdot 6^{2k} \\
&= 125 \cdot 5^{3k} - 36 \cdot 6^{2k} \\
&= (89 + 36) \cdot 5^{3k} - 36 \cdot 6^{2k} \\
&= 89 \cdot 5^{3k} + 36 \cdot 5^{3k} - 36 \cdot 6^{2k} \\
&= 89 \cdot 5^{3k} + 36(5^{3k} - 6^{2k})
\end{aligned}$$

Since  $89 \mid 89 \cdot 5^{3k}$  and  $89 \mid 36(5^{3k} - 6^{2k})$ , then  $89 \mid 5^{3(k+1)} - 6^{2(k+1)}$

Alternative Solution:

Basic Step:  $n = 0$

$$89 \mid 5^0 - 6^0 = 0$$

Inductive Step: Assume  $n = k$

$$89 \mid 5^{3k} - 6^{2k} \Rightarrow 5^{3k} - 6^{2k} = 89m, m \in \mathbb{Z} \Rightarrow 5^{3k} = 89m + 6^{2k}$$

Show  $n = k + 1$

$$89 \mid 5^{3(k+1)} - 6^{2(k+1)}$$

$$\begin{aligned}
5^{3(k+1)} - 6^{2(k+1)} &= 5^{3k+3} - 6^{2k+2} \\
&= 5^3 \cdot 5^{3k} - 6^2 \cdot 6^{2k} \\
&= 125(89m + 6^{2k}) - 36 \cdot 6^{2k} \\
&= 125 \cdot 89m + 125 \cdot 6^{2k} - 36 \cdot 6^{2k} \\
&= 125 \cdot 89m + 89 \cdot 6^{2k} \\
&= 89(125m + 6^{2k})
\end{aligned}$$

Hence,  $89 \mid 5^{3(k+1)} - 6^{2(k+1)}$

20. Consider all strings with 9 English letters.

(a) If repetitions are not allowed, how many strings contain the letter  $x$ ?

$$9 \times (25 \times 24 \times 23 \times 22 \times 21 \times 20 \times 19 \times 18)$$

(b) If repetitions are not allowed, how many strings contain both letter  $x$  and  $y$ ?

$$9 \times 8 \times (24 \times 23 \times 22 \times 21 \times 20 \times 19 \times 18)$$

(c) If repetitions are not allowed, how many strings contain both letter  $x$  and  $y$  in consecutive positions?

$$(2 \times 8) \times (24 \times 23 \times 22 \times 21 \times 20 \times 19 \times 18)$$

(d) If repetitions are allowed, how many strings contain the letter  $x$ ?

$$A = \{\text{Strings with } x\}$$

$$\overline{A} = \{\text{Strings with no } x\}$$

$$U = \{\text{All strings with 9 letters}\}$$

$$|U| = 26^9, \quad |\overline{A}| = 25^9$$

$$|A| = |U| - |\overline{A}| = 26^9 - 25^9$$

(e) If repetitions are allowed, how many strings contain either letter  $x$  or letter  $y$ ?

$$S = \{\text{All strings with either } x \text{ or } y\}$$

$$\overline{S} = \{\text{All strings with no } x \text{ and } y\}$$

$$U = \{\text{All strings with 9 letters}\}$$

$$|U| = 26^9, \quad |\overline{S}| = 24^9$$

$$|S| = |U| - |\overline{S}| = 26^9 - 24^9$$

(f) If repetitions are allowed, how many strings contain both letter  $x$  and  $y$ ?

$$A \cap B = \{\text{All strings with both } x \text{ and } y\}$$

$$A = \{\text{Strings with } x\}$$

$$B = \{\text{Strings with } y\}$$

$$A \cup B = \{\text{All strings with either } x \text{ or } y\}$$

$$|A \cup B| = |A| + |B| - |A \cap B|$$

$$\begin{aligned}
|A \cup B| &= 26^9 - 24^9, \quad |A| = 26^9 - 25^9, \quad |B| = 26^9 - 25^9 \\
|A \cap B| &= |A| + |B| - |A \cup B| = (26^9 - 25^9) + (26^9 - 25^9) - (26^9 - 24^9) \\
&= 26^9 - 25^9 + 26^9 - 25^9 - 26^9 + 24^9 \\
&= 26^9 - 2(25^9) + 24^9
\end{aligned}$$

21. Consider a set of seven digits  $\{0, 1, 3, 5, 6, 7, 9\}$ , and a four-digit number is to be constructed by using the numbers in the set without repetitions.

(a) How many different numbers can be formed?

$$7 \times 6 \times 5 \times 4 = 840$$

(b) How many such numbers are odd?

$$6 \times 5 \times 4 \times 5 = 600$$

(c) How many such numbers are greater than 5000?

$$4 \times 6 \times 5 \times 4 = 480$$

(d) How many such numbers are greater than 5300?

Case 1: 1st digit is 5, then 2nd digit must be  $\geq 3$ , and 5 cannot be used

$$1 \times 4 \times 5 \times 4 = 80$$

Case 2: 1st digit is  $> 5$

$$3 \times 6 \times 5 \times 4 = 360$$

$$\text{Total} = 80 + 360 = 440$$