

Mathematics-Module 2

1. In the figures below (not drawn to scale), right triangle $\triangle ABC$ is similar to $\triangle EFG$. What is the ratio of the area of $\triangle ABC$ to the area of $\triangle EFG$?





2. The function f is defined by a polynomial. The table below gives values for x and f(x). Which of the following must be a factor of f(x)?

х	f(x)
0	4
1	2
2	0

- A) x
- B) x 1
- C) x + 2
- D) x 2

3. In the system of equations below, a is a constant and x and y are variables. For which of the following values of a will the system have no solution?

$$x + 3y = -7$$
$$ax - 2y = 6$$

- A) $-\frac{2}{3}$
- B) $-\frac{3}{3}$
- C) 0
- D) $\frac{7}{3}$


4. What is the sum of all values of x that satisfy the following equation?

$$5x^2 - 30x - 32 = 3$$

- A) -1
- B) 0
- C) 6
- D) 7



5. In right triangle MNP below, angle M measures a° and $\cos a^{\circ} = \frac{3}{5}$. What is $\cos(90 - a^{\circ})$?



6. If 52% of 25 people polled in a coffee survey preferred Brand A and 48% preferred Brand B, which of the following inferences is a logical conclusion?

- A) At least 52% of shoppers will prefer Brand A over Brand B.
- B) The majority of shoppers will prefer Brand A over Brand B.
- C) The majority of shoppers will prefer Brand B over Brand A.
- D) The number of shoppers who prefer Brand A may be similar to the number of shoppers who prefer Brand B.

7. Ben bought a bag of assorted rubber balls and measured each one. Each had a diameter between 1.2 and 1.5 inches. What is a possible volume, rounded to the nearest cubic inch, of one of the balls? Use 3.14 for π .

8. How many gallon jugs of water should Julia buy to completely fill a container in the shape of a rectangular prism if the length is 8 feet, the width is 4 feet, and the height is 3 feet (1 gallon = 231 cubic inches)?

- A) 5
- B) 718
- C) 719
- D) 165,888

9. Subtract the polynomials: $(2x^2 + 3x + 2) - (x^2 + 2x - 3)$

- A) $x^2 + x + 5$
- B) $x^2 + x 1$
- C) $x^2 + 5x + 5$
- D) $x^2 + 5x 1$

WEEK3.2

10. The table below shows the breakdown of orders at a frozen yogurt shop. There were two flavors of yogurt (strawberry and lemon) and two sizes (small and large). If a person among those who ordered frozen yogurt is chosen at random, what is the probability that this person ordered a small serving of lemon yogurt?

	Strawberry	Lemon	Total
Small	12	9	21
Large	20	16	36
Total	32	25	57

- A) $\frac{9}{25}$
- B) $\frac{3}{7}$
- C) $\frac{7}{19}$
- D) $\frac{3}{19}$

11. If (x, y) is the solution to the following system of equations, what is the value of x + y?

$$3x + y = -2$$
$$y = -x$$

- A) -1
- B) 0
- C) 1
- D) 2

Refer to the following for questions 12 - 13:

Elise, Max, and Rylie are each offering lawn care services. Their prices for each service are listed in the table below. Prices are per square foot.

	Elise	Max	Rylie
Mowing	\$0.0025	\$0.002	\$0.003
Edging	\$0.0015	\$0.0015	\$0.002
Weeding	\$0.055	\$0.065	\$0.05

12. Rylie was asked to mow and edge a lawn as well as weed a 1,000 square foot garden. If she will spend \$12 on gas and other overhead costs, how large of a lawn does she need to be able to make a \$75 profit?

- A) 2,600 ft²
- B) $5,000 \text{ ft}^2$
- C) 6,875 ft²
- D) $7,400 \text{ ft}^2$

13. Ms. Lin wants to have her 7,500 square foot lawn mowed and edged. She also has a flower bed of x square feet to be weeded. Which of the following inequalities represents x if Elise's services are a better deal than Max's?


- A) x < 250
- B) x < 375
- C) x > 250
- D) x > 375

Mømetrix



14. A number of cash prizes were awarded for the top salespeople at ABC Communications, each in the amount of \$100, \$200, or \$400. If the prizes totaled \$1,600 and there were 7 total prizes awarded, what is the greatest possible number of \$100 prizes awarded?

15. The graph of the system of inequalities $y \le \frac{2}{3}x - \frac{3}{2}$ and $y \ge -2x - \frac{5}{2}$ has solutions in which quadrants on the xy-plane below?



- A) Quadrants I and IV
- B) Quadrants II and III
- C) Quadrants I, II, and IV
- D) Quadrants I, III, and IV

16. The equation below is used to solve for the distance, d (meters), that an object travels in a certain amount of time, t (seconds), with an initial velocity of v meters per second. Which of the following equations gives a in terms of d, v, and t?

$$d=vt+\frac{1}{2}at^2$$

- A) $\frac{2d-2vt}{t}$
- B) $\frac{2dvt}{t^2}$
- C) $\frac{2d}{t^2} \frac{2v}{t}$
- D) $\frac{2d}{t^2} 2vt$

17. The function f is defined by a polynomial. The table below gives values for x and f(x). Which of the following must be a factor of f(x)?

х	f(x)
-5	14
-3	0
-1	-6

- A) x + 3
- B) x 3
- C) x + 5
- D) x 5

Mømetrix



8. Simplify the following expression.

$$(3x+5)(4x-6)$$

- A) $12x^2 38x 30$
- B) $12x^2 + 2x 30$
- C) $12x^2 2x 1$
- D) $12x^2 + 2x + 30$

19. If (x, y) is the solution to the following system of equations, what is the value of x - y?

$$4x - y = 11$$
$$x = 3y$$

- A) 0
- B) 1
- C) 2
- D) 3

20. The equation below is used to calculate the period of a pendulum (i.e., the time it takes a pendulum to make one full oscillation). If T refers to the time in seconds, L is the length of the pendulum in meters, and g is the acceleration due to gravity in meters per second squared, which of the following equations gives L in terms of T, g, and π ?

$$T=2\pi\sqrt{rac{L}{g}}$$

- A) $(T 2\pi)^2 g$
- B) $g\sqrt{\frac{T}{2\pi}}$
- C) $\frac{T^2g}{4\pi^2}$
- D) $\frac{4\pi^2}{T^2g}$

21. If x > 0 and $x^2 - 4x = 12$, what is the value of x?

22. In the function below, b is a constant. If f(2) = 0, what is the value of f(-1)?

$$f(x)=\frac{3}{2}x+b$$

- A) $-\frac{9}{2}$
- C) $-\frac{3}{1}$
- D) $\frac{3}{2}$