

**PHILHEALTH CIRCULAR**No. 2026-0007**TO : ALL ACCREDITED HEALTH CARE PROVIDERS,  
PHILHEALTH MEMBERS, PHILHEALTH REGIONAL  
OFFICES AND ALL OTHERS CONCERNED****SUBJECT : Clinical and Pharmaceutical Standards for PhilHealth  
Yaman Ng Kalusugan Program (YAKAP)****I. RATIONALE**

The Universal Health Care Act (Republic Act No. 11223) mandates equitable access for all Filipinos to quality and cost-effective promotive, preventive, curative, rehabilitative, and palliative services delivered through a whole-of-system approach. It requires continuous quality improvement in health services and system performance, ensuring that care at all levels—including primary care—is safe, evidence-based, and appropriate. Section 37 of the National Health Insurance Act of 2013 (Republic Act No. 7875, as amended) further mandates PhilHealth to implement a comprehensive Quality Assurance Program to ensure that services provided to members, including prescriptions, are necessary, appropriate, and consistent with accepted medical standards and ethics.


PhilHealth expands its primary care through the Yaman ng Kalusugan Program (YAKAP), which integrates primary care services, the PhilHealth Guaranteed and Accessible Medications for Outpatient Treatment (PhilHealth GAMOT) benefit, and select cancer screening tests. Strengthening YAKAP through clear clinical standards supports effective implementation by promoting consistent, high-quality primary care, improved disease management, and timely referral across the health system.

**II. OBJECTIVES**

This PhilHealth Circular shall establish clinical and pharmaceutical standards to ensure that primary and outpatient care services under YAKAP are of good quality and lead to better health outcomes.

This PhilHealth Circular aims to:

- A. Provide a reference framework for reviewing and revising benefit payment designs, establishing accreditation requirements, and guiding other YAKAP-related issuances;
- B. Guide the payment and monitoring processes, and systems design; including the development of performance indicators and monitoring tools to ensure consistent and measurable provider performance; and



- C. Define the commitments, accountability, and expected standards of practice for all YAKAP providers to ensure consistent quality in healthcare delivery and adherence to PhilHealth policies.


### III. SCOPE

This PhilHealth Circular provides the clinical and pharmaceutical standards governing the delivery of primary and outpatient care services under the YAKAP. It outlines the required standards of care, prescribing and dispensing practices, and quality assurance mechanisms to ensure safe, effective, and coordinated primary care across the continuum of services. This PhilHealth Circular covers all stakeholders involved in the delivery of primary care services.


### IV. DEFINITION OF TERMS

For the purpose of this PhilHealth Circular, the following terms are defined:

- A. **Antimicrobial Stewardship** – a multidisciplinary, multi-intervention, and coordinated approach to improve the appropriate use of antimicrobials by promoting the selection of the optimal antimicrobial regimen to ensure the right drug, right administration, right time, and right duration to minimize harm to the patients.
- B. **Authorized Physician** – accredited physician with a PhilHealth GAMOT Application account.
- C. **Empanelment** – the establishment of a formal care relationship between beneficiaries and their chosen YAKAP Clinics after completion of the First Patient Encounter (FPE) and signing of the Mutual Care Agreement (MCA) or YAKAP Empanelment Slip (YES), thereby creating mutual accountability and ongoing clinical relationships. Once empaneled, the health care team assumes responsibility for coordinating comprehensive services for their assigned patients, ensuring continuity of care with support from the broader care team.
- D. **First Patient Encounter (FPE)** – the initial episode of patient contact for the year, whereby a primary care clinic takes and/or updates the basic health data of an eligible beneficiary to identify their health risks. The FPE is not yet a consultation.
- E. **Inappropriate Polypharmacy** – the condition when one or more medicines are prescribed that are not or no longer needed. This occurs when (a) there is no evidence based indication, the indication has expired or the dose is unnecessarily high; (b) one or more medicines fail to achieve the therapeutic objectives they are intended to achieve; (c) one, or the combination of several medicines cause ADRs, or put the patient at a high risk of ADRs or because (d) the patient is not willing or able to take one or more medicines as intended.



- F. Medical Prepayment Review** – the process of reviewing and evaluating clinical data before claims payment to determine compliance to PhilHealth policies and widely accepted medical practice.
- G. Patient Navigation** – the function of a primary care provider as the first point of contact for patients to coordinate and direct the individuals to the appropriate health care provider of a particular level of care, to obtain health services needed to manage a wide range of health needs.
- H. Pharmaceutical Care** – a responsible provision of drug therapy for the purpose of achieving definite outcomes that improve the patient's quality of life.
- I. Polypharmacy** – often defined as the routine use of five or more medications at the same time.
- J. Primary Care** – the initial-contact, accessible, continuous, comprehensive, and coordinated care that is accessible at the time of need, including a range of services for all presenting conditions, and the ability to coordinate referral to other health care providers in the health care delivery system, when necessary.
- K. Primary Care Providers (YAKAP Providers)** – licensed and accredited health care institutions or professionals accredited by PhilHealth to deliver outpatient primary care services, including consultation, lab tests, and medicines. They could be any of the following:
- 1. GAMOT Facilities** – PhilHealth-engaged drug dispensing facilities responsible for providing Guaranteed and Accessible Medications for Outpatient Treatment benefit to eligible members.
  - 2. Primary Care Clinics** – PhilHealth-accredited public or private healthcare providers offering primary care services, in accordance with PhilHealth Circular No. 2024-0013 and its succeeding revisions. They are currently referred to as YAKAP Clinics.
  - 3. Primary Care Physicians (YAKAP Physicians)** – PhilHealth-accredited physicians under the YAKAP Program who are responsible for the delivery of medical services in accordance with established clinical, prescribing, and quality assurance standards.
  - 4. Cancer Screening Facilities** – health facilities that are duly licensed by the Department of Health and accredited by PhilHealth as providers of PhilHealth benefit packages for selected outpatient cancer screening tests.
- L. Rational Drug Use** – the principle that patients shall receive medications appropriate to their clinical needs, in doses that meet their individual requirements, for an adequate period of time, and at the lowest cost to them and their community.




- M. Therapeutic Duplication** – the practice of prescribing multiple medications for the same indication or purpose without a clear distinction of when one agent should be administered over another.

## **V. POLICY STATEMENTS**

### **A. Core Functions of Primary Care Providers**


YAKAP providers shall perform the four core functions of primary care. They shall serve as the first point of contact and provide comprehensive, accessible, and continuous care to PhilHealth beneficiaries.

1. YAKAP Clinics shall deliver, at the minimum, the following services:
  - a. First Patient Encounter (FPE);
  - b. Empanelment;
  - c. Initial assessment, collection of basic health data, and risk profiling;
  - d. Primary care consultations, laboratory tests and medicines;
  - e. Referral to higher level facilities and other primary care services such as tuberculosis treatment, maternity care, oral health, cancer screening among others; and
  - f. Creation and maintenance of patient/client health records.
2. As first-contact providers, YAKAP Clinics shall serve as the initial and continuous access point for clients seeking primary care services. They shall also function as patient navigators, ensuring that PhilHealth beneficiaries are referred to appropriate services based on their healthcare needs.
3. The YAKAP Clinics shall provide comprehensive primary care including promotive, preventive, and curative services that address the health needs of their catchment population. They must be capable of managing the most common health conditions across all life stages.
4. As providers of coordinated care, YAKAP Clinics shall ensure that PhilHealth beneficiaries receive integrated, person-centered care across different settings and levels of the health system.
  - a. YAKAP Clinics shall establish and maintain a care coordination, patient navigation, and referral system with GAMOT providers, cancer screening providers, and other health care providers within their catchment area to facilitate timely access to appropriate services and continuity of care.
  - b. Patient navigation and referrals for cancer screening tests shall be guided by the beneficiaries' health risk profiles, consistent to the risk criteria established by PhilHealth through PhilHealth Circular Nos. 2025-0024 (Quality Standards on the Screening for Breast Cancer as Reference of the Corporation), 2025-0025 (Quality Standards on the Screening for Colorectal Cancer as Reference of the Corporation), 2025-0026 (Quality Standards on the Screening for Liver Cancer as Reference of the Corporation), 2025-0027 (Quality




Standards on the Screening for Lung Cancer as Reference of the Corporation), and future issuances.

5. YAKAP Clinics shall ensure continuity of care through a series of coherent and relevant patient encounters, consultations, and follow-ups. These interactions must be directed toward achieving sustained improvements in the client's health and well-being.
  - a. YAKAP Clinics shall have follow-up consultations with their clients based on the risk, diagnosis, and clinical practice guidelines.
  - b. Cancer screening centers, GAMOT facilities, and other referral facilities (such as maternity care clinics, TB-DOTS centers among others) shall refer back their clients to YAKAP Clinics. These shall be documented through a feedback report or its equivalent and shall be integrated into the patients' primary care records.
- B. YAKAP Clinics shall maintain and update clients' records containing complete history, diagnoses, medications, laboratory and imaging results, medications, referrals, and follow-up plans. All members of the health care team shall document each client encounter accurately and promptly. Clinical diagnoses shall be properly translated into corresponding ICD-10 code/s. In the future, YAKAP Clinics shall transition to a standardized electronic medical records (EMR) to streamline documentation, and ensure effective coordination and continuity of care.
- C. YAKAP physicians shall adhere to the current clinical practice guidelines, protocols, and standards of care including but not limited to the Omnibus Health Guidelines, National Antibiotic Guidelines, Philippine Package of Essential Non-communicable Disease Interventions, and PhilHealth-issued quality policies. When necessary, PhilHealth shall issue guidelines for specific conditions to ensure uniform implementation of this PhilHealth Circular.
- D. Medicines under PhilHealth GAMOT facilities shall be dispensed only by duly registered and licensed pharmacists, and upon presentation of a valid prescription issued by YAKAP physicians.
- E. Prescribing and Dispensing Standards and Rational Drug Use  
YAKAP providers shall uphold to the principles of rational drug use and antimicrobial stewardship.
  1. Primary care physicians shall prescribe medications in accordance with all applicable laws and regulations, including the Generics Act of 1988 (Republic Act No. 6675) and the Comprehensive Dangerous Drugs Act of 2002 (Republic Act No. 9165). They shall align the prescriptions on the patient's needs and clinical condition, the drugs' approved therapeutic indications (Annex A: Table of GAMOT Drugs and Indications).
  2. Primary care physicians shall ensure that patients receive medications only when clinically indicated, in appropriate doses, for an adequate



duration, at the lowest possible cost, and administered correctly by the appropriate person.

3. In support of antimicrobial stewardship through rational antimicrobial use and adherence to evidence-based clinical guidelines, prescribing limits for oral antimicrobials shall NOT exceed fourteen (14) days per prescription. Cases requiring extended therapy shall only be undertaken following proper clinical assessment and evaluation by authorized primary care physicians. The use and duration of other antimicrobial agents shall be in accordance with applicable clinical guidelines.
4. Medications prescribed and claimed under PhilHealth GAMOT must be intended solely for the patient's active or current medical condition. Prescriptions issued upon request or designated as standby medications for non-active conditions, and/or those not related to the patient's current diagnosis, shall not be eligible for claims under the GAMOT Program.
5. Primary care physicians shall prescribe therapeutically sound and cost-effective medicines consistent with clinical guidelines and the Philippine National Formulary.
6. Primary care physicians shall evaluate potential drug-drug interactions when prescribing multiple medications, especially in cases of polypharmacy, to ensure patient safety, prevent adverse drug events, and avoid inappropriate polypharmacy.
7. GAMOT facilities and its pharmacists, as essential providers of pharmaceutical care, shall rationally dispense drugs ensuring the safe, effective, and appropriate provision of medicines and health services (including patient counseling, if necessary) tailored to the patients' needs.
8. GAMOT facilities through its pharmacists shall ensure the following:
  - a. The "Five Rights" of dispensing and medication administration by verifying the "Right Patient, Right Drug, Right Dose, Right Route, and Right Time" before dispensing any medication.
  - b. The completeness and compliance of GAMOT prescriptions and dispensing practices with PhilHealth Circular No. 2025-0013 - Nationwide Implementation of PhilHealth Guaranteed and Accessible Medications for Outpatient Treatment (PhilHealth GAMOT) specifically Section V.C.4.c. on prescription requirements and Section V.5 on dispensing guidelines.
9. PhilHealth shall define the limits in the quantity of dispensed medicines under PhilHealth GAMOT (Annex B – PhilHealth GAMOT Dispensing Limits) to ensure that patients have regular contact with pharmacists and prescribers, which allows for monitoring of treatment effectiveness, managing side effects, and re-evaluating the need for the medication.




10. For medication exceeding the dispensing limit, the patient shall be instructed to follow up with their primary care physicians to determine whether to discontinue or extend the treatment. Extension of treatment should comply with current clinical practice guidelines and standards of care.
  11. Any YAKAP clinic or GAMOT facility who prepares the prescription or submits a GAMOT claim exceeding what is medically warranted, or engages in inappropriate polypharmacy by prescribing non-essential or unnecessary medications for the purpose of taking advantage of the benefit, shall be placed under appropriate monitoring processes and be subjected to actions deemed necessary in accordance to prevailing policies, guidelines and regulatory directives.
- F. Drug management by YAKAP clinics or GAMOT facilities is consistent with regulatory and safety requirements in the following areas: drug selection and procurement, prescription, dispensing, and drug disposal.
- G. PhilHealth shall use the quality standards and outcomes of care as the basis in setting the performance-based payment and for monitoring the quality of primary care services.
- H. When necessary, PhilHealth may implement oversight mechanisms—such as medical pre-payment review, medical post-audit and pre-authorization requirements—to ensure that YAKAP benefits are provided appropriately, consistently, and in accordance with established clinical and pharmaceutical standards. It shall likewise employ payment recovery mechanisms to affected claims in accordance with existing policies.
- I. Monitoring and Evaluation
1. PhilHealth shall set criteria and indicators (Annex C: Quality Indicators for YAKAP) and develop tools to measure the performance of providers under the Health Care Provider Performance Assessment System (HCPPAS) in accordance with this PhilHealth Circular.
  2. These standards shall be regularly reviewed and updated as necessary. Any updates, as a result of the review, shall be disseminated in another PhilHealth Circular.
- J. PhilHealth shall set operational guidelines and procedures, information dissemination, and monitoring activities to enforce this PhilHealth Circular.

K. List of Annexes

The following annexes shall be posted in PhilHealth website.

1. Annex A: Table of GAMOT Drugs and Indications
2. Annex B: PhilHealth GAMOT Dispensing Limits
3. Annex C: Quality Indicators for YAKAP



## VI. PENALTY CLAUSE

Any violation of this PhilHealth Circular shall be dealt with and penalized in accordance with pertinent provisions of Republic Act No. 11223 (Universal Health Care Act), R.A. No. 7875 as amended by R.A. Nos. 9241 and 10606 (National Health Insurance Act of 2013), and R.A. No. 10918 (Philippine Pharmacy Act) and their respective Implementing Rules and Regulations.

## VII. SEPARABILITY CLAUSE

In the event that any part or provision of this PhilHealth Circular is declared unauthorized or invalid by any Court of Law or competent authority, those provisions not affected by such declaration shall remain valid and effective.


## VIII. REPEALING CLAUSE

1. Section V.E.9 and Annex B of this PhilHealth Circular amend Section V.C.4.f of PhilHealth Circular 2025-0013 - Nationwide Implementation of Guaranteed and Accessible Medication for Outpatient Treatment (PhilHealth GAMOT).
2. Annex A of this PhilHealth Circular amends Annex A.1 of PhilHealth Circular 2025-0013.

All other PhilHealth Circulars, issuances, rules, and regulations, or parts thereof, that are inconsistent with this PhilHealth Circular are hereby repealed, amended, or modified accordingly.

## IX. DATE OF EFFECTIVITY

This PhilHealth Circular shall take effect fifteen (15) days after its publication in the Official Gazette or in any newspaper of general circulation. Three (3) certified copies shall be deposited in the Office of the National Administrative Register (ONAR) at the University of the Philippines Law Center.

  
**EDWIN M. MERCADO, MD, MHA, MMSc**  
President and Chief Executive Officer

Date signed: 06/03/2024

DC:  Date: 6/4/26  
Clinical and Pharmaceutical Standards for PhilHealth Yaman Ng Kalusugan Program  
(YAKAP)




## Annex A: Table of GAMOT Drugs and Indications

Table of GAMOT Drugs and Indications

| Drug           | Indications                                                                                                                                                                                                                                                                                                                                                 |
|----------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Albendazole    | Management of filariasis; treatment of intestinal helminths                                                                                                                                                                                                                                                                                                 |
| Amoxicillin    | Treatment of acute bacterial exacerbation of chronic bronchitis, acute otitis media, community acquired pneumonia, eradication of <i>H. pylori</i> , uncomplicated typhoid fever, mild leptospirosis, buccal cellulitis, pericoronitis and dental prophylaxis; second line treatment for exudative tonsillitis                                              |
| Azithromycin   | Suspected uncomplicated genital chlamydial infections (e.g., urethritis, cervicitis, trachoma); low-risk community-acquired pneumonia (CAP-LR) in adults, when <i>Legionella</i> , <i>Chlamydia</i> or <i>Mycoplasma</i> is suspected, or when there is hypersensitivity to the beta-lactams; non-severe CAP in child with hypersensitivity to amoxicillin. |
| Cefixime       | Second line therapy for uncomplicated typhoid fever, salmonella gastroenteritis, uncomplicated gonococcal infections of the cervix, urethras, rectum and conjunctiva. Step down therapy for severe/complicated typhoid fever                                                                                                                                |
| Cefuroxime     | Treatment of infections caused by methicillin sensitive staphylococci, group B streptococci, <i>H. influenzae</i> (type A and B), <i>E. coli</i> , <i>Enterobacter</i> , <i>Salmonella</i> , and <i>Klebsiella</i> ; treatment of susceptible infections of the upper and lower respiratory tract, otitis media                                             |
| Ciprofloxacin  | Restricted use of treatment of severe infections of the GI tract including MDR typhoid fever and <i>Shigella</i> dysentery, chancroid; febrile neutropenia, alternative treatment of acute uncomplicated cystitis; meningococcal meningitis prophylaxis.                                                                                                    |
| Clarithromycin | Management of low risk community-acquired pneumonia in adults (CAP-LR) due to <i>Legionella</i> , <i>Chlamydia</i> or <i>Mycoplasma</i> ; non-severe PCAP with hypersensitivity to amoxicillin; <i>Helicobacter pylori</i> infection. In combination with other drugs. Alternative in penicillin allergic patients.                                         |
| Clindamycin    | Skin and soft tissue infections caused by community-associated methicillin-resistant <i>Staphylococcus aureus</i> (MRSA); alternative for anaerobic infections. Alternative to macrolides for severe penicillin allergy; recurrent pharyngitis                                                                                                              |
| Clotrimazole   | Treatment of acute bacterial otitis externa caused by organisms susceptible to the action of ciprofloxacin, such as <i>Pseudomonas aeruginosa</i> , <i>Staphylococcus aureus</i> , and <i>Enterobacteriaceae</i> .<br><br>NOTE: NOT indicated for the treatment of otitis media                                                                             |
| Clotrimazole   | Management of anogenital and vulvovaginal candidiasis; ringworm; skin infections, including pityriasis versicolor and dermatophytosis (e.g., tinea corporis, tinea pedis, tinea cruris); protozoal infections (e.g., trichomoniasis)                                                                                                                        |
| Cloxacillin    | Treatment of bacterial infections caused by susceptible strains of penicillinase-producing staphylococci                                                                                                                                                                                                                                                    |
| Co-amoxiclav   | Upper respiratory tract infection: exudative or diffuse erythematous, recurrent pharyngitis, peritonsillar abscess (Quincy), deep neck                                                                                                                                                                                                                      |

MASTER COPY

DC:  Date: 6/4/26



| Drug                                            | Indications                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |
|-------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                                                 | abscess/retropharyngeal abscess, parapharyngeal space infection, acute bacterial rhinosinusitis (ABRS), acute otitis media, acute otitis media (clinical failure after 3 days), severe lower respiratory tract infection, community-acquired pneumonia, febrile neutropenia                                                                                                                                                                                                                                                                                                                                                                               |
| Cotrimoxazole (Sulfamethoxazole + Trimethoprim) | Treatment of uncomplicated typhoid fever, internal hordeolum, dacrocystitis, pertussis, and pathogen-specific infections such as pneumocystis jirovecii                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| Doxycycline                                     | Treatment of leptospirosis; juvenile periodontitis; gastroenteritis; acute bacterial exacerbation of chronic bronchitis (ABECB); blepharitis with associated acne rosacea; chlamydia and mycoplasma infections; second-line treatment of acute bacterial rhinosinusitis (ABRS) or for patients with severe penicillin allergy; pneumonia due to Burkholderia pseudomallei                                                                                                                                                                                                                                                                                 |
| Erythromycin                                    | Treatment of superficial ocular infections involving the conjunctiva or cornea caused by organisms susceptible to erythromycin; prophylaxis of ophthalmia neonatorum from both N. gonorrhoeae and C. trachomatis.<br>Alternative to penicillin in hypersensitive patients such as in secondary prevention of rheumatic fever and tonsillitis; conjunctivitis, pharyngitis, pertussis, chlamydia infections and community-acquired pneumonia.                                                                                                                                                                                                              |
| Fluconazole                                     | Treatment of candidiasis (esophageal, oropharyngeal, peritoneal, urinary tract, vaginal); Systemic candida infections (e.g., candidemia, disseminated candidiasis, and pneumonia); Cryptococcal meningitis; Antifungal prophylaxis in allogeneic bone marrow transplant recipients.                                                                                                                                                                                                                                                                                                                                                                       |
| Ketoconazole                                    | Treatment of tinea corporis, tinea cruris, tinea versicolor, cutaneous candidiasis, seborrheic dermatitis                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
| Mebendazole                                     | Management of common roundworm infections                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
| Metronidazole                                   | Eradication of H. pylori; Treatment of acute intestinal amebiasis (amebic dysentery) and amebic liver abscess;<br>Treatment of bacterial septicemia;<br>Treatment (adjunctive therapy) of bone and joint infections;<br>Treatment of CNS infections, including meningitis and brain abscess;<br>Treatment of gynecologic infections including endometritis, endomyometritis, tubo-ovarian abscess, or postsurgical vaginal cuff infection;<br>Treatment of intra-abdominal infections, including peritonitis, intraabdominal abscess and liver abscess;<br>Treatment of lower respiratory tract infections, including pneumonia, empyema and lung abscess |
| Nitrofurantoin                                  | Treatment of acute, uncomplicated cystitis in non-pregnant women, asymptomatic bacteriuria; recurrent urinary tract infection in non-pregnant women (for asymptomatic bacteriuria in pregnancy, urine culture is necessary).                                                                                                                                                                                                                                                                                                                                                                                                                              |

| Drug                       | Indications                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
|----------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Oseltamivir                | Prophylaxis of influenza A and B (not a substitute for flu vaccination); treatment of uncomplicated acute illness due to influenza A and B infection in patients who have been symptomatic for no more than 2 days.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
| Tobramycin                 | Treatment of external infections of the eye                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| Tobramycin + Dexamethasone | Steroid-responsive ocular inflammation with suspected or confirmed bacterial infection.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |
| Clopidogrel                | Prevention of atherothrombotic events in: peripheral arterial disease, or within 35 days of MI, or within 6 months of ischemic stroke; in acute coronary syndrome without ST-segment elevation (unstable angina or non-Q wave MI), including patients undergoing stent placement following percutaneous coronary intervention (in combination with aspirin); in acute MI with ST-segment elevation, and in combination with ASA in medically-treated patients eligible for thrombolytic therapy; in patients with atrial fibrillation where oral anticoagulation is unsuitable; neurologic indications; transient ischemic attack and acute ischemic cerebral infarction (for early specific treatment or secondary prevention of ischemic stroke). |
| Aspirin                    | Primary prevention of acute MI and stroke in patients with risk factors; secondary prevention of thrombotic cardiovascular or cerebrovascular disease, and following bypass surgery; management of acute MI and acute ischemic stroke.<br><br>Pain and inflammation associated with musculoskeletal and joint disorders; juvenile rheumatoid arthritis                                                                                                                                                                                                                                                                                                                                                                                              |
| Budesonide + Formoterol    | Treatment of asthma in patients $\geq 12$ years of age where combination therapy is indicated; maintenance treatment of airflow obstruction associated with chronic obstructive pulmonary disease (COPD), including chronic bronchitis and emphysema                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
| Fluticasone + Salmeterol   | For asthma patients aged 4 years and older; maintenance treatment of airflow obstruction, and reduction of exacerbations in patients with COPD<br><br>NOTE: This should be used only if the asthma is deemed to be not well-controlled with a long-term asthma control medicine (e.g., inhaled corticosteroids).                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| Ipratropium                | Chronic Asthma                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| Montelukast                | Prophylaxis and treatment of asthma (as an adjunctive therapy in combination with a corticosteroid); symptomatic relief of seasonal allergic rhinitis in patients with asthma<br><br>NOTE: There is a place for montelukast in children aged 2–5 years with intermittent viral-induced asthma, where leukotrienes play a minor role.                                                                                                                                                                                                                                                                                                                                                                                                                |
| Prednisone                 | Management of autoimmune disease; croup; short-term suppression of inflammation in allergic disorders; Pneumocystis carinii pneumonia; atopic dermatitis; dermatomyositis or polymyositis; nephrotic syndrome (idiopathic or related to lupus erythematosus); immune-mediated thrombocytopenia; inflammatory bowel disease; eye                                                                                                                                                                                                                                                                                                                                                                                                                     |

MASTER  
COPY

DC: *JP* Date: 6/4/26

| Drug                                     | Indications                                                                                                                                                                                                                                                                    |
|------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                                          | inflammation; rheumatic disease; immunosuppression; acute severe asthma (reliever); severe persistent asthma not responding to high-dose inhaled steroids, long-acting agonists, and methylxanthines (controller); adrenal insufficiency                                       |
| Salbutamol                               | Prophylaxis and treatment of asthma; bronchospasm in patients with reversible obstructive airway disease; prevention of exercise-induced bronchospasm                                                                                                                          |
| Ipratropium + Salbutamol                 | Treatment of COPD in patients who are currently on a regular bronchodilator, who continue to have evidence of bronchospasm and require a second bronchodilator                                                                                                                 |
| Tiotropium                               | Maintenance treatment of bronchospasm associated with chronic obstructive pulmonary disease (COPD), including chronic bronchitis and emphysema; reduction of COPD exacerbations                                                                                                |
| Aluminum Hydroxide + Magnesium Hydroxide | Symptomatic relief of symptoms related to hyperacidity from heartburn, hiatal hernia, upset stomach, peptic ulcer, peptic esophagitis, or gastritis.                                                                                                                           |
| Butamirate                               | Chronic, non-productive, irritative cough; symptomatic treatment of cough of various origins                                                                                                                                                                                   |
| Celecoxib                                | Relief of the signs and symptoms of osteoarthritis, ankylosing spondylitis, and rheumatoid arthritis; and short-term (=7 days) management of moderate to severe acute injury (post-operative, musculoskeletal or soft tissue), or following dental extraction                  |
| Cetirizine                               | Perennial and seasonal allergic rhinitis; other allergic symptoms such as hay fever, conjunctivitis, and chronic idiopathic urticaria                                                                                                                                          |
| Colchicine                               | Management of acute gout; treatment of familial Mediterranean fever in adults and children 4 years and older; prophylaxis and treatment of acute gout flares                                                                                                                   |
| Chlorphenamine                           | Used for symptomatic relief of allergic conditions including urticaria and angioedema, rhinitis, and conjunctivitis, and in pruritic skin disorders. For symptomatic treatment of cough and the common cold.                                                                   |
| Diphenhydramine                          | Relief of allergic symptoms caused by histamine release, including rhinitis, cough, conjunctivitis and allergic dermatoses; moderate to severe angioedema and anaphylaxis (adjunct to epinephrine)                                                                             |
|                                          | Management of Parkinsonism; drug-induced dystonic reaction (Extrapyramidal Syndrome).                                                                                                                                                                                          |
| Ferrous Salt (Iron Preparations)         | Treatment of iron-deficiency anemia and in people on hemodialysis who are receiving erythropoietin (supplement and prophylaxis); hematinic.                                                                                                                                    |
| Folic Acid + Iron Ferrous                | Prevention of iron and folic acid deficiencies, especially in pregnancy; nutritional supplement during pregnancy. NOTE: Low doses of folic acid in combination preparations are inadequate for the treatment of megaloblastic anemia, overdose, and therapy with deferoxamine. |
| Ibuprofen                                | Acute pain and inflammation in rheumatic diseases and other musculoskeletal disorders; mild-to-moderate pain, including dysmenorrhea and headache; pain in children; acute migraine attack.                                                                                    |

MASTER  
COPY

DC: *[Signature]* Date: 6/4/26

| Drug                    | Indications                                                                                                                                                                                                                                                                                                                                                                                   |
|-------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Lagundi (Vitex Negundo) | For the relief of mild to moderate cough due to common colds and flu                                                                                                                                                                                                                                                                                                                          |
| Loratadine              | Relief of nasal and non-nasal symptoms of seasonal allergic rhinitis; hay fever; chronic idiopathic urticaria, and other allergic dermatological disorder                                                                                                                                                                                                                                     |
| Mefenamic Acid          | Symptomatic relief of acute pain and inflammation; symptomatic relief of dysmenorrhea; menorrhagia due to dysfunctional causes or the presence of an IUD when organic pelvic pathology has been excluded; premenstrual syndrome                                                                                                                                                               |
| Naproxen                | Rheumatoid arthritis; osteoarthritis; ankylosing spondylitis; juvenile idiopathic arthritis; for the relief of signs and symptoms of tendonitis and bursitis; acute gout; management of pain due to inflammation (e.g., period pain, migraine); management of primary dysmenorrhea; inflammatory joint pain                                                                                   |
| Omeprazole              | Management of duodenal ulcer, gastric ulcer, and NSAID-associated gastric and duodenal ulcers and erosions, symptomatic GERD, reflux esophagitis, acid-related dyspepsia; H. pylori eradication in peptic ulcer disease prophylaxis of acid aspiration before surgery, or aspiration of gastric contents during general anesthesia.                                                           |
| Oral Rehydration Salts  | Replacement of fluid and electrolyte losses in mild to moderate dehydration due to acute diarrhea or vomiting; replacement of continuing loss from vomiting or diarrhea.                                                                                                                                                                                                                      |
| Paracetamol             | Management of mild-moderate pain, fever.                                                                                                                                                                                                                                                                                                                                                      |
| Zinc                    | Adjunct to oral rehydration therapy in acute and persistent diarrhea                                                                                                                                                                                                                                                                                                                          |
| Dapagliflozin           | As an add-on treatment to metformin (MET) for adults with Type 2 Diabetes Mellitus (T2DM) who have or are at a high risk of having cardiovascular disease and/or diabetic nephropathy                                                                                                                                                                                                         |
| Gliclazide              | Management of type 2 diabetes mellitus                                                                                                                                                                                                                                                                                                                                                        |
| Metformin               | Management of type 2 diabetes mellitus                                                                                                                                                                                                                                                                                                                                                        |
| Atorvastatin            | Adjunct to diet for the reduction of elevated total cholesterol, LDL cholesterol, apolipoprotein B, and triglycerides, and elevation HDL cholesterol in patients with primary hypercholesterolemia and combined hyperlipidemia<br><br>NOTE: Atorvastatin has not been studied in conditions where the major lipoprotein abnormality is elevation of chylomicrons (Fredrickson Types I and V). |
| Fenofibrate             | Hypertriglyceridemia; dyslipidemia associated with type 2 diabetes; second-line treatment in hypercholesterolemia                                                                                                                                                                                                                                                                             |
| Rosuvastatin            | Prevention of cardiovascular events in patients at high risk of a first cardiovascular event; mixed dyslipidemia, or homozygous familial hypercholesterolemia in patients who have not responded adequately to diet and other appropriate measures; (adults) treatment of primary hypercholesterolemia, e.g., heterozygous familial and non-familial                                          |

MASTER  
COPY

DC: *JG* Date: *6/4/26*

| Drug                            | Indications                                                                                                                                                                                                                                                                                                                                                                                              |
|---------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                                 | hypercholesterolemia; (children and adolescents) treatment of heterozygous familial hypercholesterolemia                                                                                                                                                                                                                                                                                                 |
| Simvastatin                     | For prevention of cardiovascular events in patients with high cardiovascular risk due to atherosclerotic cardiovascular disease or DM; primary hypercholesterolemia; homozygous familial hypercholesterolemia, or combined hyperlipidemia in patients who have not yet responded adequately to diet or other appropriate measures; (children and adolescents) heterozygous familial hypercholesterolemia |
| Gabapentin                      | Management of postherpetic neuralgia (PHN) in adults; neuropathic pain.                                                                                                                                                                                                                                                                                                                                  |
| Amlodipine                      | Management of hypertension                                                                                                                                                                                                                                                                                                                                                                               |
| Atenolol                        | Treatment of hypertension, alone or in combination with other agents; management of angina pectoris; secondary prevention post-myocardial infarction; for acute coronary syndrome                                                                                                                                                                                                                        |
| Captopril                       | Management of mild-to-moderate essential hypertension (alone, or with thiazide-diuretic therapy) and severe hypertension resistant to other treatment; CHF with left ventricular dysfunction following MI; diabetic nephropathy (microalbuminuria >30 mg/day) in type 1 diabetes                                                                                                                         |
| Clonidine                       | Hypertension (alone or used concomitantly with other antihypertensive agents)                                                                                                                                                                                                                                                                                                                            |
| Diltiazem                       | Prophylaxis and treatment of angina                                                                                                                                                                                                                                                                                                                                                                      |
| Enalapril                       | Management of hypertension; heart failure; acute coronary syndrome; post-myocardial infarction maintenance                                                                                                                                                                                                                                                                                               |
| Enalapril + Hydrochlorothiazide | Treatment of hypertension                                                                                                                                                                                                                                                                                                                                                                                |
| Hydrochlorothiazide             | Management of mild hypertension (alone), or in moderate or severe hypertension (in combination with other drugs); edema.                                                                                                                                                                                                                                                                                 |
| Isosorbide Dinitrate (ISDN)     | Prophylaxis and treatment of angina                                                                                                                                                                                                                                                                                                                                                                      |
| Isosorbide Mononitrate (ISMN)   | Prevention of angina pectoris                                                                                                                                                                                                                                                                                                                                                                            |
| Losartan                        | Treatment of hypertension; treatment of diabetic nephropathy in patients with type 2 diabetes mellitus (noninsulin dependent, NIDDM) and a history of hypertension; risk reduction of stroke in patients with HTN and left ventricular hypertrophy (LVH)                                                                                                                                                 |
| Losartan + Hydrochlorothiazide  | Treatment of hypertension; to reduce the risk of stroke in patients with hypertension and left ventricular hypertrophy (LVH)                                                                                                                                                                                                                                                                             |
| Methyldopa                      | For gestational hypertension and chronic hypertension in pregnant women.                                                                                                                                                                                                                                                                                                                                 |
| Metoprolol                      | Management of hypertension; angina pectoris; acute coronary syndrome; post-myocardial infarction maintenance                                                                                                                                                                                                                                                                                             |


MASTER  
COPY

Dr: *JP* Date: 6/4/26

| Drug                              | Indications                                                                                                                                                                                                                                                             |
|-----------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Tamsulosin                        | Treatment of sign and symptoms of BPH; relief of symptoms of obstruction in BPH                                                                                                                                                                                         |
| Telmisartan                       | Cardiovascular risk reduction in patients $\geq 55$ years of age unable to take ACE inhibitors and who are at high risk of major cardiovascular events (e.g., MI, stroke, death); treatment of hypertension, alone or in combination with other antihypertensive agents |
| Telmisartan + Hydrochlorothiazide | Treatment of hypertension                                                                                                                                                                                                                                               |
| Valsartan                         | Treatment of primary hypertension, alone or in combination with other antihypertensive agents; reduction of cardiovascular mortality in left ventricular dysfunction post-myocardial infarction; treatment of HF, NYHA Class II–IV                                      |
| Valsartan + Hydrochlorothiazide   | Treatment of hypertension                                                                                                                                                                                                                                               |

Source: Electronic Philippine National Formulary (ePNF) accessed through <https://pnf.doh.gov.ph/> last March 30, 2026


Note to the YAKAP Providers: For ease of patients' journey and transaction, YAKAP facilities should translate their diagnosis up to 4th character subdivision of the ICD-10 whenever available; which should be reflected when writing GAMOT prescription and requesting for cancer screening tests.



## Annex B: PhilHealth GAMOT Dispensing Limits

Table 1. Dispensing Limits of Oral Medications

| Molecule                                         | Use Case                                                                                                                                                                                         | Dispensing Limit                                                                                                                                                  |
|--------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Aluminum Hydroxide + Magnesium Hydroxide         | Medications for acute, self-limiting conditions prescribed as needed or for a short period of time. (Including NSAIDS per PC No. 2025-0013)                                                      | 7 days                                                                                                                                                            |
| Butamirate                                       |                                                                                                                                                                                                  |                                                                                                                                                                   |
| Celecoxib                                        |                                                                                                                                                                                                  |                                                                                                                                                                   |
| Cetirizine                                       |                                                                                                                                                                                                  |                                                                                                                                                                   |
| Chlorphenamine                                   |                                                                                                                                                                                                  |                                                                                                                                                                   |
| Diphenhydramine                                  |                                                                                                                                                                                                  |                                                                                                                                                                   |
| Ibuprofen                                        |                                                                                                                                                                                                  |                                                                                                                                                                   |
| Loratadine                                       |                                                                                                                                                                                                  |                                                                                                                                                                   |
| Mefenamic Acid                                   |                                                                                                                                                                                                  |                                                                                                                                                                   |
| Naproxen                                         |                                                                                                                                                                                                  |                                                                                                                                                                   |
| Oral Rehydration Salts (ORS 75-replacement)      |                                                                                                                                                                                                  |                                                                                                                                                                   |
| Paracetamol                                      |                                                                                                                                                                                                  |                                                                                                                                                                   |
| Vitex Negundo (Lagundi)                          |                                                                                                                                                                                                  |                                                                                                                                                                   |
| Omeprazole                                       | Medications for symptom relief/manage inflammation in chronic conditions                                                                                                                         | 14 days                                                                                                                                                           |
| Prednisone                                       |                                                                                                                                                                                                  |                                                                                                                                                                   |
| Albendazole                                      | Oral antimicrobials against infections<br><br>Note: Prescribing limits for oral antimicrobials shall NOT exceed fourteen (14) days per prescription. (Section V.E.3 of this PhilHealth Circular) | Anti-infectious medications shall always be dispensed in full, within the claiming period. (Partial filling is not allowed - per PC 2025-0013 Section V. C.5.g.4) |
| Amoxicillin                                      |                                                                                                                                                                                                  |                                                                                                                                                                   |
| Amoxicillin + Clavulanic acid (Co-Amoxiclav)     |                                                                                                                                                                                                  |                                                                                                                                                                   |
| Azithromycin                                     |                                                                                                                                                                                                  |                                                                                                                                                                   |
| Cefixime                                         |                                                                                                                                                                                                  |                                                                                                                                                                   |
| Cefuroxime                                       |                                                                                                                                                                                                  |                                                                                                                                                                   |
| Ciprofloxacin                                    |                                                                                                                                                                                                  |                                                                                                                                                                   |
| Clarithromycin                                   |                                                                                                                                                                                                  |                                                                                                                                                                   |
| Clindamycin                                      |                                                                                                                                                                                                  |                                                                                                                                                                   |
| Cloxacillin                                      |                                                                                                                                                                                                  |                                                                                                                                                                   |
| Doxycycline                                      |                                                                                                                                                                                                  |                                                                                                                                                                   |
| Erythromycin                                     |                                                                                                                                                                                                  |                                                                                                                                                                   |
| Fluconazole                                      |                                                                                                                                                                                                  |                                                                                                                                                                   |
| Mebendazole                                      |                                                                                                                                                                                                  |                                                                                                                                                                   |
| Metronidazole                                    |                                                                                                                                                                                                  |                                                                                                                                                                   |
| Nitrofurantoin                                   |                                                                                                                                                                                                  |                                                                                                                                                                   |
| Oseltamivir                                      |                                                                                                                                                                                                  |                                                                                                                                                                   |
| Sulfamethoxazole + Trimethoprim (Co-trimoxazole) |                                                                                                                                                                                                  |                                                                                                                                                                   |
| Zinc                                             | Mineral supplement<br>Adjunct to oral rehydration therapy                                                                                                                                        | 14 days                                                                                                                                                           |
| Amlodipine                                       | Medications for chronic disease management. (Maximum of 1 month                                                                                                                                  | 30 days                                                                                                                                                           |
| Aspirin                                          |                                                                                                                                                                                                  |                                                                                                                                                                   |
| Atenolol                                         |                                                                                                                                                                                                  |                                                                                                                                                                   |
| Atorvastatin                                     |                                                                                                                                                                                                  |                                                                                                                                                                   |






| Molecule                          | Use Case                                       | Dispensing Limit |
|-----------------------------------|------------------------------------------------|------------------|
| Captopril                         | provision per PC 2025-0013 Section V. C.5.g.1) |                  |
| Clonidine                         |                                                |                  |
| Clopidogrel                       |                                                |                  |
| Colchicine                        |                                                |                  |
| Dapagliflozin                     |                                                |                  |
| Diltiazem                         |                                                |                  |
| Enalapril                         |                                                |                  |
| Enalapril + Hydrochlorothiazide   |                                                |                  |
| Fenofibrate                       |                                                |                  |
| Folic Acid + Iron Ferrous         |                                                |                  |
| Gabapentin                        |                                                |                  |
| Gliclazide                        |                                                |                  |
| Hydrochlorothiazide               |                                                |                  |
| Iron (Ferrous Salt)               |                                                |                  |
| Isosorbide Dinitrate              |                                                |                  |
| Isosorbide Mononitrate            |                                                |                  |
| Losartan                          |                                                |                  |
| Losartan + Hydrochlorothiazide    |                                                |                  |
| Metformin                         |                                                |                  |
| Methyldopa                        |                                                |                  |
| Metoprolol                        |                                                |                  |
| Montelukast                       |                                                |                  |
| Rosuvastatin                      |                                                |                  |
| Simvastatin                       |                                                |                  |
| Tamsulosin                        |                                                |                  |
| Telmisartan                       |                                                |                  |
| Telmisartan + Hydrochlorothiazide |                                                |                  |
| Valsartan                         |                                                |                  |
| Valsartan + Hydrochlorothiazide   |                                                |                  |

Table 2. Dispensing Limits of Medications with Special Preparation or Forms

| Molecule               | Use Case                                  | Preparation/Form     | Dispensing Limit      |
|------------------------|-------------------------------------------|----------------------|-----------------------|
| Budesonide+ Formoterol | Medication for chronic disease management | Dry powder inhaler s | 1 inhaler per 15 day  |
|                        |                                           | Metered dose inhaler | 1 inhaler per 30 days |
| Clotrimazole           | Medication requiring full course therapy  | Cream                | 1 tube per 7 days     |
| Erythromycin           | Medication requiring full course therapy  | Ophthalmic ointment  | 1 tube per 14 days    |

MASTER COPY  
 DC: [Signature]  
 Date: 6/4/26

| <b>Molecule</b>            |  | <b>Use Case</b>                           | <b>Preparation/Form</b>             | <b>Dispensing Limit</b>                     |
|----------------------------|--|-------------------------------------------|-------------------------------------|---------------------------------------------|
| Fluticasone + Salmeterol   |  | Medication for chronic disease management | Metered dose inhaler                | 1 inhaler per 30 days                       |
|                            |  |                                           | Dry powder inhaler with 28 doses    | 1 inhaler per 7 days                        |
|                            |  |                                           | Dry powder inhaler with 60 doses    | 1 inhaler per 14 days                       |
| Ipratropium                |  | Medication for symptom relief             | Nebulizing solution (nebule)        | 20 nebulas every 5 days                     |
| Ipratropium + Salbutamol   |  | Medication for symptom relief             | Nebulizing solution (nebule)        | 20 nebulas every 5 days                     |
|                            |  |                                           | Metered dose inhaler with 200 doses | 1 inhaler per 30 days                       |
| Ketoconazole               |  | Medication requiring full course therapy  | Cream                               | 1 tube per 14 days                          |
|                            |  |                                           | Shampoo                             | 1 bottle per 14 days or 1 sachet per 3 days |
| Salbutamol                 |  | Medication for symptom relief             | Nebulizing solution (Nebule)        | 20 nebulas every 5 days                     |
|                            |  |                                           | Dry powder inhaler (capsule-based)  | 20 capsules every 5 days                    |
|                            |  |                                           | Metered dose inhaler with 200 doses | 1 inhaler per 30 days                       |
| Tiotropium                 |  | Medication for chronic disease management | Dry powder inhaler (capsule-based)  | 30 capsules every 21 days                   |
| Tobramycin                 |  | Medication requiring full course therapy  | Ophthalmic ointment                 | 1 tube per 14 days                          |
|                            |  |                                           | Ophthalmic solution                 | 1 bottle per 14 days                        |
| Tobramycin + Dexamethasone |  | Medication requiring full course therapy  | Ophthalmic ointment                 | 1 tube per 14 days                          |
|                            |  |                                           | Ophthalmic suspension               | 1 bottle per 14 days                        |



## Annex C: Quality Indicators for YAKAP

**Table 1. Quality Indicators for YAKAP Clinics**

| KPI                                                 | Formula                                                                              | Remarks                                                                                                                                                                                                    |
|-----------------------------------------------------|--------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 1. FPE rate                                         | No. of patients with FPE /No. of registered members                                  | <ul style="list-style-type: none"> <li>Annual rate.</li> </ul>                                                                                                                                             |
| 2. Consultation rate                                | No. of patients with consult /No. of patients with FPE                               | <ul style="list-style-type: none"> <li>Annual rate.</li> </ul>                                                                                                                                             |
| 3. Hypertension rate                                | No. of patients diagnosed with HPN / No. of patient with consult and FPE             | <ul style="list-style-type: none"> <li>Annual rate.</li> <li>Based on final diagnosis (ICD-10)</li> </ul>                                                                                                  |
| 4. Diabetes Mellitus (DM) rate                      | No. of patients diagnosed with DM / No. of patient consult and FPE                   | <ul style="list-style-type: none"> <li>Annual rate.</li> <li>Based on final diagnosis (ICD-10)</li> </ul>                                                                                                  |
| 5. Tuberculosis screening rate (chest x-ray result) | No. of patients screened for TB / No. of patient consult and FPE                     | <ul style="list-style-type: none"> <li>Annual rate.</li> <li>Based on chest x-ray result.</li> </ul>                                                                                                       |
| 6. COPD and Asthma rate                             | No. of patients with COPD or Asthma / No. of patients with consult and FPE           | <ul style="list-style-type: none"> <li>Annual rate</li> </ul>                                                                                                                                              |
| 7. Chronic disease rate                             | No. of patients with chronic diseases / No. of patients with consult and FPE         | <ul style="list-style-type: none"> <li>Annual rate.</li> <li>Chronic conditions are: Hypertension, diabetes mellitus, chronic obstructive pulmonary disease, asthma, and pulmonary tuberculosis</li> </ul> |
| 8. Acute disease rate                               | No. of patients with acute diseases / No. of patients with consult and FPE           | <ul style="list-style-type: none"> <li>Annual rate.</li> <li>Acute conditions are: acute respiratory tract infection, gastroenteritis, headache, musculoskeletal pain</li> </ul>                           |
| 9. Referral rate                                    | No. of patients referred to a specialist / No. of patient consult and FPE            | <ul style="list-style-type: none"> <li>Annual rate.</li> </ul>                                                                                                                                             |
| 10. Dental service referral rate                    | No of referral to dental service / No. of patient consult and FPE                    | <ul style="list-style-type: none"> <li>Annual rate.</li> </ul>                                                                                                                                             |
| 11. Optometry service referral rate                 | No. of referral to optometry service / No. of patients eligible with consult         | <ul style="list-style-type: none"> <li>Annual rate</li> <li>Eligible: 0-15 age, visual acuity done during consult.</li> </ul>                                                                              |
| 12. Visual screening rate                           | No of patients with visual screening (ie Snellen's chart) / No. of patients with FPE | <ul style="list-style-type: none"> <li>Annual rate</li> </ul>                                                                                                                                              |
| Hospital admission rate                             | No. of patients admitted to a hospital / No. of patient with consult and FPE         | <ul style="list-style-type: none"> <li>Annual rate.</li> </ul>                                                                                                                                             |

MASTER COPY

DC: 200 Date: 10/4/20

| KPI                               | Formula                                                                             | Remarks                                                                                                                                     |
|-----------------------------------|-------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------|
| 14. Breast CA Screening Rate      | No. of patients screened (Mammo or UTZ) / No. of identified high risk patients      | <ul style="list-style-type: none"> <li>Annual rate</li> <li>Lab result with finding</li> <li>High risk: Based on FPE and consult</li> </ul> |
| 15. Liver CA Screening Rate       | No. of patients screened (upper abd UTZ) / No. of identified high risk patients     | <ul style="list-style-type: none"> <li>Annual rate</li> <li>Lab result with finding</li> <li>High risk: Based on FPE and consult</li> </ul> |
| 16. Cervical CA screening Rate    | No. of patients screened thru PAP Smear / No. of identified high risk patients      | <ul style="list-style-type: none"> <li>Annual rate</li> <li>Lab result with finding</li> <li>High risk: Based on FPE and consult</li> </ul> |
| 17. Lung CA Screening Rate        | No. of patients screened (Low-dose Chest CT) / No. of identified high risk patients | <ul style="list-style-type: none"> <li>Annual rate</li> <li>Lab result with finding</li> <li>High risk: Based on FPE and consult</li> </ul> |
| 18. Colorectal CA Screening Rate  | No. of patients screened (FOBT/ FIT) / No. of identified high risk patients         | <ul style="list-style-type: none"> <li>Annual rate</li> <li>Lab result with finding</li> <li>High risk: Based on FPE and consult</li> </ul> |
| 19. Overall cancer screening rate | No. of patients screened / No. of identified high risk patients                     | <ul style="list-style-type: none"> <li>Annual rate</li> <li>Screened for: Breast, Liver, Cervical, Lung, and Colorectal Cancer</li> </ul>   |

**Table 2. Quality Indicators for Cancer Screening Facilities**

| KPI                                           | Formula                                                                 | Remarks                                                                                     |
|-----------------------------------------------|-------------------------------------------------------------------------|---------------------------------------------------------------------------------------------|
| Lung CA screening rate (Chest CT Scan)        | No. of lung cancer screening tests / No. of referred patients by YAKAP  | <ul style="list-style-type: none"> <li>Annual rate</li> <li>Based on lab results</li> </ul> |
| 2. Liver CA screening rate (UTZ or AFP)       | No. of Liver CA screening tests / No. of referred patients by YAKAP     | <ul style="list-style-type: none"> <li>Annual rate</li> <li>Based on lab results</li> </ul> |
| 3. Colorectal CA screening rate (Colonoscopy) | No. of Colorectal CA screening tests / No. of referred patient by YAKAP | <ul style="list-style-type: none"> <li>Annual rate</li> <li>Based on lab results</li> </ul> |
| 4. Breast CA screening rate (Mammo or UTZ)    | No. of Breast CA screening tests / No. of referred patient by YAKAP     | <ul style="list-style-type: none"> <li>Annual rate</li> <li>Based on lab results</li> </ul> |
| 5. Overall screening rate                     | No. of patients screened / No. of patients referred back to YAKAP       | <ul style="list-style-type: none"> <li>Annual rate</li> <li>Based on lab results</li> </ul> |

MASTER  
COPY

DC: *JP* Date: 6/4/26

**Table 3. Quality Indicators for GAMOT Facilities:**

| KPI                                                 | Formula                                                                                                                          | Remarks                                                                                                                                                                                                                                            |
|-----------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 1. Top 10 drugs dispensed                           | Frequency of generic name of drug dispensed                                                                                      | <ul style="list-style-type: none"> <li>Monthly count</li> <li>Overall drugs dispensed</li> </ul>                                                                                                                                                   |
| 2. Top 10 certified drugs dispensed                 | Frequency of generic name of certified drugs dispensed                                                                           | <ul style="list-style-type: none"> <li>Monthly count</li> <li>Drugs indicated in Certificate of Availability</li> </ul>                                                                                                                            |
| 3. Top 10 uncertified drugs dispensed               | Frequency of generic name of uncertified drugs dispensed                                                                         | <ul style="list-style-type: none"> <li>Monthly count</li> <li>GAMOT Drugs that are not indicated in Certificate of Availability</li> </ul>                                                                                                         |
| 4. Top 10 certified drugs stocked-out               | Frequency of generic name of certified drugs stocked-out                                                                         | <ul style="list-style-type: none"> <li>Monthly count</li> <li>Drugs indicated in Certificate of Availability</li> </ul>                                                                                                                            |
| 5. Reasons for stocked-out                          | Reasons for stocked-out and its frequency                                                                                        | <ul style="list-style-type: none"> <li>Monthly count</li> <li>Drugs indicated in Certificate of Availability</li> </ul>                                                                                                                            |
| 6. Anti-hypertensive maintenance drug dispense rate | No. of patients dispensed with 1 month supply of anti-hypertensive drug / No. of patients prescribed with anti-hypertensive drug | <ul style="list-style-type: none"> <li>Monthly rate</li> </ul>                                                                                                                                                                                     |
| 7. Anti-diabetes maintenance drug dispense rate     | No. of patients dispensed with 1 month supply of anti-diabetes drug / No. of patients prescribed with anti-diabetes drug         | <ul style="list-style-type: none"> <li>Monthly rate</li> </ul>                                                                                                                                                                                     |
| 8. Single drug prescription dispense rate           | No. of patients dispensed with complete single drug prescription / No. of patients with single drug prescription                 | <ul style="list-style-type: none"> <li>Monthly rate</li> </ul>                                                                                                                                                                                     |
| 9. Multiple drug prescription dispense rate         | No. of patients dispensed with complete multiple drug prescription / No. of patients with multiple drug prescription             | <ul style="list-style-type: none"> <li>Monthly rate</li> </ul>                                                                                                                                                                                     |
| 10. No. of prescriptions completely filled          | No. of prescriptions completely filled / Total No. of prescriptions for the month                                                | <ul style="list-style-type: none"> <li>Monthly rate</li> </ul>                                                                                                                                                                                     |
| 11. No. of prescriptions partially filled           | No. of prescriptions partially filled / Total No. of prescriptions for the month                                                 | <ul style="list-style-type: none"> <li>Monthly rate</li> </ul>                                                                                                                                                                                     |
| 12. No. of prescriptions rejected                   | No. of prescriptions rejected / Total No. of prescriptions for the month                                                         | <ul style="list-style-type: none"> <li>Monthly rate</li> <li>Rejected: if the GAMOT facility has questions on possible inappropriateness of prescriptions in terms of diagnosis</li> <li>Drugs indicated in Certificate of Availability</li> </ul> |

MASTER  
COPY

DC: *[Signature]* Date: 6/4/20

| KPI                       | Formula                                 | Remarks                                                                                                                     |
|---------------------------|-----------------------------------------|-----------------------------------------------------------------------------------------------------------------------------|
| 13. Reasons for rejection | Reasons for rejection and its frequency | <ul style="list-style-type: none"> <li>• Monthly count</li> <li>• Drugs indicated in Certificate of Availability</li> </ul> |

Note: Certificate of Availability of GAMOT as described among the documentary requirements of GAMOT Facilities in PhilHealth Circular 2025-0013

