

**Exercise 3.** Show that Proposition 3.4.3 would be false if  $p$  were allowed to be infinite. (Hint: construct a Borel subset  $A$  of  $[a, b]$  such that  $\|\chi_A - f\|_\infty \geq 1/2$  holds whenever  $f$  is a step function.)

**Proposition 3.4.3.** *Suppose that  $[a, b]$  is a closed bounded interval and that  $p$  satisfies  $1 \leq p < +\infty$ . Then the subspace of  $L^p([a, b])$  determined by the step functions on  $[a, b]$  is dense in  $L^p([a, b])$ .*

*Proof.* Let  $0 < \alpha < 1$ , there exists a Borel subset  $A$  of  $[a, b]$  such that:

- $\lambda(A) = \alpha \cdot (b - a)$
- $A$  does not contain any nonempty closed interval

Let  $f \in \mathcal{L}^\infty([a, b])$  be a step function

$$f = \sum_{i=0}^n b_i \chi_{[a_i, a_{i+1}]}$$

with  $a = a_0 < \dots < a_n = b$  real numbers in  $[a, b]$  and  $b_i$  complex numbers for all  $i$ . Suppose that  $\|\chi_A - f\|_\infty < \alpha/3$ . For all  $0 \leq i \leq n$ , there exists  $x_0 \in [a_i, a_{i+1}]$  such that  $x_0 \notin A$ , since  $A$  does not contain any nonempty closed interval. From this we deduce that  $\alpha/3 > \|\chi_A - f\|_\infty \geq |b_i|$  for all  $i$ . Both  $f$  and  $\chi_A$  are integrable on  $[a, b]$  and we have

$$\begin{aligned} \int_{[a,b]} \chi_A \, d\lambda &\leq \int_{[a,b]} |\chi_A - f| \, d\lambda + \int_{[a,b]} |f| \, d\lambda \\ \alpha \cdot (b - a) &\leq (b - a) \frac{\alpha}{3} + \sum_{i=0}^n |b_i| \cdot (a_{i+1} - a_i) \\ &\leq 2(b - a) \frac{\alpha}{3} \end{aligned}$$

from which we deduce that  $\alpha \leq 2\alpha/3$ , which is a contradiction. Therefore no step function can approach  $\chi_A$  arbitrarily, so that the subspace of  $L^\infty([a, b])$  determined by the step functions is not dense in  $L^\infty([a, b])$ .

Let us show the existence of a set  $A$  with the expected properties. Since the Lebesgue measure is invariant by translation, it is enough to show the existence of such a set in the interval  $[0, b - a]$ . Given a Borel set  $B$  and some real number  $x > 0$ ,  $x \cdot B$  is also a Borel set, so we can further reduce the interval to  $[0, 1]$ .

Let  $\beta$  be a real such that  $0 < \beta < 1$ , and let  $\sum_k u_k$  be a series of positive real numbers such that  $\sum_{k=0}^\infty u_k = 1 - \beta$ . Let  $(A_n)_{n \in \mathbb{N}}$  be a sequence of closed sets defined recursively by  $A_0 = [0, 1]$ , and, for each  $n$ ,  $A_{n+1}$  be the union of  $2^{n+1}$  closed intervals  $I_n$  obtained by removing from each of the  $2^n$

closed intervals  $I_n$  composing  $A_n$  an open interval of length  $u_n/2^n$  centered on  $I_n$ .

For all  $n$ ,  $A_n$  is closed and measurable, and  $\lambda(A_{n+1}) = 1 - (u_0 + u_1 + \dots + u_n)$ . Moreover,  $A_{n+1} \subset A_n$ , and  $\lambda(A_0) = 1$  is finite, so

$$\lambda\left(\bigcap_{n=0}^{\infty} A_n\right) = \lim_{n \rightarrow \infty} \lambda(A_n) = 1 - \lim_{n \rightarrow \infty} \sum_{k=0}^n u_k = \beta$$

Let  $A = \bigcap_{n \in \mathbb{N}} A_n$ .  $A$  is measurable, has measure  $\beta$ , and does not include any nonempty closed interval: suppose that  $[a, b] \subset A$ , then  $[a, b] \subset A_n$  for all  $n$ . Since  $A_n$  is the union of  $2^n$  disjoint intervals  $I_n$ , each of measure  $(1 - u_0 - \dots - u_n)/2^n$ , we have

$$0 \leq \lambda([a, b]) \leq \frac{1}{2^n} \left(1 - \sum_{k=0}^n u_k\right)$$

And since the series  $\sum_k u_k$  of positive reals converges, we deduce that  $\lambda([a, b]) = 0$ , so that  $[a, b] = \emptyset$ .

□