

**PROGRAMMING USING C**

---

**WEEK 6 ONE DIMENSIONAL ARRAYS**

**NAME : JAGADISH S A**

**CLASS : AIML B**

**REG NO : 241501071**

Given an array A of sorted integers and another non negative integer k, find if there exists 2 indices i and j such that  $A[i] - A[j] = k$ ,  $i \neq j$ .

Input Format

1. First line is number of test cases T. Following T lines contain:
2. N, followed by N integers of the array
3. The non-negative integer k

Output format

Print 1 if such a pair exists and 0 if it doesn't.

Example

Input:

```
1
3 1 3 5
4
```

Output:

```
1
```

Input:

```
1
3 1 3 5
qq
```

Answer: (penalty regime: 0 %)

```
1 #include<stdio.h>
2 int main()
3 {
4     int t;
5     scanf("%d",&t);
6     while(t--){
7         int n;
8         scanf("%d",&n);
9         int a[n];
10        for(int i=0;i<n;i++){
11            scanf("%d",&a[i]);
12        }
13        int k;
14        scanf("%d",&k);
15        int flag =0;
16        for(int i=0;i<n;i++){
17            for(int j=i+1;j<n;j++){
18                if(a[i]-a[j]==k || a[j]-a[i]==k){
19                    flag=1;
20                    break;
21                }
22            }
23            if(flag)break;
24        }
25        printf("%d\n",flag);
26    }
27    return 0;
28 }
```

	Input	Expected	Got	
✓	1 3 1 3 5 4	1	1	✓
✓	1 3 1 3 5 99	0	0	✓

Passed all tests! ✓

Sam loves chocolates and starts buying them on the 1st day of the year. Each day of the year,  $x$ , is numbered from 1 to  $Y$ . On days when  $x$  is odd, Sam will buy  $x$  chocolates; on days when  $x$  is even, Sam will not purchase any chocolates.

Complete the code in the editor so that for each day  $N_i$  (where  $1 \leq x \leq N \leq Y$ ) in array `arr`, the number of chocolates Sam purchased (during days 1 through  $N$ ) is printed on a new line. This is a function-only challenge, so input is handled for you by the locked stub code in the editor.

#### Input Format

The program takes an array of integers as a parameter.

The locked code in the editor handles reading the following input from `stdin`, assembling it into an array of integers (`arr`), and calling `calculate(arr)`.

The first line of input contains an integer,  $T$  (the number of test cases). Each line  $i$  of the  $T$  subsequent lines describes the  $i$ th test case as an integer,  $N_i$  (the number of days).

#### Constraints

$$1 \leq T \leq 2 \times 10^5$$

$$1 \leq N \leq 2 \times 10^6$$

$$1 \leq x \leq N \leq Y$$

#### Output Format

For each test case,  $T_i$  in `arr`, your `calculate` method should print the total number of chocolates Sam purchased by day  $N_i$  on a new line.

#### Sample Input 0

Answer: (penalty regime: 0 %)

```
1 #include<stdio.h>
2 int main()
3 {
4     int t;
5     scanf("%d",&t);
6     while(t--){
7         int n,c=0;
8         scanf("%d",&n);
9         for(int i=0;i<=n;i++){
10             if(i%2!=0)
11                 c=c+i;
12         }printf("%d\n",c);
13     }
14 }
```

	Input	Expected	Got	
✓	3	1	1	✓
	1	1	1	
	2	4	4	
	3			
✓	10	1296	1296	✓
	71	2500	2500	
	100	1849	1849	
	86	729	729	
	54	400	400	
	40	25	25	
	9	1521	1521	
	77	25	25	
	9	49	49	
	13	2401	2401	
	98			

Passed all tests! ✓

The number of goals achieved by two football teams in matches in a league is given in the form of two lists. Consider:

- Football team A, has played three matches, and has scored { 1 , 2 , 3 } goals in each match respectively.
- Football team B, has played two matches, and has scored { 2, 4 } goals in each match respectively.
- Your task is to compute, for each match of team B, the total number of matches of team A, where team A has scored less than or equal to the number of goals scored by team B in that match.
- In the above case:
- For 2 goals scored by team B in its first match, team A has 2 matches with scores 1 and 2.
- For 4 goals scored by team B in its second match, team A has 3 matches with scores 1, 2 and 3.

Hence, the answer: {2, 3}.

Complete the code in the editor below. The program must return an array of  $m$  positive integers, one for each  $\text{maxes}[i]$  representing the total number of elements  $\text{nums}[j]$  satisfying  $\text{nums}[j] \leq \text{maxes}[i]$  where  $0 \leq j < n$  and  $0 \leq i < m$ , in the given order.

It has the following:

`nums[nums[0],...,nums[n-1]]`: first array of positive integers

`maxes[maxes[0],...,maxes[m-1]]`: second array of positive integers

Constraints

- $2 \leq n, m \leq 105$
- $1 \leq \text{nums}[j] \leq 109$ , where  $0 \leq j < n$ .
- $1 \leq \text{maxes}[i] \leq 109$ , where  $0 \leq i < m$ .

Input Format For Custom Testing

Input from stdin will be processed as follows and passed to the function.

The first line contains an integer  $n$ , the number of elements in `nums`.

The next  $n$  lines each contain an integer describing  $\text{nums}[j]$  where  $0 \leq j < n$ .

The next line contains an integer  $m$ , the number of elements in  $\text{maxes}$ .

The next  $m$  lines each contain an integer describing  $\text{maxes}[i]$  where  $0 \leq i < m$ .

Sample Case 0

Sample Input 0

4

1

4

2

4

2

3

5

Sample Output 0

2

4

Explanation 0

We are given  $n = 4$ ,  $\text{nums} = [1, 4, 2, 4]$ ,  $m = 2$ , and  $\text{maxes} = [3, 5]$ .

1. For  $\text{maxes}[0] = 3$ , we have 2 elements in  $\text{nums}$  ( $\text{nums}[0] = 1$  and  $\text{nums}[2] = 2$ ) that are  $\leq \text{maxes}[0]$ .
2. For  $\text{maxes}[1] = 5$ , we have 4 elements in  $\text{nums}$  ( $\text{nums}[0] = 1$ ,  $\text{nums}[1] = 4$ ,  $\text{nums}[2] = 2$ , and  $\text{nums}[3] = 4$ ) that are  $\leq \text{maxes}[1]$ .

Thus, the function returns the array  $[2, 4]$  as the answer.

Answer: (penalty regime: 0 %)

```
1 #include<stdio.h>
2 int main()
3 {
4     int s1,s2,ans;
5     scanf("%d",&s1);
6     int ta[s1];
7     for(int i=0;i<s1;i++)
8         scanf("%d",&ta[i]);
9     scanf("%d",&s2);
10    int tb[s2];
11    for(int i=0;i<s2;i++)
12        scanf("%d",&tb[i]);
13    for(int j=0;j<s2;j++){
14        ans=0;
15        for(int i=0;i<s1;i++){
16            if(tb[j]>=ta[i])
17                ans++;
18        }printf("%d\n",ans);
19    }
20 }
```

	Input	Expected	Got	
✓	4	2	2	✓
	1	4	4	
	4			
	2			
	4			
	2			
	3			
	5			
✓	5	1	1	✓
	2	0	0	
	10	3	3	
	5	4	4	
	4			
	8			
	4			
	3			
	1			